



Use of Organic Inputs in Management of Alfisols and Ultisols for Sustainable Maize Production in Western Kenya

J. Achieng¹, G. Ouma^{2*} and G. Odhiambo³

¹Kenya Agricultural Research Institute, P. O. Box 169-50100, Kakamega, Kenya.

²Bondo University College, P. O. Box 210, Bondo, Kenya.

³Maseno University, Private Bag, Maseno, Kenya.

Authors' contributions

This work was part of a PhD thesis. Author JA was the student and he designed the study, managed the experiment, collected all the data and conducted statistical analysis. He also wrote the protocol and the first draft of the manuscript. Author GO was the principal supervisor while author GO was the second supervisor. Besides supervision, they managed the literature searches. All the authors read and approved the final manuscript.

Research Article

Received 4th May 2013
Accepted 24th June 2013
Published 24th July 2013

ABSTRACT

Maize (*Zea mays* L.) is an important food crop in western Kenya, mostly grown by smallholder farmers in complex and risky cropping systems. Trends in population growth in the country indicate that the demand for maize is projected to increase 3-4% annually. Production is however hampered by the predominance of fragile ecosystems (acidic soils), low soil fertility and low use of chemical inputs. The average grain yield is less than 1.0 t ha⁻¹ instead of a reachable 5.0 t ha⁻¹, leading to vicious cycle of food insecurity. An on-farm experiment was conducted in two soil types (Alfisols and Ultisols) in Kakamega, western Kenya, between February and September 2007 to test effects of various organic inputs (Farmyard manure, Tithonia biomass and Desmodium cover crop) in combination with inorganic fertilizers (N, P, K, Mg, B and Zn) on yield of maize. The design was Randomized Complete Block Design, replicated 5 times and the data was subjected to ANOVA and DMRT tests. Soil analysis before planting indicated that pH was 5.0 and 5.4 in Alfisols and Ultisols, respectively. Both soils were deficient in N and P but adequate in exchangeable bases (K, Ca and Mg). Maize grain yield was higher in Ultisols compared to Alfisols. In Alfisols, organic inputs in combination with 30 kg N ha⁻¹ gave maize grain yield

*Corresponding author: Email: goumaoindo@gmail.com;

improvement of nearly 100% over farmer's practice (non improved maize variety, wider plant spacing, inorganic fertilizer applied at the rate of 20 N + 20 P ha⁻¹, one weeding). When inorganic fertilizer (60 kg N + 60 kg P ha⁻¹) was applied to maize/banana intercrop, maize yield increased by about 40%. In contrast, in Ultisols, organic inputs increased maize grain yield by between 85% and 115%, while maize/banana intercrop (plus 60 kg N + 60 kg P ha⁻¹) increased maize grain yield by only 4% over the farmer's practice. Banana intercrop reduced maize population resulting in low maize grain and stover yield. Use of either farmyard manure, tithonia biomass or desmonium cover crop in combination with 30 kg of N ha⁻¹ (from inorganic source) can enhance maize production among smallholder farmers in acidic soils who ordinarily cannot afford to purchase adequate quantities of inorganic fertilizers.

Keywords: Alfisols; Ultisols; maize; organic inputs; sustainable production; Western Kenya.

1. INTRODUCTION

Maize (*Zea mays* L.) is the most important staple food crop in Kenya. It is estimated that about 1.6 million hectares of land are under the crop annually with 80% being grown by smallholder farmers [1,2], majority of whom are in western part of the country. However, the production does not match the population and industrial demand that increases at about 4.7% annually. Low soil fertility has been identified as one of the major constraints to maize production among smallholder farmers. Exacerbating the low soil fertility in the region is the high rate of acidity in many soils due to high rainfall, depletion of soil carbon and continuous cultivation. Predominant soils in western Kenya include Alfisols and Ultisols [3]. Alfisols are known to have undergone moderate leaching, are susceptible to surface compaction, erosion and acidify rapidly under continuous cultivation [4]. They have low water retention capacity [4] and are deficient in N and P [5,6]. On the other hand, Ultisols are strongly leached, highly acidic, have multiple nutrient deficiencies of N, P, K, Ca and Zn and P-fixation tendencies; they are susceptible to erosion and compaction [7].

In Kenya, acid soils cover about 13% of prime agricultural land, mostly concentrated in western part of the country [8]. Acid soils significantly limit crop production through toxic levels of aluminum (Al) and manganese (Mn), as well as sub-optimal levels of P [9]. For optimum growth and grain yield, maize requires soils with $\geq 2.0\%$ N [10], ≥ 13 mg P kg⁻¹ [11] and pH range of 5.5-6.5. Studies have shown that crop yields in acidic, un-limed soils have declined even with application of adequate amounts of inorganic fertilizers [12]. Use of organic inputs, especially *Tithonia diversifolia* and *Desmodium* sp., can raise the pH [13] and improve fertility of the acidic soils [14]. Soils fallowed to *Tithonia* improved soil's physical properties and had higher mineral values like N, P, K Ca and Mg [15]. *Tithonia* has a faster rate of decomposition because of its low lignin (6.5%) and polyphenol (1.6%) content [15], but has high nitrogen (3.5%), phosphorus (0.4%) and potassium (4.1%) [16]. *Tithonia* is an invasive annual weed growing abundantly in western Kenya mainly along roads, hedges and on uncultivated fields. *Desmodium* is a nitrogen-fixing shrub legume characterized by the following attributes: fast growing, easily degradable biomass, compatibility with maize crop and has no allelopathic effect on cereals [16, 17]. Several *desmodium* sp. are used as cover crops in maize and sorghum fields to repel *Chillo partellus* (stem borer) as well as suppression of witch-weed. They are also used as living mulch, green manure, besides being good animal fodder. Although organic inputs cannot meet crop P requirements, they can increase P availability by reducing its adsorption capacity in the soil [18]. Nutrient content of some commonly used organic materials is presented in Table 1.

The objective of this experiment was to study the effect of organic inputs and their combinations with inorganic fertilizers on growth and yield of maize intercropped with banana in Alfisols and Ultisols in Kakamega, western Kenya.

Table 1. Nutrient content of some organic materials

| Material | N (%) | P (%) | K (%) |
|-----------------------|--------------|--------------|--------------|
| Cow manure | 0.4-0.6 | 0.2 | 0.2-0.5 |
| Goat manure | 1.4 | 0.2 | 0.3-1.0 |
| Sheep manure | 0.7 | 0.3 | 0.4 |
| Poultry manure | 1.1-1.5 | 0.8-1.3 | 0.5-2.7 |
| Tithonia biomass | 3.97 | 0.3 | 4.6 |
| Desmodium biomass | 3.44 | 0.15 | - |
| Mucuna biomass | 3.56 | 0.17 | - |
| Leucaena leucocephala | 3.74 | 0.26 | 3.37 |

Source: [19].

2. MATERIALS AND METHODS

2.1 Description of the Study Site

The study was carried out in Kakamega within four administrative divisions of Ikolomani, Ileho, Shinyalu and Lurambi, western Kenya. The region has one of the most densely populated rural areas in the world with 500 to 1200 people km² [20,21] with average household farm size of 0.7 ha of which about one half is planted to maize in combination with beans and sometimes, banana. The remaining portion is usually either under tea or sugarcane. The whole area is characterized by low input-low output and low incomes. Maize grain deficit lasts for at least four months every year, resulting in high levels of hunger and extreme poverty [22]. The experiment was carried out during the long rains season of 2007, beginning February to September.

The soils in the study site are mainly Ultisols or Alfisols. Although there are pockets of fertile soils in the County, about 85% of the land is poor as a result of leaching from high rainfall and intensive cultivation with inadequate measures to maintain soil fertility. Annual rainfall ranges from 1200 to 1800 mm with a bimodal distribution allowing two cropping seasons. Long rains are experienced from March to June, peaking in April/May while the short rains begin in July, peaking in September. The mean maximum and minimum temperatures range from 28°C to 32°C and 11°C to 13°C, respectively. Mean altitude of the area is 1200 m.

2.2 Field Experiment

Soil sampling was done in representative locations within the study site covering Alfisols and Ultisols prior to planting. Air-dried 2 mm sieved samples were subjected to chemical analysis at National Agricultural Research Laboratories, Nairobi, as described by Okallebo [23]. Total N and available P were determined by micro Kjeldahl and molybdenum blue colorimetry methods, respectively. Exchangeable K, Ca and Mg were extracted using ammonium acetate. Soil pH was measured using a soil water ratio of 1:2.5 (w/v).

On-farm experiment was established on selected ten farms (five each in Alfisols and Ultisols). Land preparation was done by hand-hoeing to a fine tilth and planting was done at the on-set of rain. Each experimental plot was planted with maize variety H614 at a spacing

of 75 cm between the rows and 30 cm along the rows; two seeds per hole but later thinned to one plant per hill after emergence. The treatments of the experiment were: control/farmer's practice (non-improved maize variety, wider plant spacing, fertilizer rate of 20 N/P ha⁻¹, one weeding); no weeding + no fertilizer, weeded-no fertilizer, 60 kg N ha⁻¹, 60/60 kg N/P ha⁻¹, 60/60/40 kg N/P/K ha⁻¹, 5 tons FYM + 30 kg N ha⁻¹, 5 tons tithonia + 30 kg N ha⁻¹, desmodium + 30/30 kg N/P ha⁻¹, banana + 60/60 kg N/P ha⁻¹, optimal fertilizer, and modified farmers' practice (one weeding + 20/20 kg N/P ha⁻¹). The treatments were arranged in a randomized complete block design with five replications in each soil type. Each experimental unit measured 19.5 m². Data was subjected to ANOVA using SAS [24] Version 8 and Duncan's Multiple Range Test (DMRT) at 5% level was used to separate the means.

Tissue-culture banana (cv. Kanyinjia) was planted as an intercrop at the same time with maize at a spacing of 3m x 3m. *Desmodium intortum* (green leaf) seed, also planted at the same time with maize, was drilled midway between maize rows at the rate of 1kg per ha. Desmodium cover crop was pruned two times during the season and leaves left on the soil surface to decompose and generate organic matter. Tithonia twigs were cut from the nearby roadsides, carried to the Research Station, sun dried and stored in a dry place in gunny bags. Tithonia and farmyard manure were incorporated on the plots 7 days before planting at the rate of 5 t ha⁻¹ dry weight of each. Basal fertilizer was applied at the time of planting followed by side-dressing five weeks after emergence of maize. Inorganic N fertilizer was sourced from Urea, P was from Di-Ammonium Phosphate, K was from Potassium Chloride, Mg was from Magnesium Chloride, Zn was from Zinc Chloride while B was from Borate Chloride. Hand weeding was done two times except on farmers' practice treatment that received one weeding.

Maize was harvested at physiological maturity as indicated by the formation of black layer at the pedicel of 50% of the sampled kernels. At this stage, mean plant height was obtained by randomly selecting 5 plants from each plot, which were then cut at the soil surface level and their length measured with a tape measure from the cut surface to the tip of the longest tassel branch. Maize was harvested from four inner rows from a net plot of 13.5 m². Number of plants and ears from the net plot were counted. An ear was described as any biological structure on a maize plant having a cob (with or without grains). All clean ears were hand shelled, weighed and recorded at the farm. Grain yield, adjusted to 13% moisture content was calculated by the formula:

$$\text{Grain yield (t ha}^{-1}\text{)} = \text{FW} * (100 - \text{MC})/87 * (10,000/\text{NP}) * \text{SP}/1,000$$

Where: FW = Grain field weight from the net plot (kg),
MC = Grain moisture content at harvesting,
NP = Net plot (m²),
SP = Shelling percentage of the ear (estimated at 0.85)

Another set of five plants was randomly selected and cut at the soil surface for purposes of estimating plant biomass. The plants were de-husked, cobs were removed and the remaining parts tied together, labeled and taken to KARI-Kakamega Laboratories. They were then chopped into small pieces, put in paper bags and dried in an oven at 70°C until a constant weight was reached. The final weight represented the final biomass for each plot.

3. RESULTS

3.1 Soil Chemical Properties before Planting Maize

Soil analysis before planting indicated significant differences between Alfisols and Ultisols (Table 2). Alfisols was more acidic compared to Ultisols. Other than available P that was higher in Alfisols, all other nutrients (carbon, nitrogen, potassium, calcium and magnesium) were higher in Ultisols.

Table 2. Soil characteristics of maize fields in Alfisols and Ultisols in the administrative divisions of the study site

| Division | *Soil Type | Soil pH (H ₂ O) | Total C (%) | Total N (%) | Extr. P (mg kg ⁻¹) | Exch. K (cmol kg ⁻¹) | Exch. Ca (g kg ⁻¹) | Exch. Mg (g kg ⁻¹) |
|-----------------|------------|----------------------------|-------------|-------------|--------------------------------|----------------------------------|--------------------------------|--------------------------------|
| Ileho | Alf. | 5.0 | 1.09 | 0.12 | 4.78 | 0.35 | 0.75 | 0.07 |
| Lurambi | Alf. | 4.9 | 1.08 | 0.11 | 7.85 | 0.42 | 1.07 | 0.10 |
| Shinyalu | Ult. | 5.4 | 2.39 | 0.22 | 5.00 | 0.55 | 1.65 | 0.18 |
| Ikolomani | Ult. | 5.3 | 1.45 | 0.15 | 5.23 | 0.41 | 1.14 | 0.14 |
| Mean (Alfisols) | | 5.0 | 1.07 | 0.12 | 6.18 | 0.38 | 0.81 | 0.08 |
| Mean (Ultisols) | | 5.4 | 1.92 | 0.18 | 5.14 | 0.48 | 1.39 | 0.16 |

*Soil type: Alf. = Alfisols, Ult. = Ultisols

3.2 Effect of Organic Inputs on Growth Parameters of Maize

3.2.1 Maize population at physiological maturity

In Alfisols, highest plant population (37,778 per hectare) was recorded in optimal fertilizer treatment (Table 3). This was about 85% of the potential population. Farmer's practice (control) had 62% of the potential plant population. All other treatments gave populations higher than farmer's practice but significantly less than that in optimal fertilizer. In Ultisols, N (60 kg) treatment gave the highest plant population (39,556); which was 89% of maximum expected while farmer's practice gave 62% of maximum expected (Table 4). Organic inputs: FYM, tithonia and desmodium gave populations that were significantly similar to that of NP treatment. Banana intercrop gave 68% of potential population.

Table 3. Effects of organic inputs and inorganic fertilizer combinations on attributes and maize grain yield in Alfisols of Kakamega, Kenya

| Treatment | Plants (ha ⁻¹) | Ears (ha ⁻¹) | Plant Height (m) | Maize Yield (t ha ⁻¹) | Stover Yield (t ha ⁻¹) |
|--------------------------------------|----------------------------|--------------------------|------------------|-----------------------------------|------------------------------------|
| Farmer's practice (Control) | 27,704 b | 17,778 de | 2.3 c | 1.9 e | 6.0 f |
| Unweeded, no fertilizer | 28,593 ab | 15,407 e | 2.3 c | 2.0 e | 7.6 ef |
| Weeded, no fertilizer | 31,407 ab | 22,222 bcde | 2.3 c | 2.9 cde | 10.2 cdef |
| N (60 kg) | 31,407 ab | 21,037 cde | 2.4 bc | 4.1 bcd | 13.9 bc |
| NP (60 kg N + 60 kg P) | 30,815 ab | 26,222 bc | 2.5 abc | 4.2 bc | 13.7 bc |
| NPK (60 kg N + 60 kg P + 40 kg K) | 36,000 ab | 29,481 ab | 2.6 abc | 4.7 b | 16.5 b |
| FYM + 30 kg N | 30,815 ab | 22,815 bcde | 2.6 abc | 3.9 bcd | 12.2 bcde |
| Tithonia + 30 kg N | 34,815 ab | 25,630 bcd | 2.4 bc | 3.8 bcd | 13.1 bcd |
| Desmodium + 30 kg N + 30 kg P | 33,926 ab | 26,963 bc | 2.5 abc | 3.4 bcd | 11.9 bcde |
| ¹ Optimal fertilizer | 37,778 a | 35,407 a | 2.8 ab | 6.6 a | 24.3 a |
| Modified farmer's practice | 31,852 ab | 22,518 bcde | 2.5 bc | 2.7 de | 8.6 def |
| Banana intercrop + 60 kg N + 60 kg P | 29,630 ab | 19,259 cde | 2.9 a | 2.7 de | 11.6 bcde |
| Mean | 32,235 | 24,047 | 2.5 | 3.7 | 12.5 |
| CV (%) | 17.1 | 19.5 | 9.4 | 24.1 | 24.0 |

Means followed by the same letter within the same column are not significantly different at $P \leq 0.05$, using DMRT.

¹Optimal fertilizer = Optimal fertilizer: (N, P, K, Mg, Zn and B ((NPK was broadcast at planting at the rate of 200, 60 and 120 kg ha⁻¹ of each)), respectively; Mg 20 kg ha⁻¹, Zn 20 kg ha⁻¹ and B 5 kg ha⁻¹)

3.2.2 Number of ears

In Alfisols, optimal fertilizer treatment gave the highest number of ears (35,407) (Table 3). In relation to optimal fertilizer treatment, unweeded- no fertilizer and control gave 43% and 50% ears, respectively. Among the organic inputs, desmodium gave number of ears significantly similar to that of NP (60 kg N + 60 kg P). FYM gave number of ears similar to that recorded in weeded, no fertilizer treatment. Banana intercrop treatment gave nearly half of the ears in relation to optimum fertilizer treatment.

Table 4. Effects of organic inputs and inorganic fertilizer combinations on attributes and maize grain yield in Ultisols of Kakamega, Kenya

| Treatment | Plants (ha ⁻¹) | Ears (ha ⁻¹) | Plant Height (m) | Maize Yield (t ha ⁻¹) | Stover Yield (t ha ⁻¹) |
|--------------------------------------|----------------------------|--------------------------|------------------|-----------------------------------|------------------------------------|
| Farmer's practice (control) | 27,654 de | 20,741 c | 2.4 b | 2.5 c | 14.1 c |
| Unweeded, no fertilizer | 32,444 bcd | 25,333 bc | 2.5 b | 3.9 bc | 13.9 c |
| Weeded, no fertilizer | 33,111 abcd | 26,222 bc | 2.6 b | 4.6 bc | 16.1 bc |
| N (60 kg) | 39,556 a | 29,926 ab | 2.6 b | 5.4 b | 22.5 b |
| NP (60 kg N + 60 kg P) | 36,296 abc | 29,037 ab | 2.8 ab | 5.4 b | 20.5 bc |
| NPK (60 kg N + 60 kg P + 40 kg K) | 33,556 abcd | 27,704 abc | 2.6 b | 5.6 b | 21.8 b |
| FYM + 30 kg N | 35,111 abc | 27,259 bc | 2.6 b | 5.3 b | 20.1 bc |
| Tithonia + 30 kg N | 35,185 abc | 27,852 abc | 2.7 ab | 5.0 bc | 22.4 b |
| Desmodium + 30 kg N + 30 kg P | 36,593 abc | 25,778 bc | 2.6 b | 4.7 bc | 21.6 b |
| ¹ Optimal fertilizer | 37,111 ab | 34,518 a | 3.2 a | 9.1 a | 30.8 a |
| Modified farmer's practice | 26,111 e | 24,259 bc | 2.9 ab | 4.0 bc | 20.3 bc |
| Banana intercrop + 60 kg N + 60 kg P | 30,371 cde | 20,741 c | 2.8 ab | 2.6 c | 19.3 bc |
| Mean | 34,060 | 27,084 | 2.7 | 4.8 | 20.5 |
| CV (%) | 12.4 | 17.4 | 11.9 | 14.1 | 22.4 |

Means followed by the same letter within the same column are not significantly different at $P \leq 0.05$, using DMRT.

¹Optimal fertilizer = Optimal fertilizer: (N, P, K, Mg, Zn and B ((NPK was broadcast at planting at the rate of 200, 60 and 120 kg ha⁻¹ of each)), respectively; Mg 20 kg ha⁻¹, Zn 20 kg ha⁻¹ and B 5 kg ha⁻¹)

In Ultisols, optimal fertilizer treatment gave the highest number of ears (34,518) (Table 4). In relation to optimal fertilizer, control and banana intercrop treatments reduced ear number by nearly 40% while modified farmer's practice + fertilizer reduced it (ears) by about 30%. Whereas in relation to optimum fertilizer, desmodium and FYM reduced ear yield by nearly 25%, tithonia and NPK reduced the ears by about 20%. N and NP in relation to optimum fertilizer reduced ear yield by nearly 15%.

3.2.3 Maize plant height

In Alfisols, banana intercrop gave tallest maize plants (2.9 m) (Table 3). Control, unweeded-no fertilizer and weeded-no fertilizer treatments gave plants shorter by nearly 30% compared to banana intercrop treatment. NP, NPK, FYM and desmodium treatments were equal in height though nearly 10% shorter than the banana intercrop treatment. N, tithonia and modified farmer's practice + fertilizer gave plants nearly 22% shorter compared to banana intercrop. In Ultisols, optimal fertilizer gave tallest plants (3.2 m) followed by NP, tithonia, farmer's practice + fertilizer and banana intercrop which were nearly 13% shorter (Table 4). All the other treatments were not significantly different in height though they were about 21% shorter than banana intercrop treatment.

3.2.4 Maize grain yield

In Alfisols, farmer's practice (control) gave lowest maize yield of 1.9 t ha⁻¹ (Table 3). Optimal fertilizer gave maize yield of 6.6 t ha⁻¹, an increment of nearly 250% over the control. NPK and NP had a grain yield increment of 150% and 120% over control. Organic inputs (FYM, tithonia and desmodium) and N gave significantly similar yield with about 100% increment over control. Banana intercrop and modified farmer's practice + fertilizer gave 42% yield improvement over control. In Ultisols, control gave the lowest grain yield of 2.5 t ha⁻¹ (Table 4). Optimal fertilizer gave 9.1 t ha⁻¹, an increment of nearly 260% over control. Treatments N, NP, NPK and FYM gave grain yield increment of about 115% over control. Tithonia, desmodium, unweeded-no fertilizer and weeded-no fertilizer all gave grain yield improvement of nearly 84% while banana intercrop had a 4% increment.

3.2.5 Maize stover yield

In Alfisols, control treatment recorded the lowest stover yield (6.0 t ha⁻¹) (Table 3.) while optimal fertilizer gave 24.3 t ha⁻¹, an increment of about 300%. NPK treatment increased stover yield by nearly 175% over control while N and NP gave an increment of about 130%. Organic inputs (tithonia, FYM and desmodium) and banana treatment doubled stover yield over control. In Ultisols, control treatment gave stover yield of 14.1 t ha⁻¹ compared to optimal fertilizer that gave 30.8 t ha⁻¹, an increment of about 118% over the control (Table 4). Treatments N, NPK, tithonia and desmodium yielded about 56% above the control while NP, FYM, weeded-no fertilizer and banana treatments had an improvement of only 38% above control.

4. DISCUSSION

In our study, both Alfisols and Ultisols had low levels of N that may not be adequate for proper growth of maize (Table 2). Maize requires high levels of N (2%) to perform well [9]. Both soils were also low in P as they both fell below the critical level of 13.0 mg P kg⁻¹ required by maize for normal growth [10]. However, both soils had enough K with values ranging over the critical limit of 0.2 cmol K [25]. Abundance of rainfall in the area is partly responsible for leaching away of nitrates resulting in deficiencies of N. However, P is less prone to leaching. Its deficiency results rather from retrogression process in clay soils. Despite high rainfall in the area, the soils are moderate in Ca and Mg but generally higher in Ultisols compared to Alfisols. These exchangeable bases (K, Ca and Mg) are less prone to leaching because they are normally attached to the negatively charged surfaces of the soil colloids and not in soil solution as nitrates and phosphates. The soil analysis indicated that Ultisols are more productive (Table 2) and hence favour maize production compared to Alfisols. This probably explains why in the study, plant population, plant height, grain and stover yield was higher in Ultisols.

Plant population was higher in Ultisols compared to Alfisols. However in both soils, cover crop (desmodium) and tithonia mulch significantly increased plant population. A similar observation was reported in a wheat crop [26]. Besides increasing soil fertility, mulches and cover crops help in retaining soil moisture, reduce soil temperatures and conserve soil. They also provide a more stable environment for seedling establishment and growth than un-mulched soil [27,28].

Organic inputs with added minimal doses of inorganic fertilizer increased plant height in both soil types compared to non-fertilized treatments. This confirms a similar report by [14,29].

Tithonia biomass is rich in N and P [15,30]. Organic fertilizers exert strong influence on plant growth and development. Luxuriant growth resulting from fertilizer application leads to larger dry matter production, owing to better utilization of solar radiation and more nutrients. The significant increase in plant height reflects the effect of organic fertilizer nutrients. An optimum plant height is positively correlated with productivity of the plant [31].

Even though maize grain and stover yield showed higher improvement in Ultisols due to organic inputs, both soils recorded substantial increase in yield compared to non-fertilized treatments. A similar observation was reported by [32] in Ultisols of Cameroon. Desmodium cover crop, with minimal addition of N had a dramatic effect on maize plant height and grain yields in both soils. [33] reported a similar observation.

5. CONCLUSION

Integrated use of desmodium, tithonia biomass and farmyard manure with minimal addition of inorganic N fertilizer increased maize grain and yield components. The implication is that organic inputs could play significant role in enhancing and replenishing the depleted nutrients and sustain maize production in fragile soils. Combining organic and inorganic fertilizers appear to be advantageous because they both support the plant at early growth stage as well as late in the development stage. Moreover, nutrients from organic sources are not susceptible to leaching. Tithonia biomass that is cheaply and abundantly available is rich in N and P. Several desmodium spp are useful as living mulch and as green manure, as they are able to replenish fertility due to their ability to fix atmospheric N besides being good animal fodder as well as stem borer repellent. Intercropping maize and banana reduced maize grain yield. However, it is not likely that farmers will stop the cropping practice. There is need to study socio-economic benefits of the system before a recommendation can be made. Adoption of integrated soil improvement approaches remains the only feasible option for smallholder farmers to improve maize production in fragile environments as long as the prices of imported inorganic fertilizers remain un-affordable.

ACKNOWLEDGEMENTS

This study was made possible by financial contribution from BIOTA-subproject under Prof. Dr. Mathias Becker of Bonn University and the Director of Kenya Agricultural Research Institute, Nairobi, Kenya. The financial assistance was highly appreciated.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Mills B, Hassan R, Mwangi P. Maize research priorities in Kenya. Measuring research induced economic surplus changes with multiple spatially separated markets. A contribution to KARI maize programme priority setting exercise. Kenya Agricultural Research Institute, Nairobi, Kenya; 1994.

2. Mureithi JG, Macharia PN, Kiiya WW, Gitahi FM, Nekesa CO. Identifying suitable green manure legumes for the major agro-ecological zones in Kenya: Legume screening and database development. In: Mureithi JG, Gachene CKK, Wamuongo JW, Eilitta M. (Eds.), Development and up-scaling of green manure legume technologies in Kenya. Kenya Agricultural Research Institute. Nairobi, Kenya. 2006;9-42.
3. Jaetzold R, Schmidt H. Farm Management Handbook of Kenya. Vol. 2A. Ministry of Agriculture, Kenya, in Cooperation with the German Agricultural Team (GAT) of the German Agency for Technical Cooperation (GTZ); 1982.
4. Kang BT, Juo ASR. Management of low activity clay soils in Tropical Africa for food crop production. In: Beinroth FH, Neel H, Eswaran H. (Eds.). Proceedings of the Fourth International Soil Classification Workshop, Kigali, Rwanda. 1983;450-70.
5. Lal R. Role of mulching techniques in tropical soils and water management. Technical Bulletin 1. IITA, Ibadan, Nigeria. 1974.
6. Cottenie A, Kang BT, Kiekens L, Sajjapongso A. Micronutrients status. In: Greenland DJ. (Ed.). Characterization of soils in relation to their classification and management for crop production. Examples of some areas of the humid tropics. London: Oxford University Press. 1981;149-63.
7. Kang BT, Fox RL. Management of soils for continuous production and controlling nutrient status. In: Greenland DJ. (Ed.). Characterization of soils in relation to their classification and management for crop production. Examples of some areas of the humid tropics. London: Oxford University Press. 1981;202-13.
8. Kanyanjua SM, Ileri L, Wambua S, Nandwa SM. Acidic soils in Kenya: Constraints and remedial options. KARI Technical Note No. 11. Kenya Agricultural Research Institute, Nairobi, Kenya; 2002.
9. Kochian LV, Hoekenga OA, Pineros MA. How do crop plants tolerate acid soils? Mechanisms of aluminum tolerance and phosphorus efficiency. Ann. Rev. Plant Biol. 2004;55:459-93.
10. Landon JR. In Landon JR. (Ed.). "Booker tropical science manual: A handbook for soil survey and agricultural land evaluation in the tropics and subtropics". Longman Science and Technical Publication, Harlow, UK; 1991.
11. Schnier HF, Recke H, Muchena FN, Muriuki AW. Towards a practical approach to fertilizer recommendations for food crop production in smallholder farms in Kenya. Nutr Cycl in Agroecosystems. 1996;47:213-26.
12. Lungu OIM, Dynoodt RFP. Acidification from long-term use of urea and its effect on selected soil properties. African Journal of Food Agriculture Nutrition and Development. 2008;8(1):63-76.
13. Tanimu J, Iwuafor ENO, Odunze AC, Tian G. Effect of Incorporation of Leguminous Cover Crops on Yield and Yield Components of Maize. World Journal of Agricultural Sciences. 2007;3(2):243-249.
14. Giller KE, Cadish G, Ehaliotis C, Adams E, Sakala WD, Mafongoya PL. Building soil nitrogen capital in Africa. P. 151-192. In R. J. Buresh, P. A. Sanchez, and F. Calhoun (Eds.). Replenishing soil fertility in Africa. SSSA Spec. Publ. No. 51. SSSA, Madison, WI; 1997.
15. Ojeniyi SO, Adedina SA, Agbede TM. Soil productivity improving attributes of Mexican Sunflower (*Tithonia diversifolia*) and Siam weed (*Chromolaena odorata*). Emirates Journal of Food and Agriculture. 2012;24(3):243-47.

16. Babajide PA, Akambi WB, Olabode OS, Olaniyi JO, Ajibola AT. Influence of pre-application handling techniques of *Tithonia diversifolia* residues on sesame in southwestern Nigeria. *Journal of Animal and Plant Sciences*. 2012;15(2):2135-46.
17. Jama B, Palm CA, Buresh RJ, Niang' A, Gachengo C, Nziguheba G, Amadalo B. *Tithonia diversifolia* as a green manure for soil fertility improvement in western Kenya: A review. *Agroforestry Systems*. 2000;49:201-21.
18. Nziguheba G, Palm CA, Buresh RJ, Smithson PC. Soil phosphorus fractions and absorption as affected by organic and inorganic sources. *Plant and Soil*. 1998;198:159-68.
19. Gachene CKK, Kimaru G. (Ed.). *Soil Fertility and Land Productivity: A Guide for Extension Workers in the Eastern Africa Region*. RELMA, Nairobi; 2003.
20. Diwani TNA. Implications of resource management on soil fertility in common farm types in Kakamega, western Kenya. PhD Thesis, University of Bonn, Germany; 2009.
21. Hoekstra D, Corbett JD. Sustainable agricultural growth for highlands of East and Central Africa: Prospects to 2020. International Food Policy Research Institute, Washington, DC; 1995.
22. Tittonell P, Vanlauwe B, Leffelaar PA, Rowe EC, Giller KE. Exploring diversity in soil fertility management of smallholder farms in western Kenya. I. Heterogeneity at region and farm scale. *Agriculture, Ecosystems and Environment*. 2005. In press.
23. Okallebo JR, Gathua KW, Woome PL. Laboratory methods of soil and plant analysis: A working manual. TSBF. Soil Science Society of East Africa Technical Publication No. 1. Nairobi, Kenya. 1993.
24. SAS. Statistical Analytical Systems, SAS Institute Inc., Cary, NC. USA; 2007.
25. Gikonyo EW, Smithson PC. Effects of farmyard manure, Potassium and their combinations on maize yields in the high and medium rainfall areas of Kenya. In B. Andre, (Ed.), "Managing Nutrient Cycles to Sustain Soil Fertility in Sub Saharan Africa". Academy Science Publishers (ASP), Nairobi; 2004.
26. Ahmed ZI, Ansar M, Iqbal M, Minhas NM. Effect of planting geometry and mulching on moisture conservation, weed control and wheat growth under rain-fed conditions. *Pakistan Journal of Botany*. 2007;39(4):1189-95.
27. Doshouyu T, Enpig MW, Wqingshan S. A study of the overall effect of straw mulching and ploughing-in and techniques for its applications. *Ningxia Journal of Agriculture, Forestry Science and Technology*. 1995;(2):10-14.
28. Osuiji GE. Tillage and mulching effect on seed zone soil environment and cowpea seedling growth in humid tropics. *Soil Use and Management*. 1990;6(3):152-56.
29. Olubukola SA, Aderemi OO, Adewoyin E, Tinuke D, Akintoye HA, Oladipupo AJ. Comparing the use of *Tithonia diversifolia* and compost as soil amendments for growth and yield of *Celosia argentea*. *New York Science Journal*. 2010;3(6):133-37.
30. Olabode OS, Ogunyani Sola, Akambi WB, Adesina GO, Babajide PA. Evaluation of *Tithonia diversifolia* for soil improvement. *World Journal of agricultural Sciences*. 2007;3(4):503-07.
31. Saeed IM, Abbasi R, Kazim M. Response of maize to nitrogen and phosphorus fertilization under agro-climatic conditions of Rawalokol, Azad Jammu and Kaslim and Kashmir. *Pakistan Journal of Biological Sciences*. 2001;4:949-52.
32. Kaho F, Yemefack M, Nguimgo BAK, Zonkeng CG. The effect of short rotation *Desmodium distortum* planted fallow on the productivity of Ultisols in center Cameroon. *Tropicultura*. 2004;22(2):49-55.

33. Midega CAO, Pittchard J, Salifu D, Pickett JA, Khan ZR. Effects of mulching, N-fertilization and intercropping with *Desmodium uncinatum* on *Striga hermonthica* infestation in maize. Crop Protection. 2013;44:44-49.

© 2013 Achieng et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sciencedomain.org/review-history.php?iid=236&id=2&aid=1726>