









# tRIBS v5.2: A multi-resolution, parallel platform for tributary hydrology in forest applications

L. W. Raming<sup>1,2</sup><sup>¶</sup>, E. R. Vivoni<sup>1,2</sup>, G. Mascaro<sup>1,2</sup>, C. J. Cederstrom<sup>1</sup>,  
A. Ko<sup>1</sup>, A. P. Schreiner-McGraw<sup>1</sup>, and C. Lizarraga-Celaya<sup>1</sup>

<sup>1</sup> School of Sustainable Engineering and the Built Environment, Arizona State University, Tempe, AZ, USA, 85287. <sup>2</sup> Center for Hydrologic Innovations, Arizona State University, Tempe, AZ, USA, 85287. <sup>¶</sup> Corresponding author

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

## Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Open Journals](#) 

## Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

## License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

## Summary

Distributed hydrologic models provide earth scientists and engineers with tools to test and explore hypotheses related to the movement and storage of water within a landscape (Fatichi et al., 2016; Grayson et al., 2002; Keller et al., 2023). The Triangulated Irregular Network (TIN)-based Real-Time Integrated Basin Simulator denoted as tRIBS (Ivanov et al., 2004a, 2004b), is an example of such a process-based distributed model and has been used to address a wide range of problems from hillslope scale processes in ecohydrology (e.g., Mahmood & Vivoni, 2011) to flood management of large watersheds (e.g., Cázares-Rodríguez et al., 2017). Yet, in spite of the extensive use and application of tRIBS to current topics in hydrology, engineering, and the earth sciences, the code has been essentially maintained as a proprietary software. Here, we document the release of tRIBS v5.2, an updated open source code base and its application for forested watersheds that serve as tributaries to larger river systems. This release includes improvements in hydrologic processes with new functionality for simulating channel transmission losses (Schreiner-McGraw & Vivoni, 2018) and reservoir routing (Cázares-Rodríguez et al., 2017). Additionally, it features updated documentation, improved infrastructure for sustainable code development and employment, and improved computational efficiency. These additions provide a robust and sustainable code base, enhancing access and applications of the model.


## Statement of Needs

### Model Description

tRIBS is written in C++ and uses an object oriented design founded on a hydrologically conditioned TIN mesh (see Tucker et al., 2001 and; Vivoni et al., 2004). Building on the work of Garrote & Bras (1995), tRIBS is a continuous hydrologic model simulating the coupled dynamics between the vadose and saturated zones (Vivoni et al., 2007). Accounting for these key hydrologic processes while using computationally efficient methods (Figure 1), tRIBS actively tracks both the evolution of wetting fronts and moisture losses, allowing for continuous simulation throughout wet and dry periods (Ivanov et al., 2004a). With the addition of a single-layer snowpack module (Rinehart et al., 2008), tRIBS can also be applied in cold and mountainous forest environments (Figure 2).

Furthermore, the unstructured TIN mesh in tRIBS provides a multiresolution approach to distributed hydrologic modeling (Figure 1 and Figure 3). As a consequence, tRIBS allows for detailed control in resolving hydrologic dynamics across multiple scales (Vivoni et al., 2004), maximizing model fidelity to physical processes, while minimizing computational expenses. This multi-scaling behavior when paired with parallelization (Vivoni et al., 2011) allows for hyper-resolution modeling (Wood et al., 2011) of hydrologic dynamics at unprecedented scales,

42 from simulations rendered at a point (Vivoni et al., 2010) to 21,000 km<sup>2</sup> watersheds simulated  
43 for a period of 10 years at a nominal cell resolution of ~88 m (Ko et al., 2019).




**Figure 1:** Conceptual overview of tRIBS end-to-end workflow highlighting key processes. Asterisks indicate new features or processes available in tRIBS v5.2. Soil and vegetation parameters may be provided in a raster with continuous values or in a classification table.

## 44 Updates and Modifications

45 Building on core tRIBS functionality, as described by Ivanov et al. (2004a), Rinehart et  
46 al. (2008), and Vivoni et al. (2011), tRIBS v5.2 provides two new process additions: (1)  
47 reservoir routing using the level-pool method (Cázarez-Rodríguez et al., 2017), and (2) channel  
48 transmission losses (Schreiner-McGraw & Vivoni, 2018). In addition, the code base has been  
49 restructured with mechanisms for improved maintainability, robustness, performance, and  
50 integration. This includes updates for code compatibility with newer compilers (Clang and  
51 GCC), the introduction of a CMake build system providing flexibility for compiling serial and  
52 parallel versions, and modernization of the model version control system and documentation.  
53 Additionally, we refactored the snow module (Rinehart et al., 2008), resulting in a reduction  
54 of redundant code and enhanced code organization. Memory leaks associated with parallel

operations were fixed, allowing for increased scalability. Finally, we included Docker images for both tRIBS v5.2 and the auxiliary program MeshBuilder. The Docker image for MeshBuilder facilitates an end-to-end workflow that utilizes METIS (Karypis & Kumar, 1998), enabling rapid and easy partitioning of a watershed domain for parallel simulations (Vivoni et al., 2011). These and other features of tRIBS v5.2 can be explored using two newly updated benchmark scenarios. This first benchmark is a point-scale simulation of the Happy Jack SNOTEL site in northern Arizona, USA (Figure 2). The second is a basin-scale simulation of the Big Spring watershed located in the headwaters of Sycamore Creek in northern Arizona (Figure 3). Both benchmarks are hosted on Zenodo, see Figure 2 and Figure 3 for more details.




**Figure 2:** A point-scale (i.e. a single Voronoi cell) tRIBS simulation of snow water equivalent (SWE) at the Happy Jack SNOTEL site in northern Arizona, USA. Top panel shows the time series of observed (black) and simulated SWE (blue). Bottom panel compares the observed and simulated peak SWE from 2002 to 2017. Dashed black line is a one-to-one relation. The color bar indicates the time difference in the occurrence of the peak SWE for each water year. Zenodo repository for this simulation with additional details can be found at: <https://zenodo.org/records/10909507>.

## Conclusion

Embracing the FAIR principles (Findability, Accessibility, Interoperability, and Reusability; Wilkinson et al., 2016) and recognizing the importance of free and open source software in hydrology (Kabo-bah et al., 2012), here we document the release of tRIBS v5.2. This version represents years of cumulative efforts with major code improvements related to maintainability, robustness, performance, and integration as well as new process based functionality. The benchmarks provided exemplify tRIBS v5.2 applications in forested tributary watersheds of larger river systems. We anticipate that tRIBS v5.2 will be a valuable asset in addressing a wide range of problems for the broader hydrology community.

## Big Spring, Arizona, USA: Map of Mean Evapotranspiration Rate



**Figure 3:** An example of a basin-scale tRIBS simulation showing a spatial map of mean hourly evapotranspiration rates averaged over the course of a 4-year simulation period. Big Spring basin is a tributary to Sycamore Creek in northern Arizona, USA. Zenodo repository for this simulation with additional details can be found at: <https://zenodo.org/records/10909729>.

## Acknowledgements

We thank the tRIBS developers at the Massachusetts Institute of Technology, New Mexico Institute of Mining and Technology, Los Alamos National Laboratory, and Arizona State University, including Valeriy Y. Ivanov, Scott M. Rybarczyk, Greg E. Tucker, Sue Mniszewski, Pat Fasel, and Alex J. Rinehart. We also thank Rafael L. Bras, Dara Entekhabi, and Everett P. Springer, for guidance on model development. Lastly, we are grateful for the support of Elvy Barton and Bruce Hallin for encouraging further development and application of tRIBS to new and pressing problems. Over the years, tRIBS model development has been funded by: Army Research Office, National Science Foundation, National Oceanic and Atmospheric Administration, National Aeronautics and Space Administration, Los Alamos National Laboratory, Salt River Project, and Arizona State University. The most current contribution documented here was facilitated by the Arizona Water Innovation Initiative.

Cázares-Rodríguez, J. E., Vivoni, E. R., & Mascaro, G. (2017). Comparison of Two Watershed Models for Addressing Stakeholder Flood Mitigation Strategies: Case Study of Hurricane Alex in Monterrey, México. *Journal of Hydrologic Engineering*, 22(9), 05017018. [https://doi.org/10.1061/\(ASCE\)HE.1943-5584.0001560](https://doi.org/10.1061/(ASCE)HE.1943-5584.0001560)

Fatichi, S., Vivoni, E. R., Ogden, F. L., Ivanov, V. Y., Mirus, B., Gochis, D., Downer, C. W., Camporese, M., Davison, J. H., Ebel, B., Jones, N., Kim, J., Mascaro, G., Niswonger, R., Restrepo, P., Rigon, R., Shen, C., Sulis, M., & Tarboton, D. (2016). An overview of current applications, challenges, and future trends in distributed process-based models in hydrology. *Journal of Hydrology*, 537, 45–60. <https://doi.org/10.1016/j.jhydrol.2016.03.026>

- 94 Garrote, L., & Bras, R. L. (1995). A distributed model for real-time flood forecasting using  
95 digital elevation models. *Journal of Hydrology*, 167(1), 279–306. [https://doi.org/10.1016/](https://doi.org/10.1016/0022-1694(94)02592-Y)  
96 [0022-1694\(94\)02592-Y](https://doi.org/10.1016/0022-1694(94)02592-Y)
- 97 Grayson, R. B., Blöschl, G., Western, A. W., & McMahon, T. A. (2002). Advances in the  
98 use of observed spatial patterns of catchment hydrological response. *Advances in Water*  
99 *Resources*, 25(8), 1313–1334. [https://doi.org/10.1016/S0309-1708\(02\)00060-X](https://doi.org/10.1016/S0309-1708(02)00060-X)
- 100 Ivanov, V. Y., Vivoni, E. R., Bras, R. L., & Entekhabi, D. (2004a). Catchment hydrologic  
101 response with a fully distributed triangulated irregular network model. *Water Resources*  
102 *Research*, 40(11). <https://doi.org/10.1029/2004WR003218>
- 103 Ivanov, V. Y., Vivoni, E. R., Bras, R. L., & Entekhabi, D. (2004b). Preserving high-resolution  
104 surface and rainfall data in operational-scale basin hydrology: A fully-distributed physically-  
105 based approach. *Journal of Hydrology*, 298(1), 80–111. [https://doi.org/10.1016/j.jhydrol.](https://doi.org/10.1016/j.jhydrol.2004.03.041)  
106 [2004.03.041](https://doi.org/10.1016/j.jhydrol.2004.03.041)
- 107 Kabo-bah, A., Yuebo, X., & Odoi, J. (2012). The Future of Free and Open Source Software  
108 (FOSS) for Hydrology and Water Resources Management. *Hydrol Current Res*, 3(136), 2.  
109 <https://doi.org/10.4172/2157-7587.1000136>
- 110 Karypis, G., & Kumar, V. (1998). A Fast and High Quality Multilevel Scheme for Partitioning  
111 Irregular Graphs. *SIAM Journal on Scientific Computing*, 20(1), 359–392. [https://doi.org/](https://doi.org/10.1137/S1064827595287997)  
112 [10.1137/S1064827595287997](https://doi.org/10.1137/S1064827595287997)
- 113 Keller, A. A., Garner, K., Rao, N., Knipping, E., & Thomas, J. (2023). Hydrological  
114 models for climate-based assessments at the watershed scale: A critical review of existing  
115 hydrologic and water quality models. *Science of The Total Environment*, 867, 161209.  
116 <https://doi.org/10.1016/j.scitotenv.2022.161209>
- 117 Ko, A., Mascaro, G., & Vivoni, E. R. (2019). Strategies to Improve and Evaluate Physics-  
118 Based Hyperresolution Hydrologic Simulations at Regional Basin Scales. *Water Resources*  
119 *Research*, 55(2), 1129–1152. <https://doi.org/10.1029/2018WR023521>
- 120 Mahmood, T. H., & Vivoni, E. R. (2011). A climate-induced threshold in hydrologic response  
121 in a semiarid ponderosa pine hillslope. *Water Resources Research*, 47(9). [https://doi.org/](https://doi.org/10.1029/2011WR010384)  
122 [10.1029/2011WR010384](https://doi.org/10.1029/2011WR010384)
- 123 Rinehart, A. J., Vivoni, E. R., & Brooks, P. D. (2008). Effects of vegetation, albedo, and  
124 solar radiation sheltering on the distribution of snow in the Valles Caldera, New Mexico.  
125 *Ecohydrology*, 1(3), 253–270. <https://doi.org/10.1002/eco.26>
- 126 Schreiner-McGraw, A. P., & Vivoni, E. R. (2018). On the Sensitivity of Hillslope Runoff and  
127 Channel Transmission Losses in Arid Piedmont Slopes. *Water Resources Research*, 54(7),  
128 4498–4518. <https://doi.org/10.1029/2018WR022842>
- 129 Tucker, G., Lancaster, S. T., Gasparini, N. M., Bras, R. L., & Rybarczyk, S. M. (2001).  
130 An object-oriented framework for distributed hydrologic and geomorphic modeling using  
131 triangulated irregular networks. *Computers & Geosciences*, 27(8), 959–973. [https://doi.](https://doi.org/10.1016/S0098-3004(00)00134-5)  
132 [org/10.1016/S0098-3004\(00\)00134-5](https://doi.org/10.1016/S0098-3004(00)00134-5)
- 133 Vivoni, E. R., Entekhabi, D., Bras, R. L., & Ivanov, V. Y. (2007). Controls on runoff generation  
134 and scale-dependence in a distributed hydrologic model. *Hydrology and Earth System*  
135 *Sciences*, 11(5), 1683–1701. <https://doi.org/10.5194/hess-11-1683-2007>
- 136 Vivoni, E. R., Ivanov, V. Y., Bras, R. L., & Entekhabi, D. (2004). Generation of Triangulated  
137 Irregular Networks Based on Hydrological Similarity. *Journal of Hydrologic Engineering*,  
138 9(4), 288–302. [https://doi.org/10.1061/\(ASCE\)1084-0699\(2004\)9:4\(288\)](https://doi.org/10.1061/(ASCE)1084-0699(2004)9:4(288))
- 139 Vivoni, E. R., Mascaro, G., Mniszewski, S., Fasel, P., Springer, E. P., Ivanov, V. Y., &  
140 Bras, R. L. (2011). Real-world hydrologic assessment of a fully-distributed hydrological



- 141 model in a parallel computing environment. *Journal of Hydrology*, 409(1), 483–496.  
142 <https://doi.org/10.1016/j.jhydrol.2011.08.053>
- 143 Vivoni, E. R., Rodríguez, J. C., & Watts, C. J. (2010). On the spatiotemporal variability of soil  
144 moisture and evapotranspiration in a mountainous basin within the North American monsoon  
145 region. *Water Resources Research*, 46(2). <https://doi.org/10.1029/2009WR008240>
- 146 Wilkinson, M. D., Dumontier, M., Aalbersberg, Ij. J., Appleton, G., Axton, M., Baak, A.,  
147 Blomberg, N., Boiten, J.-W., Silva Santos, L. B. da, Bourne, P. E., Bouwman, J., Brookes,  
148 A. J., Clark, T., Crosas, M., Dillo, I., Dumon, O., Edmunds, S., Evelo, C. T., Finkers,  
149 R., ... Mons, B. (2016). The FAIR Guiding Principles for scientific data management and  
150 stewardship. *Scientific Data*, 3(1), 160018. <https://doi.org/10.1038/sdata.2016.18>
- 151 Wood, E. F., Roundy, J. K., Troy, T. J., Beek, L. P. H. van, Bierkens, M. F. P., Blyth,  
152 E., Roo, A. de, Döll, P., Ek, M., Famiglietti, J., Gochis, D., Giesen, N. van de, Houser,  
153 P., Jaffé, P. R., Kollet, S., Lehner, B., Lettenmaier, D. P., Peters-Lidard, C., Sivapalan,  
154 M., ... Whitehead, P. (2011). Hyperresolution global land surface modeling: Meeting a  
155 grand challenge for monitoring Earth's terrestrial water. *Water Resources Research*, 47(5).  
156 <https://doi.org/10.1029/2010WR010090>

DRAFT