

*viis; luteo-aurantid, fasciá latá albidá, fusco grandimaculatá, prope foramina ornatá.*

*Hab. — ?*

<sup>1954</sup> B. M. HALIOTIS CRUENTA. *Hal. testá ovatá, anticè subattenuatá, spirá elevatiusculá, spiraliter peculiariter undato- et corrugato-striatá; foraminibus subapproximatis, octonis perviis, sanguineo albi-punctato et albido sanguineo-punctato pulcherrimè variegatá.*

*Hab. New Zealand.*

<sup>13. 1954</sup> HALIOTIS INCISA. *Hal. testá ovatá, medio subdepressá, spiraliter incisá, subtuberculiferá; foraminibus amplis, quaternis perviis; albidá et purpureo-viridi marmoratá, albido minutissimè rufo-punctatá.*

*Hab. — ?*

The next paper was also communicated by Mr. Lovell Reeve, and contained "Descriptions of fifty-four new species of *Mangelia*, from the collection of H. Cuming, Esq.":—

The *Mangelia* are nearest allied to those aberrant species of *Pleurotoma* in which the predominant character of that genus, the fissure in the upper extremity of the lip, becomes modified into a somewhat obscure sinus. Their general aspect is that of a more or less fusiform *Marginella*, without plaits or polished exterior; distinguished, on the other hand, by a row of faint wrinkle-like denticulations on the inner surface of the lip and columella, and a gutter-like sinus in the lip at its junction with the body-whorl.

MANGELIA SICULA. *Mang. testá subfusiformi, spirá acuminatá, anfractibus rotundatis, gibbosiuseculis, concentricè costatis, levibus; aperturá brevi, ovatá, sinu subconspicuo; intus extusque castaneo-fuscá, labro flavicante, fusco-lineato.* B M

*Hab. Sicily.*

MANGELIA VEXILLUM. *Mang. testá oblongo-ovatá, anfractibus superne depressis, nodosis, costis e nodis descendentes, superficie totá decussatim striatá, quasi subtilissimè decussatá; aurantio-luteá, fasciis albicantibus angustis undique cingulatá.*

*Hab. Ilo Ilo, isle of Panhay, Philippines (found under stones); Cuming.*

MANGELIA LYRA. *Mang. testá trigono-fusiformi, anfractibus superne angulatis, longitudinaliter costatis, costis subobliquis, ad angulum incrassatis; castaneo-fuscá, lineis albicantibus decussatis, labro albicante.*

*Hab. Island of Ticao, Philippines (found on the sands); Cuming.*

MANGELIA ANTILLARUM. *Mang. testá subpyriformi-ovatá, levigatá, longitudinaliter concentricè costatá, costis fortibus, subobtusis, distantibus; cinereo-carned, costis fasciá rubidá tinctis, labro albicante, rubido fasciatim tincto.*

*Hab. West Indies.*



**MANGELIA MARGINELLOIDES.** *Mang. testâ pyriformi-ovatâ, spirâ brevi, acutâ; anfractibus supernè angulatis, longitudinaliter multicostratis, costis tenuibus, crebris, suturis descendentibus; lacteo-cæruleâ aut cinereo-fuscâ, lineis rubidis cingulatâ, anfractûs ultimi parte supra angulum maculâ grandi nigricante conspicuè tinctâ.*

*Hab.* Island of Burias, Philippines (found in sandy mud at the depth of seven fathoms); Cuming.

**MANGELIA FUNICULATA.** *Mang. testâ trigono-fusiformi, subelongatâ, anfractibus supernè depressis, longitudinaliter tenuicostatis, lævibus; cinereo-fuscâ, costis labroque albicantibus.*

*Hab.* Islands of Ticao and Masbate, Philippines (found on the sands); Cuming.

29, 1958 **MANGELIA CAVERNOSA.** *Mang. testâ oblongo-ovatâ, spirâ angulato-turritâ; anfractibus supernè angulatis, intra costis cavernosis, costis fortibus, obtusis, suturis descendentibus; albâ, aurantio-fusco hic illic sparsim maculatâ.*

*Hab.* Island of Ticao, Philippines (found on the sands); Cuming.

**MANGELIA CYLINDRICA.** *Mang. testâ cylindræo-fusiformi, apice acutâ, longitudinaliter subtiliter costatâ, transversim elevato-striatâ; pellucido-albâ, fuscescente tinctâ, aurantio-fusco infra suturas maculatâ.*

*Hab.* Cagayan, island of Mindanao, Philippines (found in sandy mud at the depth of twenty-five fathoms); Cuming.

**MANGELIA CAPILLACEA.** *Mang. testâ ovato-fusiformi; spiræ suturis profundis; anfractibus supernè depressis, transversim subtilissimè et creberrimè elevato-striatis, longitudinaliter costatis, angustis, suberectis, supernè leviter mucronatis; fuscescente, lineis subtilibus fuscescentibus zonatâ.*

*Hab.* Island of Burias, Philippines (found among coarse sand at the depth of seven fathoms); Cuming.

**MANGELIA GRACILIS.** *Mang. testâ gracili-fusiformi, transversim subtilissimè striatâ, longitudinaliter costatâ, costis angustis; albâ, castaneo-fusco indistinctè zonatâ et maculatâ.*

*Hab.* Island of Ticao, Philippines (found under stones at low water); Cuming.

**MANGELIA LAMELLATA.** *Mang. testâ fusiformi-ovatâ, spiræ suturis profundis, subcavernosis, anfractibus transversim elevato-striatis, striis fortibus, subdistantibus, longitudinaliter costatis, costis angustis, erectis, lamellæformibus, supernè leviter mucronatis; albidâ, fusco pallidissimè zonatâ.*

*Hab.* Isle of Burias, Philippines (found among coarse sand at the depth of seven fathoms); Cuming.

**MANGELIA VITTATA.** *Mang. testâ oblongo-ovatâ, subconicâ, spirâ brevi, obtusâ; anfractibus longitudinaliter costatis, costarum in-*



*terstitiis striis fortibus clathratis; albidá, olivaceo-fusco latifasciatá, costarum parte olivaceo-fusca albipunctatá.*

*Hab.* Island of Ticao, Philippines (found on the sands); Cuming.

**MANGELIA ZONATA.** *Mang. testá abbreviato-fusiforimi, spirá breviusculá, turritá suturis profundis; anfractibus longitudinaliter costatis, costis e suturis descendentibus; lævigatá; albá, zoná conspicuá aurantio-fusca cingulatá.*

*Hab.* Island of Ticao, Philippines (found on the sands); Cuming.

**MANGELIA INTERRUPTA.** *Mang. testá oblongo-ovatá, spirá brevi, anfractibus supernè nodosis, costis e nodis descendentibus; albá, lineis nigris subtilissimis transversis interruptis, creberrimè dispositis, inter costas ornatá.*

*Hab.* Island of Ticao, Philippines (found on the sands); Cuming.

**MANGELIA RETICULATA.** *Mang. testá fusiformi-ovatá, spirá breviusculá, apice acuminatá; anfractibus supernè perpendiculariter compressis, deinde tumidiusculis et longitudinaliter costatis, costarum interstitiis exiliter fusco-punctatis.*

*Hab.* Island of Ticao, Philippines (found on the reefs); Cuming.

**MANGELIA PULCHELLA.** *Mang. testá fusiformi-ovatá, subcylindraceá, spirá breviusculá, apice acuminatá; anfractibus supernè rotundatis, longitudinaliter multicostatis, costis tenuibus gracilibus, concentricè dispositis, anfractuum superficie totá exilissimè reticulatá; luteo-albicante, maculis quadratis parvis rufis costarum interstitiis fasciatim cingulatá.*

*Hab.* Island of Ticao, Philippines (found on the sands); Cuming.

**MANGELIA FUSIFORMIS.** *Mang. testá fusiformi, anfractibus supernè subangulatis, ad angulum nodosis, costis tenuibus subsuperciliariis e nodis descendentibus, transversim creberrimè striatis; luteo-albicante, punctis perpaucis aurantio-fuscis tinctá.*

*Hab.* Island of Corrigidor, Philippines (found among coarse sand at the depth of ten fathoms); Cuming.

**MANGELIA LYRICA.** *Mang. testá fusiformi, utrinque acuminatá, anfractibus longitudinaliter concentricè costatis, transversim elevato-striatis; fuscescente, aurantio-fusco pallidè et indistinctè fasciatá.*

*Hab.* Island of Burias, Philippines (found among coarse sand at the depth of seven fathoms); Cuming.

**MANGELIA GIBBOSA.** *Mang. testá ovato-conicá, spirá brevissimá; anfractibus supernè gibbosis et nodulosis, longitudinaliter costatis, lævigatis; cinereo-albicante, lineis aurantio-fuscis exilibus cingulatá, dorso supernè nigricante tincto.*

*Hab.* Island of Ticao, Philippines (found on the reefs); Cuming.

**MANGELIA MACULATA.** *Mang. testá subfusiformi, basi truncatá, longitudinaliter costatá, costis tenuibus subdistantibus, concentricè dispositis; sinu latiusculo; albá, maculis subquadratis aurantio-fuscis inter costas.*



*Hab.* Island of Ticao, Philippines (found under stones at low water); Cuming.

**MANGELIA TURRICULA.** *Mang. testá fusiformi-turritá, suturis profundis; anfractibus plano-depressis, longitudinaliter concentricè costatis; albicante, lineis subtilissimis aurantio-fuscis obsoletè cingulatá.*

*Hab.* Island of Ticao, Philippines (found on the sands); Cuming.

**MANGELIA COLUMBELLOIDES.** *Mang. testá ovatá, spirá brevi, acutá; anfractibus supernè leviter rotundatis, transversim elevato-striatis, longitudinaliter tenuicostatis, labro medio tumido, intus fortiter denticulato; nived, costis eximè aurantio-fusco punctatis.*

*Hab.* Baclayon, island of Bohol, Philippines (found on mud banks); Cuming.

**MANGELIA CONOHELICOIDES.** *Mang. testá ovato-conicá, spirá brevissimá, acutá; anfractibus transversim elevato-striatis, longitudinaliter multiliratis, liris tenuibus, ante suturas evanidis; luteo-albicante, dorso maculá grandi aurantio-fusá interdum tincto.*

*Hab.* Daleguete, isle of Zebu, Philippines (found under stones); Cuming.

**MANGELIA TENEBROSA.** *Mang. testá subfusiformi, spirá acuminato-turritá; anfractibus supernè plano-depressis, suturis profundis, transversim striatis, longitudinaliter costatis, costis distantibus; intus extusque castaneo-fusá.*

*Hab.* Cagayan, island of Mindanao (found in sandy mud at the depth of twenty-five fathoms); Cuming.

**MANGELIA NOVÆ HOLLANDIÆ.** *Mang. testá ovatá, infernè attenuatá, subconicá, spirá brevi; anfractibus supernè rotundatis, longitudinaliter obliquè costatis; livido-cinereá, apice basique nigricantibus, labro albido, aperturae fauce fusá.*

*Hab.* Swan River.

**MANGELIA LIVIDA.** *Mang. testá subfusiformi-ovatá, apice acuminatá; anfractibus longitudinaliter concentricè costatis, costis tenuibus; livido-carnéa.*

*Hab.* Island of Ticao, Philippines (found on the reefs); Cuming.

**MANGELIA ABYSSICOLA.** *Mang. testá oblongo-ovatá, utrinque conspicuè attenuatá, longitudinaliter costatá, lineis elevatis cingulatá; albidá, fusco zonatá.*

*Hab.* Island of Mindanao, Philippines (found in sandy mud at the depth of twenty-five fathoms); Cuming.

**MANGELIA BICOLOR.** *Mang. testá oblongá, concentricè costatá, costarum interstitiis subtilissimè striatis; supra albidá, infra plumbeá.*

*Hab.* Island of Ticao, Philippines; Cuming.

**MANGELIA FUNEBRIS.** *Mang. testá oblongá, concentricè costatá,*



*costarum interstitiis lævibus; albidâ fasciatâ plumbeâ latâ cingulatâ.*

*Hab.* Island of Ticao (found under stones at low water); Cuming.

**MANGELIA STROMBOIDES.** *Mang. testâ fusiformi-ovatâ, spirâ subturritâ; anfractibus supernè angulatis, longitudinaliter costatis, costis ad angulum nodosis, interstitiis subtilissimè striatis; albidâ.*

*Hab.* Island of Bohol, Philippines.

**MANGELIA PALLIDA.** *Mang. testâ ovatâ, concentricè costatâ, costis subobtusis; albâ.*

*Hab.* Island of Ticao, Philippines; Cuming.

**MANGELIA PESSULATA.** *Mang. testâ cylindræco-oblongâ, spirâ breviusculâ, longitudinaliter eximè subobliquè costatâ, costarum interstitiis subtilissimè striatis; niveâ.*

*Hab.* Philippine Islands; Cuming.

**MANGELIA RIGIDA.** *Mang. testâ ovatâ, longitudinaliter fortiter tuberculato-costatâ, transversim subtilissimè striatâ; fuscescente.*

*Hab.* Ægean Sea; Forbes.

**MANGELIA ELEGANS.** *Mang. testâ oblongâ, spirâ breviusculâ, acuminatâ, anfractibus supernè angulatis, longitudinaliter costatis, costis angustis, ad angulum mucronatis, costarum interstitiis pulcherrimè elevato-striatis; lutescente-albâ, exilissimè fusco-zonatâ.*

*Hab.* Island of Mindoro, Philippines; Cuming.

**MANGELIA LINEATA.** *Mang. testâ ovatâ, spirâ acuminatâ; concentricè obtuso-costatâ, lævigatâ; carneo-fuscescente, lineis saturatioribus undique cingulatâ.*

*Hab.* — ?

**MANGELIA PLANILABRUM.** *Mang. testâ fusiformi, utrinque acuminatâ, lævissimâ, anfractibus supernè depressis, longitudinaliter costatis; labro planulato, supra subangulato; cinereo-purpurascete, albizonato.*

*Hab.* Island of Ticao, Philippines; Cuming.

**MANGELIA HORNBECKII.** *Mang. testâ ovatâ, spirâ breviusculâ, acutâ, suturis profundis, subcavernosis, longitudinaliter costatâ, costis prominentibus, transversim subtilissimè striatâ, albâ.*

*Hab.* Island of St. Thomas, West Indies; Dr. Hornbeck.

**MANGELIA CASTANEA.** *Mang. testâ oblongâ, spirâ acuminatâ, concentricè tenuicostatâ, costarum interstitiis striatis; castaneo-fuscâ.*

*Hab.* Island of Burias, Philippines; Cuming.

**MANGELIA PUSILLA.** *Mang. testâ fusiformi-ovatâ, lævigatâ, longitudinaliter costatâ, costis solidiusculis obtusis; albidâ, castaneo copiosè tinctâ et lineatâ.*

*Hab.* — ?



**MANGELIA MARMOROSA.** *Mang. testá ovatá, solidiusculá, spirá breviusculá; longitudinaliter costatá, costis obesis, striis elevatis, decussatis; sinu amplo; albá, aurantio-fusco perparce maculatá.*  
*Hab.* — ?

**MANGELIA CASTA.** *Mang. testá ovato-turritá, longitudinaliter costatá, costis tenuibus subdistantibus, lævigatá; aperturá brevi, basi truncatá; carneo-fuscá, obscurè fasciatá, costis albicantibus.*  
*Hab.* — ?

**MANGELIA OBELISCUS.** *Mang. testá subulatá, hexagoná, basi truncatá, striis pulcherrimis elevatis undique creberrimè cingulatá, longitudinaliter costatá, costis distantibus, sequentibus; aperturá minutá, ovatá; sordidè albá, aperturæ fauce lutescente.*

*Hab.* Islands of Corrigidor, Philippines (found among coarse sand at the depth of ten fathoms); Cuming.

**MANGELIA BALTEATA.** *Mang. testá elongatá, subfusiformi, longitudinaliter costatá costis angustis, distantibus; albá, zoná fuscá conspicuá cingulatá.*  
*Hab.* — ?

**MANGELIA ASTRICTA.** *Mang. testá subfusiformi-ovatá, spirá breviusculá, suturis profundis; longitudinaliter costatá, costis crebriusculis; albidá, zoná fuscá angustá cingulatá.*  
*Hab.* — ?

**MANGELIA BADIA.** *Mang. testá fusiformi-ovatá, concentricè plicato-costatá, transversim fortiter striatá; castaneo-fuscá.*  
*Hab.* — ?

**MANGELIA PELLUCIDA.** *Mang. testá ovatá, utrinque attenuatá, lævigatá, pellucidá, nitidá, longitudinaliter crebricostatá; albidá, basi fuscá.*  
*Hab.* — ?

**MANGELIA ANGULATA.** *Mang. testá fusiformi-ovatá, anfractibus medio peculiariter angulatis, suturis profundis, longitudinaliter costatis, costis angulos super mucronatis; albidá, fuscescente lineatá.*

*Hab.* Bay of Manila (found in sandy mud at the depth of four fathoms); Cuming.

**MANGELIA PURA.** *Mang. testá oblongo-ovatá, spirá subturritá, suturis profundis; anfractibus concentricè costatis; albidá, maculis perpaucis aurantio-fuscis.*  
*Hab.* — ?

**MANGELIA SOLIDA.** *Mang. testá cylindraceo-ovatá, utrinque attenuatá, solidá, undique creberrimè granoso-clathratá; aperturá longiusculá; purpurascete.*

*Hab.* Island of Burias, Philippines (among sand at the depth of seven fathoms); Cuming.

**MANGELIA DERELICTA.** *Mang. testá ovatá, longitudinaliter fortiter concentricè costatá, transversim subobsoletè striatá; fuscescente.*  
*Hab.* — ?



MANGELIA ZEBUENSIS. *Mang. testá ovato-oblongá, spirá acuminatá, basi subattenuatá, concentricè fortiter costatá, transversim creberrimè striatá; fuscescente.*

*Hab.* Island of Zebu, Philippines (found in sandy mud at the depth of four fathoms); Cuming.

MANGELIA CINCTA. *Mang. testá subfusiformi-ovatá, spirá turrítá, suturis subprofundis; anfractibus supernè angulatis, longitudinaliter costatis, costis distantibus; albidá, anfractibus fasciá latá fuscescente supernè cinctis.*

*Hab.* Island of Bohol, Philippines (found under stones at low water); Cuming.

MANGELIA DIGITALIS. *Mang. testá fusiformi-ovatá, solidá, undique creberrimè granoso-clathratá; albicante, zonis duabus purpureis angustis cingulatá.*

*Hab.* — ?

MANGELIA NANA. *Mang. testá, ovatá, spirá brevi, turrítá, apice acutá; anfractibus supernè angulatis, longitudinaliter obliquè costatis, interstitiis cavis, subtiliter striatis.*

*Hab.* Island of Mindanao, Philippines (found in sandy mud at the depth of twenty-five fathoms); Cuming.

July 28, 1846.

No business was transacted.



Reeve, Lowell Augustus. 1846. "Descriptions of fifty-four New Species of Mangelia, from the collection of Hugh Cuming, Esq." *Proceedings of the Zoological Society of London* 14(07-14), 59–65.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/46217>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/387981>

**Holding Institution**

Natural History Museum Library, London

**Sponsored by**

Natural History Museum Library, London

**Copyright & Reuse**

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.