



Biomass Coverage in Forage Fish Taxonomy Group




Biomass Coverage Over Time in Forage Fish Taxonomy Group
1/3 of Stocks with highest average biomass




Biomass Coverage Over Time in Forage Fish Taxonomy Group

1/3 of Stocks with medium average biomass




Biomass Coverage Over Time in Forage Fish Taxonomy Group

1/3 of Stocks with lowest average biomass




Peruvian anchoveta North-Central Peruvian coast – Forage Fish Taxonomy Group

Stock represents 24% of average absolute surplus production




Herring ICES 1–2–4a–5–14a (Norwegian Spring Spawners) – Forage Fish Taxonomy Group

Stock represents 4% of average absolute surplus production




Japanese sardine Pacific Coast of Japan – Forage Fish Taxonomy Group

Stock represents 6% of average absolute surplus production




Sardine North West Africa Zone C – Forage Fish Taxonomy Group


Stock represents 8% of average absolute surplus production



Peruvian anchoveta North–Central Peruvian coast
in Forage Fish Taxonomy Group




Herring ICES 1-2-4a-5-14a (Norwegian Spring Spawners) in Forage Fish Taxonomy Group



First and last years with recorded surplus production are labeled

Japanese sardine Pacific Coast of Japan in Forage Fish Taxonomy Group



First and last years with recorded surplus production are labeled

Sardine North West Africa Zone C in Forage Fish Taxonomy Group

