

Inflammation in stroke: the role of cholinergic, purinergic and glutamatergic signaling

Abraham Martín, María Domercq and Carlos Matute

Ther Adv Neurol Disord

2018, Vol. 11: 1–14

DOI: 10.1177/
1756286418774267

© The Author(s), 2018.
Reprints and permissions:
[http://www.sagepub.co.uk/
journalsPermissions.nav](http://www.sagepub.co.uk/journalsPermissions.nav)

Abstract: The inflammatory response is a major factor in stroke pathophysiology and contributes to secondary neuronal damage in both acute and chronic stages of the ischemic injury. Recent work in experimental cerebral ischemia has demonstrated the involvement of neurotransmitter signaling in the modulation of neuroinflammation. The present review discusses recent findings on the therapeutic potential and diagnostic perspectives of cholinergic, purinergic and glutamatergic receptors and transporters in experimental stroke. It provides evidence of the role of neurotransmission signaling as a promising inflammatory biomarker in stroke. Finally, recent molecular imaging studies using positron emission tomography of cholinergic receptors and glutamatergic transporters are outlined along with their potential as novel anti-inflammatory therapy to reduce the outcome of cerebral ischemia.

Keywords: cerebral ischemia, cholinergic, glutamatergic, inflammation, purinergic, stroke

Received: 1 March 2018; accepted in revised form: 26 March 2018.

Introduction

The ischemic injury is a complex system of pathological processes including excitotoxicity, oxidative stress, inhibition of protein synthesis, programmed cell death and inflammation, among others, which occurs within minutes until hours and even days and months after the brain vessel occlusion.¹ This pathological process is initiated as a result of the impairment of energy levels that maintain ionic gradients,² inducing a loss of membrane potential in neurons and glia, a process known as anoxic depolarization (AD).³ AD spreads as a self-propagating wave-like depolarization across the susceptible brain parenchyma due to the release of K⁺, glutamate and adenosine triphosphate (ATP).⁴ As a result, increased entry of Ca²⁺ into the cell is thought to initiate a cascade of cytoplasmic and nuclear events, including activation of proteolytic enzymes, production of reactive oxygen species (ROS), lipid peroxidation and membrane damage, the inhibition of protein synthesis, cerebral edema formation, cellular DNA fragmentation, and activation of apoptotic cell death which together lead to primary ischemic damage and the subsequent activation of inflammation.⁵ The inflammatory

response plays a pivotal role in exerting both beneficial and detrimental effects on the acute ischemic injury and the functional recovery after stroke.⁶ In this review, we focus on the evidence regarding inflammatory biomarkers and cells that control the neuroinflammatory reaction and related mechanisms after stroke. In particular, the recent findings on the potential therapeutic and diagnostic perspectives of cholinergic, purinergic and glutamatergic biomarkers for the use in both therapy and diagnosis of stroke outcome will be also discussed.

Inflammation after brain ischemia

Inflammation is a complex response to necrotic cells and the subsequent generation of ROS during the secondary injury following stroke.⁷ Early after ischemia, the release of a repertoire of different proinflammatory prostaglandins, cytokines and chemokines results in the activation of microglia, the resident immune cell population of the brain.⁸ Once activated, microglia leads to the induction of adhesion molecules, integrins and selectins in both the brain vasculature and immune cells through the release of proinflammatory cytokines.⁹

Correspondence to:
Abraham Martín
Experimental Molecular
Imaging, Molecular
Imaging Unit, CIC
biomaGUNE, Pº Miramon
182, San Sebastian, Spain
amartin@cicbiomagune.es

María Domercq
Carlos Matute
Department of
Neurosciences, University
of the Basque Country,
Barrio Sarriena s/n, Leioa,
Spain
Achucarro Basque Center
for Neuroscience-UPV/
EHU, Zamudio, Spain
Instituto de Salud
Carlos III, Centro de
Investigación Biomédica
en Red de Enfermedades
Neurodegenerativas
(CIBERNED), Leioa, Spain

Hence, the overexpression of all these molecules mediates leukocyte adhesion to the vascular endothelium leading to subsequent entry into the brain tissue.¹⁰ In addition, leukocyte infiltration is amplified by the disruption of the blood–brain barrier (BBB) through the release of cytotoxic agents such as metalloproteinases by microglia and other infiltrated leukocytes.⁷ Microglia and infiltrated macrophages can be classified into at least two subsets with distinct molecular phenotypes and functions depending on the activation pathway. The ‘classically activated’ proinflammatory microglial cells and macrophages play a central role in host defense against pathogens, but they can also damage healthy cells as neurons and glial cells. In contrast, anti-inflammatory phenotypes or ‘alternatively activated’ cells downregulate inflammation and promote tissue remodeling or repair and angiogenesis after stroke.¹¹ Thus, a plethora of pathways and mediators might determine the final fate of these cells under a pathological situation in the brain.¹² Both *ex vivo* and *in vivo* studies have suggested that many neurotransmitter receptors including cholinergic, purinergic and glutamatergic receptors or transporters are overexpressed in microglia under pathological situations, behaving as modulators of the inflammatory response¹² (Table 1). Therefore, the precise management of neuroreceptors as inflammatory biomarkers may ultimately promote novel therapeutic and diagnostic strategies for treating ischemic damage.

Cholinergic receptors

Cholinergic receptors show neuronal protective effects which have been related to the modulation of immune cells by the cholinergic anti-inflammatory pathway.⁴⁴ The mechanism for inhibition of cytokine release is attributable to acetylcholine (ACh) through the inflammatory reflex of the vagus nerve.^{45–47} Macrophages and other immune cells express acetylcholine receptors (AChRs), which are able to transduce an intracellular signal that inhibits cytokine synthesis.⁴⁸ To date, the most characterized cholinergic receptors as inflammatory modulators are those formed by the $\alpha 7$ subunit of the AChR. Studies in mice have shown that activation of these receptors is required for acetylcholine inhibition of macrophage tumor necrosis factor (TNF) release, becoming an essential key for the regulation of inflammation.⁴⁸ In addition, pharmacological stimulation of $\alpha 4\beta 2$ nicotinic receptors promotes inhibition of amyloid toxicity to cortical neurons

and modulation of inflammation after cerebral ischemia in rats.^{23,49}

Nicotinic acetylcholine receptors (nAChRs) are ligand-gated ion channels composed of five subunits ($\alpha 2$ – $\alpha 10$ and $\beta 2$ – $\beta 4$) and the most abundant nAChRs in the mammalian brain are heteromeric receptors containing $\alpha 4$ and $\beta 2$ subunits and homomeric $\alpha 7$.⁵⁰ The expression of nAChRs has been described in a large variety of neural cells,⁵¹ and non-neuronal cells such as microglia, astrocytes, oligodendrocyte precursor cells and endothelial cells,^{52–56} where they can decrease the extent of cell death and enhance synaptic plasticity.⁵⁷ Therefore, these receptors are potential therapeutic targets for several neurodegenerative disorders such as Parkinson’s disease, schizophrenia, depression and Alzheimer’s disease.⁵⁸ nAChRs have been proposed as promising novel candidates for the treatment of neuroinflammation following stroke.⁵⁷

As previously discussed, the cholinergic anti-inflammatory pathway is controlled by vagus nerve stimulation and particularly *via* the $\alpha 7$ -nAChRs expressed on innate immune cells, evidencing the effect of the peripheral cholinergic system.⁴⁸ Some studies have shown that stimulation of the vagus nerve attenuates cerebral ischemia injury and reperfusion.^{59,60} A brief stimulation of the vagus nerve after both permanent and cerebral ischemia displayed a reduction in the protein levels of $\alpha 7$ receptors followed by a reduction on inflammation, apoptosis and neuroprotection through the $\alpha 7$ -nAChR/JAK2 anti-inflammatory pathway.^{59,60} Accordingly, the pharmacological activation of $\alpha 7$ receptors with selective agonists confirmed reduction of the brain injury and neuroprotection after intracerebral hemorrhage in rodent models through reduction of the inflammatory response.^{13,14} In the latter study, the use of methyllycaconitine as a potent and selective $\alpha 7$ -nAChR antagonist reversed the anti-inflammatory potential of the agonist PHA-543613 after intracerebral hemorrhage in mice.¹⁴ Following cerebral ischemia, several preclinical studies have observed that both the local upregulation and the pharmacological activation of nicotinic receptors protects the brain against ischemic injury, suggesting the protective central cholinergic effect and the potential role of these receptors as promising inflammatory biomarkers.^{15–23}

Table 1. Inflammatory biomarkers targeting cholinergic, purinergic and glutamatergic systems in stroke.

Name of the target	Neurotransmitter system	Biological activity	References
$\alpha 7$	Cholinergic	Neuroprotective and anti-inflammatory action	Duris <i>et al.</i> , ¹³ Krafft <i>et al.</i> , ¹⁴ Han <i>et al.</i> , ¹⁵ Zou <i>et al.</i> , ¹⁶ Han <i>et al.</i> , ¹⁷ Guan <i>et al.</i> , ¹⁸ Shimohama <i>et al.</i> , ¹⁹ Kalappa <i>et al.</i> , ²⁰ Fujiki <i>et al.</i> , ²¹ Colás <i>et al.</i> , ²²
$\alpha 4\beta 2$	Cholinergic	Inhibition of inflammation	Martin <i>et al.</i> , ²³
A ₁	Purinergic	Modulation of interleukin release	Burnstock and Boeynaems ²⁴
A _{2A}	Purinergic	Control of immune cell infiltration after cerebral ischemia	Verkhatsky <i>et al.</i> , ²⁵ Puchalowicz <i>et al.</i> , ²⁶ Vuorimaa <i>et al.</i> , ²⁷
P2X7	Purinergic	Neuroprotective activity and modulation of inflammation	Mayne <i>et al.</i> , ²⁸ Li <i>et al.</i> , ²⁹ Yu <i>et al.</i> , ³⁰ Dai and Zhou, ³¹ Trodec <i>et al.</i> , ³² Collo <i>et al.</i> , ³³ Skape <i>et al.</i> , ³⁴
P2X4	Purinergic	Control of the inflammatory reaction	Suzuki <i>et al.</i> , ³⁵ Chu <i>et al.</i> , ³⁶ Monif <i>et al.</i> , ³⁷
P2Y12	Purinergic	Modulation of the inflammatory reaction and platelet aggregation	Melani <i>et al.</i> , ³⁸ Vazquez-Villoldo <i>et al.</i> , ³⁹ Cavaliere <i>et al.</i> , ⁴⁰ Li <i>et al.</i> , ⁴¹
System xc ⁻	Glutamatergic	Control of oxidative stress underlying inflammation	Liu <i>et al.</i> , ⁴² Patel <i>et al.</i> , ⁴³

A novel approach proposed the use of positive allosteric modulators of $\alpha 7$ nAChRs that converts endogenous agonists of $\alpha 7$ nAChRs such as ACh into potent neuroprotective agents in postischemic neuronal injury in cortical and subcortical brain regions.²⁰ In another study, the use of donepezil, an acetylcholinesterase inhibitor used for the treatment of Alzheimer's disease, displayed upregulation of nAChRs that attenuated the cerebral brain infarction volume after cerebral ischemia in rats and mice.^{21,61} Moreover, the use of other cholinesterase inhibitors such as huperzine A and galantamine has shown their potential anti-inflammatory and neuroprotective effects after cerebral ischemia in rodents.^{62–64} The pharmacological use of nicotine after a rat model of global ischemia increased the neuronal survival of CA1 pyramidal neurons accompanied by a reduction of microglial cells, TNF α and interleukin

(IL)-1 β in the region of the infarction. In addition, pretreatment with α -bungarotoxin, a selective $\alpha 7$ nAChR antagonist, could prevent the inhibitory effects of nicotine on cultured microglial proliferation, suggesting the role of nicotine in microglial activation through the activation of nicotinic receptors.¹⁸ Therefore, these results suggest that cholinergic agonists may be of clinical relevance for the treatment of stroke.

The use of selective agonists of $\alpha 7$ nAChR such as PHA 568487 has also shown very promising results, reducing the ischemic brain injury and inflammatory response after experimental stroke in rodents.^{15,17,22} Following permanent cerebral ischemia in mice, treatment with PHA 568487 showed a reduction of functional deficits at the acute stage of cerebral ischemia.¹⁵ Furthermore, PHA treatment reduced lesion volume, decreased




Figure 1. Temporal evolution of system xc⁻, α4β2 and α7 receptor expression after cerebral ischemia in rats. nAChR, nicotinic acetylcholine receptor.

the number of CD68+ and M1 macrophages, and increased the number of M2 or anti-inflammatory microglia or macrophages at days 3 and 14 after permanent middle cerebral artery occlusion (MCAO) in mice.¹⁵ This study suggested that α7 receptors might decrease the inflammatory response through control over microglia or macrophage polarization after cerebral ischemia. Nevertheless, a recent study showed that the daily treatment of PHA 568487 during the first week after transient cerebral ischemia in rats displayed a nonsignificant decrease of the expression marker values for both proinflammatory and anti-inflammatory microglia markers.²² Moreover, this study observed fewer values of selectins, adhesion molecules and infiltrated T lymphocytes after treatment with PHA, suggesting a possible role of α7 nAChRs in the regulation of leukocyte infiltration into the ischemic tissue.²² Likewise, the infiltration of leukocytes after stroke can also be influenced by the disruption of the BBB; however, activation of α7 receptors showed similar levels of BBB disruption after MCAO in rats.²² In contrast, Zou and colleagues examined the effect of α7 nAChRs activation with PHA after cerebral ischemia in mice showing an improvement of the BBB integrity.¹⁶

In spite of all these discrepancies, α7 nAChRs play a promising key role in the inflammatory reaction following cerebral ischemia in rodents. For this reason, *in vivo* imaging of these receptors with a positron emission tomography (PET) technique might be of great importance to further our understanding of the role of α7 receptors in brain diseases such as stroke. During the last few years, promising radiotracers for imaging these receptors have been synthesized;^{65–69}

however, only a PET imaging study has been carried out to evaluate the role of α7 nAChRs in neuroinflammation. This study used the selective orthosteric α7 nAChR agonist PET radioligand, [¹¹C]NS14492 to monitor the expression of α7 receptors during the following month after cerebral ischemia in rats. PET imaging with [¹¹C]NS14492 described an overexpression of these receptors as a response to the ischemic injury that was identified in microglia and infiltrated macrophages at day 7 after ischemia (Figures 1 and 2). Finally, PET imaging of neuroinflammation with [¹⁸F]DPA-714, a specific radioligand for the translocator protein (18KDa) (TSPO),⁷⁰ showed a reduction in the radioligand uptake as a result of treatment with PHA 568487 on day 7 after cerebral ischemia, supported by a reduction in activated microglia and macrophages²² (Figure 2). Hence, the anti-inflammatory activity exerted by the cholinergic system following stroke has been attributed to α7 nAChRs, although α4β2 nicotinic receptors have also been involved in the inflammatory reaction underlying cerebral ischemia in rats.²³ The expression of α4β2 nicotinic receptors is increased after microglia or macrophage and astrocyte activation during cerebral ischemia evolution.²³ *In vivo* PET imaging with 2[¹⁸F]-fluoro-A85380, a selective radiotracer for α4β2 nAChRs, showed a radioligand uptake increase at day 7 after ischemia followed by a progressive decrease later on (Figure 1). The α4β2 expression pattern is in accordance with the uptake increase of [¹¹C]PK11195, a PET radiotracer for imaging inflammation.⁷¹ Furthermore, treatment with the α4β2 antagonist dihydro-β-erythroidine hydrobromide (DhβE) caused an increase in [¹¹C]PK11195




Figure 2. Magnetic resonance imaging (MRI) and positron emission tomography (PET) with [^{11}C]NS14492 and [^{18}F]DPA-714, selective radiotracers for $\alpha 7$ nicotinic acetylcholine receptors (nAChRs) and inflammation respectively. (A) [^{11}C]NS14492-PET and MRI coregistered images at control (day 0), day 3 and day 7 after middle cerebral artery occlusion (MCAO). (B) MRI-T₂W and PET images of [^{18}F]DPA-714 before (day 1) and at day 7 after MCAO in controls and in rats treated with the $\alpha 7$ antagonist PHA.

binding after cerebral ischemia in rats, evidencing the potential role of $\alpha 4\beta 2$ nAChRs in the regulation of the neuroinflammatory response after stroke²³ (Figure 1).

Purinergic receptors

In the nervous system, ATP and its derivatives act as extracellular signaling molecules through a large variety of receptors known as purinergic receptors.¹² The effects of purines and pyrimidines are mediated through an extended family of purinergic receptors, which are classified as metabotropic P1 adenosine receptors (A_1 , A_{2A} , A_{2B} and A_3), metabotropic P2Y and ionotropic P2X purinoreceptors.⁷² Several investigations have reported the role of purinergic signaling following neurodegenerative diseases, epilepsy, neuropsychiatric disorders and stroke.⁷³ Likewise, several studies have proposed ATP, adenosine and purinergic receptors as promising biomarkers for stroke therapy.⁷⁴ Cerebral ischemia leads to

the high increase of concentrations of both ATP and adenosine that can stimulate the purinergic receptors.⁷⁵ Indeed, the effect of ATP and adenosine after cerebral ischemia may be due to the interaction of purinergic receptors expressed on neurons, glial cells and peripheral inflammatory cells such as lymphocytes and neutrophils.^{24–27,76}

P1 receptors

Infusion of adenosine to the rat brain is protective against cerebral ischemia as it reduces the infarct volume and improves the neurological outcome⁷⁷ due to mainly the response of the adenosine A_1 receptors.⁷⁴ Moreover, these receptors modulate the expression of IL-10 release by immune cells after neonatal hypoxic ischemic brain injury.⁷⁸ Despite these findings, the role of A_1 receptors on inflammatory reaction after stroke has been scarcely explored to date. However, the A_{2A} adenosine receptors are expressed in innate (microglia, macrophages, mast cells and neutrophils) but also

in adaptive immunity (lymphocytes) cells, supporting its control of the neuroinflammatory response.²⁴ Following cerebral ischemia, activation of A_{2A} receptors reduced processes related to the infiltration of peripheral inflammatory cells such as chemotaxis, rolling, adhesion and transmigration.⁷⁹ Accordingly, low doses of the selective A_{2A} agonist CGS21680 decreased the number of infiltrated granulocytes, microgliosis, astrogliosis and improved myelin organization in the injured lesion after cerebral ischemia in rats.⁸⁰ In agreement with this effect, treatment with CGS21680 attenuated both the infiltration of neutrophils and TNF α production after intracerebral hemorrhage.²⁸ Activation of these receptors with BAY 60-6583 inhibits tissue plasminogen activator (tPA)-induced hemorrhagic transformation, thus reducing brain swelling and lesion volume after experimental stroke.²⁹ In addition, treatment with BAY 60-6583 induced a decrease in metalloproteinase-9 activation, which is mainly involved in the permeabilization of the BBB.²⁹ Therefore, restoration of the BBB by A_{2A} receptors can likely reduce the infiltration of leukocytes into the infarcted tissue after stroke. Conversely, selective inactivation of A_{2A} receptors in bone marrow-derived cells (BMDCs) protected against brain injury by reducing the production of proinflammatory cytokines such as IL-1 β , IL-6 and IL-12 in the brain.³⁰ Hence, all these findings support the bidirectional modulation of inflammation by A_{2A} receptors after cerebral ischemia.³¹ Finally, activation of adenosine A₃ receptors displayed an anti-inflammatory effect and a depletion of the infiltration or migration of macrophages and microglia after cerebral ischemia in rats.⁸¹

P2 receptors

P2 receptors are classified in P2X ionotropic and P2Y metabotropic receptors. P2X receptors are a family of seven receptors (P2X₁₋₇) permeable to cations (Na⁺, K⁺ and Ca²⁺) and expressed on the surface of cells. P2Y receptors are coupled to G protein and a total of nine subtypes P2Y_{1,2,4,6,11,12,13,14} have been cloned to date in mammalian species.⁷² An increase in the extracellular concentrations of ATP after brain injury results in direct damage to neurons and oligodendrocytes through Ca²⁺ intracellular loading, a result of P2X7 activation.^{82,83} Thus, P2X7 antagonists reduce neuronal damage and infarct size after transient focal ischemia^{32,83,84} and ameliorate oligodendroglial and axonal damage after

white matter ischemia.⁸² Furthermore, ATP might also be crucial for the regulation of microglia after pathological situations.³³ In the early stages of ischemia, low ATP levels can induce the recognition and migration of microglia to the lesion.³⁴ In fact, activation of microglia from their resting state is marked by overexpression of P2X7 receptors in the peri-infarct region, which promotes the release of neurotrophic factors and the enhancement of neuronal survival.^{35,85} However, during the later stages of cerebral ischemia the ATP levels increase, promoting the proliferation of microglia and cell death.³⁶ Under these conditions, microglia upregulate the expression of proinflammatory cytokines such as IL-1 β , IL-6 and TNF α , enhancing the inflammatory response after cerebral ischemia.³⁷ In addition, *de novo* expression of P2X7 receptors observed in both activated and reactive microglia suggests a differential role of these receptors in core and neighboring regions of the brain infarction.³⁸ The P2 unselective antagonist Reactive Blue reduced ischemic brain damage by blocking activated microglia in the core of the lesion. Conversely, the same treatment increased the expression of P2X7 receptors in remote areas, promoting the restoration and defense of the tissue.³⁸ Therefore, these temporal and spatial contrary processes should be taken into account in future therapeutic approaches targeting these receptors for treating stroke.

Microglial cells have also been characterized by the overexpression of P2X4 receptors that contribute to the control of the fate and the survival of the microglia.³⁹ Following brain ischemia, P2X4 receptors are upregulated in microglia or infiltrated macrophages and P2 antagonists can decrease ischemic cell death, reducing P2X4 receptor expression.⁴⁰ During hypoxic situations, expression of these receptors mediates activation of the amoeboid microglial cells and the release of proinflammatory cytokines induced by the increase in ATP levels.⁴¹ Therefore, P2X4 receptor expression on microglial cells is associated with the progression of neuroinflammation following cerebral ischemia, despite the underlying mechanisms not being clearly deciphered to date.⁸⁶

P2Y₁₂ receptors have also been shown to be expressed on microglial cells, suggesting their role in inflammation.^{87,88} The oral administration of the P2Y₁₂ antagonist ticagrelor promotes protection against stroke damage through the inhibition

of microglia activation, infiltration of blood-derived cells and the expression of proinflammatory mediators such as IL-1, monocyte chemoattractant protein 1 (MCP-1) and nitric oxide synthase (iNOS). This study also showed that ticagrelor inhibited adenosine diphosphate (ADP)-induced chemotaxis in primary cultured microglia.⁸⁸ Furthermore, ticagrelor exerts antithrombotic actions and the P2Y₁₂ receptor is also expressed on circulating platelets, mediating its aggregation and activation.⁸⁹ Moreover, the use of clopidogrel, a P2Y₁₂ receptor inhibitor with antiplatelet action, has been used in therapy for the secondary prevention of ischemic stroke.⁴² These findings suggest that P2Y₁₂ antagonist can conduct anti-inflammatory, neuroprotective and antiplatelet activity after cerebral ischemia.⁷⁴

Glutamate transporters

Although glutamate is the main excitatory neurotransmitter in the central nervous system (CNS), activation of glutamate receptors induces neuronal death after stroke.⁹⁰ Under physiological situations glutamate is stored intracellularly, however under pathological conditions such as cerebral ischemia the levels of extracellular glutamate can dramatically be increased, promoting the excitotoxicity mechanism through the influx of calcium into the neurons.⁹¹ The clearance of extracellular glutamate levels through a family of transporter proteins called excitatory amino acid transporters (EAATs) has been mainly attributed to the glutamate-scavenging action of astrocytes.⁹² In the normal CNS, astrocytes express EAAT1, EAAT2 and glutamine synthetase (GS) that first transport glutamate into intracellular milieu, and second convert the glutamate into glutamine.⁹³

Cystine glutamate antiporter

Another transporter involved in glutamate homeostasis is the cystine glutamate antiporter, also known as system xc⁻.⁹⁴ It is a heterodimer composed of a heavy chain subunit (4F2hc) and a light chain subunit (xCT).⁹⁵ System xc⁻ mediates the cellular import of cystine and the release of glutamate to the extracellular space in exchange. Cystine is intracellularly converted to cysteine, the rate-limiting substrate for glutathione aporoduction and oxidative protection. Thus, the expression and function of system xc⁻ are modulated in different pathologies secondary to the induction of oxidative stress.^{96–98}

After cerebral ischemia, overexpression of the cystine glutamate antiporter contributes to the increase in the extracellular glutamate concentration that promotes ischemic neuronal death through activation of extrasynaptic N-methyl-D-aspartate (NMDA) receptors⁹⁵ (Figure 3). This study shows that pharmacological inhibition of system xc⁻ displays reduced neuronal death after *in vitro* ischemia.⁹⁵ Furthermore, *in vivo* PET imaging of system xc⁻ activity with the radiotracer [¹⁸F]FSPG showed an increase in the PET signal uptake from 5 min to 5 h after ischemia, showing the increase in the function of these transporters during the subacute stage of stroke.⁹⁵ [¹⁸F]FSPG is a fluorine-18-labeled L-glutamate derivative taken by the system xc⁻ due to the lack of discrimination between its natural substrate cystine and glutamate for the inward transport.^{43,99} Recently, a PET study with this radiotracer has demonstrated the overexpression of system xc⁻ on microglial cells following experimental autoimmune encephalomyelitis (EAE) in rats.¹⁰⁰ Additionally, the depletion of microglia with clodronate showed a reduction in the [¹⁸F]FSPG PET signal in the spinal cord, confirming the link between microglial activation and cysteine or glutamate antiporter activity in EAE rats.¹⁰⁰ Another clinical study used PET with [¹⁸F]FSPG to detect inflammatory lesions provoked by the activation of macrophages in diseases such as sarcoidosis.¹⁰¹ Several studies support these findings showing that the expression of system xc⁻ is particularly abundant in microglia, macrophages and other immune cells.^{102,103} In fact, the release of glutamate from both resting and activated microglia by system xc⁻ can induce a regional increase in glutamate levels that leads to excitotoxic oligodendrocyte death contributing to the pathogenesis of white matter disorders.¹⁰⁴

Upregulation of system xc⁻ occurs during the following week after ischemia reperfusion¹⁰⁴ (Figures 1 and 4). This stands in accordance with the expression of TSPO, a marker of microglial activation, using the radiotracer [¹⁸F]DPA-714.¹⁰⁵ Indeed, the expression of the cystine/glutamate antiporter takes place in activated microglia and infiltrated macrophages during the first week, and marginally in astrocytes at 1 month after ischemia onset. Additionally, the inhibition of system xc⁻ with sulfasalazine and S-4-CPG resulted in a decrease in inflammatory response through the inactivation of both microglia and infiltrated macrophages, a decrease in proinflammatory markers (CCL2,




Figure 3. The release of glutamate during brain ischemia triggers neuronal death by overactivation of NMDA receptors. Different mechanisms contribute to glutamate homeostasis alterations. Astrocytic glutamate transporters play a key role in maintaining synaptic glutamate levels. However, extrasynaptic glutamate is mainly regulated by the cystine glutamate antiporter, also known as system x_c^- .⁹⁴ Because N-methyl-D-aspartate (NMDA) receptors involved in neuronal death are typically extrasynaptic, it has been proposed that glutamate release by the cystine or glutamate antiporter activates extrasynaptic N-methyl-D-aspartate receptors (NMDARs).⁹⁵ Cells involved in glutamate release by cystine glutamate antiporter during ischemic insults remain to be determined. EAAT, excitatory amino acid transporter.

TNF and iNOS) and an increase in the anti-inflammatory marker arginase after experimental stroke in rats.¹⁰⁵ These results showed that the blocking of the source of glutamate release on microglia and macrophages during cerebral ischemia evolution may be a relevant therapeutic intervention to halt progression of the brain damage after ischemia. Furthermore, these results fully support that system x_c^- might play a key role in the modulation of inflammatory response following stroke.

Conclusion

Inflammation plays an important role at different stages of the cerebral postischemic injury. Consequently, the use of anti-inflammatory strategies in stroke therapy might offer a wider therapeutic window than current treatments. The inflammatory reaction following stroke involves a large variety of signaling pathways and mediators that can determine stroke outcome. During the last decade, both *in vivo* and *ex vivo* studies have described the role of neurotransmitter receptors and transporters, including cholinergic, purinergic and glutamatergic on the modulation of the inflammatory reaction after

cerebral ischemia. It became evident that the expression of these receptors and transporters in neurons, glial cells and infiltrated immune cells in the ischemic brain promotes the release of a battery of signal molecules that enhance the inflammatory response after cerebral ischemia. Moreover, the activation or inhibition of these neuroreceptors and transporters has shown promising therapeutic responses in animal models of stroke. Recently, molecular imaging studies using PET of these neuroreceptors have described their link with the inflammatory reaction after ischemia and their potential role as novel imaging biomarkers of neuroinflammation. Therefore, the data reviewed here suggest that cholinergic, purinergic and glutamatergic agents may be useful both as biomarkers for neuroinflammation and as a treatment to attenuate the deleterious consequences of the inflammatory response after stroke. In our opinion, among the different candidates proposed here, the activation of nicotinic receptors might become a promising strategy for treating stroke in the near future. Despite this, future clinical studies are needed to support all these findings and their true potential to ameliorate the neurological care of stroke.




Figure 4. Magnetic resonance imaging (MRI) [T₂W] and positron emission tomography (PET) with $[^{18}\text{F}]$ FSPG and $[^{18}\text{F}]$ DPA-714, markers of cystine glutamate antiporter activity and inflammation respectively. (A) $[^{18}\text{F}]$ FSPG-PET and MRI coregistered images at control (day 0), day 3 and day 7 after middle cerebral artery occlusion (MCAO). (B) PET images of $[^{18}\text{F}]$ DPA-714 at day 7 after cerebral ischemia in vehicle (MCAO), SAS (MCAO+SAS) and S-4-CPG (MCAO+ S-4-CPG) treated rats. SAS and S-4-CPG are inhibitors of cystine glutamate antiporter.

Funding

We acknowledge financial support by MINECO SAF2014-54070-JIN (A.M.) and SAF2016-75292-R (C.M.).

Conflict of interest statement

The authors declare that there is no conflict of interest.

References

1. Dirnagl U, Iadecola C and Moskowitz MA. Pathobiology of ischaemic stroke: an integrated view. *Trends Neurosci* 1999; 22: 391–397.
2. Martin RL, Lloyd HG and Cowan AI. The early events of oxygen and glucose deprivation: setting the scene for neuronal death? *Trends Neurosci* 1994; 17: 251–257.
3. Katsura K, Kristian T and Siesjö BK. Energy metabolism, ion homeostasis, and cell damage in the brain. *Biochem Soc Trans* 1994; 22: 991–996.
4. Somjen GG. Mechanisms of spreading depression and hypoxic spreading depression-like depolarization. *Physiol Rev* 2001; 81: 1065–1096.
5. Zhao SC, Ma LS, Chu ZH, *et al.* Regulation of microglial activation in stroke. *Acta Pharmacol Sin* 2017; 38: 445–458.
6. Bonaventura A, Liberale L, Vecchie A, *et al.* Update on inflammatory biomarkers and treatments in ischemic stroke. *Int J Mol Sci* 2016; 17: pii: E1967.

7. Kawabori M and Yenari MA. Inflammatory responses in brain ischemia. *Curr Med Chem* 2015; 22: 1258–1277.
8. Kreutzberg GW. Microglia: a sensor for pathological events in the CNS. *Trends Neurosci* 1996; 19: 312–318.
9. Becker KJ. Inflammation and acute stroke. *Curr Opin Neurol* 1998; 11: 45–49.
10. Hallenbeck JM. Significance of the inflammatory response in brain ischemia. *Acta Neurochir Suppl* 1996; 66: 27–31.
11. Hu X, Leak RK, Shi Y, *et al.* Microglial and macrophage polarization-new prospects for brain repair. *Nat Rev Neurol* 2015; 11: 56–64.
12. Kettenmann H, Hanisch UK, Noda M, *et al.* Physiology of microglia. *Physiol Rev* 2011; 91: 461–553.
13. Duris K, Manaenko A, Suzuki H, *et al.* Alpha7 nicotinic acetylcholine receptor agonist PNU-282987 attenuates early brain injury in a perforation model of subarachnoid hemorrhage in rats. *Stroke* 2011; 42: 3530–3536.
14. Krafft PR, McBride D, Rolland WB, *et al.* Alpha7 nicotinic acetylcholine receptor stimulation attenuates neuroinflammation through JAK2-STAT3 activation in murine models of intracerebral hemorrhage. *Biomed Res Int* 2017; 2017: 8134653.
15. Han Z, Shen F, He Y, *et al.* Activation of alpha-7 nicotinic acetylcholine receptor reduces ischemic stroke injury through reduction of pro-inflammatory macrophages and oxidative stress. *PLoS One*. 2014; 9: e105711.
16. Zou D, Luo M, Han Z, *et al.* Activation of alpha-7 nicotinic acetylcholine receptor reduces brain edema in mice with ischemic stroke and bone fracture. *Mol Neurobiol* 2017; 54: 8278–8286.
17. Han Z, Li L, Wang L, *et al.* Alpha-7 nicotinic acetylcholine receptor agonist treatment reduces neuroinflammation, oxidative stress, and brain injury in mice with ischemic stroke and bone fracture. *J Neurochem* 2014; 131: 498–508.
18. Guan YZ, Jin XD, Guan LX, *et al.* Nicotine inhibits microglial proliferation and is neuroprotective in global ischemia rats. *Mol Neurobiol* 2015; 51: 1480–1488.
19. Shimohama S, Greenwald DL, Shafroon DH, *et al.* Nicotinic alpha 7 receptors protect against glutamate neurotoxicity and neuronal ischemic damage. *Brain Res* 1998; 779: 359–363.
20. Kalappa BI, Sun F, Johnson SR, *et al.* A positive allosteric modulator of alpha7 nAChRs augments neuroprotective effects of endogenous nicotinic agonists in cerebral ischaemia. *Br J Pharmacol* 2013; 169: 1862–1878.
21. Fujiki M, Kobayashi H, Uchida S, *et al.* Neuroprotective effect of donepezil, a nicotinic acetylcholine-receptor activator, on cerebral infarction in rats. *Brain Res* 2005; 10: 1–2.
22. Colás LDM, Ramos P, Palma A, *et al.* In vivo imaging of $\alpha 7$ nicotinic receptors as a novel method to monitor neuroinflammation after cerebral ischemia. *Glia*. Epub ahead of print 12 March 2018. DOI: 10.1002/glia.23326.
23. Martin A, Szczupak B, Gomez-Vallejo V, *et al.* In vivo PET imaging of the alpha4beta2 nicotinic acetylcholine receptor as a marker for brain inflammation after cerebral ischemia. *J Neurosci* 2015; 35: 5998–6009.
24. Burnstock G and Boeynaems JM. Purinergic signalling and immune cells. *Purinergic Signal* 2014; 10: 529–564.
25. Verkhratsky A, Krishtal OA and Burnstock G. Purinoceptors on neuroglia. *Mol Neurobiol* 2009; 39: 190–208.
26. Puchalowicz K, Baranowska-Bosiacka I, Dziedzic V, *et al.* Purinergic signaling and the functioning of the nervous system cells. *Cell Mol Biol Lett* 2015; 20: 867–918.
27. Vuorimaa A, Rissanen E and Airas L. In vivo PET imaging of adenosine 2A receptors in neuroinflammatory and neurodegenerative disease. *Contrast Media Mol Imaging* 2017; 2017: 6975841.
28. Mayne M, Fotheringham J, Yan HJ, *et al.* Adenosine A2A receptor activation reduces proinflammatory events and decreases cell death following intracerebral hemorrhage. *Ann Neurol* 2001; 49: 727–735.
29. Li Q, Han X, Lan X, *et al.* Inhibition of tPA-induced hemorrhagic transformation involves adenosine A2b receptor activation after cerebral ischemia. *Neurobiol Dis* 2017; 108: 173–182.
30. Yu L, Huang Z, Mariani J, *et al.* Selective inactivation or reconstitution of adenosine A2A receptors in bone marrow cells reveals their significant contribution to the development of ischemic brain injury. *Nat Med* 2004; 10: 1081–1087.
31. Dai SS and Zhou YG. Adenosine 2A receptor: a crucial neuromodulator with bidirectional effect in neuroinflammation and brain injury. *Rev Neurosci* 2011; 22: 231–239.

32. Troadec JD, Thirion S, Petturiti D, *et al.* ATP acting on P2Y receptors triggers calcium mobilization in primary cultures of rat neurohypophysial astrocytes (pituicytes). *Pflugers Arch* 1999; 437: 745–753.
33. Collo G, Neidhart S, Kawashima E, *et al.* Tissue distribution of the P2X7 receptor. *Neuropharmacology* 1997; 36: 1277–1283.
34. Skaper SD, Facci L, Culbert AA, *et al.* P2X(7) receptors on microglial cells mediate injury to cortical neurons in vitro. *Glia* 2006; 54: 234–242.
35. Suzuki T, Hide I, Ido K, *et al.* Production and release of neuroprotective tumor necrosis factor by P2X7 receptor-activated microglia. *J Neurosci* 2004; 24: 1–7.
36. Chu K, Yin B, Wang J, *et al.* Inhibition of P2X7 receptor ameliorates transient global cerebral ischemia/reperfusion injury via modulating inflammatory responses in the rat hippocampus. *J Neuroinflammation* 2012; 9: 69.
37. Monif M, Burnstock G and Williams DA. Microglia: proliferation and activation driven by the P2X7 receptor. *Int J Biochem Cell Biol* 2010; 42: 1753–1756.
38. Melani A, Amadio S, Gianfriddo M, *et al.* P2X7 receptor modulation on microglial cells and reduction of brain infarct caused by middle cerebral artery occlusion in rat. *J Cereb Blood Flow Metab* 2006; 26: 974–982.
39. Vazquez-Villoldo N, Domercq M, Martin A, *et al.* P2X4 receptors control the fate and survival of activated microglia. *Glia* 2014; 62: 171–184.
40. Cavaliere F, Florenzano F, Amadio S, *et al.* Up-regulation of P2X2, P2X4 receptor and ischemic cell death: prevention by P2 antagonists. *Neuroscience* 2003; 120: 85–98.
41. Li F, Wang L, Li JW, *et al.* Hypoxia induced amoeboid microglial cell activation in postnatal rat brain is mediated by ATP receptor P2X4. *BMC Neurosci* 2011; 12: 111.
42. Liu F, Tantry US and Gurbel PA. P2Y12 receptor inhibitors for secondary prevention of ischemic stroke. *Expert Opin Pharmacother* 2015; 16: 1149–1165.
43. Patel SA, Warren BA, Rhoderick JF, *et al.* Differentiation of substrate and non-substrate inhibitors of transport system xc(-): an obligate exchanger of L-glutamate and L-cystine. *Neuropharmacology* 2004; 46: 273–284.
44. Pavlov VA, Wang H, Czura CJ, *et al.* The cholinergic anti-inflammatory pathway: a missing link in neuroimmunomodulation. *Mol Med* 2003; 9: 125–134.
45. Tracey KJ. Physiology and immunology of the cholinergic antiinflammatory pathway. *J Clin Invest* 2007; 117: 289–296.
46. Tracey KJ. The inflammatory reflex. *Nature* 2002; 420: 853–859.
47. Borovikova LV, Ivanova S, Zhang M, *et al.* Vagus nerve stimulation attenuates the systemic inflammatory response to endotoxin. *Nature* 2000; 405: 458–462.
48. Wang H, Yu M, Ochani M, *et al.* Nicotinic acetylcholine receptor alpha7 subunit is an essential regulator of inflammation. *Nature* 2003; 421: 384–388.
49. Ueno K, Togashi H, Matsumoto M, *et al.* Alpha4beta2 nicotinic acetylcholine receptor activation ameliorates impairment of spontaneous alternation behavior in stroke-prone spontaneously hypertensive rats, an animal model of attention deficit hyperactivity disorder. *J Pharmacol Exp Ther* 2002; 302: 95–100.
50. Hogg RC, Raggenbass M and Bertrand D. Nicotinic acetylcholine receptors: from structure to brain function. *Rev Physiol Biochem Pharmacol* 2003; 147: 1–46.
51. Murakami K, Ishikawa Y and Sato F. Localization of alpha7 nicotinic acetylcholine receptor immunoreactivity on GABAergic interneurons in layers I–III of the rat retrosplenial granular cortex. *Neuroscience* 2013; 252: 443–459.
52. Rogers SW, Gregori NZ, Carlson N, *et al.* Neuronal nicotinic acetylcholine receptor expression by O2A/oligodendrocyte progenitor cells. *Glia* 2001; 33: 306–313.
53. Sharma G and Vijayaraghavan S. Nicotinic cholinergic signaling in hippocampal astrocytes involves calcium-induced calcium release from intracellular stores. *Proc Natl Acad Sci U S A* 2001; 98: 4148–4153.
54. De Simone R, Ajmone-Cat MA, Carnevale D, *et al.* Activation of alpha7 nicotinic acetylcholine receptor by nicotine selectively up-regulates cyclooxygenase-2 and prostaglandin E2 in rat microglial cultures. *J Neuroinflammation* 2005; 2: 4.
55. Hawkins BT, Eggleton RD and Davis TP. Modulation of cerebral microvascular

- permeability by endothelial nicotinic acetylcholine receptors. *Am J Physiol Heart Circ Physiol* 2005; 289: H212–H219.
56. Sharma G and Vijayaraghavan S. Nicotinic receptor signaling in nonexcitable cells. *J Neurobiol* 2002; 53: 524–534.
57. Neumann S, Shields NJ, Balle T, *et al.* Innate immunity and inflammation post-stroke: an alpha7-nicotinic agonist perspective. *Int J Mol Sci* 2015; 16: 29029–29046.
58. de Jonge WJ and Ulloa L. The alpha7 nicotinic acetylcholine receptor as a pharmacological target for inflammation. *Br J Pharmacol* 2007; 151: 915–929.
59. Jiang Y, Li L, Liu B, *et al.* Vagus nerve stimulation attenuates cerebral ischemia and reperfusion injury via endogenous cholinergic pathway in rat. *PLoS One* 2014; 9: e102342.
60. Lu XX, Hong ZQ, Tan Z, *et al.* Nicotinic acetylcholine receptor alpha7 subunit mediates vagus nerve stimulation-induced neuroprotection in acute permanent cerebral ischemia by a7nAChR/JAK2 pathway. *Med Sci Monit* 2017; 23: 6072–6081.
61. Wang T, Lv P, Jin W, *et al.* Protective effect of donepezil hydrochloride on cerebral ischemia/reperfusion injury in mice. *Mol Med Rep* 2014; 9: 509–514.
62. Wang ZF, Wang J, Zhang HY, *et al.* Huperzine A exhibits anti-inflammatory and neuroprotective effects in a rat model of transient focal cerebral ischemia. *J Neurochem* 2008; 106: 1594–1603.
63. Odorcyk FK, Nicola F, Duran-Carabali LE, *et al.* Galantamine administration reduces reactive astrogliosis and upregulates the anti-oxidant enzyme catalase in rats submitted to neonatal hypoxia ischemia. *Int J Dev Neurosci* 2017; 62: 15–24.
64. Odorcyk FK, Sanches EF, Nicola FC, *et al.* Administration of huperzia quadrifariata extract, a cholinesterase inhibitory alkaloid mixture, has neuroprotective effects in a rat model of cerebral hypoxia-ischemia. *Neurochem Res* 2017; 42: 552–562.
65. Hillmer AT, Li S, Zheng MQ, *et al.* PET imaging of alpha7 nicotinic acetylcholine receptors: a comparative study of [(18)F]ASEM and [(18)F]DBT-10 in nonhuman primates, and further evaluation of [(18)F]ASEM in humans. *Eur J Nucl Med Mol Imaging* 2017; 44: 1042–1050.
66. Horti AG. Development of [(18)F]ASEM, a specific radiotracer for quantification of the alpha7-nAChR with positron-emission tomography. *Biochem Pharmacol* 2015; 97: 566–575.
67. Hillmer AT, Zheng MQ, Li S, *et al.* PET imaging evaluation of [(18)F]DBT-10, a novel radioligand specific to alpha7 nicotinic acetylcholine receptors, in nonhuman primates. *Eur J Nucl Med Mol Imaging* 2016; 43: 537–547.
68. Rotering S, Deuther-Conrad W, Cumming P, *et al.* Imaging of alpha7 nicotinic acetylcholine receptors in brain and cerebral vasculature of juvenile pigs with [(18)F]NS14490. *EjNMMI Res* 2014; 4: 43.
69. Ettrup A, Mikkelsen JD, Lehel S, *et al.* 11C-NS14492 as a novel PET radioligand for imaging cerebral alpha7 nicotinic acetylcholine receptors: in vivo evaluation and drug occupancy measurements. *J Nucl Med* 2011; 52: 1449–1456.
70. Martin A, Boisgard R, Theze B, *et al.* Evaluation of the PBR/TSPO radioligand [(18)F]DPA-714 in a rat model of focal cerebral ischemia. *J Cereb Blood Flow Metab* 2010; 30: 230–241.
71. Rojas S, Martin A, Arranz MJ, *et al.* Imaging brain inflammation with [(11)C]PK11195 by PET and induction of the peripheral-type benzodiazepine receptor after transient focal ischemia in rats. *J Cereb Blood Flow Metab* 2007; 27: 1975–1986.
72. Burnstock G. Purine and pyrimidine receptors. *Cell Mol Life Sci* 2007; 64: 1471–1483.
73. Burnstock G. Purinergic signalling: therapeutic developments. *Front Pharmacol* 2017; 8: 661.
74. Pedata F, Dettori I, Coppi E, *et al.* Purinergic signalling in brain ischemia. *Neuropharmacology* 2016; 104: 105–130.
75. Melani A, Corti F, Stephan H, *et al.* Ecto-ATPase inhibition: ATP and adenosine release under physiological and ischemic in vivo conditions in the rat striatum. *Exp Neurol* 2012; 233: 193–204.
76. Brodie C, Blumberg PM and Jacobson KA. Activation of the A2A adenosine receptor inhibits nitric oxide production in glial cells. *FEBS Lett* 1998; 429: 139–142.
77. Kitagawa H, Mori A, Shimada J, *et al.* Intracerebral adenosine infusion improves neurological outcome after transient focal ischemia in rats. *Neurol Res* 2002; 24: 317–323.

78. Winerdal M, Winerdal ME, Wang YQ, *et al.* Adenosine A1 receptors contribute to immune regulation after neonatal hypoxic ischemic brain injury. *Purinergic Signal* 2016; 12: 89–101.
79. Hasko G, Linden J, Cronstein B, *et al.* Adenosine receptors: therapeutic aspects for inflammatory and immune diseases. *Nat Rev Drug Discov* 2008; 7: 759–770.
80. Melani A, Corti F, Cellai L, *et al.* Low doses of the selective adenosine A2A receptor agonist CGS21680 are protective in a rat model of transient cerebral ischemia. *Brain Res* 2014; 10: 59–72.
81. Choi IY, Lee JC, Ju C, *et al.* A3 adenosine receptor agonist reduces brain ischemic injury and inhibits inflammatory cell migration in rats. *Am J Pathol* 2011; 179: 2042–2052.
82. Domercq M, Perez-Samartin A, Aparicio D, *et al.* P2X7 receptors mediate ischemic damage to oligodendrocytes. *Glia* 2010; 58: 730–740.
83. Arbeloa J, Perez-Samartin A, Gottlieb M, *et al.* P2X7 receptor blockade prevents ATP excitotoxicity in neurons and reduces brain damage after ischemia. *Neurobiol Dis* 2012; 45: 954–961.
84. Amadio S, D'Ambrosi N, Cavaliere F, *et al.* P2 receptor modulation and cytotoxic function in cultured CNS neurons. *Neuropharmacology* 2002; 42: 489–501.
85. Kharlamov A, Jones SC and Kim DK. Suramin reduces infarct volume in a model of focal brain ischemia in rats. *Exp Brain Res* 2002; 147: 353–359.
86. Cheng RD, Ren JJ, Zhang YY, *et al.* P2X4 receptors expressed on microglial cells in post-ischemic inflammation of brain ischemic injury. *Neurochem Int* 2014; 67: 9–13.
87. Amadio S, Parisi C, Montilli C, *et al.* P2Y(12) receptor on the verge of a neuroinflammatory breakdown. *Mediators Inflamm* 2014; 2014: 975849.
88. Gelosa P, Lecca D, Fumagalli M, *et al.* Microglia is a key player in the reduction of stroke damage promoted by the new antithrombotic agent ticagrelor. *J Cereb Blood Flow Metab* 2014; 34: 979–988.
89. Kashiwazaki D, Kuwayama N, Akioka N, *et al.* The roles and issues of P2Y12 percent inhibition assessed by VerifyNow assay for patients undergoing neurointervention: a prospective study. *J Stroke Cerebrovasc Dis* 2014; 23: 1830–1836.
90. Chamorro A, Dirnagl U, Urra X, *et al.* Neuroprotection in acute stroke: targeting excitotoxicity, oxidative and nitrosative stress, and inflammation. *Lancet Neurol* 2016; 15: 869–881.
91. Krzyzanowska W, Pomierny B, Filip M, *et al.* Glutamate transporters in brain ischemia: to modulate or not? *Acta Pharmacol Sin* 2014; 35: 444–462.
92. Rothstein JD, Martin L, Levey AI, *et al.* Localization of neuronal and glial glutamate transporters. *Neuron* 1994; 13: 713–725.
93. Martinez-Hernandez A, Bell KP and Norenberg MD. Glutamine synthetase: glial localization in brain. *Science*. 1977; 195: 1356–1358.
94. Baker DA, Xi ZX, Shen H, *et al.* The origin and neuronal function of in vivo nonsynaptic glutamate. *J Neurosci* 2002; 22: 9134–9141.
95. Soria FN, Perez-Samartin A, Martin A, *et al.* Extrasynaptic glutamate release through cystine/glutamate antiporter contributes to ischemic damage. *J Clin Invest* 2014; 124: 3645–3655.
96. Bannai S and Kitamura E. Transport interaction of L-cystine and L-glutamate in human diploid fibroblasts in culture. *J Biol Chem* 1980; 255: 2372–2376.
97. Lo M, Wang YZ and Gout PW. The x(c)-cystine/glutamate antiporter: a potential target for therapy of cancer and other diseases. *J Cell Physiol* 2008; 215: 593–602.
98. Conrad M and Sato H. The oxidative stress-inducible cystine/glutamate antiporter, system x (c) (-): cystine supplier and beyond. *Amino Acids* 2012; 42: 231–246.
99. Koglin N, Mueller A, Berndt M, *et al.* Specific PET imaging of xC- transporter activity using a (1)(8)F-labeled glutamate derivative reveals a dominant pathway in tumor metabolism. *Clin Cancer Res* 2011; 17: 6000–6011.
100. Martin A, Vazquez-Villoldo N, Gomez-Vallejo V, *et al.* In vivo imaging of system xc- as a novel approach to monitor multiple sclerosis. *Eur J Nucl Med Mol Imaging* 2016; 43: 1124–1138.
101. Chae SY, Choi CM, Shim TS, *et al.* Exploratory clinical investigation of (4S)-4-(3-18F-fluoropropyl)-L-glutamate PET of inflammatory and infectious lesions. *J Nucl Med* 2016; 57: 67–69.

102. Sato H, Kuriyama-Matsumura K, Hashimoto T, *et al.* Effect of oxygen on induction of the cystine transporter by bacterial lipopolysaccharide in mouse peritoneal macrophages. *J Biol Chem* 2001; 276: 10407–10412.
103. Qin S, Colin C, Hinnens I, *et al.* System Xc- and apolipoprotein E expressed by microglia have opposite effects on the neurotoxicity of amyloid-beta peptide 1–40. *J Neurosci* 2006; 26: 3345–3356.
104. Domercq M, Sanchez-Gomez MV, Sherwin C, *et al.* System xc- and glutamate transporter inhibition mediates microglial toxicity to oligodendrocytes. *J Immunol* 2007; 178: 6549–6556.
105. Domercq M, Szczupak B, Gejo J, *et al.* PET imaging with [(18)F]FSPG evidences the role of system xc(-) on brain inflammation following cerebral ischemia in rats. *Theranostics* 2016; 6: 1753–1767.

Visit SAGE journals online
[journals.sagepub.com/
home/tan](http://journals.sagepub.com/home/tan)

