



AMERICAN JOURNAL OF PHARMTECH RESEARCH

Journal home page: <http://www.ajptr.com/>

Analytical method development and validation of Niraparib and Abiraterone using RP-HPLC method by using QbD method

Mounica B, Vinutha Kommineni, Sreedevi P

Sri Venkateswara college of Pharmacy, Madhapur, Hyderabad

ABSTRACT

A robust RP-HPLC technique was set up for the estimation of Abiraterone & Niraparib. SST parameters, including resolution, theoretical plates, & tailing factor, were within acceptable boundaries, ensuring the technique's reliability. technique verification followed ICH rules, demonstrating high assay accuracy with recoveries within 98-102%. The linearity scale was 10-50 µg per ml for Abiraterone & 2-10 µg per ml for Niraparib, with R^2 of 0.999. Precision, ID precision, & robustness tests confirmed the technique's consistency across different conditions. LOD & LOQ were established, showcasing the technique's sensitivity. This verified RP-HPLC technique is suitable for regular analysis of Abiraterone & Niraparib in 1 formulations.

Keywords: Niraparib, Abiraterone, QbD, RP HPLC.

*Corresponding Author Email: Vinutha08.ch@gmail.com

Received 10 November 2024, Accepted 15 December 2024

Please cite this article as: Vinutha K *et al.*, Analytical method development and validation of Niraparib and Abiraterone using RP-HPLC method by using QbD method. American Journal of PharmTech Research 2024.

INTRODUCTION

Niraparib is a small molecule poly ADP ribose polymerase 1 and 2 inhibitors prescribed for treating ovarian cancer in adult patients. It is also used to treat primary peritoneal cancer and fallopian tube cancer for patients who are showing partial or complete response to platinum-based chemotherapy. Niraparib works by irreversible multiple breaks in the double strand with Breast cancer susceptibility protein (BRCA) type 1 and 2 mutations leading to the death of the cells.^(1,2)

According to Global Cancer Statistics 2020, prostate cancer is the second most commonly occurring cancer in men and the fifth most common cause of cancer-related death. Abiraterone (ART) acetate is used to treat metastatic castration-resistant prostate cancer and metastatic high-risk castration sensitive prostate cancer. ART acetate is recommended in combination with prednisone or prednisolone. In vivo, ART is an active metabolite of ART acetate. Both ART acetate and ART are potent, selective, and irreversible inhibitors of 17 α -hydroxylase/C17, 20-lyase (CYP17), a key enzyme used in testosterone biosynthesis. By blocking CYP17, ART leads to the termination of androgen biosynthesis in testicular, adrenal, and prostatic tumour tissues. ART chemically known as (3b)-17-(pyridin-3-yl) androsta-5,16-dien-3-ol.^(3;4)

Several analytical techniques, including high-performance liquid chromatography (HPLC)-fluorescence⁽¹¹⁾, HPTC⁽⁸⁾, ultraviolet (UV)^(5,6), UPLC⁽⁷⁾ and liquid chromatography-mass spectrometry (LC-MS/MS)⁽⁹⁾ have been reported to quantify Abiraterone and Niraparib. In the extensive literature review, it was confirmed that no analytical Liquid Chromatographic Method was reported for estimation of Abiraterone and Niraparib.

Hence the present work intended to establish an analytical Liquid Chromatographic Method for the estimation of Niraparib and Abiraterone.

MATERIALS AND METHOD

Chemicals and Reagents

Niraparib, Abiraterone (ART) acetate, Sodium hydroxide (NaOH), water, Phosphate buffer, Orthophosphoric acid (OPA) and Potassium dihydrogen phosphate (KH₂PO₄) acetonitrile (ANC) and trifluoroacetic acid chemicals are used in this research.

Instruments

HPLC System: Shimadzu (LC-2010C HT) with a ultraviolet (UV)-visible detector, ultra-sonicator and nylon filter (0.45 μ m) was used

Detector: photodiode array, Software Empower 3, 2695 separation modules 996 PDA

Methods

Preparation of mobile phase

Mix a 300ml TFA PH: 3 (30%) & 700 ml ACN (70%) & remove gases in ultra sonication water bath for 5 mins. Filter using 0.45micron vacuum filtration.

Standard Solution Preparation:

Accurately weigh and transfer 50 mg of Abiraterone and 10 mg of Niraparib working standards add into a separate 25 ml clean dry volumetric flasks add Diluents and sonicate to dissolve it completely and make volume up to the mark with the same solvent. (Stock solution)

Further pipette 0.3ml & 0.3 ml of the above stock solutions into a 20ml volumetric flask and dilute up to the mark with Diluents

Design Space

Method optimization was performed with Box-Behnken statistical design comprising the CMPs, which include five significant factors (the composition of the determined flow rate, organic phase ratio, column, detector wavelength and column temperature), encompassing five levels and include five significant responses (tailing factor of 1st and 2nd drug, retention time of 1st and 2nd drug and resolution. Seventeen experimental runs were established. Design-Expert software was used to validate the model after the data was examined. A positive correlation was found between the quadratic model and the experimental data used for design space navigation. It was verified that the predicted values from the practical responses were obtained within the design space of the optimized results, and they were found to be satisfactory.



Method development and optimization Using Box-Behnken experimental design

Table 1: Experimental runs and responds values

| Std | Run | Block | Factor 1 | Factor 2 | Factor 3 | Response 1 | Response 2 | Response 3 | Response 4 | Response 5 |
|-----|-----|---------|----------|----------|----------|------------|------------|------------|------------|------------|
| 15 | 1 | Block 1 | 0.90 | 50.00 | 35.00 | 1.06 | 1.01 | 2.122 | 2.76 | 5.09 |
| 2 | 2 | Block 1 | 1.00 | 40.00 | 35.00 | 1.07 | 1.02 | 2.118 | 2.72 | 5.05 |
| 8 | 3 | Block 1 | 1.00 | 50.00 | 45.00 | 1.08 | 1.03 | 2.119 | 2.73 | 5.06 |
| 1 | 4 | Block 1 | 0.80 | 40.00 | 35.00 | 1.02 | 1 | 2.06 | 2.4 | 5 |
| 12 | 5 | Block 1 | 0.90 | 60.00 | 45.00 | 1.15 | 1.19 | 2.13 | 2.64 | 4.9 |
| 4 | 6 | Block 1 | 1.00 | 60.00 | 35.00 | 1.09 | 1.04 | 2.12 | 2.74 | 5.07 |
| 7 | 7 | Block 1 | 0.80 | 50.00 | 45.00 | 1.04 | 1.01 | 2.08 | 2.6 | 5.01 |
| 5 | 8 | Block 1 | 0.80 | 50.00 | 25.00 | 1.04 | 1.01 | 2.08 | 2.6 | 5.01 |
| 6 | 9 | Block 1 | 1.00 | 50.00 | 25.00 | 1.08 | 1.03 | 2.119 | 2.73 | 5.06 |
| 13 | 10 | Block 1 | 0.90 | 50.00 | 35.00 | 1.06 | 1.01 | 2.122 | 2.76 | 5.09 |
| 9 | 11 | Block 1 | 0.90 | 40.00 | 25.00 | 1.16 | 1.21 | 2.132 | 2.65 | 5.01 |
| 14 | 12 | Block 1 | 0.90 | 50.00 | 35.00 | 1.06 | 1.01 | 2.122 | 2.76 | 5.09 |
| 3 | 13 | Block 1 | 0.80 | 60.00 | 35.00 | 1.05 | 1.02 | 2.09 | 2.7 | 5.02 |
| 17 | 14 | Block 1 | 0.90 | 50.00 | 35.00 | 1.06 | 1.01 | 2.122 | 2.76 | 5.09 |
| 16 | 15 | Block 1 | 0.90 | 50.00 | 35.00 | 1.06 | 1.01 | 2.122 | 2.76 | 5.09 |
| 10 | 16 | Block 1 | 0.90 | 60.00 | 25.00 | 1.09 | 1.04 | 2.12 | 2.74 | 5.07 |
| 11 | 17 | Block 1 | 0.90 | 40.00 | 45.00 | 1.16 | 1.21 | 2.132 | 2.65 | 5.01 |

Table 2: ANNOVA of quadratic model

| Ource | Sum of Squares | df | Mean Square | F Value | p-value |
|------------------|----------------|----|-------------|---------|---------|
| Model | 0.025 | 9 | 2.786E-003 | 7.57 | 0.0071 |
| Flow rate | 3.613E-003 | 1 | 3.613E-003 | 9.82 | 0.0165 |
| Organic mp ratio | 1.125E-004 | 1 | 1.125E-004 | 0.31 | 0.5975 |
| Column Temp | 4.500E-004 | 1 | 4.500E-004 | 1.22 | 0.3053 |

Figure 1: Tailing factor of Niraparib**Figure 2: Tailing factor of Abiraterone****Figure3: Retention of Niraparib**

Design-Expert® Software

Rt 1st drug
 ● Design Points
 2.132
 2.06

X1 = A: Flow rate
 X2 = B: Organic mp ratio

Actual Factor
 C: Column Temperature = 35.00




Figure 4: Retention of Abiraterone

Design-Expert® Software

Rt 2nd drug
 ● Design Points
 2.70
 2.4

X1 = A: Flow rate
 X2 = B: Organic mp ratio

Actual Factor
 C: Column Temperature = 35.00





Figure 5: Resolution of Niraparib and Abiraterone

Design-Expert® Software

Resolution
 ● Design Points
 5.09
 4.9

X1 = A: Flow rate
 X2 = B: Organic mp ratio

Actual Factor
 C: Column Temperature = 35.00



METHOD VALIDATION

LINEARITY

Six concentrations of the niraparib and abiraterone acetate working standard ranging from 10 to 50 ppm were prepared and analysed. The calibration plot was constructed by plotting the chromatographic peak areas versus concentrations and values of observed concentrations were determined. The correlation coefficient obtained was 0.999 which is in the acceptance limit. The linearity range was found to lie from 10µg/ml to 50µg/ml of Abiraterone and 2µg/ml to 10µg/ml Niraparib and chromatograms are shown below.




Figure 1: Calibration graphs of Niraparib and Abiraterone

ACCURACY

A series of solutions were prepared with known standard concentrations of niraparib and abiraterone acetate of 50%, 100%, and 150% (5, 10, and 15 µg/mL) in triplicate and the % recovery was calculated.

Table 3: Accuracy (recovery) data for Abiraterone

| %Concentration Abiraterone | Area* | Amount Added (mg) | Amount Found (mg) | % Recovery | Mean Recovery |
|-------------------------------|----------|-------------------|-------------------|---------------|---------------|
| 50% | 1629425 | 25 | 24.95 | 99.81 | 99.8 |
| 100% | 3258850 | 50 | 49.90 | 99.81 | |
| 150% | 48883005 | 75 | 74.77 | 99.70 | |

Table 4: Accuracy (recovery) data for Niraparib


| %Concentration Niraparib | Area* Amount Added (mg) | Amount Found (mg) | % Recovery | Mean Recovery |
|--------------------------|-------------------------|-------------------|------------|---------------|
| 50% | 3258855 | 4.95 | 100.09 | 100.03 |
| 100% | 65107010 | 9.84 | 99.98 | |
| 150% | 97725515 | 14.92 | 100.05 | |

PRECISION:

The precision of the developed technique, expressed in % relative standard deviation (RSD), was calculated by performing repeatability and intermediate precision studies. The developed analytical QbD -based method was validated by the precision studies.

Table 5: Results of Precision for Abiraterone and Niraparib

| Injection | Areas of Abiraterone | Areas of Niraparib |
|---------------------------|----------------------|--------------------|
| Injection-1 | 3258850 | 651770 |
| Injection-2 | 3248850 | 641770 |
| Injection-3 | 3158850 | 650770 |
| Injection-4 | 3248850 | 651770 |
| Injection-5 | 3238850 | 651770 |
| Injection-6 | 3258850 | 651770 |
| Average | 3235517 | 649936.7 |
| Standard Deviation | 38297.08 | 4020.779 |
| %RSD | 1.18 | 0.61 |

**Figure 2: Chromatogram for Precision****LIMIT OF LOD AND LOQ:**

The lowest concentration of the sample was prepared with respect to the base line noise and measured the signal to noise ratio. The estimated value for LOD was found to be 0.024 ppm, and the estimated value for LOQ was 0.08 ppm.




Figure 3: Chromatogram of niraparib and abiraterone

Table 6: Result of LOQ

| Drug name | Baseline noise(μ V) | Signal obtained(μ V) | S/N ratio | CONC. |
|-------------|--------------------------|---------------------------|-----------|-----------------|
| Abiraterone | 54 | 538 | 9.96 | 0.08 μ g/ml |




Figure 4: Chromatogram of niraparib and abiraterone

Table 7: Result of LOD

| Drug name | Baseline noise(μ V) | Signal obtained | S/N ratio | Conc. |
|-------------|--------------------------|-----------------|-----------|------------------|
| Abiraterone | 54 | 161 | 2.98 | 0.024 μ g/ml |
| Niraparib | 54 | 160 | 2.96 | 0.024 μ g/ml |

Assay of formulations

Twenty tablets of commercial brands of niraparib and abiraterone acetate were chosen and their average weight was determined. The tablets were crushed to fine powder, 250 mg was weighed, and the powder was dissolved in 200 mL of acetonitrile (in a 250 mL volumetric flask). Then, it was shaken for 20 min and ultrasonicated for 20 min. After that, it was allowed to cool at room

temperature, and the solution was diluted up to mark with diluents (1000 mcg mL^{-1}). The final obtained solution was then diluted to 10 mcg mL^{-1} with acetonitrile/phosphate buffer (20:80 %v/v) and subsequently injected to an HPLC system for estimation in triplicate.

Table 8: Assay of Niraparib and Abiraterone

| Drugs | Label Claim (mg) | % Assay |
|---------------------------|------------------|---------|
| Abiraterone and Niraparib | 500mg/100mg | 99.95 |

CONCLUSION

The AQbD approach proved effective for holistic process understanding and analytical method development. By identifying CMPs and optimizing key parameters, the developed method achieved reliable results for niraparib and abiraterone. This approach can be extended to other drug analyses, promoting quality and efficiency in pharmaceutical research. The proposed method was found to be linear, with concentrations of 2-100 ppm having r^2 : 0.998. The remarkable % recovery (within 98-102%) of the drug reflects that the excipients existing in the tablet formulation have no impediment in the quantitation of the drug. The optimized conditions by AQbD of the anticipated method established that the proposed study was cost-effective, extremely robust, and stability indicating. Therefore, the developed stability-indicating method was economical, accurate, and precise. It can be successfully implemented for the routine analysis of niraparib abiraterone acetate in bulk and pharmaceutical formulations.

REFERENCES

1. Van Andel L, Zhang Z, Lu S, Kansra V, Agarwal S, Hughes L, Tibben MM, Gebretensae A, Rosing H, Schellens JH, Beijnen JH. Liquid chromatography-tandem mass spectrometry assay for the quantification of niraparib and its metabolite M1 in human plasma and urine. *Journal of Chromatography B*. 2017 Jan 1;1040:14-21.
2. Beg S, Malik AK, Afzal O, Altamimi AS, Kazmi I, Al-Abbasi FA, Almalki WH, Barkat MA, Kawish SM, Pradhan DP, Rahman M. Systematic development and validation of a RP-HPLC method for estimation of abiraterone acetate and its degradation products. *Journal of Chromatographic Science*. 2021 Jan;59(1):79-87.
3. Kamani VG, Sujatha M. an Innovative Stability Indicating Hplc Method With Impurity Profiling of Niraparib-an Anticancer Drug in Pharmaceutical Formulations. *Rasayan J. Chem.*. 2022;15:2976-83.
4. Scott LJ. Niraparib: first global approval. *Drugs*. 2017 Jun;77:1029-34.

5. Chandra Reddy BJ, Sarada NC. Development and validation of a novel RP-HPLC method for stability-indicating assay of Abiraterone acetate. *Journal of Liquid Chromatography & Related Technologies*. 2016 Apr 20;39(7):354-63.
6. Yahya BA, Bawazir AS, Sangshetti JN, Baig SS, Shaikh SS. QbD-based development and validation of an efficient RPHPLC method for estimation of abiraterone acetate in bulk, tablet, and in-house-developed nano-formulation. *Analytical chemistry letters*. 2021 Jan 2;11(1):112-30.
7. Goud VM, Rani BS, Sharma JV, Sirisha P. Development and Validation for estimation of Abiraterone acetate in Bulk and Pharmaceutical Dosage Form by UPLC. *Research Journal of Pharmacy and Technology*. 2019;12(6):3029-32.
8. Belleville T, Noé G, Huillard O, Thomas-Schoemann A, Vidal M, Goldwasser F, Alexandre J, Blanchet B. A HPLC-fluorescence method for the quantification of abiraterone in plasma from patients with metastatic castration-resistant prostate cancer. *Journal of Chromatography B*. 2015 May 1;989:86-90
9. Dai L, Lv P, He Y, Wang X, Chen L, Dai J. A Novel LC-MS Method for the Determination of Abiraterone in Rat Plasma and its Application to Pharmacokinetic Studies. *Current Pharmaceutical Analysis*. 2022 Jul 1;18(6):590-7.
10. Kumar SV, Rudresha G, Gurav S, Zainuddin M, Dewang P, Kethiri RR, Rajagopal S, Mullangi R. Validated RP-HPLC/UV method for the quantitation of abiraterone in rat plasma and its application to a pharmacokinetic study in rats. *Biomedical Chromatography*. 2013 Feb; 27(2):203-7.

AJPTR is

- **Peer-reviewed**
- **bimonthly**
- **Rapid publication**

Submit your manuscript at: editor@ajptr.com

