

1903. Z Watson, Hugh, Bracondale, The Avenue, Cambridge.  
 1908. Weaver, G. H., 31, Devonshire Road, Palmer's Green, London, N.  
 1900. Webb, Walter F., 202, Westminster Road, Rochester, N.Y., U.S.A.  
 1902. Weeks, Wm. H., jr., 508, Willoughby Avenue, Brooklyn, N.Y., U.S.A.  
 1895. Welch, Robert John, M.R.I.A., 49, Lonsdale Street, Belfast.  
 1907. Wheat, Silas C., 987, Sterling Place, Brooklyn, N.Y., U.S.A.  
 1886. Whitwell, Wm., F.L.S., Brookside, Darley Knowle, Warwickshire.  
 1911. Williams, James M. M., Imperial House, Pontlottyn, Cardiff.  
 1889. Williams, John M., 31, Grove Park, Liverpool.  
 1906. Winkworth, John F., 290, Burdett Road, London, E.  
 1890. Wood, Albert, Midland Lodge, Sutton Coldfield, Warwickshire.  
 1910. Woodcock, R., Fauvic, Jersey.  
 1901. Z Woodruffe-Peacock, Rev. E. A., F.L.S., etc., Cadney, Brigg, Lincs.  
 1911. Woods, Rev. F. H., B.D., Bainton Rectory, Driffeld.  
 1898. Woods, Henry, M.A., F.G.S., 39, Barton Road, Cambridge.  
 1886. Z Woodward, Bernard B., F.L.S., etc., 4, Longfield Rd., Ealing, W.  
 1903. Worsdale, R., 102, Dudley Terrace, Dudley Road, Grantham.  
 1906. Wragge, Clement L., F.R.G.S., etc., Perth, Western Australia.  
 1895. Wright, Charles East, Woodside, Rockingham Road, Kettering.

**Succinea elegans** Risso new to the Orkneys.—Amongst a few common species of shells which I received recently from the Orkneys are examples of *Succinea elegans*. These have been verified by Mr. J. W. Taylor, and the Recorder tells me that it is a new record for vice-county III.—J. R. LE B. TOMLIN (*Read before the Society*, Nov. 8, 1911).

**Vertigo angustior** Jeff. in Hertfordshire.—In February, 1911, I noticed that in the earth cast up by moles in the low-lying pastures which border the large reservoir at Wilstone, near Tring, there were many bleached shells of mollusca. A section exposed by a ditch which traversed one of the meadows showed that the thin layer of alluvium overlying the chalk was crowded with shells. The associated species suggested that the construction of the reservoir about a hundred years ago had entirely altered the character of the place, transforming swampy ground into comparatively dry meadow land, and, incidentally, destroying the habitat of the mollusca. An hour's search in the mole-hills yielded two specimens of *Vertigo angustior*, many *V. pygmaea*, *V. antivertigo*, *Jamnia muscorum*, *Vallonia excentrica*, *Hygromia hispida*, *Succinea elegans* and *Cochlicopa lubrica*. In smaller numbers, but conspicuous owing to their larger size, were *Helix nemoralis* and *Helicigona arbustorum* var. *alpestris*, whilst *Succinea putris* and *Limnaea truncatula* were not uncommon. I found two specimens of *Cacilioides acicula* and a few *Euconulus fulvus*, *Carychium minimum* and *Pisidium casertanum*. — CHAS. OLDHAM (*Read before the Society*, Sept. 13th, 1911).



Oldham, Charles. 1912. "Vertigo angustior Jeff. in Hertfordshire." *Journal of Conchology* 13(9), 264–264. <https://doi.org/10.5962/p.406047>.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/98594>

**DOI:** <https://doi.org/10.5962/p.406047>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/406047>

**Holding Institution**

Smithsonian Libraries and Archives

**Sponsored by**

Biodiversity Heritage Library

**Copyright & Reuse**

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.