

Tableau des distances génétiques intraspécifiques

Gruhn et al. : Le genre Resinicum dans les Outre-Mer, avec la description de sept espèces nouvelles

Taxon	% moyen	% minimum	% maximum
Resinicum austroasianum	0.0046	0.0046	0.0046
Resinicum bicolor	NA	NA	NA
Resinicum confertum	NA	NA	NA
Resinicum friabile	0.0095	0.0000	0.0202
Resinicum luteolus	0.0014	0.0000	0.0036
Resinicum grandisporum	0.0095	0.0000	0.0255
Resinicum lateastrocystidium	0.0000	0.0000	0.0000
Resinicum monticola	0.0020	0.0020	0.0020
Resinicum mutabile	0.0050	0.0000	0.0132
Resinicum nouraguense	NA	NA	NA
Resinicum rimulosum	0.0105	0.0105	0.0105
Resinicum saccharicola	0.0203	0.0022	0.0617
Resinicum tenue	0.0000	0.0000	0.0000
Resinicum yunnanense	0.0124	0.0124	0.0124
Skvortzovia furfuracea	0.1667	0.0000	0.2578
Rickenella fibula	NA	NA	NA
Resinicum simile	NA	NA	NA
Resinicum gastrodiae	0.0000	0.0000	0.0000
Resinicum borbonicum	0.0000	0.0000	0.0000
Resinicum guttatum	NA	NA	NA
Resinicum mahorense	NA	NA	NA