

# GammaPBHPlotter: A public code for calculating the complete Hawking evaporation gamma-ray spectra from primordial black holes

John Carlini<sup>1</sup> and Ilias Cholis<sup>1</sup>

<sup>1</sup> Department of Physics, Oakland University, Rochester, Michigan, 48309, USA

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

## Software

- [Review](#)
- [Repository](#)
- [Archive](#)

Editor: [Open Journals](#)

## Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

## License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

## Summary

We present GammaPBHPlotter, a public Python package for calculating and plotting the Hawking radiation gamma-ray spectra of primordial black holes in the mass range of  $5 \times 10^{14}$  to  $10^{18}$  grams. This tool allows users to compute the monochromatic and mass-averaged spectra of black holes over a range of parameters. We include the primary/direct Hawking emission, the secondary emission from the decay and hadronization of unstable particles, the final state radiation, and the in-flight annihilation gamma-ray emission components.

## Statement of Need

Hawking radiation (Hawking, 1974) remains an unobserved property of black holes. As the temperature of black holes is inversely proportional to the square of their mass, conventional stellar mass black holes are expected to emit too little radiation to ever be detected. However, primordial black holes (PBHs) that could have formed from the collapse of primordial perturbations in the early universe can provide detectable signals (Carr et al., 2016). PBHs with mass less than  $10^{14}$  grams would have evaporated via Hawking radiation long before the present age of the universe. Upcoming gamma-ray telescopes such as e-ASTROGAM (Tavani & others, 2018) and AMEGO-X (Caputo & others, 2022) will be sensitive enough in the MeV range to detect the Hawking spectra of PBHs lying between this lower bound and  $10^{19}$  grams. We have developed GammaPBHPlotter, as an open source and user friendly means of simulating gamma-ray spectra produced from different PBH mass-distributions. By simulating in-flight annihilation, and final state radiation, this package additionally allows for more accurate and comprehensive simulations than existing software.

## Hawking Spectra

### Modeling the emission components

The gamma-ray spectrum of a PBH within the relevant mass range consists of four primary components; direct/primary Hawking radiation, secondary radiation, final-state radiation, and in-flight annihilation.

Direct Hawking radiation accounts for all kinematically allowed elementary particles formed at the event horizon (Hawking, 1974), including gamma-ray photons. Secondary radiation originates from the decay of unstable particles and contributes significantly at lower energies. We rely on BlackHawk (Arbey & Auffinger, 2021) to evaluate the gamma-ray primary and secondary spectral components. BlackHawk uses PYTHIA (Sjöstrand et al., 2015) for the

37 modeling of the hadronization and decay processes leading to the secondary spectra. Final-  
38 state radiation originates from relativistic electrons and positrons and has a differential spectrum  
39 given by Eq.~1,

$$\frac{dN_{\gamma}^{\text{FSR}}}{dE_{\gamma}} = \frac{\alpha}{2\pi} \int dE_{e^+} \frac{dN_{e^+}}{dE_{e^+}} \left( \frac{2}{E_{\gamma}} + \frac{E_{\gamma}}{E_{e^+}^2} - \frac{2}{E_{e^+}} \right) \left[ \ln \left( \frac{2E_{e^+} + (E_{e^+} - E_{\gamma})}{m_{e^+}^2} \right) - 1 \right],$$

40 where  $\alpha = 137.037$  is the fine structure constant,  $E_{e^+}$  is the kinetic energy of a given positron  
41 ( $e^+$ ),  $E_{\gamma}$  is the energy of the emitted photon,  $m_{e^+} = 0.511$  MeV is the rest mass of the  
42 electron, and  $\frac{dN_{e^+}}{dE_{e^+}}$  the differential spectrum of emitted electrons/positrons. In addition to the  
43 previously mentioned components, gamma-rays can be produced through pair-annihilation of  
44 positrons with interstellar medium electrons. This is known as in-flight annihilation and its  
45 differential spectrum is (Keith et al., 2022),


$$\begin{aligned} \frac{dN_{\gamma}^{\text{IA}}}{dE_{\gamma}} &= \frac{\pi\alpha^2 n_H}{m_e} \int_{m_e}^{\infty} dE_{e^+} \frac{dN_{e^+}}{dE_{e^+}} \int_{E_{\min}}^{E_{e^+}} \frac{dE}{dE/dx} \frac{P_{E_{e^+} \rightarrow E}}{(E^2 - m_e^2)} \\ &\times \left( -2 - \frac{(E + m_e)(m_e^2(E + m_e) + E_{\gamma}^2(E + 3m_e) - E_{\gamma}(E + m_e)(E + 3m_e))}{E_{\gamma}^2(E - E_{\gamma} + m_e)^2} \right). \end{aligned}$$

46 We take  $n_H = 1 \text{ cm}^{-3}$  as the density of interstellar medium hydrogen (and by extension  
47 electrons).  $E_{e^+}$  is again the initial positron total energy,  $E$  is the final positron total energy,  
48  $dE/dx$  is the rate of positron energy lost per path via the Bethe-Bloch formula (Bethe  
49 & Ashkin, 1953),  $E_{\gamma}$  is the resulting photon energy from annihilation, and  $P_{E_{e^+} \rightarrow E}$  is the  
50 probability of a particular positron of a given initial and final energy to decay. This probability  
51 matrix can be calculated as (Keith et al., 2022),

$$P_{E_{e^+} \rightarrow E} = \exp \left( -n_H \int_E^{E_{e^+}} \sigma_{\text{ann}}(E') \frac{dE'}{dx} dE' \right),$$

52 where  $\sigma_{\text{ann}}$  is the cross section of annihilation for positrons of a given energy.


53 In Fig.~1, we give the individual gamma-ray spectral components as well as their sum for a  
54 PBH of mass  $3 \times 10^{15}$  grams.



**Figure 1:** The total gamma-ray spectrum of a  $3 \times 10^{15}$  grams PBH as well as its components.

## PBH Mass Distribution

Users can calculate the gamma-ray spectra from four types of PBH mass distributions. Those are, i) a monochromatic distribution with a mass to be set in the range of  $5 \times 10^{13}$  to  $1 \times 10^{19}$  grams, ii) a Gaussian distribution of PBH masses originating from a Gaussian distribution of density perturbations (Biagetti et al., 2021), iii) a more realistic non-Gaussian PBH mass distribution from (Biagetti et al., 2021) and iv) a log-normal distribution of PBH masses. In Fig.~2, we give the gamma-ray spectra from monochromatic and Gaussian PBH mass-distributions.



**Figure 2:** The total gamma-ray spectrum per PBH, from a PBH of mass  $3 \times 10^{15}$  grams (blue line) and from a Gaussian distribution of density perturbations leading to a distribution of a mean mass of  $3 \times 10^{15}$  grams.  $\sigma$  refers to the standard deviation of initial density perturbations (Biagetti et al., 2021).

## Software content

GammaPBHPlotter was written in Python version 3.9 and is capable of running on Windows, Linux, and Mac. The main code uses five modules in its routine. Those being colorama (Hartley, 2022), NumPy (Harris et al., 2020), Matplotlib (Hunter, 2007), tqdm, (Costa-Luis & developers, 2024) and SciPy (Virtanen et al., 2020). The package is available via PyPI and the archived release is cited as (Carlini & Cholis, 2025).

We acknowledge the use of BlackHawk (Arbey & Auffinger, 2021). This material is based upon work supported by the U.S. Department of Energy, Office of Science, Office of High Energy Physics, under Award No. DE-SC0022352.

## References

- Arbey, A., & Auffinger, J. (2021). Physics beyond the standard model with BlackHawk v2.0. *European Physical Journal C*, 81, 910. <https://doi.org/10.1140/epjc/s10052-021-09702-8>
- Bethe, H., & Ashkin, J. (1953). Passage of radiations through matter. In E. Segrè (Ed.), *Experimental nuclear physics* (pp. 253–253). John Wiley & Sons.
- Biagetti, M., De Luca, V., Franciolini, G., Kehagias, A., & Riotto, A. (2021). The formation probability of primordial black holes. *Physics Letters B*, 820, 136602. <https://doi.org/10.1016/j.physletb.2021.136602>
- Caputo, R., & others. (2022). All-sky medium energy gamma-ray observatory eXplorer mission concept. *Journal of Astronomical Telescopes, Instruments, and Systems*, 8(4), 044003. <https://doi.org/10.1117/1.JATIS.8.4.044003>
- Carlini, J., & Cholis, I. (2025). *GammaPBHPlotter: Code and user manual* (Version v1.1.2). Zenodo. <https://doi.org/10.5281/zenodo.17527115>

- 85 Carr, B., Kühnel, F., & Sandstad, M. (2016). Primordial black holes as dark matter. *Physical*  
86 *Review D*, 94(8), 083504. <https://doi.org/10.1103/PhysRevD.94.083504>
- 87 Costa-Luis, C. O. da, & developers, tqdm. (2024). *Tqdm: A fast, extensible progress bar for*  
88 *python and CLI*. <https://doi.org/10.5281/zenodo.11107065>
- 89 Harris, C. R., Millman, K. J., Walt, S. J. van der, Gommers, R., Virtanen, P., Cournapeau, D.,  
90 Wieser, E., Taylor, J., Berg, S., Smith, N. J., Kern, R., Picus, M., Hoyer, S., Kerkwijk,  
91 M. H. van, Brett, M., Haldane, A., Fernández del Río, J., Wiebe, M., Peterson, P., ...  
92 Oliphant, T. E. (2020). Array programming with NumPy. *Nature*, 585(7825), 357–362.  
93 <https://doi.org/10.1038/s41586-020-2649-2>
- 94 Hartley, J. (2022). *Colorama*. <https://pypi.org/project/colorama/>
- 95 Hawking, S. W. (1974). Black hole explosions? *Nature*, 248(5443), 30–31. [https://doi.org/](https://doi.org/10.1038/248030a0)  
96 [10.1038/248030a0](https://doi.org/10.1038/248030a0)
- 97 Hunter, J. D. (2007). Matplotlib: A 2D graphics environment. *Computing in Science &*  
98 *Engineering*, 9(3), 90–95. <https://doi.org/10.1109/MCSE.2007.55>
- 99 Keith, C., Hooper, D., Linden, T., & Liu, R. (2022). Sensitivity of future gamma-ray telescopes  
100 to primordial black holes. *Physical Review D*, 106(4), 043003. [https://doi.org/10.1103/](https://doi.org/10.1103/PhysRevD.106.043003)  
101 [PhysRevD.106.043003](https://doi.org/10.1103/PhysRevD.106.043003)
- 102 Sjöstrand, T., Ask, S., Christiansen, J. R., Corke, R., Desai, N., Ilten, P., Mrenna, S., Prestel,  
103 S., Rasmussen, C. O., & Skands, P. Z. (2015). An introduction to PYTHIA 8.2. *Computer*  
104 *Physics Communications*, 191, 159–177. <https://doi.org/10.1016/j.cpc.2015.01.024>
- 105 Tavani, M., & others. (2018). Science with e-ASTROGAM: A space mission for MeV–GeV  
106 gamma-ray astrophysics. *Journal of High Energy Astrophysics*, 19, 1–106. [https://doi.org/](https://doi.org/10.1016/j.jheap.2018.07.001)  
107 [10.1016/j.jheap.2018.07.001](https://doi.org/10.1016/j.jheap.2018.07.001)
- 108 Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D.,  
109 Burovski, E., Peterson, P., Weckesser, W., Bright, J., Walt, S. J. van der, Brett, M.,  
110 Wilson, J., Millman, K. J., Mayorov, N., Nelson, A. R. J., Jones, E., Kern, R., Larson, E., ...  
111 Contributors, S. 1.0. (2020). SciPy 1.0: Fundamental algorithms for scientific computing  
112 in python. *Nature Methods*, 17(3), 261–272. <https://doi.org/10.1038/s41592-019-0686-2>