

SPECIES

To Cite:

Chryso RB, Ragul S. Occurrence of ragged sea hare, *Bursatella leachii* Blainville, 1817 (Gastropoda: Heterobranchia: Aplysiidae) from Vellar Estuary, Tamil Nadu, India. *Species* 2023; 24: e49s1544
doi: <https://doi.org/10.54905/diss/v24i73/e49s1544>

Author Affiliation:

Centre of Advanced Study in Marine Biology, Annamalai University, Parangipettai, 608502, Tamil Nadu, India

*Corresponding author

Centre of Advanced Study in Marine Biology, Annamalai University, Parangipettai, 608502, Tamil Nadu, India
Email: nivinragulsiva@gmail.com

Peer-Review History

Received: 19 April 2023
Reviewed & Revised: 22/April/2023 to 01/June/2023
Accepted: 05 June 2023
Published: 9 June 2023

Peer-Review Model

External peer-review was done through double-blind method.

Species

pISSN 2319–5746; eISSN 2319–5754



© The Author(s) 2023. Open Access. This article is licensed under a [Creative Commons Attribution License 4.0 \(CC BY 4.0\)](https://creativecommons.org/licenses/by/4.0/), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

Occurrence of ragged sea hare, *Bursatella leachii* Blainville, 1817 (Gastropoda: Heterobranchia: Aplysiidae) from Vellar Estuary, Tamil Nadu, India

Reslin Berlin Chryso, Sivakumar Ragul*

ABSTRACT

This paper documents a new record of *Bursatella leachii* Blainville, 1817 from Vellar estuary, Parangipettai, South-eastern coast of Tamil Nadu. Eight specimens of *Bursatella leachii* with standard length (SL) of 43.48-55.92 mm and weight of 20.9-26.2g were collected at Vellar estuary, Parangipettai, during June 2021 - July 2022. The present record of this species from estuarine waters of Tamil Nadu, India expands and confirms the known distribution of *Bursatella leachii*, previously reported from Australia, Brazil, Canada, China, Greece, Hong Kong, Iran, Israel, Italy, Japan, Lebanon, Malta, Marshall Island, Mozambique, Panama, Philippines, Slovenia, South Africa, Tunisia, Turkey and United Kingdom.

Keywords: Ragged Sea Hare, Vellar estuary, South east coast of India.

1. INTRODUCTION

Molluscs are the second largest animal phylum on earth and include an enormous diversity of species (Gosliner et al., 2008). Along the Indian coast, Phylum Mollusca comprises of 6 orders and 53 families, 292 species of marine heterobranchia with 14 aplysiidae family (Sreeraj, 2020). Sea Hare are generally termed as 'butterflies of the ocean' because of their prominent coloration (Behera et al., 2020). *B. leachii* Blainville, (1817) a gastropod mollusc of the order Anaspide and family Aplysiidae, colloquially known as ragged sea hare and shaggy sea hare. The genus *Bursatella* contains only three species, *Bursatella hirsuta* Nimbs and Wilson, (2020) (southern and western Australia); *Bursatella leachii* Blainville, (1817) (Pantropical distribution which includes South Africa to Indo-Pacific region); *Bursatella ocelligera* (Bergh, 1902) (Philippines).

B. leachii are usually observed in intertidal and subtidal of ocean inlets and sand or muddy bottoms of estuaries. This species frequently encountered from tropical and subtropical seagrass and mangrove communities (Lowe and Turner, 1976). Clarke, (2004) stated that this species mostly depends on seagrass habitats because of their sporadic high densities and their feeding specialization on cyanobacteria. *B. leachii* is distributed worldwide in warm temperate to tropical

marine environments (Clarke, 2006; Selfati et al., 2017) and there are reports on the occurrence of this species in estuarine environments (Sethi et al., 2015). The maximum standard length (SL) recorded for this species is 250 mm (Voss, 1980). The type specimen of this species was originally collected from Italy, Levantine coast, France, Morocco, southern Spain and east Mediterranean Red Sea.

During our survey on molluscan diversity of Parangipettai coast, India we collected the above species from the Vellar estuary and confirmed the occurrence of this species in this estuary for the first time. Thus, present record of this species from Vellar estuary expands and confirms the known distribution of *Bursatella leachii*, which is previously reported from Pulicat Lake (Sethi et al., 2015) and Krishna estuary (Behera et al., 2020) in India.

2. MATERIALS AND METHODS

Monthly fish surveys were conducted in Vellar estuary, south-east India, from June 2021 to July 2022, to document cryptic species diversity. Specimens of *Bursatella* were collected using scoop net and by hand picking methods during the receding period of the incoming tides.

Five specimens of *B. leachii* from Vellar mangrove zone (11°29'25"N, 79°45'57"E) on during low tide, when the intertidal region was partially exposed. Three specimens from small seagrass bed adjacent to the fish pond (11°29'32"N, 79°46'06"E). This sand-dwelling species were procured and preserved after undertaking morphological studies.

Morphometric and meristic characters were observed and measured from live specimens anesthetized using clove oil and then euthanized and fixed in 10% formaldehyde and later preserved in 70% ethanol (R). Specimens were identified to species level and measured, based on Bebbington, 1974. Preserved specimens were deposited in the Centre of Advanced Studies in Marine Biology, Faculty of Marine Sciences, Annamalai University, Reference Museum (CASMBURM), Parangipettai, India for further future studies.




Figure 1 Dorsal head region with (A) Two oral tentacles; (B) Two olfactory tentacles (rhinophores); (C) Ocelli (blue eyespots) and (D) Dorsal view of live *Bursatella leachii* Blainville, (1817) collected from Vellar estuary (CASMBURM/230521)

3. RESULTS

Systematics

ORDER Aplysiida Lamarck, 1809

FAMILY Aplysiidae Lamarck, 1809

GENUS *Bursatella* Blainville, 1817

Bursatella leachii Blainville, 1817

Material examined

Five specimens (46.61-55.92 mm SL and weight 21.4-26.2g) (CASMBURM/230520-24), coll. R.B. Chryso on 11 June 2022, intertidal mangrove zone opposite to the Marine Biological Research station, Vellar Estuary (11°29'25"N, 79°45'57"E); Three specimens (43.48-

51.23 mm SL and weight 20.9-24.0g) (CASMBURM/230525-27), coll. S. Ragul, 02 July, 2022, a small seagrass bed (*Halodule* sp.), Vellar Estuary, (11°29'32"N, 79°46'06"E), Tamil Nadu.

Description

The body is compressed with prominent head and neck regions (Figure 1D); head part is wide and short. The body surface was covered with both simple and compound villus of various sizes, with multiple branched villi over the head, cephalic tentacles and the rhinophores region. Dorsal side of the body is blackish grey with olive-green and there were several blue rimed black blotched ocelli (blue eyespots) scattered over the body (Figure 1C).

The body is also covered with many long, fleshy white branched villi that give the species its ragged appearance. Head comprises with four tentacles: Two olfactory tentacles called rhinophores (Figure 1B) originating on the dorsal part of the head and two oral tentacles (Figure 1A), above the mouth region. The gill is protected by a pair of fleshy parapodia. The parapodia were joined high up posteriorly leaving an oval slit on the summit. The foot was rather broad for its length and its anterior margin was double.

Coloration

Colour in live specimen, blackish grey with olive-green and there were several blue rimed black blotched ocelli scattered over the body. Preserved specimen, faded brown body with black eye spots.

4. DISCUSSION

B. leachii is a complex species which has posed many taxonomic statuses with seven geographical subspecies as well as on morphological differences (Eales and Engel, 1935; Rudman, 2007; Bazzicalupo et al., 2020). Rudman, (2007) states that the colour of eye spots can vary within a local population, as can the background colour and the degree of 'woolliness' of *B. leachii*. Correspondingly the Seven subspecies i.e., *Bursatella leachii africana* (Engel, 1926: South Africa; very woolly in appearance), *Bursatella leachii guineensis* (Bebbington, 1969: Ghana; less woolly than *B. leachii Africana*; peacock blue eye spots rather than green), *Bursatella leachii lacunculata* (Gould, 1852: Brazil), *Bursatella leachii leachii* (Blainville, 1817: Indo-Pacific; green ocelli), *Bursatella leachii pleii* (Rang, 1828: West Indies), *Bursatella leachii rosea* (Engel, 1926: West Africa; pinkish in colour), *Bursatella leachii savigniana* (Audouin, 1826: Red Sea)

This species feeds on filamentous brown algae (Masterson, 2008). Correspondingly, From India Sethi et al., (2015) have reported *B. leachii* with maximum length of 75 mm and weight of 23 g from Pulicat Lake and Behera et al., (2020) recorded the average total length of 30.93 ± 1.34 mm and weight of 4.10 ± 0.74 g from Krishna estuary respectively. Ozvarol, (2014) have reported the maximum size of 150 mm along the Mediterranean coast, Turkey.

Appleton et al., (2002) isolated a novel bioactive compound malyngamide from *B. leachii* from New Zealand. Kamiya et al., (2006) derived few antimicrobial and cytotoxic proteins from sea hares. Rajaganapathi et al., (2002) have derived anti-HIV properties from purple fluid protein of Sea Hare, *B. leachii*. Capper et al., (2005) supposed that the sea hare consumes on filamentous cyanobacterium, *B. leachii* capable of sequestering secondary toxic metabolites (e.g., lyngbyatoxin-a) in the digestive gland and in bodily secretions. There is very limited knowledge of the diversity, distribution and occurrence of sea hares from the estuaries of Tamil Nadu. This present report confirms the occurrence of *Bursatella leachii* Blainville, 1817 from Vellar estuarine waters.

Acknowledgements

The authors are thankful to the Director and Dean of the Faculty of Marine Sciences, Annamalai University, for providing necessary facilities.

Author's contribution

Chryso RB collected the specimens and drafted the manuscript. Ragul S prepared the images and reviewed the draft manuscript.

Informed consent

Not applicable.

Ethical approval & declaration

In this article, as per the National Biodiversity Authority (NBA) algae regulations & the ethical regulations of Centre of Advanced Study in Marine Biology, Annamalai University, India, the authors observed the ragged sea hare, *Bursatella leachii* Blainville, 1817 (Gastropoda: Heterobranchia: Aplysiidae) from Vellar Estuary, Tamil Nadu, India. The algae ethical guidelines are followed in the study for species observation, identification & experimentation.

Conflicts of interests

The authors declare that there are no conflicts of interests.

Funding

The study has not received any external funding.

Data and materials availability

All data associated with this study are present in the paper.

REFERENCES AND NOTES

1. Appleton DR, Sewell MA, Berridge MV, Copp BR. A new biologically active malonyamide from a New Zealand collection of the sea hare *Bursatella leachii*. J Nat Prod 2002; 65: 630-631.
2. Bazzicalupo E, Crocetta F, Gosliner TM, Berteaux-Lecellier V, Camacho-García YE, Chandran BS, Valdés Á. Molecular and morphological systematics of *Bursatella leachii* de Blainville, 1817 and *Stylocheilus striatus* Quoy and Gaimard, 1832 reveal cryptic diversity in pantropically distributed taxa (Mollusca: Gastropoda: Heterobranchia). Invertebr Syst 2020; 34(5):535-568.
3. Behera PR, Jishnudev MA, Shiva P, Ghosh S. New distributional record of Ragged Sea Hare, *Bursatella leachii* Blainville, 1817 along western Bay of Bengal. J Mar Biol Assoc India 2020; 62(2):125-127.
4. Bergh R. Gastropoda opisthobranchiata. B Lunos 1902; 1:60.
5. Blainville HMD. Dictionnaire des Sciences Naturelles. Cuvier F (Editor) 1817; 5:138.
6. Capper A, Tibbetts IR, Neil JMO, Shaw GR. The fate of *Lyngbya majuscula* toxins in three potential consumers. J Chem Ecol 2005; 31:1595-1606.
7. Clarke CL. The population dynamics and feeding preferences of *Bursatella leachii* (Opisthobranchia: Anaspidea) in northeast Queensland, Australia. Rec West Aust Mus 2006; 69:11-21. doi: 10.18195/issn.0313122x.69.2006.011-021
8. Gosliner TM, Behrens DW, Valdés A. Indo-Pacific nudibranchs and sea slugs, a field guide to the world's most diverse fauna. Sea Challengers, Gig Harbor, WA/California Academy of Sciences, San Francisco, CA 2008; 426.
9. Kamiya H, Sakai R, Jimbo M. Bioactive molecules from sea hares. Prog Mol Subcell Biol 2006; 43:215-239.
10. Lowe EF, Turner RL. Aggregation and trail-following in juvenile *Bursatella leachii* plei (gastropoda: Opisthobranchia). Veliger 1976; 19:153-155.
11. Masterson J. *Bursatella leachii*. In Smithsonian Marine Station at Fort Pierce 2008. http://www.sms.si.edu/irlspec/Bursatella_leachii.html
12. Nimbs MJ, Wilson NG. Description of a new species of *Bursatella* Blainville, 1817 (Gastropoda, Aplysiida, Aplysiidae) from southern Australia. Molluscan Res 2020; 40(4):369-378.
13. Rajaganapathi J, Kathiresan K, Singh TP. Purification of anti-HIV protein from purple fluid of the sea hare *Bursatella leachii* de Blainville. Mar Biotechnol (NY) 2002; 4:447-453.
14. Rudman WB. *Bursatella leachii* subspecies. In Sea Slug Forum. Australian Museum: Sydney, NSW, Australia 2007. <http://www.seaslugforum.net/find/19181>
15. Selfati M, El-Ouamari N, Crocetta F, Mesfioui A, Boissery P, Bazairi H. Closing the circle in the Mediterranean Sea: *Bursatella leachii* Blainville, 1817 (Mollusca: Gastropoda: Anaspidea) has reached Morocco. Bioinvasions Rec 2017; 6:129-134. doi: 10.3391/bir.2017.6.2.07
16. Sethi S, Kokane MR, Otta SK, Sethi G. First record of ragged sea hare *Bursatella leachii* Blainville, 1817 (Opisthobranchia: Euopisthobranchia: Aplysiidae) in Pulicat Lake, east coast of India. J Mar Biol Assoc U K 2015; 8.
17. Sreeraj CR. Mollusca: Heterobranchia. In: Faunal Diversity of Biogeographic Zones: Coasts of India. Zool Surv India 2020; 575-594.
18. Voss GL. Seahorse life of Florida and the Caribbean. Dover Publications, Inc., NY 1980; 199.