



Annals and Magazine of Natural History

Series 7

ISSN: 0374-5481 (Print) (Online) Journal homepage: <http://www.tandfonline.com/loi/tnah13>

XLVII.—New generic names for some Entomostraca and Cirripedia

Canon A.M. Norman F.R.S.

To cite this article: Canon A.M. Norman F.R.S. (1903) XLVII.—New generic names for some Entomostraca and Cirripedia, *Annals and Magazine of Natural History*, 11:64, 367-369, DOI: [10.1080/00222930308678780](https://doi.org/10.1080/00222930308678780)

To link to this article: <http://dx.doi.org/10.1080/00222930308678780>



Published online: 29 Sep 2009.



Submit your article to this journal [↗](#)



Article views: 1



View related articles [↗](#)



Citing articles: 3 View citing articles [↗](#)

spine on their upper surface near apex and with four shorter spines arranged in pairs beneath.

Long. 7 millim.; exp. pronot. angl. 3 millim.

Hab. Dorey (Wallace, Brit. Mus.).

NABINÆ.

Nabis subcoleoptratus.

Nabis subcoleoptratus, Kirby, in Rich. Faun. Bor.-Amer. iv. p. 282 (1837).

Nabis xanthopus, MSS., Walk. Cat. Het. vii. p. 143. 22 (1873).

XLVII.—New Generic Names for some Entomostraca and Cirripedia. By Canon A. M. NORMAN, F.R.S. &c.

In preparing for publication a second edition of the Catalogue of the Crustacea in my collection and tracing the generic names of the lower orders in that class I find that the following changes are necessary.

CLADOCERA.

Genus SIMOSA *, nov. nom.

= *Simocephalus*, Schœdler.

Two genera named *Simocephalus* were published almost simultaneously. The name was used for a genus of snakes by Dr. Günther in his 'Catalogue of Columbrine Snakes,' the exact date of the publication of which, I learn from him, was October 9, 1858; and the name was applied to a genus of Cladocera by Schœdler in a paper read September 22, 1858, and as it was illustrated by a quarto plate, it is not likely that it could have been published within so short a time after the reading as October 9.

Genus RHYNCHOTALONA †, nov. nom.

In 1861 G. O. Sars instituted a genus *Harporhynchus* for the interesting species he was describing—*H. falcatus*; but that name having been previously (in 1848) employed by Cabanis for a genus of birds, in 1884 C. L. Herrick substituted the name *Leptorhynchus*—a still more unfortunate choice, for that name had been employed at least three times previously, and it is therefore necessary to supersede it.

* σιμός, snub-nosed.

† ῥυγχωτός, having a beak.

COPEPODA.

HALICYCLOPS *, nov. nom.

In 1893 Claus instituted a genus *Hemicyclops* with the type *Cyclops æquoreus*, Fischer; but the name had been previously used in a different sense by Boeck in 1872.

Genus EUTERPINA, nov. nom.

In 1873 Claus employed the name *Euterge* for a genus of marine Copepoda with the type *E. gracilis*; but that generic name was preoccupied by Swainson (1831) among the Lepidoptera.

Genus TEGASTES †, nov. nom.

I propose this name as a substitute for *Amygone*, Claus, 1860, type *A. satyrus*, Claus, since *Amygone* was not only a name given in 1785 to a larval form by O. F. Müller, of which Claus was aware when he instituted his genus, but was employed by Savigny in 1817 for a genus of Annelids.

Genus DACTYLOPUSIA, nov. nom.

Dactylopus is in use for a genus of fishes thus named by Gill in 1859, and therefore it is necessary that the same name applied to a genus of Copepods by Claus in 1862 should be changed. Type *D. Stræmii* (Baird).

CLAUSIDIUM, Kossman, 1874.

This name must be employed instead of *Hersilia*, Philippi, a name preoccupied by Savigny in Arachnida and by Dejean in 1834 for a genus of Coleoptera.

Genus PHYLLOTHYREUS ‡, nov. nom.

Instead of *Phyllophora*, H. Milne-Edwards, a name which had been three times earlier employed. The type is *Phyllothureus cornutus* (H. Milne-Edwards). I may take this opportunity of announcing this interesting species as a member of the British fauna. It has been known to me for more than forty years. It was sent to me by William Laughrin at a

* ἅλς, the sea.

† τεγαστής for στεγαστής, like τέγος for στέγος, one who covers up (*Stegastes* already in use).

‡ φύλλον, a leaf; θυρεός, an oblong shield.

time when I was employing that old coastguardsman to collect fish-parasites for me. He said that he had found it on the blue shark at Polperro.

The two Cornish specimens agree in every particular with Milne-Edwards's description and capital figures (Hist. des Crust. vol. iii. p. 472, pl. xxxviii. figs. 13, 14). I am not aware that the species has been recorded by any other author; and the type specimen was from near Tongatubu, Friendly Islands!

CONGERICOLA, J. van Beneden.

This name must be used with van Beneden's type *C. pallida*. The earlier name *Cygnus*, H. Milne-Edwards, 1840, had been preoccupied by Hübner, 1816, for a genus of Lepidoptera.

CIRRIPEdia.

Genus TRYPETESA *, nov. nom.

Type *Trypetesa (Alcippe) lampas* (A. Hancock).

Unfortunately the name *Alcippe*, which Hancock gave to this very interesting burrowing *Lepas* found off the Northumberland coast by him, was preoccupied by Blyth in 1844, and is in use for a genus of birds. The genus *Alcippe*, Hancock, is not to be found in either Scudder or Waterhouse and Sharp. In Scudder, moreover, *Alcippe*, Blyth, did not appear; and it was no doubt in consequence of this that a third genus *Alcippe* was instituted in 1896 among insects.

XLVIII.—British Land Isopoda.—Supplement.

By Canon A. M. NORMAN, F.R.S. &c.

IN the 'Annals' for January 1899 (ser. 7, vol. iii. p. 70) I published a paper containing notes on all the Land Isopoda known to belong to our fauna up to that date. Three species have since been added, which I will now notice, with some other brief remarks.

Genus TRICHONISCUS, Brandt.

Trichoniscus vividus (Koch).

Dr. Scharff has recorded this species from Cappagh, Co.

* *τρυπητής*, a borer. I have used this particular ending because *Trypetes* and *Trypeta* are already in use.

Ann. & Mag. N. Hist. Ser. 7. Vol. xi.

27