



CODEN [USA]: IAJPBB

ISSN: 2349-7750

INDO AMERICAN JOURNAL OF PHARMACEUTICAL SCIENCES

Available online at: <http://www.iajps.com>

Research Article

EXPERIMENTAL AND COMMISSION TESTS OF EFFICIENCY OF THE NEW DRUG FENBENTAL GRANULATE 10% IN CALVES NEOASCARIDOSIS

¹Anatoly Murashevich Bittirov, ¹Anvar M. Ataev, ¹Murat N. Tuganov, ²Mutalif I. Kalabekov, ²Marina K. Kurmanova, ²Safiyat A. Begieva, ³Shamil M. Kadyzhev, ⁴Leyla Kh. Eldarova, ⁵Eduard V. Beslaneev, ⁶Seda Sh. Mantaeva

¹Department of Veterinary Medicine Kabardino-Balkarian State Agrarian University named after V.M. Kokov, Nalchik, Russia., ²Department of Veterinary Sanitary Expertise and Zootechny, Kabardino-Balkarian State Agrarian University named after V.M. Kokov, Nalchik, Russia., ³Departments of Veterinary Medicine and Zootechny of the North Caucasus State Humanitarian Technological Academy, Cherkessk, Russia., ⁴Departments of Ecology and Nature Management of the Dagestan State Medical Academy, Makhachkala, Russia., ⁵Department of Commodity Science and Tourism of Kabardino-Balkarian State Agrarian University named after V.M. Kokov, Nalchik, Russia., ⁶Departments of Botany, Zoology and Ecology, Chechen State University, Grozny, Russia.

Article Received: September 2019 **Accepted:** October 2019 **Published:** November 2019

Abstract:

In the North Caucasus, neoascariidosis of calves (among nematodosis of the digestive tract of young cattle) is a poorly understood disease. However, this pathology in calves is found with an invasion extensity of 6 to 28%, which requires the development of new methods of therapy and prevention using new domestic complex preparations. In experiments with group prescribing, the new multidisperse anthelmintic drug Fenbental granulate 10% at a dose of 20 mg / kg body weight, mixed with food 1: 100, showed EE -87, 50% and IE - 91, 10% and at a dose of 30 mg / kg body weight, mixed with food 1: 100, showed EE -100% and IE - 100% is recommended as an effective of means neoascariidosis calves. Fenbental granulate 10% of at a dose of 30 mg / kg body weight during commission tests was showed EE 90, 00% with an IE of 92,40%. Albefen granulate 10% experimentally and in commission experiments is a highly effective drug and is recommended for the treatment and prevention of calves neoascariidosis by individual and group methods of with food.

Keywords: calves; invasion; neoascariidosis, Fenbental granulate 10%; extensefficiency; Intensefficiency.

Corresponding author:

Anatoly Murashevich Bittirov,

Department of Veterinary Medicine, Faculty of Veterinary Medicine,
Kabardino-Balkarian State Agrarian University named after V.M. Kokov,
Nalchik, Russia. Email: bam_58a@mail.ru

QR code



Please cite this article in press Anatoly Murashevich Bittirov et al., *Experimental And Commission Tests Of Efficiency Of The New Drug Fenbental Granulate 10% In Calves Neoascariidosis.*, Indo Am. J. P. Sci, 2019; 06(11).

INTRODUCTION:

In the North Caucasus, neoascariidosis of calves (among nematodoses of the digestive tract of young cattle) is a poorly understood disease. However, according to many authors, this pathology in calves is found with an invasion extensity of 6 to 28% [1, 4, 5, 7, 10, 13, 15, 16, 17]¹. In calves stationary content, regardless of breed, neoascariidosis appears enzootically with EI 5–12% [2, 4, 7, 8, 9, 16, 17], in of calves with of pasture content with EI 8–17% [5, 8, 10], which requires the development of new methods of therapy and prevention using new domestic complex preparations. In this regard, the development of new complex drugs for the treatment and prevention of neoascariidosis in calves and other intestinal nematodoses is an important task [1, 2, 3, 4, 5, ..., 10 - 17]². The goal is to check the anthelmintic activity of the drug Fenbental granulate 10% in calves of neoascariidosis of group method.

MATERIALS AND METHODS:

Experiments to test the anthelmintic activity of the drug Fenbental granulate 10% in neoascariidosis of calves were conducted on 24 heads. Experimental and control of calves (n = 24) were divided into 3 groups of 8 animals each. Calves 1 group (n = 8), with invasion of neoascariidosis, was given the anthelmintic drug Fenbental granulate 10% at a dose of 20 mg / kg body weight, group 2 of calves (n = 8) at a dose of 30 mg / kg body weight, once, also by group method. Calves the of 3rd group (n = 8) served as a control, as infested with neoascariidosis, not receiving a new anthelmintic drug Fenbental granulate 10% with food. According to the scheme of

the experiment, after 3, 5, 7, 10 and 15 days after a single injection of the new drug, Fenbental granulate 10%, the feces of all individuals underwent coproscopy [1-7]. Commission tests of Albefen granulate 10% were carried out on 50 of calves. The results of experimental testing of the anthelmintic composition Fenbental granulate 10% with neoascariidosis of calves were subjected to statistical processing using the "Biometrics" program.

RESULTS:

New anthelmintic drug Fenbental granulate 10% per 1 g of powder includes: fenbendazole 250 mg, albendazole 200 mg, copper chelate 100 mg, dry bentonite 450 mg. In the 1st experimental group (n = 8) with neoascariidosis of calves administered with the group method of the new drug Fenbental granulate 10% at a dose of 20 mg / kg of body weight with food 1: 100, showed a rather high extensefficiency - 87, 50% and intensefficiency (IE) – 91, 10% (Table 1). At the same time, in the 2nd group of calves (n = 8) with neoascariidosis of calves, a new anthelmintic composition Fenbental granulate 10% in a dose of 30 mg / kg of body weight showed extensefficiency 100% with intensefficiency (IE) - 100%. This dosage of the new composition Fenbental granulate 10% should be recognized as an effective therapeutic dose (Table 1). Calves of the 3rd group (invasive control, n = 8) remained infected with neoascariidosis when detecting 48,0-52,4 ekz. of eggs and larvae in 5 g of feces.

Table 1- Efficacy anthelmintic composition of Fenbental granulate 10% in neoscaridosis of calves

Group	The number of infected neoscaridos is of calves	The number of free from neoscaridosis of calves after treatment	EE, %	The number of eggs and larvae of in 5 g of feces calves, ekz.		IE, %
				Before therapy	After therapy	
1	8	7	87,50	42,6±3,8	3,8±0,7	91,10
2	8	8	100	44,3±4,1	-	100
3	8	0	0	48,0±5,3	52,4±5,5	0

The commission tests amounted extensefficiency of the new anthelmintic composition Fenbental granulate 10% at a dose of 30 mg / kg of body weight against in of neoscaridosis calves during to 90,00% with intensefficiency (IE) – 92,40% (Table 2).

Table 2- Efficacy of Fenbental granulate 10% anthelmintic composition in neoscaridosis of calves at commission tests (according to coprolaroscopy)

Group	The number of infected lambs	The number of free from nematodes of lambs after treatment	EE, %	The number of eggs and larvae of nematodes in 5 g of feces of lambs, ekz.		IE, %
				Before therapy	After therapy	
1	50	45	90,00	46,2±4,3	3,5±0,6	92,40
2	50	0	0	48,4±4,6	50,7±4,9	0

Thus, the complex anthelmintic compound Fenbental granulate 10% in a dose of 30 mg / kg body weight, mixed with 1:100 feed experimentally, in commission experiments is a highly effective drug and is recommended for the treatment and prevention in neoscaridosis of calves for group administration with feed.

DISCUSSION:

The test results of the new anthelmintic drug Fenbental granulate 10% against the in neoscaridosis of calves were obtained by us for the first time. New data were obtained on the therapeutic efficacy of Fenbental granulate 10% at a dose of 30 mg / kg of body weight of calves. At the same time, information on the need to develop methods for group treatment and prevention of in neoscaridosis of calves is consistent with the positions of many well-known authors [1,2, 3, 4, 5, 8, 9, 10, 11, 12, 13, 14, 15,16,17]³.

CONCLUSION:

For the new anthelmintic drug Fenbental granulate 10%, in the neoscaridosis of calves, a dose of 30 mg/ kg of body weight is therapeutically effective.

During the experiments, Fenbental granulate 10% at a dose of 30 mg / kg body weight in a mixture of 1:100 with food once showed a 85,2-100% effect in neoscaridosis of calves by the group method. The complex anthelmintic compound Fenbental granulate 10% in a dose of 30 mg / kg body weight, mixed with 1:100 feed experimentally, in commission and production experiments is a highly effective drug and is recommended for the treatment and prevention of in neoscaridosis of calves for group administration of with feed.

REFERENCES:

1. Zalikhanov M.Ch., Bittirov A.M., Begiev S.A. Modern biological threats and global regulation to ensure the biosafety of livestock products. In the collection: Breeding on modern populations of domestic dairy cattle as the basis for the import substitution of livestock products. Materials of the All-Russian scientific-practical conference with international participation. FSBI "Belgorod Federal Agricultural Research Center of the Russian Academy of Sciences". 2018; 245-253.
2. Bittirov A.M., Shakhbiev Kh.Kh. The results of epizootological studies and measures to combat the dominant helminthiasis of animals in the

- region of the North Caucasus. In the collection: Scientific support for the sustainable development of the agro-industrial complex in the North Caucasian Federal District. Collection of reports of the All-Russian scientific-practical conference with international participation. 2013; 592-595.
3. Bittirov A.M., Kabardiev S.Sh., Gazimagomedov M.G., Magomedov O.A., Abdulmagomedov S.Sh., Kabardiev Sh.S., Shakhmurzov M.M., Uyanaeva F.B., Bittirova A.A. Ecological and epizootic assessment of the fauna of bio- and geohelminths of sheep in the climatic zones of the North Caucasus. *Veterinary Medicine*. 2017; 9: 36-39.
 4. Shakhbiev Kh.Kh., Shakhbiev I. Kh., Bittirov A.M. Preparations "Triclafenal" in the treatment of fasciolosis in the North Caucasus and imported breeds of ruminants. In the collection: Science and Youth. All-Russian scientific and practical conf. students, young scientists and graduate students. 2016; 90-94.
 5. Kabardiev S.Sh., Bittirov A.M., Gazimagomedov MG, Magomedov O.A., Begiev S.Zh., Karpushchenko K.A., Bittirova A.A., Makhiev I.I., Kalabekov A.A. Complex antiparasitic composition "Azinal plus" - 3 for chemotherapy and prevention of trichuriasis, ankylostomaosis and echinococcosis of dogs. patent RUS 2614711 01.19. 2016.
 6. Shikhalieva, M.A., Atabieva, Zh.A., Kolodiy I.V., Bittirov A.M., Sarbasheva M.M., Bichieva M.M., Bittirov A.M. The structure of the parasitocenosis of the North Caucasus // *Veterinary Pathology*. 2012; 2 (40): 109-113.
 7. Kabardiev S.Sh., Bittirov A.M., Begieva S.A., Bittirova A.A., Magomedov O.A., Begiev S.Zh., Alieva Zh.R. A new anthelmintic drug for the treatment and prevention of osterohagiosis in cattle and small ruminants. patent for the invention RUS 2608132 26.
 8. Bittirov A.M., Kabardiev S.Sh., Begiev S.A., Karpuschenko K.A., Bittirova A.A., Begiev S.Zh., Abdulmagomedov S.Sh. Anthelmintic agent for the treatment and prevention of fasciolosis, dicrocoeliosis and paramptomatosis of cattle and small ruminants. patent for the invention RUS 2612013 01.25.201601.2016.
 9. Bittirov A.M., Kabardiev S.Sh. Veterinary and sanitary problems of regional pathology of fasciolosis of sheep and goats in the region of the North Caucasus and new methods for their elimination. Scientific-practical publication. Makhachkala: 2014.
 10. Bittirov A.M., Begiev S.Zh., Bittirova A.A., Kabardiev S.Sh., Eldarova L.Kh., Musaev Z.G. Embryotropic properties of the new composition of fenbendazole and albendazole (panaverm plus)// *Russian parasitological Journal*. 2015; 3: 86-88.
 11. Bittirov A.M., Kabardiev S.Sh., Begiev S.Zh., Bittirova A.A., Shakhbiev Kh.Kh. New complex treatment of chronic fasciolosis of domestic goats. Proceedings of the All-Russian correspondence scientific-practical conference. 2016; 111-116.
 12. Thakakhova A.A., Bittirova A.A., Berezhko V.K., Bittirov A.M. The species composition of helminthes and the contamination of sheep in the mountain tracts of Kabardino-Balkaria at an altitude of 1200-2500 m. Sea. Theory and practice of combating parasitic diseases. 2017; 18: 492-495.
 13. Uspensky A.V., Kabardiev S.Sh., Bittirov A.M. Problems of regional pathology and prevention of dangerous zoonoses in the region of the Central Caucasus. In the collection: Materials of scientific works of the Doctor of biological sciences, Prof. Bittirov A.M. "Theory and practice of innovative development of agrarian science". Dedicated to the 55th birthday. Caspian Zonal Vet. Research Institute. Makhachkala, 2014: 310-314.
 14. Bittirov A.M., Kabardiev S.Sh., Magomedov O.A., Musaev Z.G., Eldarova L.Kh., Shipshev B.M., Begiev S.Zh., Slonova E.S., Bittirova A.A. The effectiveness of new compositions based on albendazole in intestinal sheep nematodoses. Theory and practice of combating parasitic diseases. 2015; 16: 57-58.
 15. Bittirov A.M., Kabardiev S.Sh., Gazimagomedov M.G., Magomedov O.A., Begiev S.Zh., Karpuschenko K.A., Mutaev I.M., Kalabekov M.I., Bittirova A.A. Integrated method of treatment of cattle fasciolosis. patent for invention RUS 2584212 12/8/2014.

16. Bittirov A.M., Mantaeva S. Sh., Shakhbiev Kh.Kh, Shakhbiev I. Kh. et. al. Monitoring Of Intestinal Nematodoses Of Dogs In The Kabardino-Balkarian Republic And The Test Results Of The New Drug Avertfen Powder, Indo Am. J. P. Sci, 2019; 06(10); P S. Sh. Mantaeva,. 13883 – 13887.
17. Bittirov A.M., Mantaeva S. Sh. , Shakhbiev Kh.Kh, Shakhbiev I. Kh. et. al. The Of Results The Test New Multidisperse Anthelmintic Composition Prazinox At With Mono-And Of Mixtinvasion Of Echinococcosis And Multiceptosis Dogs, Indo Am. J. P. Sci, 2019; 06(10); P. 13888 – 13891.