


DETERMINATION OF FUNDAMENTAL PROPERTIES OF M-DWARFS FROM NEW MODEL ATMOSPHERES AND SPECTRA

Aishwarya Iyer¹, Michael Line¹, Philip Muirhead², Jonathan Fortney³, and Ehsan Gharib-Nezhad⁴
¹Arizona State University, School of Earth and Space Exploration, ²Boston University, Department of Astronomy, ³University of California, Santa Cruz
Department of Astronomy and Astrophysics, ⁴Nasa Postdoctoral Fellow, NASA Ames Research Center



Why M-dwarfs and Challenges Modeling them

- M-dwarfs have cooler temperatures (between 4000 - 2500 K) relative to sun-like stars, **facilitating molecule formation throughout their atmospheres** [1, 2], forming continuum opacities which are difficult to model using **standard stellar methods traditionally designed for FGK stars** [3].
- **Generational differences in line list databases cause inconsistencies in model spectra** [4].
- **Possible unknown physics in modeling M-dwarf stellar atmospheres.**

STRENGTHS OF OUR MODEL PIPELINE:

- 1) We produce a **self-consistent Radiative-Convective thermochemical equilibrium grid of models which permit physical plausibility + a Bayesian retrieval framework which will permit arbitrary species abundance determinations of M-dwarf atmospheres**, thereby stress-testing model assumptions.
- 2) Our model has the **most up-to-date opacities computed with the correlated-K method**, and are flexible to modify our model spectral resolutions for a variety of datasets.
- 3) We **leverage the broadband molecular absorption features** occurring in M-dwarf spectra over a wide wavelength range seamlessly to infer molecular abundances over the **near-infrared bandpass**—contrary to classical stellar abundance methods which focus on narrow spectral lines at higher resolutions.
- 4) We primarily **focus within the low-spectral resolution (R~120) regime**, with an advantage for neglecting non-LTE effects and microturbulence.
- 5) Our grid model retrieval component will incorporate an open source Bayesian Inference tool with an **error covariance matrix via a Gaussian Process scheme** [9], giving us **better control over model and data systematics and associated interpolation errors**, thereby **maximizing risk mitigation**.
- 6) **Our grid model pipeline will be open source** (on platforms like *Zenodo*) and allow for flexibility to include additional parameters of choice such as α/M , N/C ratio etc, in addition to providing a straightforward way of including opacity data by the user allowing for easy upgrades.

Future Work

- Benchmark our stellar model grid for M+G binaries [10, 11] with robust metallicity measurements.
- Incorporate Gaussian Process Scheme interpolator, accounting for the error covariance matrix for our fits.
- Perform Bayesian retrieval fits to estimate stellar properties and investigate model deficiencies.
- Incorporate cloud parameterizations and extend model grid to cover low-gravity cloudy M-dwarfs.

Acknowledgements:

This material is based upon work supported by the National Science Foundation under grant No. AST-2006555, SpeX Prism Library (and/or SpeX Prism Library Analysis Toolkit), maintained by Adam Burgasser at <http://www.browndwarfs.org/spexprism>, SIMBAD Catalog maintained by CDS Strasbourg and ASU Research Computing HPC and Storage resources.

Comparison with latest PHOENIX-ACES Models

We find that our models are not only consistent with PHOENIX-ACES [5], we also show improved fits using our grid-models owing to the latest opacity calculations. **Our approach leverages broadband molecular features (unlike traditional stellar abundance analysis approaches) to derive bulk stellar chemical properties.**




Figure 1: Comparison of fits and constraints between OUR GRID and PHOENIX-ACES [5] model grid. Here we use *Pymultinest* [6] along with a linear interpolation over an irregular grid to fit a low-resolution (R~120) SpeX Prism Library Spectrum [7] of an M7 dwarf VB-8 to **retrieve Teff (K), logg, log([M/H]), and Radius (sol)**, scaling with a radius-to-distance factor (R/D)² with distances acquired from GAIA [8].

Grid Model Fits to SpeX Prism Library Data




Figure 2: Preliminary Grid-model fits of M-dwarfs varying in class from M4.5—M7 from the SpeX Prism Library Database[7]. Using updated line-list opacities and with the self-consistent approach of modeling radiative-convective thermochemical equilibrium through our model grid, we are able to reasonably estimate the stellar parameters for these targets.

[1] Bochanski, John J., et al. 2010, *The Astronomical Journal* 139.6: 2679.

[2] Laughlin, Gregory et al. 1997, *The Astrophysical Journal Letters* 491.1: L51.

[3] Veyette, Mark J., et al. 2016, *The Astrophysical Journal* 828.2: 95.

[4] Gharib-Nezhad, E., Iyer, A., et al. (In revision *ApJS*, 2021)

[5] Husser, T. O., et al. 2013 *A&A*, 553, A6

[6] Buchner, Johannes. *ASCL*(2016): ascl-1606.

[7] Burgasser, Adam J. 2014, *arXiv preprint arXiv:1406.4887*.

[8] Schönrich, Ralph, et al. 2019, *MNRAS*. 487.3: 3568-3580.

[9] Czekala, Ian, et al. 2015, *The Astrophysical Journal* 812.2:128.

[10] Mann et al. 2013, *Astronomical Journal*, vol 145, 2, 52:15pp

[11] Mann et al. 2014, *Astronomical Journal*, vol 147, 6, 160:11pp