

8 Theory – COMPLETE OVERVIEW

Manor O – 29.3.2021

Fermions, Manifolds and Arbitrary Variations

Acceleration away from Extremum Curvatures	3
Fermions as Variations of a Lorentz Manifold	4

Bosons, Primes and the Coupling Constants Series

Deriving the Coupling Constants Series	5
Majestic (3) Possible meanings	9
Majestic (3) Spin (1/2) and Prime Critical Line (1/2)	11
Majestic (3) as an Electron	14
Weak Interaction Left Orientation and Spin	17
Duality of the first Three Interactions at 26	20
Proof of the Pauli Exclusion Principle	22
The Hidden Series of Masses and Generations	25

Cosmological Predictions

The Rise of the Arrow of Time	30
The Almost Homogenous Universe	32

Universe Future & Gauge Fields & Higgs & Antimatter & PW Duality

The Future of Universe - Expansion or Collapse?	34
Coupling Constant Equation and Gauge fields Prediction	37
Quark Masses mixing & mixing angles	39
The Majestic (3) and the Higgs Field	43
Anti-matter & Dirac Delta Variation	45
Spiral galaxies & Principle of Least Curvature	50
PW Duality	54
The Feynman Variation on Lorentz Manifolds	57

Gravity & Fermion Rings & Time

Gravity as a combination of couplings	62
Predicting the Graviton Mass	64
Fermions are Imperfect Rings	65

The Primorial and The Arrow	68
Universe Packets	69
Flatness	70
Quark confinement	72
Wrapped dimensions	74
Elevating Einstein Equivalence to Perfection	75
Majestic (3) Is the Electron	76
Proof: Yang Milles Conjecture	78
<u>Golden Shades of a Final Theory</u>	
8T and QFT Axioms	80
The Three Critical Theorems	82
Refuting Magnetic Monopoles	83
Most Symmetrical Interaction is the Weak	84
Hermitian Conjugation and Primes	85
Final Shot at Quantum Relativity	86
Total Variations Pairing	88
Boson & Fermion Propagations	89
The Lagrangian Variation	90
Cluster Decomposition	91
Hadronic Symmetries & Densities	92
Feynman Diagrams Variation	94
The Axis of Evil	96
Bosonic Probabilities	97
Conservation of Variation	98
Bosonic Strings & Curvature Absorptions	99
Proof: The Riemann Hypothesis	102
Visualizations	106

Fermions, Manifolds and Arbitrary Variations

Define a Lorentz manifold, which is the connected manifold with (3,1) signature:

$$s = (M, g)$$

Invoke it to be stationary by Euler Lagrange operator, $S = S_0 \times \mathbb{R}$:

$$\mathcal{L} = (s, s', t)$$

$$\frac{\partial \mathcal{L}}{\partial s} - \left(\frac{d}{dt} \right) \frac{\partial \mathcal{L}}{\partial s'} = 0$$

Develop the last equation:

$$\frac{\partial \mathcal{L}}{\partial s} \frac{\partial s}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} - \frac{\partial \mathcal{L}}{\partial s'} \frac{\partial s'}{\partial M} \frac{\partial M}{\partial g'} \frac{\partial^2 g'}{\partial t^2} = 0 \quad (1)$$

If the Lorentz manifold to be stationary and no data is attainable from the first three terms, we can require the manifold to those two conditions:

$$\frac{\partial g}{\partial t} = 0, \quad -\frac{\partial^2 g'}{\partial t^2} = 0$$

If these two are hold to be true, we have areas of extremum curvature on the manifold and negative time invariant acceleration. The demand of extrinum curvature to stay as they are overtime means the acceleration cannot affect them – if so, directed away from them. This in agreement with what we speculate as "dark energy". Notice that M is now the matric tensor, g is the Ricci flow. That is the result of parametrizing the manifold to "s" variable and inserting it to EL operator, yielding agreement with Einstein principle of equivalence.

$$\frac{\partial \mathcal{L}}{\partial s} \frac{\partial s}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} \delta g - \frac{\partial \mathcal{L}}{\partial s'} \frac{\partial s'}{\partial M} \frac{\partial M}{\partial g'} \frac{\partial^2 g'}{\partial t^2} \delta g' = 0$$

Equation reads, length to manifold, manifold to matric, matric to flow, flow to time. δg as amount of arbitrary variations, which by demands of stationarity we require to vanish. Discretizing and partitioning the term δg into a series of sub elements, we can derive the existence of fermions, i.e. showing that it must have an even amount of elements, which differ in sign and create nine threefold combination, and no more than two distinct elements.

$$\delta g_1 + \delta g_2 \dots = \sum_{i=1}^N \delta g_i$$

$$\sum_{i=1}^N \delta g_i = 0$$

Given four elements distinct:

$$\delta g_1 + \delta g_2 > 0$$

$$\delta g_3 + \delta g_4 < 0$$

If

$$\delta g_1 + \delta g_2 + \delta g_3 + \delta g_4 \neq 0$$

Then the overall series cannot vanish, by that logic we need even amounts of equal elements of pluses and minuses. The amount must be even and summed as zero, ensuring stationary Lorentz manifold. Suppose that we had three distinct elements, two pluses and minus:

$$\delta g_1 + \delta g_2 + \delta g_3 > 0$$

or

$$\delta g_1 + \delta g_2 + \delta g_3 < 0$$

Demanding the series to vanish this will exclude this result, and so there could not be three distinct elements in the series, else the overall series will not vanish to zero. As a result of those sceneries, we require the series to have an even amount of variation elements, manifesting as two distinct elements in the series, which differ in sign. If we allow those sub elements in the series to vary as well, and by the above reasoning, there are only two elements in the series, they are varying in a discrete way, or forming a group. Let it be only four elements in the series and one of the pluses just changed its nature

$$O: \delta g_1 \rightarrow \delta g_2$$

$$\delta g_1 + \delta g_1 + \delta g_2 + \delta g_2 = 0$$

To:

$$\delta g_1 + \delta g_2 + \delta g_2 + \delta g_2 = 0$$

There must be a way to bring it back to where it was, so the overall series can vanish, it takes another map, on the varying element to bring it back to where it was.

$$Y: \delta g_2 \rightarrow \delta g_1$$

Therefore, to bring an element to itself given only two varying elements in the series we need two distinct maps, which attach a varying element to itself, by a threefold combination. $\delta_{g_1}(O)\delta_{g_2}(Y)\delta_{g_1}$ For example. Even though the sub elements in the series are varying, the overall series can vanish.

Now, count all the ways of possible combinations of those elements. We are going to analyze by the integral signs. Since it is a group, there is a natural map, which change an element to itself. One built his analysis firstly on those natural maps.

So:

(1(e)1(e)1)

2(e)2(e)2

(221)

(112)


(211)

(122)

(212)

(121)

The first two combinations are by the natural maps and one used them to build the other combinations. Overall, there are eight such combinations and additional one arrow combination, which yield (333). Here is how one built it, starting from those two natural maps. (Arrows to variations, colors to pairings):



Therefore, we have Lorenzian manifold with arbitrary variations, which vanish into matter based on that idea. One does not know whether these are the actual variations, as the mathematics does not entail any details about that. Therefore, the graph could be inaccurate in elements order. The colors meant to elements pairing. Reader does not have to agree with what one did, but as one will calculate the ratios of all the forces known, one kindly asks the reader to keep reading as some truth seem to obey the reasoning one is suggesting.

Deriving the Coupling Constants Equation

Theorem (1) – nature will not allow a prime amount of variation to appear by itself. Define prime to be $(2n+1)$ variations.

1.1) Prime amounts appear in pairs.

Theorem (2): Nature will generate force if a **prime net amount of arbitrary variation will appear**. Net variations will appear when combine two amounts of prime variations. Two does not appear, as it is an even amount of variations, which vanish.

Define N_V as the series of prime net variations and the number one.

$$N_V = 2V + 1 \quad V \geq 0$$

Count all the prime pairs of variations,

$$\begin{aligned} &(3,3) \ (3,5) \ (3,7) \ (3,11), (3,13) \dots \\ &(5,3) \ (5,5) \ (5,7) \ (5,11) \ (5,13) \dots \\ &(7,3) \ (7,5) \ (7,7) \ (7,11) \ (7,13) \dots \\ &\dots \\ &(29,19)(29,23), (29,29), (29,31) \dots \end{aligned}$$

That is a tedious work, but here is the great part. **We only need to do it twice** to find what nature does repeatedly.

Since we have only two varying elements in the series, we can eliminate almost all the options, as we require obtaining **a sum that is divisible by two and after yields a number divisible by three**. By The following reasoning: Two as we have only two varying elements. Three as these elements create a certain amount of threefold combinations.

The sums satisfying the condition is (5,13) or (7,11) and (29,31).

Of course, there are more as N_V has no limit, but as one mentioned, it took two pairs to understand the principle:

Theorem (3) :

Each prime pair should have a net variation element N_V proportional to Total Variations value divided by two.

This will be vivid with actual examples:

Analyze the (7, 11) total variations pair with $N_V = (+1)$:

Total variations sum is divisible by two:

$$18/2 = 9$$

And then by three

$$9/3 = 3$$

We know that we have $N_V = (+1)$ so it can be extracted to yield:

$$F_1 = 8 + 1$$

However, even amounts of variations vanish so we can ignore the element 8 and write:

$$F_1 = 1$$

Analyze the next pair of total variations (29, 31) with $N_V = (+3)$

$$29 + 31 = 60$$

$$60/2 = 30$$

In addition, three devisible. We know we have three net variations so extract:

$$27 + 3$$

Now that is all you need to complete the series and calculate the **next element**:

Notice:

$$\begin{aligned} 27 &= 24 + (3) \\ (8 * 3) &= 24 \end{aligned}$$

Obtain the ratio:

$$[8 + 1]: [27 + 3] = [8 + 1]: [24 + (3)] + 3$$

$$[8 + 1]: [27 + 3] = [8 + 1]: [(8 * 3) + (3)] + 3$$

Next element $V = 2$ and $N_V = +5$ **so if the overall idea to be correct** we take this element, multiply by the even sum of the previous element in the series, **add extra invariant (3)**, and we know we need add to the sum the extracted N_V .

$$[(24 * 5) + (3)] + 5 = 128.$$

Next in line:

$$\begin{aligned} [(120 * 7) + (3)] + 7 &= 850 \\ [(840 * 11) + (3)] + 11 &= 9254 \end{aligned}$$

Nature is than the interplay between averages of total arbitrary variations pairs to net variations/curvature. To calculate the magnitude of an element R:

$$F_{V=0} = 8 + (1) \tag{1.1}$$

$$F_R \# = \left(8 * \prod_{V=1}^{V=R} N_V + (3) \right) + N_V = 30:128:850:9254.. \tag{1.2}$$

$$N_V = 2 \left(V + \frac{1}{2} \right); \quad V \geq 0$$

$$N_V \in \mathbb{P} \cup (+1); \quad \mathbb{P} \rightarrow \text{Set of Primes}$$

$$N_V = P_{max} \in [0, \mathbb{R}] \cup (+1); \quad P_{max} \in \mathbb{P}$$

$$\mathcal{P}_0 = 8 + (1)$$

$$\mathcal{P}_N \# = \left(2^{\mathcal{M}} * \prod_{V=1}^{V=N} \mathcal{P}_V + (\mathcal{M}) \right) + \mathcal{P}_V = 30:128:850:9254.. \quad (1.2.A)$$

Equation (1.2A) is another way of representation. \mathcal{M} As the first letter of the word 'Majestic'.
 $\#$ Sign meant for classification as a primorial function. " \mathcal{M} " as possible magnitudes. Notice the strong symmetry pattern of this equation.

Overview of reasoning:

Axiom – prime amount of arbitrary variations pair to each other

Their overall sum must be dividable by two and three

Two distinct elements, which create threefold combinations

Define generated force as prime net variation in which we associate N_V element

$\frac{\text{total variations}}{2} \propto$ to N_V element by the relative size of total pairing

Net variation function cannot contain an even, as it will vanish

We searched for the first two prime pairs and derived $8 + (1)$ and $27 + (3)$

We saw that nature multiply the even sum by the next element of N_V

We found the invariant three element.

We obtained a number to which we add the extracted net variation

We calculated the next element to be exactly 128 and the two next interactions:

$$8 + (1): (24 + (3)) + 3: (120 + (3)) + 5: (840 + (3)) + 7 \dots$$

$$(1): (30): (128): (850): (9254) \dots$$

Predictions and Conclusions

There is an infinite number of bosonic fields, which are Lorentz manifold net curvature.

The clusters of total variations grow much more rapidly than the net variations.

The larger the cluster, the weaker the interaction.

The magnitude of interactions is manifested an in infinite series of ratios

1: 30: 128: 850: 9254...by the expression:

$$F_{V=0} = 8 + (1) \quad (1.1)$$

$$F_R \# = \left(8 * \prod_{V=1}^{V=R} N_V + (3) \right) + N_V = 30: 128: 850: 9254.. \quad (1.2)$$

Possible meanings of the Majestic (3)

Option 1

The **invariant three as a cause**. Notice that all the element within the closed term (8*..) Are two and three divisible to vanish into matter. The invariant three prevents it completely and then as a result, a net variation will appear. The net variation is proportional to the right element in the bracket(8 * 3) \propto 3 and (24 * 5) \propto 5 .

Option 2

The **invariant three as a result**-There are perfect clusters of variations such (8 * 3), (24 * 5) , which experience additional net variation causing them to destabilize. The result is manifested in the invariant three. The additional variation could affect them could be external. Less likeable option. It is less likeable as we can them create mixtures (8*3) to destabilize by five net variations, and yield invariant three and all the beauty in which we attained than will be lost.

Option 3

The **Invariant three and net variation as duals**-both appear at the same time and they are related to each other by more fundamental relation, which is not attainable nor explainable. Even though we found a jewel, many questions still stand unanswered.

Why the invariant three appear as it is and do not change?

Of course that the real answer to that question is that one does not know. However, one can guess and say that three is the smallest prime. If we assume that nature is Lagrangian oriented, it might be the minimal way to destabilize the cluster of potential matter. Why add thirty seven additional variations when only three is needed? It's a logical argument not a proof, and therefore rightfully argued by reader. One was trying to argue that three is a Prime minima, that is the reason for its invariant in the series. Remember that even variations vanish, so two is not an option.

Correlating the Majestic (3) To Spin (1/2) and Matter

In the paper about primes, we have shown that they create a non-abelian group with $1/2$ as generator, by using the anti-commutation relation and vanishing of even amounts of variation. It recently become evident to one that we can represent each element in the series in the following way:

$$[(8 * 3) + (3)] \rightarrow \left[2N_1 + \frac{1}{2}\right]$$

$$[(24 * 5) + (3)] \rightarrow \left[2N_2 + \frac{1}{2}\right]$$

$$[(120 * 7) + (3)] \rightarrow \left[2N_3 + \frac{1}{2}\right]$$

Since three is a prime, and aligned on the prime ring located on critical line of $\frac{1}{2}$. The sums alongside of it are even sums such as 8, 24, 120 and so on. These expressions are interesting, as one believes they represent the notion of matter or fermions. Notice that we omitted the additional net variation, which is also prime. Meaning it is also on the Prime Ring Located on $\frac{1}{2}$. Overall:

$$[(8 * 3) + (3)] + 3 \rightarrow \left[2N_1 + \frac{1}{2}\right] + \frac{1}{2}$$

$$[(24 * 5) + (3)] + 5 \rightarrow \left[2N_2 + \frac{1}{2}\right] + \frac{1}{2}$$

$$[(120 * 7) + (3)] + 7 \rightarrow \left[2N_3 + \frac{1}{2}\right] + \frac{1}{2}$$

So the construction within the parenthesis is prime but the overall additional net is changing it, and making it: $(1/2 + 1/2) = 1$. So the overall 1: 30: 128 will have to do with certain elements that have element one. We already knows these are bosons, as we found the coupling constants series. If so, than the rest of the terms are fermions, as only $(1/2)$ is there.

So it is the Majestic three→in this paper: one half element to destabilize perfect clusters of variations and causing a net variation to appear. Notice that one chose the first option in regards to the meaning of the invariant three, as we had in part two three ideas to it possible meaning. We have proved that the Majestic three is Spin. We also proved, that bosons will propagate within variation clusters destabilized by one half, or matter. These are non-trivial statements. We only use one equation, not experiment nor inherited knowledge. **Using that framework, we can see why bosons will propagate within fermions.** Since Its invariant, all matter must have the same spin one-half

So $(2N)$ are variation clusters, the majestic three is really a destabilizing factor which is spin one half yielding matter. Because of that process, a boson will propagate from within the fermion. The nature of the boson is correlated to right element of the term: $(8 * 3) \rightarrow 3$ (weak particle), $(24 * 5) \rightarrow 5$ or a photon, so on.

Majestic (3) as the Electron

$$F_{V=0} = 8 + (1) \quad (1.1)$$

$$F_R \# = \left(8 * \prod_{V=1}^{V=R} N_V + (3) \right) + N_V = 30:128:850:9254.. \quad (1.2)$$

$$[(8 * 3) + (3)] + 3$$

$$[(24 * 5) + (3)] + 5$$

$$[(120 * 7) + (3)] + 7$$

.....

$$[(8 * 3) + (3)] \rightarrow \left[2N_1 + \frac{1}{2} \right]$$

$$[(24 * 5) + (3)] \rightarrow \left[2N_2 + \frac{1}{2} \right]$$

$$[(120 * 7) + (3)] \rightarrow \left[2N_3 + \frac{1}{2} \right]$$

$$[(8 * 3) + (3)] + 3 \rightarrow \left[2N_1 + \frac{1}{2} \right] + \frac{1}{2}$$

$$[(24 * 5) + (3)] + 5 \rightarrow \left[2N_2 + \frac{1}{2} \right] + \frac{1}{2}$$

$$[(120 * 7) + (3)] + 7 \rightarrow \left[2N_3 + \frac{1}{2} \right] + \frac{1}{2}$$

In previous paper, (part three) we called the (1/2) an element to destabilize Perfect clusters of variations and causing a net variation to appear. In this part, we can **call it the Electron**. Later in the thesis, we will prove it by putting inside the equation of the fine structure constant.

$$[(24 * 5) + (3)] + 5 \rightarrow \left[2N_2 + \frac{1}{2} \right] + \frac{1}{2}$$

$(2N \text{ variations}) \rightarrow \text{Spin } 0$

$(2N \text{ variations} + 3) \rightarrow \text{Matter with spin } (\frac{1}{2})$

$(2N \text{ variations} + 3) + N_V \rightarrow \text{Bosons with spin } (1)$

$(2N \text{ variations} + 3) + N_{V1} + N_{V2} + \dots \rightarrow \text{Bosons with higher spin integers}$

When we first discover the coupling constant equation, we only saw the analytical aspect, by and the ratio between the total variations to net variations. However, by setting the equation on the geometrical realm and examining the critical line of the primes, we can get a deeper insight to what is going on. We are able to analyze the trait of spin, we can understand why bosons have spin one and the Invariant three or spin one half Therefore, it is the electron, which causes the boson propagation from clusters of potential matter.

Sure, we knew that, but we did not have the mathematical equation to describe it. The coupling constants equation has than another powerful use; it describes what it going on in elementary level, not just the magnitude of the interactions. It was only available to us when we examined the geometrical realm.

Please notice that the electron is inside potential cluster $\left[2N_2 + \frac{1}{2} \right]$ so we would not be able to know where it is within the cluster, it blends in $[120+3] = 123$.

$$[(24 * 5) + (3)] + 5 \rightarrow [(24 * 5) + (e)] + \gamma$$

The Complete Picture:

Perfect clusters of variations $\rightarrow 2N$

Destabilize the perfect $2N$ is the majestic $(3) \rightarrow \left(\frac{1}{2}\right) \rightarrow$ electron. Blends in the potential cluster to yield in that case $\rightarrow 123$.

The result is the net variation, also Prime $N_v \rightarrow \left(\frac{1}{2}\right) = +(5)$

The overall result yields $\left[2N + \left(\frac{1}{2}\right)\right] + \left(\frac{1}{2}\right)$ probability to an emission of a boson.

$$123 + 5 = 128.$$

We have taken the third element in the series, as we are familiar with the nature of the electrons due to the great minds of the past century, but the following result would apply to each element in the series from the second and above.

$$[(24 * 5) + (3)] + 5 \rightarrow [(24 * 5) + (e)] + \gamma$$

Weak Interaction Negative Left orientation

$$F_{V=0} = 8 + (1) \quad (1.1)$$

$$F_R \# = \left(8 * \prod_{V=1}^{V=R} N_V + (3) \right) + N_V = 30:128:850:9254.. \quad (1.2)$$

$$8 + (1)$$

$$[(8 * 3) + (3)] + 3$$

$$[(24 * 5) + (3)] + 5$$

$$[(120 * 7) + (3)] + 7$$

....

$$[(8 * 3) + (3)] \rightarrow \left[2N_1 + \frac{1}{2} \right]$$

$$[(24 * 5) + (3)] \rightarrow \left[2N_2 + \frac{1}{2} \right]$$

$$[(120 * 7) + (3)] \rightarrow \left[2N_3 + \frac{1}{2} \right]$$

$$[(8 * 3) + (3)] + 3 \rightarrow \left[2N_1 + \frac{1}{2} \right] + \frac{1}{2}$$

$$[(24 * 5) + (3)] + 5 \rightarrow \left[2N_2 + \frac{1}{2} \right] + \frac{1}{2}$$

$$[(120 * 7) + (3)] + 7 \rightarrow \left[2N_3 + \frac{1}{2} \right] + \frac{1}{2}$$

Notice that each term in the series within the parenthesis is prime $\rightarrow 123, 843, 9243 \dots$ as one did not calculate the entire series he is going to assume that is would be true concerning each higher element in the series. We are leaving out the net variation in this part.


Notice that the only term which is not a prime after added the Majestic three or spin one half is the second element in the series, in which we associate with the weak interaction.

$$[(8 * 3) + (3)] = 27$$

As the series is increasing and each term inside the parenthesis is creating a higher prime than the previous element, in order of weak interaction to be of the same nature of the rest of the forces, we would need that the sum of the parenthesis to be a prime, we look for the closest higher prime:

$$[(8 * 3) + (3)] \rightarrow 29$$

So in order to be like the rest of the forces. Meaning to have a prime inside a parenthesis, it lacks a certain amount of variation. If we associate each interaction to be invariant to direction – and the Cause of such a trait could be the prime term inside the parenthesis, than the weak interaction would differ by its nature.



The fact that the term inside the parenthesis is not on the critical line of the primes, but left to it, can explain why the weak interaction is left oriented and differ by its nature by the rest in terms of its spin. We have proved that the majestic three is really a different representation of spin, which destabilizes clusters of perfect variations causing the N_V to appear, which overall yield a propagation of a Boson from the fermion, and therefore gives us the beautiful series of coupling constants. If all the Terms on the critical line of primes are yielding interactions that are invariant to direction, than one could predict the weak interaction to be spin oriented to the left by the ratio (2), since the strong interaction is also not on the critical line, such orientation could exist in its regards as well.

$$27 - 29 = -2$$

$$\left(\frac{1}{2} - 2\right) = -\frac{3}{2}$$

Mathematical Duality of Forces-Virtual Variations

We will take the equation built and first three developments:

$$8 + (1): [(8 * 3) + (3)] + 3: [(24 * 5) + (3)] + 5$$

The idea: we will allow the net variations to vary, and when they have the same value, than the expressions inside the parentheses will become scalar multiple. This will be done by using the idea of virtual variations:

$$[(24 * 5) + (3)] + 5 \rightarrow [(24 * 5) + (3)] + 3$$

Notice that now the third is a scalar multiple of the second by a factor of five:

$$[(24 * 5) + (3)] + 3$$

$$[(8 * 3) + (3)] + 3$$

Therefore, the weak and the electric are differing now by a scalar. That is simply beautiful. However, the strong force just accepted that extra two variations so it is just become:

$$8 + (1) + 2 \rightarrow 8 + (1).$$

As Even amounts of variations vanish. It does not affect it. We can try something more interesting, and that is the real purpose of the part:

$$[(24 * 5) + (3)] + 5 \rightarrow [(24 * 5) + (3)] + 2$$

$$8 + (1) + 3$$

Now this will ruin the duality and the series, the weak and the electric are not isomorphic, and the strong just got a prime amount of variations that cannot vanish. To solve that we can define a virtual exchange of variation $\rightarrow (1\nu)$.

$$[8 + (1)] + 3 - (\mathbf{1\nu}): [(24 * 5) + (3)] + \mathbf{3}$$

The real variations are (+3) but to ensure the nature of the strong force, there is a virtual exchange of one variation, marked in bold. For a very short time period, the strong is now a scalar multiple of the other two. Overall, they have the same prime amount of net variations – will mean they are at equivalence relation. For the first three forces:

$$N_V = +(3).$$

$$[8 + (1)] + 3 - (\mathbf{1\nu}): [(8 * 3) + (3)] + 3 : [(24 * 5) + (3)] + \mathbf{3}$$

We can say that there are three real exchanges and one virtual, so overall four exchanges, which causes all the forces to align on the $N_V = +(3)$. Taking the average of the Sum: $4/2 = 2$ *net*.

The converging value of the those exchanges will modify the middle element:

$$[(8 * 3) + (3)] + 3.$$

Since we want to keep the prime net variation as it is, to ensure duality, and we can't touch the invariant three, we add this (+2), the first term:

$$((8 * 3) + 2) = 26.$$

The point where they three aligned will be $24 + 2$ variations. certain agreement with this Number exist.

Proof: The Pauli Exclusion Principle

$$F_{V=0} = 8 + (1) \tag{1.1}$$

$$F_R \# = \left(8 * \prod_{V=1}^{V=R} N_V + (3) \right) + N_V = 30:128:850:9254.. \tag{1.2}$$

$$8 + (1)$$

$$[(8 * 3) + (3)] + 3$$

$$[(24 * 5) + (3)] + 5$$

$$[(120 * 7) + (3)] + 7$$

$$[(8 * 3) + (3)] + 3 \rightarrow \left[2N_1 + \frac{1}{2} \right] + \frac{1}{2}$$

$$[(24 * 5) + (3)] + 5 \rightarrow \left[2N_2 + \frac{1}{2} \right] + \frac{1}{2}$$

$$[(120 * 7) + (3)] + 7 \rightarrow \left[2N_3 + \frac{1}{2} \right] + \frac{1}{2}$$

We have seen that we can change the term outside the parenthesis, and so we can reach duality between the forces. When we did it in the first three terms, we saw that their duality is exactly on 24+2 variations, which is in agreement with what we know in other theories of GUT. We briefly mention in that paper, that we cannot touch the invariant three. This will be the subject of this part. If we for example combine:

$$[(24 * 5) + (3)] + 5 \pm \text{INTEGER} \dots$$

We can switch and change the terms outside the parenthesis, as those are net variations and they do not seem to obey to any strict rules. However, we could not touch the invariant three and now we will examine deeply the reason.

$$[(24 * 5) + (3) + (3)] + 5 = [(24 * 5) + \textit{Even}] + 5$$

$$\textit{Even} = 0$$

$$[(24 * 5) + 0] + 5 \rightarrow \text{Impossible}$$

As even amount of variations vanish. Recall that the invariant three is the cause; It is the destabilizing factor yielding a net variation. In the case of the third element, it is the Electron. So using that framework, we can see why we cannot combine two electrons or invariant three elements together. The term then becomes meaningless, a photon cannot propagate from nowhere and the coupling constant series does not makes sense anymore. So the invariant three cannot be combined, it will repel each other. The net variation however can be changed and switched, which makes the flexibility and duality of the forces. The equation is with complete agreement with our understanding; we are just examining additional meaning of it. It allows us to examine it from a deeper, more profound view. Now we can understand why fermions do not commute – because even variations vanish and so bosons will not be propagated.

If we eliminate the electron, than no boson will be propagate at all. However, consider the following:

$$[(24 * 5) + (3)] + 5 + [(24 * 5) + (3)] + 5 + .. =$$

$$[(24 * 5) + (3)] + 7 + [(24 * 5) + (3)] + 3 + .. =$$

While we cannot touch the terms inside the parenthesis, we can change and combine the net variation, there seems to be no limitation in regards to that operation, we have done it before, and showed that the forces can be scalar multiples. We can cluster the net variations, which means that many electrons can emit net variations together, That is bosons, which agrees to what we know as laser, or what we know as bosons commutation relation in QFT. However, using the 8-theory framework we can get a new and fresh insight on why those things are the way they are using the coupling constant equation. As we mentioned in part four of the paper series on coupling constants, the invariant three blends in the total cluster of the fermions, so we cannot know where he is. That is in agreement with the Heisenberg principle of uncertainty.

Strikingly Beautiful Relation of Three Generations Masses

The idea, which is followed by the last paper, is that if $8 + (1)$ to generate force, and force is extended outward, (short or long ranged) than $8 - (1)$ would be to generate mass, or arbitrary **variations converging inward**. Equipped with this idea we can search for a mathematical pattern. First, take all the masses, accurate as they can and combine them according to generation:

$$[1.9] \qquad [1320] \qquad [172,770]$$

$$[4.4] \qquad [87] \qquad [4240]$$

$$1.9 + 4.4 \approx 6\frac{1}{3}$$

$$1320 + 87 = 1407$$

$$172,770 + 4240 = 177010$$

Seemingly nothing in common, luckily we can change it. Soon one will reason why the following exactly, multiple equation one by factor of nine and divide the third family by a factor of nine.

$$6\frac{1}{3} * 9 = 57 = 50 + 7$$

$$1320 + 87 = 1407 = 1400 + 7$$

$$\frac{177010}{9} = 19,667 = 19,660 + 7$$

Also, notice that

$$50 * 28 = 1400$$

$$1400 * 14 = 19,600$$

$$(60 \text{ Mev Difference} - 0.03\%)$$

but

$$28 = 7 * 4$$

$$14 = 7 * 2$$

so to go from first to second:

$$(7 * 4) * 50 + (7)$$

And from second to third

$$(7 * 2) * 1400 + (7)$$

Notice that it is a decreasing by an even factor of two. In addition, if we go from low to high it does not make sense physically, it should be Lagrangian oriented, nature is devising by increasing amount to minimize the arbitrary variations, so if correct we should go from three to one by devising:

$$\frac{19,660 + (7)}{7 * 2} = 1400 + (7)$$

$$\frac{1400 + (7)}{7 * 4} = 50 + (7) * \frac{1}{9}$$

Next, we can predict that **total mass** for fourth to sixth families:

$$\begin{aligned} \frac{50 + (7)}{7 * 8} * \frac{1}{9} &= 0.113 \text{ mev} \\ \frac{0.113}{7 * 16 * 9} &= 0.000113 \text{ mev} \quad \text{or} \quad \frac{0.113}{7 * 16} = 0.00100 \text{ mev} \\ \frac{0.000113}{7 * 32 * 9} &= 5.95 * 10^{-8} \text{ mev} \quad \text{or} \quad \frac{0.00100}{7 * 32} = 0.0000045 \text{ mev} \end{aligned}$$

Summing 4-6 families: 0.113113 or 0.1140 Mev. We can see a converging to the value of the forth which is 55.25-55.69 lighter than first family:

$$\frac{6.3}{0.1131130595} = 55.696 \quad \text{or} \quad \frac{6.3}{0.1140} = 55.26$$

Note that we needed to readjust the scale by the factor of $8 + (1)$ as we manipulated the data, in a search for a pattern. Adjust it in the third family, by Multiplication and in the first and by division.

The following reason, T-B family has much more mass, thus much more arbitrary variation converging inward, that might be the reason it has $8 + (1)$ factor in the nominator, and in the first, the arbitrary variations are so small, we need to adjust it in the opposite direction, to increase by $8 + (1)$. Whether in the fifth family and below, additional rescales are needed is unknown, we do include two options, with the $8 + (1)$ or without it.

So according to the above reasoning and mathematical notion, one will predict infinite family is forming below the masses of the U-D masses, converging to total value of ≈ 0.113113 Mev as family's below the six are neglected due to little contribution the total sum. So overall, we can write:

$$M_{N+1} = \frac{M_N + (7)}{7 * \prod_{E=1}^r N_E} * \frac{1}{9} \quad (1.3)$$

$$M_{N+1} = \frac{M_N + (7)}{7 * \prod_{E=1}^r N_E} \quad (1.31)$$

$$N_E = 2E ; E \geq 1$$

is to a function for two multiple of variations. N_E

Overview of ideas

Mass is a variation of the manifold converging inward. Just like force but opposite in direction. Nature is eliminating the arbitrary amount of variations by devising in increasing amounts. That prediction is the rule of dark matter in our theory. It suits the fact that very quickly the families total is converging to zero. The rate in which the conserving to zero is made is unknown.

The theory provides two options. First, with the rescaling factor to each family and second option without it. Rescaling only Once. Both options agree on the value of the total mass of the fourth, which is about 56 Times lighter than first.

$$M_{N+1} = \frac{M_N + (7)}{7 * \prod_{E=1}^r N_E} * \frac{1}{9}; \quad M_{N+1} = \frac{M_N + (7)}{7 * \prod_{E=1}^r N_E}$$

As we combined the net masses of the two elements, the value should be again, decomposed to the two separate elements. There are an infinite variety of families whose mass is decreasing, thus below first generation of quarks, this could agree with so-called, dark matter. Cosmologists to decide whether the mass values predicted agree with the data.

The Rise of the Arrow of Time

$$\frac{\partial \mathcal{L}}{\partial s} \frac{\partial s}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} - \frac{\partial \mathcal{L}}{\partial s'} \frac{\partial s'}{\partial M} \frac{\partial M}{\partial g'} \frac{\partial^2 g'}{\partial t^2} = 0 \quad (1)$$

$$F_{V=0} = 8 + (1) \quad (1.1)$$

$$F_R \# = \left(8 * \prod_{V=1}^{V=R} N_V + (3) \right) + N_V = 30:128:850:9254.. \quad (1.2)$$

In our framework we have a Lorentz manifold inside an Euler- Lagrange equation. The manifold experience arbitrary variations, which vanish into, matter, we proved it in previous papers. Each time net variation appear on the manifold, a boson is manifested into our matric. That was the idea, which derived the coupling constants equation. Net variations are prime, and for each prime, there is a boson, unique boson:

$$8 + (1)$$

$$[(8 * 3) + (3)] + 3$$

$$[(24 * 5) + (3)] + 5$$

$$[(120 * 7) + (3)] + 7$$

However, how does that relate to the arrow of time? Recall that the coupling constant equation is really a built upon a ratio between total variations divided by two and net variations which are prime. We saw that the total variations grew much more rapidly than the net, and we required a Sequence, that it will go from low to high. Therefore, the arrow of time should go from low to high as well. There could not be a photon propagation without electron, which propagate from the nuclei, or cluster of so-called quarks. The sequence of The coupling constant equation is the sequence of time it allows us to build from the elementary to the massive, first arbitrary variations eliminate and vary themselves, create protons and neutrons which vary as well, propagate electrons, which vary as well, yielding photons and electromagnetism.

Nature as the interplay of total variations to net variations, which grow in number and gets weaker from one element to another, explain why the forces at a large scale are much weaker than those at smaller scale, here are much more total variations and the net is divided across the whole cluster. So starts and galaxies must appear only after the strong, weak and electromagnetic.

Nature is going from high to low, from small amount or strong variations to weak amounts of net variations over bigger clusters of total variations. Keep in mind that when one say **variation he means curvature** as we built the 8- theory upon a Lorentz manifold. However, if we look at each element in itself, like electromagnetism for example we will not see any clues for the arrow of time, as it's not telling anything about the arrow. It is only when we found the series of Coupling constants and the intimate relation of the boson to primes and and put them in a row, than and only than we can see the rise of the arrow of time. In other words, we can reason why galaxies and cluster of galaxies can form only after the strong, weak and the electric. We are also able to reason the weakness of gravity and the interactions in higher terms in the series.

$$1 > \frac{1}{30} > \frac{1}{128} > \frac{1}{850} \dots$$

$$\xrightarrow{\hspace{1.5cm}}$$

$$1 > \frac{1}{30} > \frac{1}{128} > \frac{1}{850} \dots$$

The Almost homogenous Universe

$$\frac{\partial \mathcal{L}}{\partial s} \frac{\partial s}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} - \frac{\partial \mathcal{L}}{\partial s'} \frac{\partial s'}{\partial M} \frac{\partial M}{\partial g'} \frac{\partial^2 g'}{\partial t^2} = 0 \quad (1)$$

$$F_{V=0} = 8 + (1) \quad (1.1)$$

$$F_R \# = \left(8 * \prod_{V=1}^{V=R} N_V + (3) \right) + N_V = 30:128:850:9254.. \quad (1.2)$$

The reason the universe is not completely homogenous based on the framework is that the manifold experience arbitrary variations – which than vanish into fermions. marked in black.

$$\frac{\partial \mathcal{L}}{\partial s} \frac{\partial s}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} \delta g - \frac{\partial \mathcal{L}}{\partial s'} \frac{\partial s'}{\partial M} \frac{\partial M}{\partial g'} \frac{\partial^2 g'}{\partial t^2} \delta g' = 0$$

Those variations are arbitrary amount of curvature of a manifold, and they are subject to net variations, which yielded the coupling constant equation. We saw that nature is really the interplay between total arbitrary variations to net variations. Net variations are prime in their nature, and so in the 8- theory Framework for each prime number there exist a boson.

$$8 + (1)$$

$$[(8 * 3) + (3)] + 3$$

$$[(24 * 5) + (3)] + 5$$

$$[(120 * 7) + (3)] + 7$$

The series gives rise to the arrow of time; we should see more interactions as time goes on and so, bigger and bigger structures which makes the manifold less and less homogenous. The bigger the cluster of total variations the weaker the force, as it is divided across the whole cluster.

By looking at those two equations we can see exactly why the universe or the Lorentz manifold in The 8-theory framework is not homogenous, because of those arbitrary variations and the additional net variations. The first accounts for fermions, known as quarks, the other known as bosons. Using that framework, we can see why the manifold cannot be homogenous, it is almost obvious. Of course, the question of the homogenous structure is a question in which we cannot really answer, as it has no numerical data, it's a question revolving around a theory in which the lack of Homogeny is a feature of the main axioms and equations. We can see it in the framework of the 8-theory, or any Lagrangian oriented theory, which includes arbitrary variations, which must vanish at border. The beauty and innovative part in the 8-theory is that, all life forms, galaxies, clusters of galaxies **are** those arbitrary variations.

8-Theory on Universe Expansion-Collapse

$$\frac{\partial \mathcal{L}}{\partial s} \frac{\partial s}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} - \frac{\partial \mathcal{L}}{\partial s'} \frac{\partial s'}{\partial M} \frac{\partial M}{\partial g'} \frac{\partial^2 g'}{\partial t^2} = 0 \quad (1)$$

$$\frac{\partial g}{\partial t} = 0 \text{ and } -\frac{\partial^2 g'}{\partial t^2} = 0$$

This equation describes dark energy or time invariant acceleration from areas of extremum curvature on the Lorenz manifold. We assume no data is available from the first three terms, which describe a varying matrix in spatial dimensions. To ensure universe collapse, we need to revert the signs so we will get:

$$\begin{aligned} +\frac{\partial g}{\partial t} &\rightarrow -\frac{\partial g}{\partial t} \\ -\frac{\partial^2 g'}{\partial t^2} &\rightarrow +\frac{\partial^2 g'}{\partial t^2} \end{aligned}$$

In other words, the acceleration is now directed inwards, and the new equation is:

$$\frac{\partial \mathcal{L}}{\partial s} \frac{\partial s}{\partial M} \frac{\partial M}{\partial g} \frac{\partial^2 g'}{\partial t^2} - \frac{\partial \mathcal{L}}{\partial s'} \frac{\partial s'}{\partial M} \frac{\partial M}{\partial g'} \frac{\partial g}{\partial t} = 0 \quad (1.4)$$

Therefore, we have an inward acceleration and areas of negative curving on the Manifold, which agrees with the description of a compressed Lorentz manifold. However, is it reasonable physically to make such a transformation from (1) to (2)? Suppose it is reasonable to change the direction of the acceleration. By looking at the second term:

$$+\frac{\partial g}{\partial t} \rightarrow -\frac{\partial g}{\partial t}$$

Meaning, all the galaxies, clusters of galaxies, which represent extremum curvature on the manifold, must be eliminated and revert their direction inward, toward the manifold. Such shift will be along an inward acceleration and a process of manifold compression. The process than is synonymous to going from a lower energy state, colder state, to a much higher state of energy. It is a higher state of energy as it is a process of immense masses compressing inward, toward a converging Lorentz manifold, such process will be encompassed by friction, heat and high entropy. It is not Lagrangian oriented and not likeable scenario in our framework. There is no need for calculation of hydrogen atoms per unit space when we have the mathematical equation. We can also analyze the subject of expansion or collapse by using the coupling constant equation in its third representation, the arrow of time.

$$F_{V=0} = 8 + (1) \quad (1.1)$$

$$F_R \# = \left(8 * \prod_{V=1}^{V=R} N_V + (3) \right) + N_V = 30:128:850:9254.. \quad (1.2)$$

$$1 > \frac{1}{30} > \frac{1}{128} > \frac{1}{850} \dots$$

$$\overrightarrow{1 > \frac{1}{30} > \frac{1}{128} > \frac{1}{850} \dots}$$

A universal collapse would be to revert the side of the arrow. From weaker and weaker interactions at mega scales, to go for smaller interactions much stronger:

$$\overleftarrow{1 > \frac{1}{30} > \frac{1}{128} > \frac{1}{850} \dots}$$

The physical meaning would be than, stars, galaxies and clusters of galaxies to deform and in an endless succession until we reach quarks and gluons. Such process would require immense amount of energy and it has to happen across all the spectra of the foreseeable universe. In our framework, it means less manifold net variations (positive curving) over time. Physically it does not make sense, it's not Lagrangian oriented. To go from low state of energy and aspire the highest level. There is no indication that such process could accrue in nature, without artificial intervene. As far as one knows, it comes to an agreement with the laws of thermodynamics. Nevertheless, more importantly, in our framework, there is no reason for such unnatural thing to happen.

The Coupling Constants Equation and Gauge Fields

The coupling constant equation:

$$F_{V=0} = 8 + (1) \quad (1.1)$$

$$F_R \# = \left(8 * \prod_{V=1}^{V=R} N_V + (3) \right) + N_V = 30:128:850:9254.. \quad (1.2)$$

Each term individually:

$$8 + (1)$$

$$[(8 * 3) + (3)] + 3$$

$$[(24 * 5) + (3)] + 5$$

$$[(120 * 7) + (3)] + 7$$

Let us look at the first term:

$$8 + (1)$$

Remember back in the day, when we concluded that we could ignore the eight, since even amount of variations vanish, and just write that the first element is one.

$$8 + (1): (24 + (3)) + 3: (120 + (3)) + 5: (840 + (3)) + 7 \dots$$

$$(1): (30): (128): (850): (9254) \dots$$

We also know that there are eight gluon fields. These are mediating the strong interaction and color charge. However, this could be just a coincidence. Let us examine the next term in the series:

$$[(8 * 3) + (3)] + 3$$

This term describes the nature of the weak interaction. Notice the right inside the parenthesis:

$$(8 * 3)$$

We also know that there are three gauge fields mediating the weak interaction. The massive W and Z bosons, which we correlate to $SU(2)$ and isospin. If the right term inside the parenthesis is a reflection on the number of fields mediating an interaction then we can examine the next term on the series, electromagnetism:

$$[(24 * 5) + (3)] + 5$$

That is a daring statement to make, but if the assumption to hold true, There Should be five gauge fields mediating the electric interaction. Five distinct kinds of photons. It is really an absurd statement to make, given the fact that there are no indication that there is an agreement with experiment regarding that idea. However, sometimes in theoretical physics, bold risks must be taken, and so the author of this paper will allow his belief regarding the great power of the equation to guide him and State: **The 8-theory predicts five gauge fields mediating electromagnetism.** Whether such thing could be correct, only time and experiment will tell.

Quark Mass Mixing and Mixing Angles

Take the masses of all the generations and combine them:

$$[1.9] \quad [1320] \quad [172,760]$$

$$[4.4] \quad [87] \quad [4240]$$

$$1.9 + 4.4 = 6.3$$

$$1320 + 87 = 1407$$

$$172,760 + 4240 = 177000$$

The idea by Quark mixture we mean multiplication of masses of the first and second to yield the total mass of third, times a scalar. Therefore, a total mass of the first family multiplied by the total mass of the second family, both multiplied by a scalar, will yield the total mass of the third. We can proof that is the almost case exactly for the values of the masses above:

$$6.3 * 1407 = 8864.1$$

$$\frac{177,000}{8864.1} = 19.96$$

If we can allow a slight variation of the first masses to be 6.29 Mev and not 6.3, it will be

$$6.29 * 1407 = 8850$$

$$\frac{177,000}{8850} = 20$$

Therefore, just a slight variation of 0.01 Mev and we have a beautiful number and a way to combine the total mass of the first and the second, mix them and multiply by the scalar, to reach the total mass of the third. Reader should argue that it could be just a coincidence, a choice of certain values to yield the scalar and he might be right as the masses are not measured or known as exact, they could divert.

Assuming the mixing will accrue at scalar numbers only, we can build correction angles to ensure the scalar number will hold. So if the masses of the first divert or measured at a higher value than 6.29, there will be a correction angle to retain the same scalar we obtained. The correction angles could have more than one value and they can be positive or negative. Take the mass of the up quark to be average between 1.9 to 2.2 Mev, which is 2.05 Mev.

$$\frac{1.9 + 2.2}{2} = 2.05 \text{ Mev}$$

$$2.05 + 4.4 = 6.45 \text{ Mev}$$

$$6.45 * 1407 = 9075.15$$

$$\frac{177,000}{9075.15} = 19.503$$

The correction angle to reach desired number would be:

$$19.503 + \cos(11.5) \approx 20$$

There could be many more, the correction angles are not limited in number and depend upon the masses values taken of the first, second, and the third as well. The idea behind stay the same. The correction angle will be added to yield a scalar multiple.

$$20 * (TM_1 * TM_2) \approx TM_3$$

Among all the topics can be explained by the 8-theory, and there has been quite a few, the question of Quark mixing seems to be among the hardest ones, and among the topics not within reach. This part is not a proof of any sort but a mathematical idea, the reader should rightfully argue and doubt it. One was trying to reason in the simplest and most elegant way, the weird phenomenon of Quark mixing. Whether it makes sense or not, readers should decide after analyzing the Paper.

The Coupling Constant Equation and Higgs Mechanism

$$F_{V=0} = 8 + (1) \quad (1.1)$$

$$F_R \# = \left(8 * \prod_{V=1}^{V=R} N_V + (3) \right) + N_V = 30:128:850:9254.. \quad (1.2)$$

$$8 + (1):(24 + (3)) + 3:(120 + (3)) + 5:(840 + (3)) + 7 \dots$$

$$(1):(30):(128):(850):(9254) \dots$$

Let us look at the first term describing the strong. We saw that the eight vanish since it's an even in our framework.

$$8 + (1) \rightarrow (1)$$

We also know that from physics the gluons are massless. Let us examine the second term.

$$(24 + (3)) + 3$$

We know that the bosons that mediate the weak interaction do carry mass. Moreover, we know that the symmetry of $SU(2)$ forbids mass terms in the Lagrangian, and the solution which allows us to include mass terms without ruining the symmetry is the Higgs idea. This idea works by including extra terms. In our framework, the **extra term is the majestic three**. Therefore, the Higgs field is responsible for the lack of order in our series, which could have been a beautiful Series of eight multiples. In a sense of the standard model, we can say it is "breaking the symmetry" by inserting the invariant three. So overall, we move from spin 0 – perfect clusters of variations. With the Majestic three Inserted by the Higgs Field we move to a matter with spin one-half, we did so by setting the equation on the critical line of the primes. This *three* is really a destabilizing factor than yields a net variation, which is prime as well.

For example – Electromagnetism:

Perfect clusters of variations $\rightarrow 2N$

Destabilize the perfect $2N$ is the Majestic $(3) \rightarrow \left(\frac{1}{2}\right) \rightarrow \text{electron}$.

Blends in the potential cluster to yield in that case $\rightarrow 123$.

The result is the net variation, which is also prime: $N(V) \rightarrow \left(\frac{1}{2}\right) \rightarrow +(5)$

The overall frame yields:

$$\left[2N + \left(\frac{1}{2}\right)\right] + \left(\frac{1}{2}\right) \rightarrow \mathbf{123} + \mathbf{5} = \mathbf{128}. \text{ Magnitude of an interaction.}$$

The main point of the part is that the Majestic three is a result of the Higgs field. It is the reason the majestic three appears. So overall, our framework does not contradict the Higgs Idea but support it and allow us an additional view on how the mechanism work. As the Higgs is responsible for additional terms in the Lagrangian, and in the 8-theory we see that the first elements in the series of coupling constant differ by an additional term, the Majestic three or spin $(1/2)$.

Anti-Matter & Dirac Delta Variation

$$\frac{\partial \mathcal{L}}{\partial s} \frac{\partial s}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} - \frac{\partial \mathcal{L}}{\partial s'} \frac{\partial s'}{\partial M} \frac{\partial M}{\partial g'} \frac{\partial^2 g'}{\partial t^2} = 0 \quad (1)$$

$$\frac{\partial g}{\partial t} = 0, -\frac{\partial^2 g'}{\partial t^2} = 0$$

$$\left[\frac{\partial \mathcal{L}}{\partial s} \frac{\partial s}{\partial M} \frac{\partial M}{\partial g} \right] \frac{\partial g}{\partial t} \delta g - \left[\frac{\partial \mathcal{L}}{\partial s'} \frac{\partial s'}{\partial M} \frac{\partial M}{\partial g'} \right] \frac{\partial^2 g'}{\partial t^2} \delta g' = 0 \quad (1.1)$$

$$\sum_{i=1}^N \delta g_i = 0 \quad (2.12)$$

Reader should be familiar with the procedure. Now we have seen that we can derive the nature of fermions and the quark model by allowing the series, which contain two distinct elements to vary. So overall we obtain eight threefold combinations of those elements. Therefore, even though the elements are varying the series could vanish. That is in agreement with a stationary Lorentz manifold. There could be however, another way to ensure a stationary Lorentz manifold.

Which will match each element in the series its mirrored element. That is

$$\delta g1 + \delta \exists g1 = 0$$

$$\delta g2 + \delta \exists g2 = 0$$

By mirror, it means the same but opposite in sign. So the overall sum of the Series will hold as zero. In the 8- theory framework, Quarks are regarded as arbitrary amount of curvature on a manifold. Based on this view, anti-quarks and anti-matter is arbitrary curvature with opposite direction. Same magnitude just different direction. So overall, that framework would allow the existence of anti-matter. That is in agreement with quantum field theory and with the Dirac equation for spinors. In fact, the moment of Singularity could be a result of the series not equal to zero.

$$\delta g \neq 0$$

The moment the series is not equal to zero than means that we have net curvature, or maximal curvature on the manifold, which will yield a negative extremum time invariant acceleration from it.

$$\frac{\partial \mathcal{L}}{\partial s} \frac{\partial s}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} - \frac{\partial \mathcal{L}}{\partial s'} \frac{\partial s'}{\partial M} \frac{\partial M}{\partial g'} \frac{\partial^2 g'}{\partial t^2} = 0 \quad (1)$$

In other words, the moment of asymmetry in the series yielding net curvature on the manifold could be the reason for singularity and so called among the masses "big bang". It is just an idea of course, but up until now the 8- theory was on point in regards to Issues on other theory could explain.

Dirac Delta Variation

Our main equations in the framework:

$$\frac{\partial \mathcal{L}}{\partial s} \frac{\partial s}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} - \frac{\partial \mathcal{L}}{\partial s'} \frac{\partial s'}{\partial M} \frac{\partial M}{\partial g'} \frac{\partial^2 g'}{\partial t^2} = 0 \quad (1)$$

$$F_{V=0} = 8 + (1) \quad (1.1)$$

$$F_R \# = \left(8 * \prod_{V=1}^{V=R} N_V + (3) \right) + N_V = 30:128:850:9254.. \quad (1.2)$$

The Dirac delta in our framework is an interference on the Lorenztion manifold. An arbitrary Amount of curvature δg on the manifold. Since it is not allowed and must vanish, we require $\delta g = 0$, as we did previously in this framework.

$$\begin{aligned} \delta g &\neq 0 & at & t = 0 \\ \delta g &= 0 & at & t > 0 \end{aligned}$$

So the Dirac delta in our framework describe the process in which arbitrary amount of curvature appear, and vanish into matter. However, there is no restriction with regard to Time. Arbitrary amount of curvature can appear at any time, so we must modify the idea of the Dirac in our framework.

$$\begin{aligned} \delta g &\neq 0 & at & t = Q(t) \\ \delta g &= 0 & at & t = Q(t + \Delta t) \end{aligned}$$

We also require that $\Delta t \rightarrow 0$ as just after the arbitrary amount of interference will appear, it will immediately vanish into matter. Therefore, in this framework is rich in delta functions. The difference is that the delta can appear at time that is not null. In a sense, we have more flexibility with the delta. After the delta appeared and as a result fermions were manifested into the metric. Those fermions could still vary, and experience a net curvature or net variation. As was analyzed in this paper those net curvatures were taken to be prime numbers and that was the reasoning behind the construction of the coupling constant equation. Those net variations of the manifold are another interference, but and interference which propagate from fermions, and is prime number. Therefore, in that sense it cannot turn into fermions. **Fermions vanish in even amount of variations.** The result is a propagation across the manifold Ripples on the metric all across.

$$\delta g = 0 \quad \text{at} \quad t_1 = Q(t + \Delta t)$$

At later continuation of time:

$$t_2 > t_1$$

This condition is satisfied:

$$\delta g \neq 0 \quad \text{at} \quad t_2 = Q(t + \Delta t + \Delta t)$$

Moreover, the amount of variations is either prime or one:

$$\delta g = 2 \left(n + \frac{1}{2} \right) ; n \geq 0$$

Then we have a ripple on the manifold which propagate all across, toward all directions. The Laplacian operator than is vital to description for a mathematical description of the Manifold ripples, or bosonic fields. Important point to take is that the **underlining reason for the boson propagation all across the metric is their prime number feature**. Define a bosonic ripple across the Lorentzian metric:

$$\nabla^2 = \frac{\partial^2 M_x}{\partial^2 g} + \frac{\partial^2 M_y}{\partial^2 g} + \frac{\partial^2 M_z}{\partial^2 g} \quad (1.41)$$

That is curvature propagation across all metric spatial dimensions as:

$$M_\mu \in S$$

$$S = (M, g)$$

Reasoning for Spiral Structures of Galaxies

$$\frac{\partial \mathcal{L}}{\partial s} \frac{\partial s}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} - \frac{\partial \mathcal{L}}{\partial s'} \frac{\partial s'}{\partial M} \frac{\partial M}{\partial g'} \frac{\partial^2 g'}{\partial t^2} = 0 \quad (1)$$

$$\frac{\partial g}{\partial t} = 0, -\frac{\partial^2 g'}{\partial t^2} = 0$$

$$\left[\frac{\partial \mathcal{L}}{\partial s} \frac{\partial s}{\partial M} \frac{\partial M}{\partial g} \right] \frac{\partial g}{\partial t} \delta g - \left[\frac{\partial \mathcal{L}}{\partial s'} \frac{\partial s'}{\partial M} \frac{\partial M}{\partial g'} \right] \frac{\partial^2 g'}{\partial t^2} \delta g' = 0 \quad (1)$$

$$\delta g = 0$$

Notice the first requirement:

$$\frac{\partial g}{\partial t} = 0$$

In addition, the second requirement:

$$\delta g = 0$$

Those two simple requirements combined together can allow us to a deep Insight into the structure of galaxies. In the 8-theory framework, we have a Lorenz manifold, the manifold has areas of extremum curvature that stay as they are over time. That is given by the first requirement. The manifold also experience arbitrary variations, the second requirement. Those arbitrary variations vanish into matter in agreement with a stationary Lorentz manifold. The combination of both condition than implies that in order for the areas of extremum curvature to stay as they are, the arbitrary variations cannot appear inside them. That is by the combination of the two requirements.

However, those arbitrary variations still appear in the framework. In addition, the areas of extremum curvature are a vital part of this theory. The combination of both requirement is than resulting in areas of extremum curvatures surrounded by arbitrary variations that could not affect them. The following model of the 8-theory is than intersecting with the large scale geometrical shape of galaxies. However, it is known that so called, black holes in the center of galaxies are absorbing matter and nothing can escape them. So in a Second glance the first requirement will not hold in such case. However, that is not a real problem if we assume that those black holes, which we regard as areas of extrunum curvature inside galaxies also omit matter. We know it is the case, as we call it the hawking radiation $-\delta g(H)$.

$$\frac{\partial g}{\partial t} + \delta g + (-\delta g(H)) = 0 \quad (1.442)$$

So overall those two simple requirements in our framework provide an Interesting indication to structure of large-scale matter formations in the universe. The hawking radiation is a vital part of making the two conditions hold true. For each unit of fermions absorbed or manifested inside the area of extremum curvature we require a hawking radiation Particle emitted from the area, so the first requirement will hold true.

The Principle of Least Variation

$$F_{V=0} = 8 + (1) \quad (1.1)$$

$$F_R \# = \left(8 * \prod_{V=1}^{V=R} N_V + (3) \right) + N_V = 30:128:850:9254.. \quad (1.2)$$

$$8 + (1): (24 + (3)) + 3: (120 + (3)) + 5: (840 + (3)) + 7 ...$$

$$(1): (30): (128): (850): (9254) ...$$

We derived the coupling constant as a ration between total arbitrary variations to the net variations, N_V , which are outside the parenthesis. Those net variations are a different representation of curvature on the Lorenzton manifold. Notice the numerical relations between the total to net:

$$\frac{N_V}{T_V} \rightarrow R \quad (1.5)$$

$$\frac{1}{9} = 0.111$$

$$\frac{3}{30} = 0.1$$

$$\frac{5}{128} = 0.039$$

$$\frac{7}{850} = 0.008$$

...

$$0.111 > 0.1 > 0.039 > 0.008 \dots \quad (1.51)$$

The reasoning was clear, as the coupling constant equation is multiplies each Even sum of the previous element in the next prime, and the net variations are the prime numbers sequence itself. In means that each element the net curvature is a smaller and smaller portion of the whole variation cluster, which reason why the sequence is getting weaker and weaker. Based on this equation we can vividly derive and predict the weakness of gravity. We can say that nature is aspiring to minimize the ratio of net to total. All the possible amount of curvature can and will appear and nature, but the most common and noticeable ones are those with the bigger ratio, or least amount of net variation:

$$0.111 > 0.1 > 0.039 > 0.008 \dots \rightarrow 0$$

The bosons in which we are already know of. The interactions associated with the number one, three and five. The two lowest primes and one. The 8 – theory principle, which is derived by this analysis, is the Principle of least variation or curvature as we are dealing with a Lorenz manifold. Just as Feynman did in quantum path integrations, all is taken into account. However, the most significant routes are the simplest ones. In this framework the most significant Interactions are those with the largest ratios between the net Variations to the total variations The largest ratios are those with the least curvature or Smallest prime numbers and the number one, and primes are representing manifold variations.

The Coupling Constant Equation And The Wave-Particle Duality

$$F_{V=0} = 8 + (1) \quad (1.1)$$

$$F_R \# = \left(8 * \prod_{V=1}^{V=R} N_V + (3) \right) + N_V = 30:128:850:9254.. \quad (1.2)$$

$$8 + (1):(24 + (3)) + 3:(120 + (3)) + 5:(840 + (3)) + 7 ...$$

$$(1):(30):(128):(850):(9254) ...$$

We can vary the N_V outside of the parenthesis so by doing so, reaching duality among the three first forces at 26 variations was attained.

$$[8 + (1)] + 3 - (\mathbf{1v}): \quad [(8 * 3) + (3)] + 3 : \quad [(24 * 5) + (3)] + \mathbf{3}$$

By analyzing the third element in the series, the propagation of a photon from a fermion so called the electron. Certain insight from the new framework is becoming vividly clear. In the context of wave particle duality.

$$[(24 * 5) + (3)] + 5$$

Since it is a prime net variation outside the parenthesis, it can not vanish into matter. As fermions vanish in even amounts. The ripple field of boson across the matric is mathematically described:

$$\nabla^2 = \frac{\partial^2 M_x}{\partial^2 g} + \frac{\partial^2 M_y}{\partial^2 g} + \frac{\partial^2 M_z}{\partial^2 g}$$

$$M_{x,y,z} \in S$$

$$S = (M, g)$$

Suppose, in an experiment we decide to measure the photon momenta of position. Its done by scattering an additional photon onto the photon, which already propagated Form the electron. For simplicity sake, we suppose it is one additional element that is only one photon:

$$[(24 * 5) + (3)] + 5 \rightarrow [(24 * 5) + (3)] + 5 + 5$$

$$[(24 * 5) + (3)] + 5 \rightarrow \left[2N_2 + \frac{1}{2}\right] + \frac{1}{2}$$

$$[(24 * 5) + (3)] + 5 + 5 \rightarrow \left[2N_2 + \frac{1}{2}\right] + \frac{1}{2} + \frac{1}{2}$$

Those equations are the second variation of the coupling constant equation, which is the prime critical line. By adding the additional net variation, we reach a spin that is no longer associated with boson propagation, $3/2$. Before our measurement the bosons had spin one, and by measurement with additional photon, a variance of spin has occurred, so now our Boson behave like a fermion, it has an additional half unit of spin. Overall in the 8-theory by analyzing the coupling constant equation in the second representation, it is possible to extrapolate the reason for the phenomenon of wave particle duality.

The Feynman Path Integral Variation on Varying Lorentzian Manifolds

In the 8-theory a varying Lorentz manifold is the entity of description. The Lorentz manifold is inserted to an Euler Lagrange equation and by doing so, the main equation of the framework is obtained.

$$\frac{\partial \mathcal{L}}{\partial s} \frac{\partial s}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} - \frac{\partial \mathcal{L}}{\partial s'} \frac{\partial s'}{\partial M} \frac{\partial M}{\partial g'} \frac{\partial^2 g'}{\partial t^2} = 0 \quad (1)$$

Are the conditions, which the framework is demanding to retain a stationary manifold. Those two conditions describe a time invariant acceleration directed from extrunum areas of curvature on the Lorentzian manifold. Intersection with the so-called dark energy. In addition, if no data is attainable from the first three terms, it is vividly clear that there is an agreement with Einstein equivalence principle:

The coupling constant equation was obtained by demanding a stationary manifold to experience net variation, $N_V = 2(V + 1/2)$; $V \geq 0$; $N_V \in P$ as P to be is the set of primes and the number one. Those ideas yielded an infinite series, in which each distinct Boson has a distinct prime, which is the N_V value. Even amount of variations vanish, $2V = 0$.

$$F_{V=0} = 8 + (1) \quad (1.1)$$

$$F_R \# = \left(8 * \prod_{V=1}^{V=R} N_V + (3) \right) + N_V = 30:128:850:9254.. \quad (1.2)$$

The Feynman variation on Lorentz manifold - the objective of this part is to find out what is the probability transition of a boson from initial to final state on the manifold. Bosons are associated with prime amount of net variation $N_V \rightarrow 2 \left(V + \frac{1}{2} \right)$ which propagates as ripples on the manifold, given by a variation of the Laplacian:

$$\nabla^2 = \frac{\partial^2 M_x}{\partial^2 g} + \frac{\partial^2 M_y}{\partial^2 g} + \frac{\partial^2 M_z}{\partial^2 g}$$

First, we define a manifold $s = (M, g)$ and insert it to an Euler LaGrange equation and an initial state of the manifold $S(0) = (M, g)$.

Manor O

Than we require the manifold to experience arbitrary variations, which vanish into matter.

For simplicity of notation, define:

Let the arbitrary variations appear all across the matric on the manifold.

$$\delta g \in M(x, y, z)$$


Define a ripple propagation of a boson from an initial point on the manifold:

$$q1 = M(x1, y1, z1)$$

And a final position of the matric ripple to arrive at

$$q2 = M(x2, y2, z2)$$

Intentionally left as a blank page.



The green arrow is directing the from the initial position of the ripple wave to final Position. The blue dots are the electrons omitting bosons, in that case a photon, marked in yellow. The manifold has arbitrary amount of fermions on it, which get scattered by the initial boson wave and omit a new boson. There are infentially more ways than the above drawing, its vivid. The ripple wave will scatter all the arbitrary variaions, but the highest probability of arrival will be at the path of least curvature.

Each fermion which get scattered omitting a boson with random direction of propagation. the framework has no data regrading the position of the propagation. The more arbitrary Variations getting scattered, the less probable it is to reach the final position. The following can be analyzed by the equation. The more arbitrary variaions in the Path, the more curved the matric, as there is an accelaration of it outward. The accelaration outward is causing the path to be longer and less linear, and so the time to reach the final position is longer, if only. There is no gurentte a photon will reach the final position in this framework as arbitrary variaions created in a random fashion, and in configurations which are not predictable. However, if a photon will reach it will be in the least curved path, or the path with the least fermions getting scattered.

$$P = \int_{q1(t(i))}^{q2(t(f))} dq \exp[(S0) \int_{t(i)}^{t(f)} L(s, s') dt] \quad (10)$$

Its unclear whether (10) is solvable as the arbitrary variations themselves vary their position over time and in addition, arbitrary variaons appear in random fashion in this framework. Its given by the first equation. So in a sense we can not sum all the paths if the paths vary at all times. it's a complication of the feynamn result, and at the same time just as beautiful as feynman reuslt. But if we ignore the complication, the probablity transition should be calcuated using (10).

Gravitational Coupling Constant as a Combination of Particles

$$F_{V=0} = 8 + (1) \quad (1.1)$$

$$F_R \# = \left(8 * \prod_{V=1}^{V=R} N_V + (3) \right) + N_V = 30:128:850:9254.. \quad (1.2)$$

We can also represent the equation in the form:

$$F_R \# = \left(8 * \prod_{V=1}^{V=R} N_V + (3) \right) + N_V = \frac{1}{30} : \frac{1}{128} : \frac{1}{850} : \frac{1}{9254} ..$$

Let us analyze the third element – Electromagnetism:

$$[(24 * 5) + (3)] + 5$$

$$[(24 * 5) + (3)] + 5 \rightarrow \left[2N_2 + \frac{1}{2} \right] + \frac{1}{2}$$

$(2N \text{ variations}) \rightarrow \text{Spin } 0$

$(2N \text{ variations} + 3) \rightarrow \text{Matter with spin } (\frac{1}{2})$

$(2N \text{ variations} + 3) + N_V \rightarrow \text{Bosons with spin } (1)$

$(2N \text{ variations} + 3) + N_V 1 + N_V 2 + \dots \rightarrow \text{Boson with higher spin integers}$

Given that framework, we can vividly see that gravity is belonging to the bosons with higher spin integers, as modern theories predict the gravitational interaction to have spin two. In the 8-theory framework, what does it mean? In the context of the coupling constants equation what does it mean? Since it has spin two, we can associate gravity to the category of bosons with higher spin integers, which could relate to a certain combination of elements in the coupling constant series, as the elements are getting weaker and weaker, if the gravitational coupling will not be found by keeping developing to infinity it could mean **gravitational will be found as a combination of elements in the series**. Since it is spin two there should be three net variations outside. Gravitation as a combination of elements, using the fact it has a boson with spin 2.

$$\begin{aligned} [2N_{gravity} + (3)] + N_{V1} + N_{V2} + N_{V3} \rightarrow \left[2N_{gravity} + \frac{1}{2}\right] + \frac{1}{2} + \frac{1}{2} + \frac{1}{2} = \\ [2N_{gravity} + 2] \end{aligned}$$

Using the second representation of the coupling constant equation, meaning spin. It also means that the gravitational is a lot more rare as it is requiring a combination of elements in the series to be emitted and not just a singular element.

Using the Coupling Constant Equation to Predict the Exact Mass of the Graviton

$$F_{V=0} = 8 + (1) \quad (1.1)$$

$$F_R \# = \left(8 * \prod_{V=1}^{V=R} N_V + (3) \right) + N_V = 30:128:850:9254.. \quad (1.2)$$

$$N_V = 2 \left(V + \frac{1}{2} \right); \quad V \geq 0$$

$$N_V \in \mathbb{P} \cup (+1); \quad \mathbb{P} \rightarrow \text{Set of Primes}$$

$$N_V = P_{max} \in [0, \mathbb{R}] \cup (+1); \quad P_{max} \in \mathbb{P}$$

$$8 + (1): (24 + (3)) + 3: (120 + (3)) + 5: (840 + (3)) + 7 \dots$$

In the 8-theory framework, even amounts of manifold variations vanish. That feature allowed the following shift:

$$8 + (1) \rightarrow (1)$$

We know that the strong interaction has eight gauge fields meditating it. Those meditating particles do not carry mass. We also know that gravity has spin two. By switching to the second representation of the equation, we can represent gravity as the following:

$$[(2N_{gravity}) + (3)] + N_{V1} + N_{V2} + N_{V3} = \left[2N_{gravity} + \frac{1}{2} \right] + \frac{1}{2} + \frac{1}{2} + \frac{1}{2}$$

$$[2N_{gravity} + 2] \rightarrow 2N_{gravity}$$

ince even amount of variations vanish we will be left with one term in the final form of the term. That is similar to the strong interactions but immensely weaker. Since the bosons mediating the strong interaction are massless, and we can represent it in one term given the coupling constant equation, and by the analysis gravitation has only one term as well, we can reach a mathematical prediction, which will state, that gravitons has no mass. In agreement with reality and agreement with quantum field theory. The only thing taken from what was known before was the fact that the bosons mediating the strong interaction are massless.

Indication That Fermions Are Closed Circles by the Coupling Constant Primorial Function Variation

$$F_{V=0} = 8 + (1) \quad (1.1)$$

$$F_R \# = \left(8 * \prod_{V=1}^{V=R} N_V + (3) \right) + N_V = 30:128:850:9254.. \quad (1.2)$$

$$N_V = 2 \left(V + \frac{1}{2} \right); \quad V \geq 0$$

$$N_V \in \mathbb{P} \cup (+1); \quad \mathbb{P} \rightarrow \text{Set of Primes}$$

$$N_V = P_{max} \in [0, \mathbb{R}] \cup (+1); \quad P_{max} \in \mathbb{P}$$

$$8 + (1): (24 + (3)) + 3: (120 + (3)) + 5: (840 + (3)) + 7 \dots$$

The following representation of equation (1) by replacing the invariant three with pi.

$$F_R \# = \left(8 * \prod_{V=1}^{V=R} N_V + (3) \right) + N_V \rightarrow \left(8 * \prod_{V=1}^{V=R} N_V + (\pi) \right) + N_V$$

$$8 + \left(\frac{\pi}{3} \right): (24 + (\pi)) + 3: (120 + (\pi)) + 5: (840 + (\pi)) + 7 \dots$$

That is giving up certain accuracy on the coupling constant equation in order to get an insight regarding the shape of fermions. One is going to argue that such a representation is valid as we have a varying Lorentz manifold, there could be slight variations in the invariant three over time, toward pi and vice versa. In other words, the electron is not a perfect circle, but close to it. It is a varying circle, not a perfect shape. Varying in physical theories could mean vibration.

The fact that we have a varying framework allow us to dynamically allow such slight variations without being rigid, the fact that it is not pi, could be a positive indication. Perfect shape of a circle would be problematic in a final theory, but a varying, imperfect circle seems to be much more elegant and suitable to a framework of constant variation. So according to this representation, a boson will be emitted from something close to a perfect circle, which is the electron. We gave up certain amount of accuracy and reached an astonishing insight regarding the shape of an electron. But we can go even further by representing the net variations in pi number multiples.

$$8 + \left(\frac{\pi}{3}\right) : (24 + (\pi)) + 3 : (120 + (\pi)) + 5 : (840 + (\pi)) + 7 \rightarrow$$

$$8 + \left(\frac{\pi}{3}\right) : (24 + (\pi)) + \pi : (120 + (\pi)) + (\pi + 1.82) : (840 + (\pi)) + (2\pi + 0.716)..$$

Such representation is beautiful but what does it mean? of course that the real answer is that one does not know. Two options come to mind. The first is regrading the probability to find a boson in varying area. the bigger variations clusters, the larger the area of possible emission and the less likable it is to detect the boson. The higher the net variations, the smaller the probability to find the boson. Another possible option is of magnitude. The boson propagate across larger areas and thus its energy is getting divided across the area, so overall it gets much weaker as we develop the coupling constant series into infinity. In agreement with the weakness of gravity. Since the 8-theory was born in 2021, there could be more variations to the coupling constant equation. Other indication fermions are of closed shape is the main equation:

$$\frac{\partial \mathcal{L}}{\partial s} \frac{\partial s}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} - \frac{\partial \mathcal{L}}{\partial s'} \frac{\partial s'}{\partial M} \frac{\partial M}{\partial g'} \frac{\partial^2 g'}{\partial t^2} = 0 \quad (1)$$

Which describe a varying Lorentz manifold. Fermions were proved to be arbitrary variations of the manifold. If the manifold is of finite size, i.e. closed, the elements in it should be closed as well. They are not a separate entity of the Lorentz manifold, but appear as part of the Lorentz manifold and its ever varying nature. The closeness of the manifold indicate the closeness of the elements that appear in it. There could be more ways to prove that the following is correct.

Primorial Coupling Constants Equation and the Rise of the Arrow of Time

$$F_{V=0} = 8 + (1) \quad (1.1)$$

$$F_R \# = \left(8 * \prod_{V=1}^{V=R} N_V + (3) \right) + N_V = 30:128:850:9254.. \quad (1.2)$$

$$N_V = 2 \left(V + \frac{1}{2} \right); \quad V \geq 0$$

$$N_V \in \mathbb{P} \cup (+1); \quad \mathbb{P} \rightarrow \text{Set of Primes}$$

$$N_V = P_{max} \in [0, \mathbb{R}] \cup (+1); \quad P_{max} \in \mathbb{P}$$

$$8 + (1): (24 + (3)) + 3: (120 + (3)) + 5: (840 + (3)) + 7 \dots$$

Suppose a boson was emitted from a fermion due to net variation of a certain magnitude. If the arrow of time is two sided and reversible, there must be a way to bring the photon back to the electron. However, the physics of the 20-th Century forbids us from doing that, as we don't even know where the photon is. Momentum and position are conjugate variables in quantum field theory. So once a boson is propagated into the metric, there is no possible way to bring it where it was. An additional argument is that all bosons are indistinguishable, so even if it was possible to trace and revert the photon, in a system with more than one Photon, its again beyond reach. The reason we emphasize those arguments as to the context of the arrow of time. At first, at a certain point after the singularity, there were only elements of the first Element in the coupling series on the expended manifold:

$$8 + (1)$$

If the expended manifold experience multiple net variations of the first element than it is possible to cluster those:

$$\sum_{n=1}^{n=\infty} C_n = 8 + (1)$$

We can cluster into groups of three and get:

$$(8 * 3) + (1) * 3 = 24 + (3)$$

The invariant three, in 8-theory framework is, as you already know, is the destabilizing factor yielding a net variation so overall:

$$24 + (3) \rightarrow [24 + (3)] + 3$$

Therefore, we can derive the intimate relation between the coupling constant series and the direction of time. The following procedure can be done on any additional element in the series. **In the 8-theory what is time? Time is the result of net variations being clustered to different magnitudes.** The succession of bosons with decreasing magnitude converging to zero is the direction of the arrow. The fact that each element is different than is preceding is the physical manifestation of the arrow of time. This equation encompass all the interactions according to magnitude, and so as those are different, the difference is the factor that gives rise to the arrow. If all elements in the series were identical there could not be a rise to the arrow. Using that coupling constant equation, we can reason for the chronology of events from the moment of singularity to the present moment. We can reason for electrons propagation only after protons were created. We can reason gravitational interactions only after electric interactions and we possibly can reason also, how galaxies were formed. Notice that the fourth element in the series is only 6.65 weaker than the electric. That is immensely stronger than the gravitational interaction and using that element as a building block for clustering after electric interactions and it is possible to explain how relatively fast galaxies formed in a short window of time, from the manifold being too hot to being too

Universe Packets - Stationary Manifolds

The main equation:

$$\frac{\partial \mathcal{L}}{\partial s} \frac{\partial s}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} - \frac{\partial \mathcal{L}}{\partial s'} \frac{\partial s'}{\partial M} \frac{\partial M}{\partial g'} \frac{\partial^2 g'}{\partial t^2} = 0 \quad (1)$$


In agreement with our model of the universe. Negative time invariant acceleration From areas of extremum curvatures on the manifold. Validating the Einstein equivalence principle between gravity and acceleration. Again, we assume no data is available from the first three terms, no indication They agree with a stationary Lorentz manifold. Now we can represent the equation (1) in a different way, if there are many stationary Lorentz manifolds we can write:

$$\frac{\partial \mathcal{L}}{\partial S_1} - \frac{\partial \mathcal{L}}{\partial S_2} = 0 \quad (1.52)$$

Alternatively:

$$\frac{\partial \mathcal{L}}{\partial S_1} - \sum_{n=2}^{\infty} \frac{\partial \mathcal{L}}{\partial S_n} = 0 \quad (1.53)$$

$$\frac{\partial \mathcal{L}}{\partial S_1} \frac{\partial S_1}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} - \sum_{n=2}^{\infty} \frac{\partial \mathcal{L}}{\partial S_n} \frac{\partial S_n}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} = 0 \quad (2)$$



The Principle of Least Curvature and Cosmological Flatness

In the 8-theory a varying Lorentz manifold is the entity of description. The Lorentz manifold is inserted to an Euler Lagrange equation and by doing so, the main equation of the framework is obtained.

$$\frac{\partial \mathcal{L}}{\partial s} \frac{\partial s}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} - \frac{\partial \mathcal{L}}{\partial s'} \frac{\partial s'}{\partial M} \frac{\partial M}{\partial g'} \frac{\partial^2 g'}{\partial t^2} = 0 \quad (1)$$

$$F_R \# = \left(8 * \prod_{V=1}^{V=R} N_V + (3) \right) + N_V = 30:128:850:9254.. \quad (1.2)$$

$$N_V = 2 \left(V + \frac{1}{2} \right); \quad V \geq 0$$

The framework regard (6) as the second main construction of the theory and currently has five representation and different uses of the equation. (6) represent the concept of Least variation, the most significant interactions in nature, are those with the largest Value of total to net $\frac{N_V}{T_V}$. By analyzing the ratios of the first elements in the series we Conclude the ratio $N_V/T_V \rightarrow 0$ as $N_V \rightarrow \infty$, the result is following ratios:

$$\frac{1}{9} = 0.111, \quad \frac{3}{30} = 0.1, \quad \frac{5}{128} = 0.039 \dots$$

$$0.111 > 0.1 > 0.039 > 0.008 \dots$$

The biggest ratios are those with the least $N(V)$ amount and thus they are the most noticeable on the manifold. The third representation of (6) revolves around to the arrow of time. The direction of the series is assumed to match the direction of Time. So as we increment the time $t(1) = t(1) + \Delta t$ and allowing time aspire Infinity $t \rightarrow \infty$, the manifold will experience higher number of net variations $N(V) \rightarrow \infty$. and at the same time the ratio of $\frac{N_V}{T_V} \rightarrow 0$, which means that the matrix on the manifold Is getting more and more flat. The most curved, or intense interaction than is the first, the Strong interaction due to its largest value of $\frac{N_V}{T_V}$.

Using the coupling constant equation ratio between net to total variations (curvature) it becomes vividly clear that gravitation will be aspiring to flatness, due to the immense value of N_V . Gravity will be almost not noticeable. It is also possible to derive that the manifold will become more flat followed by the direction of the arrow. Flatness than, in the 8-theory is a continuous process.

Quark Confinement

$$\frac{\partial \mathcal{L}}{\partial s} \frac{\partial s}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} - \frac{\partial \mathcal{L}}{\partial s'} \frac{\partial s'}{\partial M} \frac{\partial M}{\partial g'} \frac{\partial^2 g'}{\partial t^2} = 0 \quad (1)$$

We already proved that δg can be analyzed as fermions. Suppose we did what we did back in the day, and break it to a sequence of N distinct elements, such;

$$\delta g \rightarrow \sum_{i=1}^N \delta g_i = 0$$

$$N \in \mathbb{R}, N \rightarrow \infty$$

How would such a set of elements would behave? To answer such a question we can analyze the second term:

$$-\delta g' \rightarrow \sum_{i=1}^N -\delta g'_i = 0$$

$$\delta g \rightarrow \sum_{i=1}^N \delta g_i = \delta g_i + \delta g_{i+1} + \dots = 0$$

$$-\delta g' \rightarrow \sum_{i=1}^N -\delta g'_i = -\delta g'_i + (-\delta g'_{i+1}) + \dots = 0$$

On one hand the entire series need to vanish, we have a sequence of opposite signs of even numbers, those arbitrary variations derivative describe a acceleration outward, they will try to aspire to reach distance from each other, that is with agreement with the so-called "anti commutation" relation of fermions. Since half of those elements in the set **already have a negative sign** as we proved before:

$$(-\delta g'_{i+1}) \rightarrow -(-\delta g'_{i+1}) \rightarrow +\delta g'_{i+1}$$

Therefore, despite the minus sign, and the outward distancing from each, in accordance to the anti-commutation relation, there is a shift and the minus now varied to a plus:

$$-\delta g' \rightarrow \sum_{i=1}^N -\delta g'_i = -\delta g'_i + (-\delta g'_{i+1}) + \dots = -\delta g'_i + \delta g'_{i+1} \dots = 0$$

Therefore, that is in agreement with quantum field theory, and their treatment of fermions. They will accelerate toward each other, which can explain the phenomena of quark confinement. The equations show that arbitrary variations of distinct sign will accelerate toward each other, if the reasoning and the mathematical development one presented here is correct.

Quark Confinement and the D'alembert Variation

$$\frac{\partial \mathcal{L}}{\partial s} \frac{\partial s}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} - \frac{\partial \mathcal{L}}{\partial s'} \frac{\partial s'}{\partial M} \frac{\partial M}{\partial g'} \frac{\partial^2 g'}{\partial t^2} = 0 \quad (1)$$

$$\delta g \rightarrow \sum_{i=1}^N \delta g_i = 0$$

$$N \in \mathbb{R}, N \rightarrow \infty$$

$$-\delta g' \rightarrow \sum_{i=1}^N -\delta g'_i = 0$$

$$\sum_{i=1}^N \delta g_i - \sum_{i=1}^N \delta g'_i = 0$$

$$\sum_{i=1}^N \partial E_i / \partial t - \sum_{i=1}^N \frac{\partial E_i^2}{\partial^2 t} = 0 \quad (2.1)$$

The sum of all arbitrary variations and accelerations is taken to zero in this framework. Similar to the procedure D'alembert taken with forces and accelerations. That is an additional take on the phenomena of quark confinement, published earlier by the author.

Infinite Dimensional Multiverse

$$\frac{\partial \mathcal{L}}{\partial s} \frac{\partial s}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} - \frac{\partial \mathcal{L}}{\partial s'} \frac{\partial s'}{\partial M} \frac{\partial M}{\partial g'} \frac{\partial^2 g'}{\partial t^2} = 0 \quad (1)$$

$$\frac{\partial \mathcal{L}}{\partial S_1} - \sum_{n=2}^{\infty} \frac{\partial \mathcal{L}}{\partial S_n} = 0 \quad (1.53)$$

$$\frac{\partial \mathcal{L}}{\partial S_1} \frac{\partial S_1}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} - \sum_{n=2}^{\infty} \frac{\partial \mathcal{L}}{\partial S_n} \frac{\partial S_n}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} = 0 \quad (2)$$

Equation (2) is the second representation of the main equation. Notice that even though the main equation describe both Einstein theory of relativity (1) and time invariant acceleration away from extremum curvature on the manifold, it lacks providing the reason for such a process. Equation (2) is than used; our universe is wrapped in many similar, stationary manifolds, which are distinct. They are assumed topologically invariant. Such a construction than allow us to understand why each manifold can not have any number of dimensions, it is confined within many other manifolds. Its also more reasonable to assume that there are many stationary manifolds than to assume that there is only one stationary manifold. The (2) is more elaborated equation than equation (1). Suppose that each manifold has n-dimensions.

$$\frac{\partial \mathcal{L}}{\partial S_1} \frac{\partial S_1}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} \rightarrow N_D^{S1}$$

Take into account the number of manifolds wrapping our manifold making its matric accelerate outward and add those dimensions

$$\sum_{n=2}^{\infty} \frac{\partial \mathcal{L}}{\partial S_n} \frac{\partial S_n}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} \rightarrow \sum_{n=2}^{\infty} N_D^{S(n)}$$

So the number of dimensions in our framework is:

$$N_D^{S1} + \sum_{n=2}^{\infty} N_D^{S(n)} = T_D \quad (2.11)$$

The Equivalence Principle in Quantum Scale

$$\frac{\partial \mathcal{L}}{\partial s} \frac{\partial s}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} - \frac{\partial \mathcal{L}}{\partial s'} \frac{\partial s'}{\partial M} \frac{\partial M}{\partial g'} \frac{\partial^2 g'}{\partial t^2} = 0 \quad (1)$$

Suppose the matric has a fermion that is an arbitrary variation of the manifold given by the term $\delta g = 0$. What would be the consequence? Given by equation (2) the arbitrary variation will cause the matric to accelerate outward. That is in complete agreement of Einstein theory of gravitation, equation (2) implies:

$$\frac{\partial g}{\partial t} = \frac{\partial^2 g'}{\partial t^2}$$

Since the matric varied due to the arbitrary variation, which appeared in it, and in particular, it expended outward, the distance increased. Suppose the quark was conscious and could perform measurement, its very existence affected the matric, and the time in which a boson field will need to reach the object measured has increased because of the quark manifesting. In special relativity, the great Einstein used velocity, but here there is no velocity. There is no such thing velocity in the 8 theory. The quark may conclude that the object is moving, but what is happening is that the matric itself is varying, because of that quark. We also have in this framework the invariance of the speed of light, given by the coupling constant equation, and the fact that the propagation process is similar in all interactions.

$$[(8 * 3) + (3)] + 3 \rightarrow \left[2N_1 + \frac{1}{2}\right] + \frac{1}{2}$$

$$[(24 * 5) + (3)] + 5 \rightarrow \left[2N_2 + \frac{1}{2}\right] + \frac{1}{2}$$

General relativity implies an equivalence relation between curvature and acceleration. 8 theory implies that as well, but also in addition implies that curvature will **cause** outward acceleration of the matric by (1). Einstein had to add the cosmological constant in an artificial way, but here it's the main equation. Such a condition than allow us to understand relativity in a new and elegant way. C is invariant, and every arbitrary variation of the manifold causing an outward acceleration of the matric, the matric itself varying in such way that those arbitrary variations will eventually measure different distances and times, the measured object can be standing still but it will observed as moving, but what is happening is that the matric is expending. The entire theory of Einstein is not only contained in just one equation but expended to a new horizon.

Majestic Three is the Electron

$$\frac{e^2}{4\pi\hbar c} = \frac{1}{128}$$

$$\frac{e^2}{4\pi} \rightarrow \frac{3^2}{128}$$

Recall that arbitrary variations vanish in pairs of even numbers. That axiom in our framework related to fermions and allowed us to make a transformation regarding the strong interaction:

$$8 + (1) \rightarrow (1)$$

So we can vary (2) to prove that the majestic three is indeed an electron and solidify our theory and its validity:

$$\frac{3^2}{128} = \frac{8 + (1)}{128}$$

Even amount of variations taken to vanish so the final form of equation (2) is exactly like the equation in the beginning of the paper.

$$\frac{8 + (1)}{128} \rightarrow \frac{(1)}{128} = \frac{e^2}{4\pi\hbar c}$$

Using the Primorial Coupling Constants Function to Derive C Invariance Yang Mills Conjecture

$$F_{V=0} = 8 + (1) \quad (1.1)$$

$$F_R \# = \left(8 * \prod_{V=1}^{V=R} N_V + (3) \right) + N_V = 30:128:850:9254.. \quad (1.2)$$

$$N_V = 2 \left(V + \frac{1}{2} \right); \quad V \geq 0$$

$$N_V \in \mathbb{P} \cup (+1); \quad \mathbb{P} \rightarrow \text{Set of Primes}$$

$$N_V = P_{max} \in [0, \mathbb{R}] \cup (+1); \quad P_{max} \in \mathbb{P}$$

The second representation of the primorial function using the prime critical line:

$$[(8 * 3) + (3)] + 3 \rightarrow \left[2N_1 + \frac{1}{2} \right] + \frac{1}{2}$$

$$[(24 * 5) + (3)] + 5 \rightarrow \left[2N_2 + \frac{1}{2} \right] + \frac{1}{2}$$

$$[(120 * 7) + (3)] + 7 \rightarrow \left[2N_3 + \frac{1}{2} \right] + \frac{1}{2}$$

Notice that beside the variation cluster which get bigger, all the interactions are taking the same form. We have the destabilizing factor, which is the electron for example, or the $\frac{1}{2}$ inside the parenthesis, yielding a net variation which is also $\frac{1}{2}$ or prime, according to first representation of the primorial function. So the propagation speed of all boson of this type must be similar, precisely because there is no detail regarding the speed of propagation and because there is no difference among the bosons, they are all of the same hand. We already proved their dynamical nature by varying the net variations and as a result making them scalar multiples.

Based on the framework of the eight theory, we can make an additional prediction. All bosons of the above type, $2n() + 1$, with no consideration of their mass, will propagate across at the same speed. The same speed applies for all. Even to bosons with mass. That is the young mills problem, how can a boson that carry mass move at the speed of a boson, which do not carry mass. In the 8-theory the answer is given. Since mass is associated with $8 - 1$ variations, and boons are of the type $8 + 1$ the combination of a boson with mass will not effect on the propagation.

$$8 - 1 + 8 + 1 = 0 \quad (3.1)$$

The boson that is a mass carrier, causing the matric to converge inward, will be balanced to the other direction by its very nature. As a result, he will move on a linear, not curved trajectory and his speed will not be effected by its mass. In equation six, we took even amount of variations to vanish and so the result is zero. No curving to either direction. Of course, the ideal would be to extract the actual speed of light from the 8-theory. It is currently beyond reach. The equation does not change under any condition that means that the speed of propagation does not means under any conditions. In the case of the third element, than, speed of light is invariant to all. Therefore, the 8-theory framework suggest an elegant and simple solution to the Yang mills conjecture.

8T and QFT – Axiomatic Analysis

Quantum field theory has certain features that play a significant rule, and repeat themselves in one way or another along each epos of the theory. Among those, we can name the commutation and anti-commutation of bosons and fermions. The Dirac delta or interference known as a field, the operators of matter creating and destructing, cluster decomposition and Lorentz invariance. In addition to Feynman path integrations and diagrams. That being said, what are the mathematical axioms in which QFT is built upon? One would like to suggest those following axioms:

Axiom (1) – Nature is probabilistic

Axiom (2) – Fermions repeal, Bosons do not

Axiom (3) – There is only one set of rules

By the first axiom, we can include the Feynman diagrams and the Feynman path integrations. In addition to arbitrary amount of matters appear and disappear by operators we insert. By the second axiom the commutation and anti-commutation relation and the nature of spin and statistics. The third axiom, the Lorentz invariance and the entire set of symmetries and conservation laws, at quantum scale (Nother) and at large scale (Lorentz). Those three axioms also stand at the heart of 8-theory, so in essence the nature of those theories, their innate ideas about nature is the same. The difference is which ideas are describing the axioms and which objectives the theory is set to achieve. Quantum field theory searches for probability of certain occurrences, it does it amazingly well but lacks to provide the reason for those arbitrary numbers, such as coupling magnitudes. QFT uses integrations across the entire space-time that are impossible to solve. 8- Theory is also probabilistic in its nature, maybe even more than QFT. It has no data regarding any direction of motion, momenta, and location at any point and so on. Very little to no physical data is manifested in this theory. However, it does describe beautifully the magnitudes of the couplings, the reason each magnitude is what it is, the process of propagation and the dynamic nature of the forces.

The methods uses are partial differential equations, and the methods also uses in quantum field theory given by axiom (2), the commuting relation of fermions and bosons. It does not currently have complicated integrations over space-time or it can specify the decays as QFT. However, it does describe the dark energy in an accurate fashion given by its main equation, a varying Lorentz manifold. Gravity is within its domain of description as it was built upon the work of two of the greatest minds in science Einstein and Lorentz. It is also supported by the coupling constant equation and predict that graviton will be massless and that gravity is actually a combination of three net variations. The 8T has two arbitrary numbers less than QFT; it predicts infinite bosonic fields, which relate to Lorentz net curvature on the manifold. It also predicts infinite families below first generation, and thus does not face questions as to those arbitrary numbers.

8T and QFT both are described in terms of the Dirac delta. QFT uses the delta as a description for the wave equation, as a way to describe a complete set of states, alongside with a set of amplitudes. 8T uses the Dirac delta in more flexible manner, it applies to times that are different from zero as well, and describe how an arbitrary amount of curvature vanish into matter. Any net variation at a later continuation of time than describing a bosonic ripple field across the manifold, given by a variation of the Laplacian. While QFT is mainly physical, 8T is mainly and almost completely mathematical, the axioms at the heart of those theories are the same, the methods are similar, the 8T describe phenomena not within the realms of QFT, and QFT can calculate probabilities not within the realm of 8T. 8T is just as probabilistic as QFT, if not more. It validates Pauli Exclusion Principle and the fermionic and bosonic difference between spin and statistic, and have just one set of rules. This set of rules has three axioms:

Axiom (1): All universes are Lorentzian manifolds

Axiom (2): All Lorentzian manifolds are stationary

Axiom (3): Net Curvature on the manifold is a bosonic field. Net are Primes.

The Three Critical Theorems

"Theorem (1) – nature will not allow a prime amount of variation to appear by itself. Define prime to be $(2n+1)$ variations.

1.1) Prime amounts appear in pairs."

Theorem (1) - The physical meaning of that theorem is that bosonic fields cannot be propagated from nowhere. The 8T correlate bosonic propagation to prime net variations of the manifold, and bosons, as we know them, propagate from fermions, which vanish in even number of variations.

Theorem (1.1) – even amount of variations is the result of two prime numbers combined. So to create variation cluster vanishing into matter we need two primes to appear in a pair.

"Theorem (2): Nature will generate force if a **prime net amount of arbitrary variation will appear**. Net variations will appear when combine two amounts of prime variations.

Two does not appear, as it is an even amount of variations, which vanish."

Theorem (2): In continuation of theorem (1), after variation cluster vanished into matter, two distinct elements in threefold combination, a net variation, which is prime can propagate from within it. The feature of the bosonic propagation is their prime number amount of variations, and therefore their expansion across the entire matrix. A boson must propagate from an even amount of variations, which is matter.

Theorem (3): "Each prime pair should have a net variation element N_V proportional to Total Variations value divided by two"

Theorem (3): Each net variation is proportional to the average of the elements in the pair. There could not be net variation $N_V = +(101)$ propagating from $(7,11)$ total variation pair. It does not make sense.

The three theorems in be put in concise and simple manner:

- (1) Bosonic fields cannot propagate from nowhere
- (2) Bosonic Fields propagate from matter clusters
- (3) Bosonic fields are infinite in kind and isomorphic to prime numbers or one.

Theorem (3) was the critical theorem that eventually allowed calculating the value of the fine structure constant and validating the entire framework.

Refuting Magnetic Monopoles

Examine the term describing the electric coupling. We proved majestic three is the electron.

$$[(24 * 5) + (3)] + 5 \rightarrow [(24 * 5) + (e)] + 5$$

Define a magnet as a set of electrons, which spin around as part of a larger cluster of matter.

$$\begin{aligned} \sum_{i=1}^N e_i &\rightarrow \sum_{i=1}^N (3)_i \\ \sum_{i=1}^N e_i &\in \sum_{k=1}^M \delta g_k; \quad M > N \\ \sum_{k=1}^M \delta g_k &= 0 \end{aligned} \tag{2.12}$$

As we did in the 8T thesis, the elements in the term describing matter anti commute, appear in an even number that differ in sign and vanish to zero when summed. However, the spinning electrons are added to a positive summation:

$$\sum_{i=1}^N (3)_i > 0$$

We have two conditions that are not aligned and contradict each other. Both were proven in the 8-Theory to be correct.

$$\sum_{i=1}^N (3)_i > 0 \quad \cap \quad \sum_{k=1}^M \delta g_k = 0$$

The only way to satisfy the second term is to add an opposite spinning cluster so the term would vanish into zero, meaning spinning cluster of electrons in the opposite direction, so (2.12) would be satisfied.

$$\begin{aligned} \sum_{i=1}^T (-3)_i &< 0 \\ \sum_{i=1}^T (-3)_i + \sum_{i=1}^N (3)_i &= 0; \quad T = N \\ \sum_{k=1}^M \delta g_k &= 0 \end{aligned}$$

The Most Symmetrical Interaction is The Weak Interaction

We have proven that the majestic (3), in the case of the electric coupling is the electron. The destabilizing factor yielding a net variation. Overall, the thesis main example was the third element in the series. Therefore the weak interaction did not get enough interaction regarding a very interesting feature it possess.

$$\left(8 * \prod_{V=1}^{V=R} N_V + (3) \right) + N_V = [(8 * 3) + (3)] + 3$$

We can replace the net variation by the majestic three and the correctness of the term will retain. It could explain why the weak interaction is different in terms of its spin, and also allow us to make prediction regarding a fermion, which is analogous to the electron, which can get propagated by the boson of the weak interaction, , $N_V = +(3)$. The overall value is the same; there is a "symmetry" in such a variation, which is not attainable in any term of the coupling constant series. It could mean that the majestic three regarding the weak and the boson, which is propagated, are isomorphic to each other.

$$[(8 * 3) + (3)] + 3 \rightarrow [(8 * (3)) + 3] + (3)$$

$$\left(8 * \prod_{V=1}^{V=R} (3) + N_V \right) + (3)$$

Hermitian Conjunction and Prime Numbers

$$\sum_{i=1}^N \delta g_i = 0$$

$$N \rightarrow \infty$$

$$N = 2n; \quad n \in \mathbb{R}$$

There is no limitation concerning such measurement, we have an even amount of arbitrary variations, which differ in sign and summed as zero. Suppose we had an odd amount of arbitrary variations.

$$N = 2n + 1; \quad n \in \mathbb{R}; \quad 2n + 1 \in \mathbb{C}$$

$$\sum_{i=1}^{N+1} \delta g_i \neq 0$$

So now, the measurement of the fermion cluster become impossible as the manifold is no longer stationary. An elimination of that extra variation must be made. Nature can eliminate it by mirror projections, i.e. Hermitian conjugation. By doing so, the measurement of the fermion cluster will become possible again, or transitioned back to the real field from the complex field.

$$\sum_{i=1}^{N+1} \delta g_i + \sum_{i=1}^{N-1} \delta g_i = 0$$

$$2n + 1 + 2n - 1 = 0$$

So even amount of variation is measurement, additional variation causing the measurement to become impossible, and transition it to the complex field which makes the measurement impossible. To retain the previous state, a mirror projection will be taken.

$$2n \in \mathbb{R}$$

$$2n + 1 \in \mathbb{C}$$

$$2n + 1 + 2n - 1 \in \mathbb{R};$$

Define Hermitian as:

$$\mathcal{H} : \mathbb{C} \rightarrow \mathbb{R}$$

Final Shot at Quantum Relativity

Define an observer, distinct observer, as an arbitrary amount of curvature on the manifold. An infinite series of fermions.

$$\sum_{i=1}^N \delta g_i = 0 \quad (2.12)$$

$$N \rightarrow \infty$$

Define an additional observer, distinct, which differ in the amount of curvature it creates on the matrix. The observer is an infinite series of fermions which overall vanish into matter.

$$\sum_{r=1}^M \delta g_r = 0$$

$$M \rightarrow \infty \quad \cap \quad M! = N$$

Now, analysis of the two observers on equation (1.2). Assume they are measuring the same object, and the entire matrix is null, the entire matrix contain each observer and the measured object. The setting chosen for simplicity sake, as those things will be too complex to analyze in a real physical scenario. Defined the measured object for both observers as:

$$\sum_{k=1}^T \delta g_k = 0$$

Now for the first observer and the measured object, the total arbitrary variation summed as:

$$\sum_{i=1}^N \delta g_i + \sum_{k=1}^T \delta g_k = 0$$

Now for the second observer and the measured object, the total arbitrary variation summed as:

$$\begin{aligned} \sum_{r=1}^M \delta g_r + \sum_{k=1}^T \delta g_k &= 0 \\ \sum_{i=1}^N \delta g_i + \sum_{k=1}^T \delta g_k &\neq \sum_{r=1}^M \delta g_r + \sum_{k=1}^T \delta g_k \\ \left[\frac{\partial \mathcal{L}}{\partial s} \frac{\partial s}{\partial M} \frac{\partial M}{\partial g} \right] \frac{\partial g}{\partial t} \delta g_{ik} - \left[\frac{\partial \mathcal{L}}{\partial s'} \frac{\partial s'}{\partial M} \frac{\partial M}{\partial g'} \right] \frac{\partial^2 g'}{\partial t^2} \delta g'_{ik} &= 0 \\ \left[\frac{\partial \mathcal{L}}{\partial s} \frac{\partial s}{\partial M} \frac{\partial M}{\partial g} \right] \frac{\partial g}{\partial t} \delta g_{rk} - \left[\frac{\partial \mathcal{L}}{\partial s'} \frac{\partial s'}{\partial M} \frac{\partial M}{\partial g'} \right] \frac{\partial^2 g'}{\partial t^2} \delta g'_{rk} &= 0 \end{aligned} \quad (2.13)$$

Those observers will cause the matric to accelerate outward so the object will be observed moving. His velocity is dependent upon the amount of curvature the observer is creating, and so two different observers, different by the above definition, will measure two different distances crossed and two different times for the same object. The reason however, is not for the object itself, it's the different nature of the observers, and in particular the amount of curvature they possess. Now since we proved the yang mills conjecture we have the same propagation speed for all bosons:

$$[(8 * 3) + (3)] + 3 \rightarrow \left[2N_1 + \frac{1}{2}\right] + \frac{1}{2}$$

$$[(24 * 5) + (3)] + 5 \rightarrow \left[2N_2 + \frac{1}{2}\right] + \frac{1}{2}$$

$$[(120 * 7) + (3)] + 7 \rightarrow \left[2N_3 + \frac{1}{2}\right] + \frac{1}{2}$$

The time needed to cross the same matric which accelerated outward in different amounts is different. So, measured time which is different for each observers is quite vivid and a must by using (1.2) and the 8T framework. In fact, using such framework makes relativity notoriously complicated, as everything needs to be taken into account. Everything is causing the matric to vary; it is at a verge of impossible to do at the real world. Our best theories are radically simplified. By "everything", one means every arbitrary variations of fermion in the matric needs to be taken into account, which was not done in that analysis for simplicity sake. The majority of the paper was known to the reader. What is different is the reason of relativity and the analysis of this beautiful idea in the 8T framework, which imposes additional complications, in quantum scale.

The Coupling Constants Series and Total Variations Pairing

$$F_{V=0} = 8 + (1) \quad (1.1)$$

$$F_R \# = \left(8 * \prod_{V=1}^{V=R} N_V + (3) \right) + N_V = 30:128:850:9254.. \quad (1.2)$$

We have obtain the net variation, N_V , as part of a total variation pair, (p_1, p_2) , which we required the sum to be two and three divisible. We gave two examples for the strong interaction:

$$(p_1, p_2) = (5, 13)$$

$$(p_1, p_2) = (7, 11)$$

Two points with regard to those pairs. First, it is commutative, we can replace the elements in the pair and nature will be invariant, the coupling series will hold:

$$(p_1, p_2) \rightarrow (p_2, p_1)$$

Nature is invariant to the actual value of the elements; we can choose any two primes, as long as their sum creating an even number, two and three divisible of certain magnitude, the coupling constant will hold as well. In the 8T thesis we chosen the first pair, it could have worked exactly as well with the second pair.

$$(p_1 + p_2) = S_1$$

$$(p_3 + p_4) = S_1$$

An additional point that was not mentioned in the thesis, the coupling series will hold with any additional amount of primes clustering. We chose the simplest one, two primes in a pair. It could have been four, six or any even number of primes pairing. Any even amount of primes added will yield an even number. Of course the adjustment needed to be made regarding to the division, so we can reach the average value.

$$\frac{\sum_{i=1}^N P_i}{N} = S_{Average} \quad (2.14)$$

If we had four primes pairing, divide by four, six primes divide by six, to reach the average. Of course the average must be two and three divisible, so it could get harder and less likely to find higher numbers of primes pairing which satisfying the condition. It will be impossible to reach the smallest sum in the series with a hundred primes pairing. So for the beginning of the series there could be a limitation.

Fermionic & Bosonic Propagations

$$\left[\frac{\partial L}{\partial s} \frac{\partial s}{\partial M} \frac{\partial M}{\partial g} \right] \frac{\partial g}{\partial t} \delta g - \left[\frac{\partial L}{\partial s'} \frac{\partial s'}{\partial M} \frac{\partial M}{\partial g'} \right] \frac{\partial^2 g'}{\partial t^2} \delta g' = 0 \quad (1)$$

$$\sum_{k=1}^M \delta g_k = 0 \quad (2.12)$$

We partitioned and discretized into a series of arbitrary variations that vanish into matter. We do not have any data regarding the position on the manifold in which those arbitrary variations appear, nor can we assume they possess momenta, as we invoked stationarity on the Lorenzian. M_0 is the connected manifold.

$$M = M_0 \times R$$

In other words, arbitrary variations, which vanish into matter, can be regarded and described by scalar fields that are real, since they have an even amount of variations.

$$\sum_{k=1}^M \delta g_k \in \mathbb{R}$$

Those arbitrary variations, still a subject to additional variance. Such a variance is either prime or one in our framework. These are the variations associated with bosonic propagation. One associated with the strong and each prime with additional coupling term, weak, electric and so on. Because of their prime number feature, they are not vanishing like a fermion scalar but rather as a vector propagation all across. The propagation is associated with a variation of the ∇^2 operator to the setting of the stationary manifold. The bosonic ripple field is then described by:

$$\nabla^2 = \frac{\partial^2 M_x}{\partial^2 g} + \frac{\partial^2 M_y}{\partial^2 g} + \frac{\partial^2 M_z}{\partial^2 g}$$

In other words, it is a vector field propagating all across the matrix, due to its prime number feature, for the second element in the coupling constant series and above. Since the bosonic propagation is associated with prime amount of variations, we can associate it to a complex field, which then require a Hermitian conjunction in order to perform measurement upon. In other words, we can associate bosonic fields to complex vector fields.

$$N_V = 2V + 1; N_V \in \mathbb{P}$$

$$N_V \in \mathbb{C}$$

The Lagrangian Variation

$$\frac{\partial \mathcal{L}}{\partial s} \frac{\partial s}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} - \frac{\partial \mathcal{L}}{\partial s'} \frac{\partial s'}{\partial M} \frac{\partial M}{\partial g'} \frac{\partial^2 g'}{\partial t^2} = 0 \quad (1)$$

$$\frac{\partial g}{\partial t} = 0 \quad - \frac{\partial^2 g'}{\partial t^2} = 0$$

$$\mathcal{L} = T - V$$

How can a representation of the Lagrangian be made on the varying Lorentz manifold, is the main question one will analyze in this short assay. Of course, the real answer to that question is one does not know. However, educated guess will be made. The kinetic term could be associated to the outward matric expansion, due to $(\partial g / \partial t)$ term, which is synonymous with energy in physical theories.

$$\left(\frac{\partial g}{\partial t} \right) = \left(\frac{\partial E}{\partial t} \right) = \frac{\partial^2 E'}{\partial t^2}$$

That is the kinetic term. A Ricci Tensor overtime, yielding an energy expansion outward causing a matric acceleration on the object generating the energy. That is the main equation that was derived by putting an Einstein manifold in Euler Lagrange Equation. Now, what is the potential energy in the 8T framework? In physical theories, the potential is associated with the mass, which is certain feature of the object itself. In the 8T we do not have any objects, the objects are manifestations of discretizing and partitioning the term $\delta g \rightarrow \sum_{i=1}^K \delta g_i = 0$ in equation (1.) to vanish into fermions. How can we translate that into a potential term?

$$\left[\frac{\partial \mathcal{L}}{\partial s} \frac{\partial s}{\partial M} \frac{\partial M}{\partial g} \right] \frac{\partial g}{\partial t} \delta g - \left[\frac{\partial \mathcal{L}}{\partial s'} \frac{\partial s'}{\partial M} \frac{\partial M}{\partial g'} \right] \frac{\partial^2 g'}{\partial t^2} \delta g' = 0 \quad (1)$$

Since we would like to measure how much arbitrary variations the object we measure contain, and those arbitrary variations are two and three divisible to vanish into matter, two distinct elements which created threefold combinations, to get a measure of the amount of arbitrary variations, the action needed is the following Transformation:

$$\sum_{i=1}^K \delta g_i = 0 \rightarrow \sum_{i=1}^K |\delta g_i| = \mathcal{V}_{8T} \quad (2.12)$$

The idea, whether correct or not, was to take the absolute number of varying elements participating in the construction of the object. The operation was done via the insight we gained in previous paper, two distinct elements which differ in sign, so by eliminating the minus sign we can estimate how many arbitrary variations appear in the cluster. To sum up, The energy, causing outward acceleration minus the total amount of arbitrary variations constructed in the cluster.

$$\mathcal{L} = \left(\frac{\partial g}{\partial t} \right) - \sum_{i=1}^K |\delta g_i| \quad (3)$$

Cluster Decomposition

In quantum field theory, one learns that the connected part of the S matrix must vanish. Distinct events do not effect each other.

$$S_{\beta\alpha}^c \rightarrow 0$$

What is the equivalent of the cluster decomposition principle on the Lorentz manifold (M, g_E) with signature (3,1), invoked stationary, $M = M_0 \times R$, is the subject of this paper.

$$\frac{\partial L}{\partial s} \frac{\partial s}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} - \frac{\partial L}{\partial s'} \frac{\partial s'}{\partial M} \frac{\partial M}{\partial g'} \frac{\partial^2 g'}{\partial t^2} = 0 \quad (1)$$

Since the manifold experience arbitrary variations that vanish into matter, all across the matric, the smoothness of the matric must be taken into account. Bosonic propagation described by the delta must cross the metric before reaching a distinct event on the manifold. The result of such a construction would be that only arbitrary variations that vanished relativity closed to each other, will have an effect on each other. Suppose we had two distinct arbitrary variations, that is by discretizing and partitioning the term δg in equation (1.11), as was done in previous papers of the 8T, to proof that these are fermions:

$$\delta g = \sum_{i=1}^{\infty} \delta g_i = 0 \quad (2.12)$$

We impose two conditions equivalent to the cluster decomposition in QFT. Those conditions are synonymous with saying that distinct events will not affect each other. Consider two arbitrary variations

$$\delta g_i + \delta g_{i+1}$$

Suppose those appeared at distinct parts of the matric, M_μ is a four vector isomorphic to the arbitrary variation with the matching index δg_i :

$$M_\mu \rightarrow M(x_i, y_i, z_i, t_i)$$

$$\delta g_i \rightarrow M_\mu$$

Same for the additional variation, δg_{i+1} , a four vector M_ν , the condition than requires that:

$$M_\mu - M_\nu \leq \epsilon$$

$$\epsilon \rightarrow 0$$

In other words, two arbitrary variations must appear close to each other on the matric, at very short time interval. That is synonymous with the quantum field theory statement of the connected part of the amplitudes to vanish. The two conditions are synoptic in the four vector. The arbitrary variations should appear close on the matric spatial dimensions and at a short time interval.

The Perfect Symmetry of Hadrons

$$F_{V=0} = 8 + (1) \quad (1.1)$$

$$F_R \# = \left(8 * \prod_{V=1}^{V=R} N_V + (3) \right) + N_V = 30:128:850:9254.. \quad (1.2)$$

$$8 + (1):(24 + (3)) + 3:(120 + (3)) + 5:(840 + (3)) + 7$$

Now, in the 8T framework, up to this point, we nicknamed the left term of each element in the coupling as a variation cluster, which is divisible by two and three to perfectly vanish into matter. This variation cluster is destabilized by the majestic three, causing a net variation to appear, or in other words, to boson propagation from the fermion. Recently one noticed a very interesting fact, that the left terms of the coupling constant series for the weak interaction are identical to densest packing D_4 highest kissing number that is 24. So all the left coupling terms are actually 4D spheres, leading to a propagation of the electron. That may sound outrageous but not in the 8T, as we only have 4D manifold, three spatial and one temporal. By looking at the coupling constant in that light, we can also regard the hadron as possessing an extreme density, as it has the highest kissing number in 4D, and the electron is not bound to it but revolves around it, as the majestic is a separate term. The following apply to each other term:

$$24 * 5 \rightarrow 120$$

$$24 * 5 * 7 \rightarrow 840$$

Notice that those numbers are associated with highest kissing numbers in higher dimensions.

$$E_8 \rightarrow 240 = 120 * 2$$

$$p_{12} \rightarrow 840$$

Of course, ignoring the higher dimensions complexity and focusing on the part of the highest kissing numbers, we can reach an insight, those fermionic clusters in each term are most dense, in agreement with what we know about the structure of the fermions, and in particular the hadron. Also, notice that those higher dimensions are scalar four multiples, which as one believes, means that should appear on the manifold eventually. The highest kissing number in D_4 is the base to all other kissing numbers at those higher dimensions. By looking at the coupling constant series, then we can correlate the manifold and validate it has only four dimensions, since all higher terms are the dimension four multiples of the kissing number, 24. And thus there could not be more than four dimensions on our manifold. There are of course other manifolds, which according to the series are four dimensional as well, interacting with our own as given by the main equation of the 8T. But by coupling constant series, it is possible to derive why the manifold has exactly four dimensions, because of the kissing number of the second term and above.

In addition, the number 24 is associated with the leech lattice, which has most density within a certain dimensional range, is intimately related to this number. In the 8-Theory however there is no use of any lattices. Rather we use variations. Notice the 24 is perfectly to and three divisible to vanish into matter. There is no additional variation left alone. The hadron is perfectly compact and most dense because of that trait. Than it is destabilized by additional term, the element in which we called the majestic three. The point one was trying to make is that the perfect symmetry of the hadronic structure is preserved along each coupling term, i.e. each interaction. In addition, it is than lessen by the electron, i.e. the additional element in the third coupling term. And either the electron is also the cause of that symmetry break in all other terms or electron analogues field.

$$\frac{24 * N_V}{mod(6)} = 0; \quad (3.12)$$

$$N_V = 2V + 1; V \geq 1$$

$$N_V \in \mathbb{P}$$

If it was any other number than 24, than the symmetry of the hadrons was not perfect, as equation (1.2) will not hold. The symmetry is breaking due to an external element added by the higgs field from the second element and above, the majestic three. It is currently unclear whether this element is the same for each of the coupling terms. For the electric, it was proven the electron. However, for the weak interaction term and higher terms it could be an electron analogues particle manifested in the element three as mentioned in the previous paragraph and again, it's so important one wanted to emphasize it here as well. There are two main points two take from this short assay. The first is the perfect symmetry of the hadronic structure due to its numerical features. The second point is that the symmetry is breaking from an external element not from within the hadronic structure, due to the higgs field, inserting the majestic three.

The Feynman Diagrams

$$F_{V=0} = 8 + (1) \quad (1.1)$$

$$F_R \# = \left(8 * \prod_{V=1}^{V=R} N_V + (3) \right) + N_V = 30:128:850:9254.. \quad (1.2)$$

$$N_V = 2 \left(V + \frac{1}{2} \right); \quad V \geq 0$$

$$[(8 * 3) + (3)] + 3 \rightarrow \left[2N_1 + \frac{1}{2} \right] + \frac{1}{2}$$

$$[(24 * 5) + (3)] + 5 \rightarrow \left[2N_2 + \frac{1}{2} \right] + \frac{1}{2}$$

$$[(120 * 7) + (3)] + 7 \rightarrow \left[2N_3 + \frac{1}{2} \right] + \frac{1}{2}$$

Examine the term describing the electric coupling. We proved majestic three is the electron in the 8-Theory thesis.

$$[(24 * 5) + (3)] + 5 \rightarrow [(24 * 5) + (e)] + 5$$

$$[(24 * 5) + (3)] + 5 \rightarrow [(24 * 5) + (e)] + \gamma$$

$$e \searrow \rightarrow (\gamma) \rightarrow e \nearrow$$

$$(+3) \rightarrow (\gamma) \rightarrow (+3)$$

$$(+3) \rightarrow (+5) \rightarrow (+3)$$

$$(+3) + N_V = (+3) + 5 = 8$$

$$8 = \text{even}$$

$$\text{even} = 0$$

The electron, represented as the majestic three combined with the net variation yielding an even amount of variation that vanish. That is synonymous with saying that the electron has absorbed the photon. The conservation of variation ensure that no electron can disappear from the manifold. However, as the combination of N_V and the electron, i.e. the three yielding an even, there has to be a vanishing of certain sort into the electron. It is moved into an excited state, vanishing of curvature, $(\gamma) = (+5)$ into the receiving electron, which causes the deflection in trajectory. Using the numerical trait and insight

gained by the coupling constant series, by the 8T framework, it is possible to add an additional layer to the Feynman diagrams and interactions among bosons and fermions in what seems as a very simple and elegant manner. What can be derived about the nature of the electron using the coupling constant representation? First of all, it is bounded by the bracket, it cannot escape and behave as the net variation, i.e. the photon. Despite the fact that both elements represented by a prime. Second, the electron is represented as a prime number, three, which cannot vanish into matter, but also cannot propagate as a bosonic fields across the matrix its behavior than would propagation across the nuclei, in agreement with current understanding about the probabilistic behavior of that particle. There is no data regarding the current position, momenta, orbitals, no physical data of any sort is manifested in the 8-theory. An additional way to analyze it is to say that the electron blends in the hadronic cluster, $[(24 * 5) + (3)]$. The hadronic cluster is closed and represented in a closed term within the bracket. The summation of the term is perfectly suitable to vanish into matter.

The Axis of Evil

The main equations in our new framework:

$$\frac{\partial L}{\partial s} \frac{\partial s}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} - \frac{\partial L}{\partial s'} \frac{\partial s'}{\partial M} \frac{\partial M}{\partial g'} \frac{\partial^2 g'}{\partial t^2} = 0 \quad (1.1)$$

$$\frac{\partial g}{\partial t} = 0 \quad - \frac{\partial^2 g'}{\partial t^2} = 0$$

Describing the Lorentz manifold (M, g_E) with signature (3,1), invoked stationary, $M = M_0 \times R$. Equation (1.1) satisfy the Einstein principle of equivalence and expends it to a cause and effect relationship. Invoking a stationary manifold, any amount of curvature on it, will yield an outward acceleration of the matric. In that sense, it is different from general relativity, as there is no need to insert the cosmological constant as a separate entity. Using that equation, we built a new way to explain relativity by saying that two distinct observers will cause different accelerations of the matric, and so, by measuring the same object, will reach different times and distances.

In our theory, the manifold has a varying matric according to a varying topology. The subtle idea is that the manifold has a compact topological space that is accessible from every point given high enough energy. Such space covers every point in matric space. Such a space is what makes the theory works, it is the space keeping the manifold stationary and with the second condition causing it to accelerate outward. Since there are no coordinate to such space, it is the same everywhere, and since every point in the matric is connected to it, there could be the illusion that each point in space was the point in which something cosmologically significant has accrued at singularity. Not the whole topological space is satisfying the condition, $\partial g / \partial t = 0$ there are arbitrary variations in that space which vanish into matter on the matric, we have proved it in previous papers. Each net variation than is isomorphic to the prime numbers or to the number one, and thus we were able to prove the coupling constant series, presented in equation(1.1) and (1.2). The point of this short assay is the fact that there is an underlining space, which is invariant to matric coordinate and covers the entire matric. We know it covers the entire matric as the manifold is connected to the topological space but no spatial coordinates are given in equation (1.1). The topological space is than invariant, and the equation is really a right to left chain of the order. Notice that the chain in equation (3.12) is exactly describing the order in which things are happening in cosmological scales.

$$\frac{\partial L}{\partial s} \leftarrow \frac{\partial s}{\partial M} \leftarrow \frac{\partial M}{\partial g} \leftarrow \frac{\partial g}{\partial t} \quad (3.12)$$

Reasoning Bosonic Probabilities

$$F_{V=0} = 8 + (1) \quad (1.1)$$

$$F_R \# = \left(8 * \prod_{V=1}^{V=R} N_V + (3) \right) + N_V = 30:128:850:9254.. \quad (1.2)$$

$$8 + (1): (24 + (3)) + 3: (120 + (3)) + 5: (840 + (3)) + 7 \dots$$

Examine the term describing the electric coupling. We proved majestic three is the electron in the 8-Theory thesis. The photon is represented as net variation, which is unbound. It is free to propagate all across the manifold.

$$[(24 * 5) + (3)] + 5 \rightarrow [(24 * 5) + (e)] + 5$$

$$[(24 * 5) + (3)] + 5 \rightarrow [(24 * 5) + (e)] + \gamma$$

Suppose such a photon just propagated from the electron. i.e. the majestic three. The meaning of such an occurrence is that there is a net curvature that is unbound on the manifold. Such curvature will effect all other potential propagation toward itself. It will create a pull effect on other potential boson propagating from fermions. That is in agreement with what we know about the commutation relation of bosons, and the fact that the probability to find a boson increase if there is already a boson in a certain position of the matrix. The innovative part of this paper and the main point to take is the new setting, a in which a photon itself is a net curvature causing other curvature propagating at later time to converge to its position. When analyzed via the new framework it than becomes quite easy to understand what is going on at that fundamental level.

$$\sum_{i=1}^N \gamma_i > 0 \quad (3.13)$$

$$\sum_{i=1}^N \gamma_i = \sum_{i=1}^N \delta g_i > 0 \quad (3.13.B)$$

The point of view presented is not presented in quantum field theory framework, the methods they use to describe the commutation and anti-commutation is VOA, vertex of algebra, and there is simply no way to imagine or to grasp the intuitive reason for the such a behavior. By using an approach combining manifolds and variation, i.e. Euler Lagrange, it is possible to explain the behavior of bosons in an intuitive and simpler fashion. It is possible to state that each boson is creating a "gravitational effect", i.e. curvature on the manifold, and thus increase the probability of arrival for other bosons to itself.

The Conservation of Variation

$$F_{V=0} = 8 + (1) \quad (1.1)$$

$$F_R \# = \left(8 * \prod_{V=1}^{V=R} N_V + (3) \right) + N_V = 30:128:850:9254.. \quad (1.2)$$

$$N_V = 2 \left(V + \frac{1}{2} \right); \quad V \geq 0$$

$$8 + (1):(24 + (3)) + 3:(120 + (3)) + 5:(840 + (3)) + 7 \dots$$

$$8 + (1) + 3 - (1v) : (24 + (3)) + 3: (120 + (3)) + 3$$

In the paper about the interactions dynamic nature, we varied the first and the third interactions, i.e. the strong and the electric, in their N_V element, so all the net variations will align on the same integer. The important point, which was not mentioned, is that the net variations varying their position among the terms are confined within the manifold. In other words, it is conserved. That is also the case with the gravitational coupling, which as far as the 8T can predict, is a result of two net variations added to the original net variation. The data regarding the nature of gravity came from the second representation, i.e. the spin representation of the coupling constant equation.

$$[2N_{gravity} + 2] = \left[2N_{gravity} + \frac{1}{2} \right] + \frac{1}{2} + \frac{1}{2} + \frac{1}{2} = [(2N_{gravity}) + (3)] + N_{V1} + N_{V2} + N_{V3}$$

We can put the conservation law in rigor and construct an appropriate theorem:

Theorem (1.0) – The sum of net variations on all the coupling elements cannot escape the manifold.

Theorem (1.1): The sum of all net variations increase with time.

$$\oint_{t=0}^{t=Z} (dM)(M_0 \times R) \left(\sum_{V=0}^{\infty} N_V \right) \in M \quad (3.16)$$

$$Z \rightarrow \infty$$

If one constructed properly, one summation of the net variation to each V across the entire manifold matric, over time, must belong to the manifold itself and cannot decrease. It could be related to the second rule of thermodynamic, the entropy rise alongside the net variations overtime. Of course, the total variations grow much faster, but that was not the subject of this paper. The point was to emphasize that the sum of net variation is bounded to the manifold, despite the fact it grows with time.

Bosonic Strings - Cyclic Groups

$$F_{V=0} = 8 + (1) \quad (1.1)$$

$$F_R \# = \left(8 * \prod_{V=1}^{V=R} N_V + (3) \right) + N_V = 30:128:850:9254.. \quad (1.2)$$

$$8 + (1): (24 + (3)) + 3: (120 + (3)) + 5: (840 + (3)) + 7 ...$$

$$8 + (1)$$

$$[(8 * 3) + (3)] + 3 \rightarrow \left[2N_1 + \frac{1}{2} \right] + \frac{1}{2}$$

$$[(24 * 5) + (3)] + 5 \rightarrow \left[2N_2 + \frac{1}{2} \right] + \frac{1}{2}$$

$$[(120 * 7) + (3)] + 7 \rightarrow \left[2N_3 + \frac{1}{2} \right] + \frac{1}{2}$$

Cyclic groups in mathematics are represented by the following, if a set of elements is generated by one single element, than we have a cyclic group. Since all the bosonic fields or net variations in the 8T are generated by the same element, i.e. the majestic three, than there is in this framework an infinite cyclic group. Define the majestic three as the generator:

$$(3) \rightarrow \mathcal{M}$$

$$\mathcal{B} = \{N_1 = (+3)\mathcal{M}, N_2 = +(5)\mathcal{M} \dots\} \quad (3.16)$$

By representing the propagation in such fashion, we can state that since the bosons are propagations are part of an infinite cyclic group, the sub elements of that cyclic group are cycles themselves. We have proven the representation of the coupling constant series in the thesis:

$$[(24 * 5) + (3)] + 5 \rightarrow [(24 * 5) + (e)] + \gamma$$

$$\mathcal{B} = \{N_1 = (+3)\mathcal{M}, N_2 = +(\gamma)\mathcal{M} \dots\}$$

Therefore, that is a proof that bosonic net variations are cycles, or in physical theories, bosonic particles are in fact closed strings. That is because they are generated by the same element.

Curvature Absorptions

$$[(24 * 5) + (3)] + 5 \rightarrow [(24 * 5) + (e)] + \gamma$$

$$(3) \rightarrow \mathcal{M}$$

$$\mathcal{B} = \{N_1 = (+3)\mathcal{M}, N_2 = +(5)\mathcal{M} \dots\}$$

$$\mathcal{B} = \{N_1 = (+3)\mathcal{M}, N_2 = +(\gamma)\mathcal{M} \dots\}$$

All of this was covered in previous papers. The majestic three is a generator of a cyclic group of bosonic net curvature propagations, isomorphic to the primes or one. We made a Feynman diagram using the new framework:

$$e \searrow \rightarrow (\gamma) \rightarrow e \nearrow$$

$$(+3) \rightarrow (+5) \rightarrow (+3)$$

$$(+3) + N_V = (+3) + 5 = 8 \rightarrow 0$$

The point of this paper is that in order to understand how the manifold vary, there are to be a summation of all curvature absorptions and emissions. As an electron absorb a photon, the manifold gets more flat, as $N_2 = +(5)$ just vanished into the electron and vice versa. By looking at clusters of photons in unit matrix, it is also possible to estimate how much curvature exits on the manifold. As bosons are net variations unbound, it was derived that preciously for that reason the probability of boson arrival after a boson is propagated.

$$\sum_{i=1}^N \gamma_i > 0 \quad (3.13)$$

$$\sum_{i=1}^N \gamma_i = \sum_{i=1}^N \delta g_i > 0 \quad (3.13.B)$$

The point is, we can use space- time summation and in particular, the distribution of fermions to bosons to estimate how curved the matric, or how it varies over time. It is vividly clear that a real world estimation is at the verge of impossible, but a rough evaluation is always within reach.

$$\sum_{i=1}^N \gamma_i \rightarrow \mathcal{P}$$

$$\frac{\partial \mathcal{P}}{\partial t} - \frac{\partial M}{\partial g} = 0 \quad (3.17)$$

Light is Bending Space-Time

$$\left[\frac{\partial L}{\partial s} \frac{\partial s}{\partial M} \frac{\partial M}{\partial g} \right] \frac{\partial g}{\partial t} \delta g - \left[\frac{\partial L}{\partial s'} \frac{\partial s'}{\partial M} \frac{\partial M}{\partial g'} \right] \frac{\partial \partial g'}{\partial \partial t} \delta g' = 0 \quad (1)$$

$$\sum_{i=1}^N \delta g_i = 0 \quad (2.12)$$

$$\sum_{i=1}^M \gamma_i > 0 \quad (3.13)$$

$$\sum_{i=1}^M \gamma_i = \sum_{i=1}^M \delta g_i > 0 \quad (3.13.B)$$

By putting the Lorentz manifold in Euler- LaGrange framework and allowing arbitrary variations to appear, in which we require to vanish, in the 8T we discretized and partitioned the term (2.12) and were able to prove that arbitrary variations of the manifold vanish into matter. Each net variation or net curvature is isomorphic to a bosonic field propagation. In particular the boson associated with photon propagation is $N_V = +(5)$. Such propagation is than yielding a positive summation, i.e. a positive curvature by (1.15), so fermion clusters are flat according to the 8T framework, but bosonic propagations are curvature on the manifold. The weird and unexpected result is that bosonic fields are deflecting fermion clusters and not the opposite as believed by GR. It is an unexpected result, but up until recently we thought there are only four forces, and such thought lead to thinking that physics can be unified.

The Riemann Hypothesis – Proof

Define a Lorentz manifold

$$\mathbf{s} = (\mathbf{M}, \mathbf{g})$$

Use it to assemble a Lagrangian and require it to be stationary:

$$L = (s, s', t)$$

$$\frac{\partial L}{\partial s} - \frac{\partial L}{\partial s'} * \frac{d}{dt} = 0$$

Allow arbitrary variations of the manifold. Ensure it will vanish:

$$\omega \mathbf{s} = 0$$

Turn it to a series of arbitrary variations:

$$\omega \mathbf{s} = \omega \mathbf{s1} + \omega \mathbf{s2} + \omega \mathbf{s3} \dots$$

If there are only four elements in the series, and we require them all to vanish, than we can allocate two pluses and two minuses:

$$\omega \mathbf{s1} + \omega \mathbf{s3} > 0$$

$$\omega \mathbf{s2} + \omega \mathbf{s4} < 0$$

If

$$\omega \mathbf{s1} + \omega \mathbf{s3} + \omega \mathbf{s2} + \omega \mathbf{s4} \neq 0$$

Than the overall series cannot vanish, by that logic we need equal amounts of plus and minuses. The overall amount must be even and summed as zero.

Suppose that we had three distinct elements, two pluses and minus:

$$\omega \mathbf{s1} + \omega \mathbf{s3} + \omega \mathbf{s2} > 0$$

or

$$\omega \mathbf{s1} + \omega \mathbf{s3} + \omega \mathbf{s2} < 0$$

Demanding the series to vanish this forbid this result, and so there could not be three distinct elements in the series, else the overall series will not vanish. As a result of those sceneries, we require the series to have an even amount of variation elements, manifesting as two distinct elements in the series, which differ in sign. If we allow those sub elements in the series to vary as well, and by the above reasoning, there are only two elements in the series, they are varying in a discrete way, or forming a group. Let it be only four elements in the series and one of the pluses just changed its nature

$$0: \omega \mathbf{s1} \rightarrow \omega \mathbf{s2}$$

$$\omega \mathbf{s1} + \omega \mathbf{s1} + \omega \mathbf{s2} + \omega \mathbf{s2} = 0$$

To:


$$\omega s1 + \omega s2 + \omega s2 + \omega s2 \neq 0$$

There must be a way to bring it back to where it was, so the overall series can vanish, it takes another map, on the varying element to bring it back to where it was.

$$Y: \omega s2 \rightarrow \omega s1$$

Therefore, to bring an element to itself given only two varying elements in the series we need two distinct maps, which attach a varying element to itself, by a threefold combination.

$\omega s1(O) \omega s2(Y) \omega s1$ For example. Even though the sub elements in the series are varying, the overall series can vanish. Now, count all the ways of possible combinations of those elements. We are going to analyze by the integral signs. Since it is a group, there is a natural map, which change an element to itself. One built his analysis firstly on those natural maps. The first two combinations are by the natural maps and one used them to build the other combinations. Overall, there are eight such combinations and additional one arrow combination, which yield (333) here is how one built it, starting from those two natural maps. (Arrows to variations, colors to pairings):



Now that we have a series of $2N$ elements, varying to one another and forming threefold combinations, **which we require to vanish at end**, we can set the stage for a proof of primes. **Define:** P^m as the set of $\{2, 3\}$ as "minimal primes". In addition, all the other primes to be in a set of P_h as meant "prime higher".

Define $P_h = \{2n + 1\}$ not divisible by P^m as "prime higher" set – $2n$ taken as amount of Lorentz manifold arbitrary variations.

$\{2n + 1\}$ as an odd amount of variations not divisible by minimal primes

$$P_t = P_h + P^m ; \text{ to be the set of all primes}$$

Define a functor V on P_h :

$$V: \text{set} \rightarrow \text{ring}$$

Analyze any multiplication or addition combination of P_h on the ring. Let the ring exist on a Lorentz manifold, a topological space.

Multiplication:

Define T to be a number aspiring infinity: $T \rightarrow \infty$ Multiply an **even or odd** series aspiring infinity of distinct higher primes to obtain:

$$[(2n_1 + 1)(2n_2 + 1)(2n_3 + 1) \dots (2n + 1)] =$$

$$2 \left[T((n_1 n_2 \dots)) + (n_1 + n_2 + n_3 \dots) + \frac{1}{2} \right]$$

$$= 2([T((n_1 n_2 \dots)) + N(s) + 1/2])$$

$$N(s) = (n_1 + n_2 + n_3 \dots) = 0$$

As sums of even amounts of arbitrary variations vanish. Since all the elements are two multiples, they all vanish. Final form:

$$2 \left([T(n_1 n_2 \dots)] + \frac{1}{2} \right)$$

Addition

Add any infinite **even series** of distinct higher primes to obtain

$$(2n_1 + 1) + (2n_2 + 1) + (2n_3 + 1) \dots = [2(n_1 + n_2 \dots) + \text{even}] =$$

$$[2(n_1 + n_2 \dots)]$$

as even = 0.

Prime cannot form, as even amount of variations vanish exactly to zero. That is the reason the paper begins with deriving fermions, their anti-commutation relation. Even amount of distinct higher primes added will never form a prime.

Add any infinite **odd series** of distinct higher primes to obtain

$$(2n_1 + 1) + (2n_2 + 1) + (2n_3 + 1) \dots =$$

$$[2(n_1 + n_2 \dots) + \text{odd}] =$$

$$[2(n_1 + n_2 \dots) + (\text{even} + 1)] \quad (10)$$

However, even amounts of arbitrary variations vanish:

$$\text{even} = 0$$

$$[2(n_1 + n_2 \dots) + 1] \text{ or:}$$

$$2[n_1 + n_2 \dots + 1/2] \quad (11)$$

Category transformations

Define a functor on "Primes higher" ring

$$G: \text{ring} \rightarrow \text{group}$$

All "primes higher" are forming a closed non-abelian group with 1/2 as generator. The condition to group forming is to have an odd amount of primes under addition and eliminating even amounts of arbitrary variations taken as an axiom. Define additional functor

$$G': \text{group} \rightarrow \text{set}$$

Add the sets:

$$P_h + P^m = P_t ;$$

Define a functor on P_t :

$$G'': \text{set} \rightarrow \text{group}$$

All primes are forming a non-abelian group of generator $1/2$. Minimal primes are part of the group by nature of the proof, defined technically to be prime. Primes are forming a non-abelian group under addition and multiplication. The condition to satisfy is to have an odd amount of primes under operation of addition. No matter how far into infinity we will go, the framework of vanishing of even amount of variations will ensure that all primes take the same form – aligned on $\frac{1}{2}$. Setting the stage and **examining primes not as numbers, but rather as arbitrary variations of a manifold**, which vanish in pairs of even variations, we are able to show primes to form a non-abelian closed group under $2(n+1/2)$. Final functor on the total group of primes:

$$\text{Riemann: Group} \rightarrow \text{ring}$$

All primes are forming an infinite ring on the critical line of $1/2$ and only there.

End of proof.

Visualization - Photon Emission

$$8 + (1): (24 + (3)) + 3: (120 + (3)) + 5: (840 + (3)) + 7 \dots$$


$$[(24 * 5) + (3)] + 5 \rightarrow [(24 * 5) + (e)] + \gamma \quad (4)$$

$$(e) = (3) \rightarrow \mathcal{M}$$

$$\mathcal{B} = \{N_1 = (+3)\mathcal{M}, N_2 = +(5)\mathcal{M} \dots\} \quad (4.1)$$

$$\mathcal{B} = \{N_1 = (+3)\mathcal{M}, N_2 = +(\gamma)\mathcal{M} \dots\}$$

Equation (4) to describe the process of emission due to the invariant three, proven the electron, assumed the electron for each higher term in the coupling series. equation (4.1) describe the invariant three as the generator of a cyclic group, meaning all bosonic propagations are sub elements of that group and so we prove they are closed cycles themselves. Therefore, we can draw the interaction between two electrons and a photon emission in the following way:




As was proven, they cannot move at the same orientation of the distortion due to their prime number feature, combined together there will be a vanishing and so the coupling series than would not make sense. The end conclusion would than imply that the boson propagated from nowhere which is impossible.

Interference

$$[(24 * 5) + (3)] + 5 \rightarrow [(24 * 5) + (e)] + \gamma$$



The image above represent a net curvature on the Lorentz manifold, in that specific case, it's the photon associated with $N_V = (+5)$ net variations, and total 128 variations. Suppose that we perform the two slits experiment and open an additional route for net curvature. this is the visualization of what could happen according to our new theory:



There are two ways to explain. The first is to say that two opposite but similar in magnitude curvature occupying the same space will have a segment of mutual cancelation. If we define ripple operators \emptyset from a starting area to another area, the mutual area of both will be the amount of interference.


$$\emptyset: A \rightarrow B$$

$$\emptyset: A' \rightarrow B$$

Interference will accrue at the manifold segment that is mutual to both starting point. Define the interference operator:

$$\approx: A \cap A' \quad (3.18)$$


Quark Visualization



Imagine constant variation so the overall construction is curvature varying, according to the combination where will be a pairing according the graph presented in the 8 theory thesis or the group suggested by the particle physicist Gell Mann. Each arrow in the visual is a representation of the gluon, or the first element in the coupling constant primordial function.

Visualization of the Multiverse

$$\frac{\partial \mathcal{L}}{\partial S_1} \frac{\partial S_1}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} - \sum_{n=2}^{\infty} \frac{\partial \mathcal{L}}{\partial S_n} \frac{\partial S_n}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} = 0 \quad (2)$$



Strong Interaction – The Electron

$$F_{V=0} = 8 + (1) \quad (1.1)$$

$$F_R \# = \left(8 * \prod_{V=1}^{V=R} N_V + (3) \right) + N_V = 30:128:850:9254.. \quad (1.2)$$

$$N_V = 2 \left(V + \frac{1}{2} \right); \quad V \geq 0$$

$$N_V \in \mathbb{P} \cup (+1); \quad \mathbb{P} \rightarrow \text{Set of Primes}$$

$$N_V = P_{max} \in [0, \mathbb{R}] \cup (+1); \quad P_{max} \in \mathbb{P}$$

$$8 + (1): (24 + (3)) + 3: (120 + (3)) + 5: (840 + (3)) + 7 \dots$$

$$[(24 * 5) + (3)] + 5 \rightarrow [(24 * 5) + (e)] + \gamma$$

The main argument of this short essay is that it is possible to regard each higher coupling terms as the strong interaction being destabilized in ever-growing fermion formations. It's the electron that has so much significance in the coupling constants series. Back in the day, when author derived the coupling series, in the thesis he believed that each term would have unique destabilizer, but now it seems very clear that such an assumption is quite likely wrong and eventually will lead to complexity that is not needed. Another way to state it is that three is isomorphic to itself. What is varying is the size of the fermionic cluster and the magnitude of the net curvature. The shift in understanding manifested itself in toward the end of the thesis but still it is important to clarify to avoid confusion among readers. It is also possible to represent the coupling, as you already know, in the form of spin representations by setting it on the prime critical strip.

$$[(8 * 3) + (3)] + 3 \rightarrow \left[2N_1 + \frac{1}{2} \right] + \frac{1}{2}$$

$$[(24 * 5) + (3)] + 5 \rightarrow \left[2N_2 + \frac{1}{2} \right] + \frac{1}{2}$$

$$[(120 * 7) + (3)] + 7 \rightarrow \left[2N_3 + \frac{1}{2} \right] + \frac{1}{2}$$

To solidify the statements made in previous papers, the variance in that representation is the fermionic clusters, represented in the right of each term, and the net variation, or net curvature that is prime or one. The conclusion if one is correct is the electron is destabilizing larger and larger fermionic cluster yielding an infinite succession of net curvature on the manifold, which causes the endless process of clustering. One prefer that version, as it is simpler than to assume that each term would have a unique destabilizer. As the fermionic cluster gets much more massive in rate, the net curvature than becomes less significant, preciously the idea behind the principle of least variation.

Virtual Curvatures

In calculus of variations, we have the procedure of the following for the vanishing of virtual displacements within a massive cluster. Such a procedure makes description of motion rather simple, as we do not need to describe the innate motion of a static body. Similar in a sense to the Laplace operator.

$$\sum_{i=1}^N F_i dr_i = 0 \quad (3.19)$$

What would be the equivalent statement in the 8-theory? As we do not use force in the innate description of the theory, all we have is net curvature, N_V , on the Lorentz manifold, which was invoked stationary by the Lagrangian operator. We also did not use radius per se, it is different from the Riemann line element in which we associate curvature. One will suggest the following analogue for the equation (1.1):

$$\sum_{i=1}^N \delta g_i \partial L_i = 0 \quad (3.19)$$

The sum of all arbitrary variations per varying manifold unit length is summed as zero. As we say variations, we mean curvature, so the sum of arbitrary curvatures is taken to zero. We can similarly use that construction in the same manner and for the same purposes used in calculus of variations, to avoid describing the inner motions of a static body.

Curvature Knots

$$F_{V=0} = 8 + (1) \quad (1.1)$$

$$F_R \# = \left(8 * \prod_{V=1}^{V=R} N_V + (3) \right) + N_V = 30:128:850:9254.. \quad (1.2)$$

$$N_V = 2 \left(V + \frac{1}{2} \right); \quad V \geq 0$$

$$N_V \in \mathbb{P} \cup (+1); \quad \mathbb{P} \rightarrow \text{Set of Primes}$$

$$N_V = P_{max} \in [0, \mathbb{R}] \cup (+1); \quad P_{max} \in \mathbb{P}$$

Suppose that instead of a prime number as in equations (1.3) and (1.31) describing the weak and the electric, we would have a number that is odd, which could be a composite of odd number primes. Define the odd number function:

$$\Phi_n = 2n + 1 \quad (3.2)$$

$$n \in \mathbb{R}; \quad \Phi_n \notin \mathbb{P}$$


$$\mathbb{P} \rightarrow \text{set of primes}$$

So Φ_n is a series of odd numbers that replace only the external N_V in the coupling constant series. The new series is now described by:

$$\left[2N_1 + \frac{1}{2} \right] + \Phi_{n1} \quad (32.1)$$

$$\left[2N_2 + \frac{1}{2} \right] + \Phi_{n2} \quad (3.2)$$

Since Φ_n is not a prime it cannot act as a bosonic ripple field on the matrix tensor. Since it is on an even number, divisor of modulo (6) it cannot vanish into matter. It is a composite of prime, or a composite of net curvature, and because it is a composite, which is stable on the matrix tensor, we will have a curvature which is time- invariant, not matter like nor boson like. In other words, a knot. The main point is if one is correct, a knot is composite of net curvature, associated with odd numbers. That is an expansion of the 8T, which did not analyze the odd numbers, but rather referred only to prime numbers and even numbers, isomorphic to primes and evens respectively. Since odds are not on the prime critical line the expressions on terms (2.1) and (2.11) would not have spin one, but neither spin one-half, that is to say they cannot be associated with a particle of any sort. According to the size of the odd numbers we should be able to observe those knots on the matrix tensor. Below an example to such knot.



Matric Tensor Fluctuations and the Birth of Universes

$$\frac{\partial \mathcal{L}}{\partial s} \frac{\partial s}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} - \frac{\partial \mathcal{L}}{\partial s'} \frac{\partial s'}{\partial M} \frac{\partial M}{\partial g'} \frac{\partial^2 g'}{\partial t^2} = 0 \quad (1)$$

$$\frac{\partial g}{\partial t} = 0, \quad -\frac{\partial^2 g'}{\partial t^2} = 0$$

$$\frac{\partial L}{\partial S_1} - \sum_{n=2}^{\infty} \frac{\partial L}{\partial S_n} = 0 \quad (1.53)$$

$$\frac{\partial L}{\partial S_1} \frac{\partial S_1}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} - \sum_{n=2}^{\infty} \frac{\partial L}{\partial S_n} \frac{\partial S_n}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} = 0 \quad (2)$$

The matric tensor experience arbitrary variations that vanish into matter. We describe the process of arbitrary variations vanishing into matter in the thesis, by the variation of the Dirac Delta function.

$$\begin{aligned} \delta g &\neq 0 & at & t = Q(t) \\ \delta g &= 0 & at & t = Q(t + \Delta t) \\ \delta g_1 + \delta g_2 \dots &= \sum_{i=1}^N \delta g_i \\ \sum_{i=1}^N \delta g_i &= 0 \end{aligned}$$


There is always a chance net curvature will appear at later continuation of time. That is bosonic fields given by the primordial coupling series:

$$\begin{aligned} \delta g &= 0 & at & t = Q(t + \Delta t) \\ \delta g &\neq 0 & at & t_2 = Q(t + \Delta t + \Delta t) \\ \Delta t &\rightarrow 0 \end{aligned}$$

Moreover, the amount of net curvature is either prime or one:

$$\delta g = N_V = 2 \left(V + \frac{1}{2} \right); V \geq 0$$

Now that we presented the 8T foundation, we can visualize the birth of new universe. By assuming a segment of the matrix tensor to experience a certain amount of curvature it could lead to a departing from the original manifold. One can try to put it in visual means. This idea is synonymous with the vacuum fluctuations in QFT.



The main point of this short assay is that the net curvature led to a departing from the original matrix tensor to a new entity. The outer shell of this new manifold will accelerate due to other manifolds wrapping around it given by equation (2). That is in agreement with QFT prediction of infinite universes. The entire evolution of the universes from singularity to complete flatness is given by the main equation (1). The stage and actual flattening moment is different in each manifold. That is an elegant way to eliminate the question – why 13.7B years?

EMT Symmetry

suppose that the electron has absorbed a discrete amount of net curvature, its energy increased. Since we are familiar with the equivalence relation between mass and energy, as presented by Albert Einstein, energy increase is synonymous with mass increase. Suppose its mass increased in such way that now instead of the electron, it is a Muon or a Tau.

$$[(24 * 5) + (3)] + 5 \rightarrow [(24 * 5) + (\mu^-)] + \gamma$$

In addition, the Tau:

$$[(24 * 5) + (3)] + 5 \rightarrow [(24 * 5) + (\tau^-)] + \gamma$$

Mass is curvature converging inward, so if the electron has absorbed net curvature its mass increased. That is supported by the Quark masses series of the 8T. Those higher generation particle according to coupling series are representing a symmetry. The magnitude will stay as it is, invariantly of the actual particle, we can call it the EMT symmetry, first letter of each generation particle name. What will vary as a result of the particle varied is energy of the photon emitted. The heavier the particle, the more energy the emitted net curvature should contain. That is again implied by equivalence between mass and energy. Such a construction allow us to make two predictions regarding the energy of the net curvature, i.e. the photon in the case of the third coupling term:

- (1) The Energy of the photon emitted is proportional to Lepton generation.
- (2) The coupling constants series is invariant to generation – what is varying is the energy of the net curvature.

The Coupling Constants Series and Probability

First, we can represent the original equation, which regard Bosonic fields to be net curvature on the varying Lorentz manifold. Those Bosons are isomorphic to prime numbers or one - $\mathbb{P} \cup (+1)$, and propagating from matter clusters destabilized by the majestic three, which is the electron, from the second element and above. Associate a probability of certain sort to the first element, $N_V = (+3)$. the majestic three and the invariant multiplier eight will be presented as a constants, \mathcal{M}, K .

$$F_{V=0} = 8 + (1) \quad (1.1)$$

$$F_R \# = \left(8 * \prod_{V=1}^{V=R} N_V + (3) \right) + N_V = 30:128:850:9254.. \quad (1.2)$$

$$N_V = 2 \left(V + \frac{1}{2} \right); \quad V \geq 0$$

$$N_V \in \mathbb{P} \cup (+1); \quad \mathbb{P} \rightarrow \text{Set of Primes}$$

$$N_V = P_{max} \in [0, \mathbb{R}] \cup (+1); \quad P_{max} \in \mathbb{P}$$

$$8 + (1): (24 + (3)) + 3: (120 + (3)) + 5: (840 + (3)) + 7 \dots$$

Correlating the net curvature element to a certain probability.

$$N_V = (+3) \rightarrow P(A)$$

$$P(A) < 1$$

Now, for simplicity sake assume that the probability is the same for all each higher element in the series. As we do not really know what is the probability of such an event, it is possible to assume that is the case. We can represent the equation in means of probability.

$$P_A \# = \left(K * \prod_{A=3}^{A=2n+1} P(A) + \mathcal{M} \right) + P(A) \quad (3.3)$$

$$A \in \mathbb{P};$$

For each higher term than there is a dependence, the next element in the series can only arise after a previous probability was satisfied, as it is a series. So the longer we develop, the smaller the probability to detect the boson as it is depended upon longer chain of events, with probability smaller than one. We can represent it in a simpler fashion by ignoring the constants:

$$P_A \# = \left(K * \prod_{A=3}^{A=2n+1} P(A) \right) \quad (3.33A)$$

Let $A \rightarrow \infty$

$$P_A \# = \left(K * \prod_{A=3}^{A=2n+1} P(A) \right) \rightarrow 0 \quad (3.1)$$

Such a representation of the primorial series than makes it easier to understand how hard it will be to detect those higher term coupling bosons, and why they have not found up to this day. However, it scientists have detected gravitational waves they should be able to detect the next elements in the coupling series, as they are about seven, and seventy two weaker than the electric. Therefore, despite each term is an individual element which have a unique boson isomorphic to \mathbb{P} for the second and above, there is an implicit dependence given by the fact that is a mathematical series and each even sum is a scalar multiple of the next prime. If we represent the series from an angle of the arrow of time, the higher the coupling term, the more time it will need to develop it. Weakest interactions appear than after longer periods of time, and the strongest most common ones appear at the beginning. We can make a prediction:

- (1) The probability of locating the boson of the third term is significantly higher than the sixth term.

Asymptotic Freedom


Bosons were proven discrete amount of net curvature on the matrix tensor, we can represent them by the term in equation (1.3):

$$\sum_{i=1}^M \delta g_i > 0 ; M \rightarrow \infty \quad (3.13)$$

$$\sum_{i=1}^M \delta g_i \in (+1) \cup \mathbb{P}$$

$$\mathbb{P} \rightarrow \text{Set of Primes}$$

Now, we have used the visualization of the sea of gluons on the Quark triplet in the following way.



In the context of asymptotic freedom, when we indulge in high energy collisions, that is synonymous with trying to roll the quark triplet uphill. It is possible to try as the bosons are just net curvature unbound as given by (1), however since each boson is a curvature of certain magnitude it increase the probability of arrival to its position, therefore we have a "sea" of gluons. For example, in the third coupling term presented in equations (3) to (3.1):

$$\sum_{i=1}^M \gamma_i = \sum_{i=1}^M \delta g_i > 0 \quad (3.13.B)$$

$$[(24 * 5) + (3)] + 5 \rightarrow [(24 * 5) + (e)] + \gamma$$

Taken from that point of analysis, asymptotic freedom is a result of curvature converging to a point, or the existence of gluons on the quark triplet. If the number of bosons is ever increasing on the quark triplet, so does the overall curvature of the magnitude. To roll a quark uphill an infinite curve is at the verge of impossible. The attempt to roll the quark triplet elements uphill will eventually lead to a the quark reaching the minima, lowest point on the curve. Similar to other physical phenomena aspiring minima. Overall the 8T from birds eye overview, allow us to explain phenomena which is considered "advanced" such as Pauli Principle, asymptotic freedom, Spin, the commutator, the reason for the coupling magnitudes, dark energy and probability of arrival in rather simple and elegant way. All we need is just two equations, (1) and the coupling constants series.

Manifold Jumps and Pharrell Transport

$$\frac{\partial \mathcal{L}}{\partial s} \frac{\partial s}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} - \frac{\partial \mathcal{L}}{\partial s'} \frac{\partial s'}{\partial M} \frac{\partial M}{\partial g'} \frac{\partial^2 g'}{\partial t^2} = 0 \quad (1)$$


$$\frac{\partial \mathcal{L}}{\partial s_1} - \sum_{n=2}^{\infty} \frac{\partial \mathcal{L}}{\partial s_n} = 0 \quad (1.53)$$

$$\frac{\partial \mathcal{L}}{\partial s_1} \frac{\partial s_1}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} - \sum_{n=2}^{\infty} \frac{\partial \mathcal{L}}{\partial s_n} \frac{\partial s_n}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} = 0 \quad (2)$$

The implicit assumption of equation of (2) is that in order of the universe packet to flatten each other the curvatures on the manifolds must be interfacing. That is synonymous with stating that the universe packet must have topologically invariant manifolds, or manifolds in which the extremum curvature distribution is identical on the metric tensor. That is because in the manifolds flatten each other due to the interaction of those areas, if they are not interacting the flattening, i.e. dark energy would not be correct. It is possible to prove that if there were only two manifolds, which are not interacting with each other via those areas, equation (2) will not be correct; the universe would not be flat as we measure it today. The requirement of the universe packet then imposes a symmetry in a sense that only topologically invariant manifolds are "allowed" on the packet. We do not know whether it is actually the case but so it seems by equation (2) and the "thought experiment" of only two manifolds interacting in the packet, assumed different topology. Another point to mention is same topology does mean same matter distribution on each manifold. Distinct manifold can have a dust of gas of certain curvature,

which is equivalent to the mass of a certain galaxy on another manifold. Those universes differ from each other in a distance measure which is not known, can could vary as other topologically invariant manifolds enter the packet. Between each manifold pair there is the same base space, Ricci flow, given by the fourth term of (2). everything written up to this point was already covered in 8T previous papers. From here we have a completely new paper.

Since the manifolds have the same curvature distribution, they have the same energy given by the term of the Ricci flow, if you can switch from the metric tensor of one manifold to its flow, and the flow is the same for all the manifolds in the packet, then you can jump or get into the metric tensor of another manifold. In other words, the Ricci flow is the kernel of the entire manifold packet. That is by equation (1.2) and the fact that each manifold, which flatten each other interact by the areas of extremum curvatures $\partial g / \partial t = 0$. So to switch from manifold to manifold, it will require an immense amount of energy, and such an energy level would lead to a deformation of the metric tensor to the kernel, the Ricci flow, and from the Ricci flow we can reach again the metric tensor of a distinct manifold. 8T than regard the metric tensor of each manifold to be a map to another metric tensor. below are two illustrations, the first is the universe packet, the metric is the dark part and the flow is the white part between each two manifolds. the second illustration is an interaction of areas of extremum curvatures between two distinct manifolds, between each two there is the Ricci flow, which is the same for both. The jump from manifold to manifold can be done via this base space.



Those universes differ from each other in a distance measure, which is not known. As the illustration above suggest, they are very close. The packet could vary as other topologically invariant manifolds enter the packet, also known as cosmological singularity. The main point of this short assay between each manifold pair, and actually all the manifolds in the packet is the same base space, Ricci flow, given by the fourth term of (1.2), which allows the jumps, as presented in equation (1.3). Its currently unclear whether there are infinite manifold packets or just one manifold packet which is infinite. It is also unclear whether the question of distance is applicable in the base space, The Ricci flow, as it pure energy oriented.

Manifold Volcanos and Curvature Eruptions

Imagine that instead of having just one electron in the primordial, we will have an entire surface full of electrons. Each of them is emitting a net curvature of prime magnitude, and they all emit that magnitude at the same temporal segment on the surface of the matric tensor M.

$$\left[(24 * 5) + \sum_{i=1}^N (e_i) \right] + \gamma_i \quad (3.41)$$

$$\odot : \sum_{i=1}^N (e_i) \rightarrow \gamma_i$$

The result of this construction is an immense eruption of net curvature off the manifold, similar to a volcano eruption in geo-physics, its concentrated amount of net curvature eruptions due to a positive summation of electrons that emit together, \odot as a time operator of all elements in the matric tensor. The eruption could be linearly polarized. In physics it is also known as "lasers".

The volcano is the summation of electrons, and the magma is the timed eruptions of photons. It is the same main equation just a different variation – applicable to many particles propagation. The energy of the eruption ray is proportion to the electron summation on the surface, which emit together and to inversely proportional to the area scattered by the eruption ray. The volcano is the electrons on the surface and the magma is the photons, in their concentrated from can melt and cut steal. An analogy makes it easier to describe. So overall the "geo-surface" of the matric tensor is flat, due to the net curvature being relativity small portions and due to the fact arbitrary amount of curvature vanish into matter. The "geo-surfaces" or matric tensors in the 8T have dormant volcanos, which could suddenly become active, causing curvature eruptions of immense magnitude at a timed moment, analogous to magma eruptions.

8T versus MT

8-Theory, revolves around varying curvature to the M-theory that is considered to be an elevated version of string theory, and includes additional dimension and unification of so called five distinct string theories. The two theories differ in noticeable and subtle difference. The first difference is that the M-theory also describe alongside the first three interactions, the interaction of gravity. In the 8T, all interactions **are** distinct amounts of gravity. That is discrete amount of curvature on the matric tensor. That is given by the Primorial coupling constants series and in the main equation, which is in agreement with the equivalence principle.

$$\frac{\partial g}{\partial t} = \frac{\partial^2 g'}{\partial t^2}$$

Second difference is the subject of description, 8T only aspire to describe a varying manifold. It does not include any particles motion or any particles of any sort, and all the particles were derived with no a-priori data regarding their nature. That was done in the three critical theorems that yielded the primorial in March 2021. M-Theory aspire to describe the behavior, vibration and motion of different "strings" or infinitesimal quantities, in three and higher dimensions. Such bold entities of description have not yielded testable predictions to date. Such an analysis is also has an implicit axiom – understating the way those infinitesimal things vary can tell us something about physics. The beginning of the M-theory is describe by the five distinct kinds of strings, and that is the subject of description in birds eye view. A third difference is the number of arbitrary numbers appearing in the theory. 8-Theory has three arbitrary numbers less than any other theory. The number of Bosons is infinite and isomorphic to prime numbers. The number of families is also infinite given by the Quark masses series, which provides us with additional prediction of fourth family below first generation, causing the matric tensor to have additional amounts of light mass particles. The third number is the number of dimensions, as the universe is part of a packet, each with its own set of finite dimensions; the overall number of dimensions is infinite as well. Those are distinct and do not get mixed into one manifold. It is the reason the manifold is flat and the reason each manifold can not be infinite in dimension, as it is confined by others. M- theory does the opposite and describe nature by additional arbitrary number which is 11D. If it is 11D, there has to be a reason it has to be that way. Why not 13D? What makes 11D special? The answer is – nothing. As a number of dimensions, it is good as any other. The fact that seemingly certain traits of Quantum physics are in agreement with this number does not make it special, it could work for a higher dimensional number of certain sort. Another way to put it, this number could be part of a subgroup of numbers. Another arbitrary number of the M-theory is the five "distinct" kind of strings, and the overall emphasis on those strings, makes the theory very weak. As one wrote above, it is building upon the implicit assumption that those strings, and in particular their shape, are important. So 8T has three arbitrary numbers less, MT has two arbitrary numbers more. A forth difference is that 8T is described in terms of spaces, extra spaces. The Matric space and the Riccy flow space, which is the base space. The relation among the two is described by a fiber bundle, since all the manifolds are topologically invariant; it is possible to jump from one manifold into under by switching to the Ricci flow. This space does not obey the rules of distance, and is compact. M-theory describe physics in terms of additional dimensions. So overall, it is much longer description, as you have to describe a-lot more according to each extra dimension. One theory describe spaces, which are two. The other dimensions which are infinite. The fifth difference and the last one, is the number of testable predictions as part of the length of description. 8T has description of dark energy, the equivalence principle, The Primorial coupling constants series and all the known Bosons to be prime amounts of net curvature, fermions as arbitrary variations that vanish. It includes the Quark masses series, curvature knots, matric tensor deformations to the base space, the duality of the thirst forces at 26, and it does so using **only one equation** (1.53). It is that simple in can be encompassed in one equation.


$$\frac{\partial \mathcal{L}}{\partial s_1} - \sum_{n=2}^{\infty} \frac{\partial \mathcal{L}}{\partial s_n} = 0 \quad (1.53)$$

M-theory does need a larger amount of numerical description. Alongside, the number of testable predictions given its main equations and power of predictions – is as far as one knows, stand at zero or very close to it at the levels of energy we can reach for today. Another way to state it, it needs a lot more time and space (on paper) to describe the M-theory, and it gives little to no testable predictions. It was the best we had up-until recently, but according to the analysis and comparing to the new 8T, it seems to have been surpassed. .

Universe Packet Density

$$\frac{\partial \mathcal{L}}{\partial s} \frac{\partial s}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} - \frac{\partial \mathcal{L}}{\partial s'} \frac{\partial s'}{\partial M} \frac{\partial M}{\partial g'} \frac{\partial^2 g'}{\partial t^2} = 0 \quad (1)$$

$$\frac{\partial \mathcal{L}}{\partial s_1} \frac{\partial s_1}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} - \sum_{n=2}^{\infty} \frac{\partial \mathcal{L}}{\partial s_n} \frac{\partial s_n}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} = 0 \quad (2)$$



Up to this point in the thesis, we assumed that there is only one packet of stationary Lorentz manifolds, which grow in number. Each manifold has a distinct arrow of time, which is a unique moment of singularity or a unique age. The older the universe the flatter it should be, as it was a subject of pressure from other manifolds for longer temporal periods. That is currently the only option presented in the 8T thesis. However, it now becomes evident that it could be wrong. There could be a limitation of the number of stationary manifolds that composes the packet. Such that if that limit is reached, any metric tensor fluctuations volatile enough will ignite a manifold, which will join a distinct packet. Similar to wave packets, which comes in an infinite number. As far as one can see, the current equations of the 8T indicate that the universe has a "sphere packing" structure, an unknown number of thin layers stacked or compressed together in a packet.

If the number is infinite then we have one packet of stationary manifolds. If there exist a limit, there are multiple. Another interesting point, if the number of manifolds in the packet is finite, then the degree of acceleration outward from areas of extremum curvatures is also finite, which is what we required for a stationary manifold. If the number of manifolds increases without a density limit, then the outward acceleration should increase overtime, as more stationary manifolds are in the packet. That seems more correct as we know that the so-called "dark energy" is time invariant. Therefore, that could imply that there is a limitation of density in the packet. We can define this density limit by varying the main equation:

$$\frac{\partial \mathcal{L}}{\partial s_1} \frac{\partial s_1}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} - \sum_{n=2}^{n < K} \frac{\partial \mathcal{L}}{\partial s_n} \frac{\partial s_n}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} = 0 \quad (2.A)$$

$$K \in \mathbb{R}$$

We can parameterize the manifolds presented in (2.1) to put the idea of stationary manifold packets, which are distinct in rigor.

$$s_1 + \sum_{n=2}^{n < K} s_n = Y_1 \quad (2.B)$$

Moreover, the new structure of the multiverse is the summation of all the universe packets:

$$Y_1 + \sum_{i=2}^{\infty} Y_i = \vartheta \quad (2.C)$$

In the 8T we assumed there is just one infinite packet, and the dark energy could be an adiabatic variant, which vary very slowly. This paper analyzed the structure of the multiverse by imposing a limitation on the density of the packet, leading to infinite number of distinct packets as described by equations (2.B) and (2.C).

The Commuter

In QFT one of the most important ideas which emphasize the difference between fermions to bosons is the mathematical expression commuting/anti commuting relations for bosons and fermions respectively. The term is presented in the following form:

$$[A_i, B_i] \pm = 0$$

Fermions anti commute, summed as zero when combined and bosons commute, the only they to be summed as zero as if they are subtracted from one another. The actual way of QFT representation is not important in this paper. The idea of the commuting anti-commuting relations of bosons and fermions is in perfect agreement with the 8T. As was presented in the thesis, the arbitrary variations term is associated with fermions. We require the term to vanish, so when partitioned we needed an even amount of two distinct elements which differ in sign.

$$\sum_{k=1}^M \delta g_k = 0 \quad (2.12)$$

On the other hand, the bosons were regarded as net curvature of discrete prime amounts as described by the primorial, which add up to a positive summation, so they only way to eliminate them is to subtract from one another. That is in agreement with QFT idea of commutation relation. The term describing Bosons is (3.13.B):

$$\sum_{i=1}^M \gamma_i = \sum_{i=1}^M \delta g_i > 0 \quad (3.13.B)$$

Overall, one of the most important ideas in Quantum Field Theory is in perfect intersection with the 8T. We can visualize it and reason why that is from an angle of curvature on the matric tensor using the main equation and primorial. We can even use the commuter on the two terms.

$$[\delta g_i, \delta g'_i] \pm = 0 \quad (1.6)$$

The first term in the commuting relation (1.5) is describing the partitioned terms, the second is the acceleration. Fermions will accelerate toward each other, in agreement with vanishing curvature. Bosons will accelerate to a joint point on the matric tensor. That is because each bosons is a net curvature that increase the probability of arrival to itself. As was analyzed before, 8T and QFT does not contradict one another.

The Curvature Code - 8T

$$\sum_{k=1}^M \delta g_k = 0 \quad (2.12)$$

$$\sum_{i=1}^M \gamma_i = \sum_{i=1}^M \delta g_i > 0 \quad (3.13.B)$$

Suppose we allocate to the terms, additional terms according to each variable in the main equation. Now as a result we have those fourfold terms for fermions and Bosons accordingly:

$$[\delta \ell \delta s \delta M] \delta g = 0$$

$$[\delta \ell \delta s \delta M] \delta g > 0$$

We do not know what are the values of the three terms inside the bracket, however since we know to associate the conditions in equations (2.12) and (3.13B) to be equal to zero and larger than zero accordingly, these are in essence constraint to the rest of the unknown chained terms. For fermions, we can deduce:

$$[\delta \ell \delta s \delta M] = 0$$

Because of the $\sum_{i=1}^N \delta g_i = 0$ auxiliary condition, which impose a constraint on the chained terms. The fermions will receive the form of points, which are flat and are infinitesimal in length, on the matrix tensor of the manifold. Now analyze the Bosons, with the auxiliary condition $\sum_{i=1}^M \delta g_i > 0$, the chained term is:

$$[\delta \ell \delta s \delta M] > 0$$

Bosons will receive the form of non-local propagation on the matrix tensor of the manifold. The opposite of infinitesimal scales, that is because they cannot vanish into matter, and isomorphic to prime numbers. Similar to how we presented the process of emission. Summing up, we do not know what are the chained three terms are, but we have proven the ideas two-pillar ideas of the 8T: (2.12) and (3.13.B) in which we can use, as auxiliary conditions. Those auxiliary conditions are used on the chained three terms, which we do not know, and thus they are the key to solve the entire chain. Those two conditions are the vital key to the curvature code – the language of nature.

Degrees of Freedom - 8T

We have derived the main equation (1) by EL operator. The following way:

$$\ell = (s, s', t)$$

We can state that the 8T analysis in that form has one degree of freedom.

$$\ell = (s_1, s'_1, t)$$

since we have proven the second representation in equation (1.2), and thus we can represent the EL operator as a system of differential equations with an infinite degrees of freedom. Those differential equations describe a system of stationary manifolds. That is a different way to state that we are dealing with an infinite dimensional universe, using the original operator.

$$\ell = (s_1 \dots s_n, s'_1 \dots s'_n, t_{1 \rightarrow n}) \quad (1.61)$$

the time operator is of course present in each manifold, but since each manifold has a unique moment of singularity, each manifold is getting flattened in different temporal moment, we have to index the time parameter, so to indicate that the arrow is in different stages for each manifold. Such a representation allows us to eliminate the question regarding the arbitrary number of 13.7B billion years. Equation (1.61) is another way to represent the structure of the multiverse, infinite manifolds that are stationary, and interact with each other. Since each manifold is part of the packet, it is confined by it and cannot escape the variation of the manifold than can be presented only within the domain of the packet. Such an analysis also eliminate the question of three dimensional universe, by representing infinite degrees of freedom, we can elevate the universe to infinite dimensions. We can represent the packet in a discrete way, for example:

$$s_1 \rightarrow \text{dimen.} (1 \rightarrow 3) + t_1$$

$$s_2 \rightarrow \text{dimen.} (4 \rightarrow 6) + t_2$$

$$s_3 \rightarrow \text{dimen.} (K \rightarrow K + 2) + t_K$$

$$K \in \mathbb{R}$$

Since we already presented a symmetry regarding the universe packet, we can change the index of the summation with no effect. Residents of the "second manifold" regard themselves as first, and thus count their dimensions as first to third, if we are residents of the "first" manifold, we count our three as the first to the third, and "theirs" as fourth to six. Each resident of distinct manifold regard "his" dimensions as the lowest, i.e. first to third plus a unique arrow.

8T- Curvature Spectra's

we have seen the multiplier of each term from the second and above, is reflecting the number of so-called "fields" of each interaction. The first coupling term has eight gluon fields:

$$8 + (1)$$

The second term has three fields, the massive W and Z bosons, in accordance to the right multiplier, marked in black:

$$[(8 * \mathbf{3}) + (3)] + 3$$

Since all we have in the 8T its curvature, author would like to coin the term – "Curvature spectra", that is each interaction has Bosonic, net curvatures which differ from one another in certain orientation. It is currently unclear which kind of a physical difference it is, it could be a difference in a orientation of the curvature, or a more obvious difference related to mass or both. The features of the W and Z bosons differ from one another supporting the idea of the spectra. Therefore, it is possible to represent the right multiplier as means of a spectrum that is to parametrize it.

$$\sum_{i=1}^{i=N_V} \Psi_i = N_V$$

Therefore, in the 8T, instead of having a certain finite number of fields, we have an infinite amount of curvature orientations, all appear on the matrix tensor and are isomorphic to prime numbers and one. The curvature spectra is parametrized and counting the number of orientations which in physical theory account the different kinds of particles associated with each coupling term. The new elevated form of the primordial is:

$$F_R \# = \left(2^{\mathcal{M}} * \prod_{i=1}^{i=N_V} \Psi_i + (\mathcal{M}) \right) + N_V \quad (1.2B)$$

The Ghost Neutrino

From experiment, we know that the electron does not propagate by itself but rather with another ghost particle, the electron neutrino. What kind of numerical traits in the 8T this particle possess? In other words, we need to add it the coupling term of the electric without changing the magnitude of the coupling. If reader is familiar with the 8T two symmetry breaking – inward to generate mass:

$$8 - (1)$$

Moreover, outward to generate a ripple on the matrix tensor given by the term:

$$8 + (1)$$

The answer is clear the ghost particle, the electron neutrino cannot be associated with neither symmetry breaking classes. It has to be a particle which has no effect on the coupling term, we can represent it but it will vanish. The answer then is that the electron neutrino is represented by the following numerical trait that associate with vanishing in the 8T:

$$\nu_e \rightarrow 8n ;$$

$$n \in \mathbb{R}$$

$$[(24 * 5) + 8 + (3)] + 5 \rightarrow [(24 * 5) + \nu_e + (e)] + \gamma$$

$$[(24 * 5) + \nu_e + (e)] + \gamma = 128$$

$$\nu_e = 0$$

$$[(24 * 5) + \nu_e + (e)] + \gamma \rightarrow [(24 * 5) + (3)] + 5$$

Based on the 8T we can predict that the electron neutrino will be massless, in order for the coupling term to stay as it is. The same apply to each higher generation neutrino according to the EMT symmetry. The fact it has no mass does not mean it cannot exert pressure. The photon is massless, it can exert pressure. If the photon will propagate on a tiny mass measuring scale it will cause the measuring scale to differ from zero due to the pressure it exerts, and effective mass as it is raw energy. Summing up, it is possible to represent the electron neutrino by using a perfect symmetry multiplier that does not affect the coupling term. The fact it is a perfect symmetry multiplier means the electron neutrino has no mass. That is in agreement with experiment.

Alternative Explanation for Dark Matter

we presented the Quark masses series, and predicted an infinite series of families with total mass aspiring zero. Mass is considered arbitrary amount of curvature converging inward, with a symmetry break of the $8 - (1)$ variations. That is the inverse to the primordial, associated with curvature diverging, or $8 + (1)$ variations. In the case of mass generation, nature is devising in increasing amounts to eliminate the arbitrary amounts of curvature. We predicted the total mass of the fourth to be 0.113 Mev, 55-56 times lighter than first. The advantage of this idea is that we no longer need to explain why there are three families.

$$19,600 \rightarrow 1400 \rightarrow 56 \rightarrow 0.113 \dots$$

$$19,600 * 9 \rightarrow 1400 \rightarrow \frac{56}{9} =$$


$$176,400 \rightarrow 1400 \rightarrow 6.3$$

The two versions are presented in the thesis as it is unknown whether the factor of nine is repeating itself for the fifth family and below. Keeping that in mind, assuming this idea is wrong, what alternative explanation can we offer for the issue of dark matter? Notice that according to the main equation (1) or (1.2) we have an infinite packet of universe which interact at areas of extremum curvatures, that means that there two distinct manifolds (if we regard each manifold to be somewhat of a thin liar), whose extremum curvature interact with our own. Since we are familiar with the equivalence principle between mass and energy, the dark energy as given by equation, can be regarded as dark mass. Those masses of distinct manifolds may have an additional gravitational interaction. If each manifold has distinct subspaces, which are newer manifolds that rose from the original manifold, those subspaces may interact with the original manifold that means a distinct set of mass, interacting with our own. The advantage of this idea is that, there could not be any additional trait of matter if the matter is own distinct (yet very close to our own) manifold. It seems to be suitable to the fact that dark matter do not do anything other than to exhort gravity. The weakness of the original idea is that if there is a fourth family below first, it could behave like original matter, omit and absorb light, which is not in agreement with what we speculate. However, if it is matter on a distinct space, or a infinite spaces of the packet, than the features of dark matter could be explained easier. Summing up, the alternative explanation of dark matter is gravitational effect from a distinct manifold, which interact at areas of extremum curvature. There is advantage to taking the point of view, as it could agree with the features of dark matter behavior. However, using that viewpoint, we still need to explain why there are only three fermions generations. The explanation is not part of this new idea, which is the disadvantage comparing to the original idea. The gravitational effect of dark matter should not be strong, as we have immense fermion clusters, according to the primordial, the ratio of net to total should be very small, aspiring zero, so if dark matter would be explained that route, the gravitational magnitude effect it should have should be weak, that is compared to the first elements in the primordial.

$$\frac{N_V}{T_V} \rightarrow R$$

$$0.111 > 0.1 > 0.039 > 0.008 \dots \rightarrow 0$$

We can take the original illustration and modify it



The Canonical Equations of Curvature Spikes

Suppose we would like to present a simple way to create an analog for the canonical equations of motion, presented by Hamilton. How can we do it in a simple way on a varying Lorentz manifold, with four chained terms in the differential equation? This is an interesting question, and the real answer is one does not know. However, here is an educated guess. The idea is to use the terms in equations (1.48) and (1.49) to derive something fundamental about the momenta of Fermions and Bosons. Suppose we replace the known variable of Hamilton by:

$$\partial q_i \rightarrow \partial g_i$$

And we know from the equivalence principle that

$$\partial g_i = \partial g'_i$$

We can present the canonical equation of curvature spikes, if one intuition is correct:

$$\dot{p}_i = \frac{\partial \ell}{\partial g_i}$$

In addition, since we know it vanish into zero, we cannot present it as being in the denominator. We can than represent the canonical equation of curvature spikes for Fermions:

$$\partial g_i = \frac{\partial \ell}{\dot{p}_i} = 0 \quad (1.7)$$

And since we can derive the beautiful result, which comes to an agreement with previous results of the 8T, fermions momenta will vanish to zero. That is another way to state that they will accelerate toward one another. Therefore, configurations of Fermions must appear stationary, similar to Quark Triplets in Hadrons for example. Notice that the emphasis is not on constant rate but rather on the momenta of arbitrary variation set. We can do the exact same thing for Bosons, since we know that they are isomorphic to prime numbers that cannot vanish into matter, the canonical equation of curvature spikes for Bosons is the following:

$$\partial g_i = \frac{\partial \ell}{\dot{p}_i} > 0 \quad (1.71)$$

Meaning that the Bosonic configuration must have some net momenta, we cannot find a Boson at rest. That is in agreement with the 8T ideas. Fermions have opposite signs, demands by stationary Lorentz manifold. Bosons are all positive summations, net curvature on the metric tensor isomorphic to prime numbers, they cannot cancel one another as Fermions do. That is the similar to the idea construction in that led to the 8T commutator for fermions, plus, and Bosons minus respectively:

$$[\delta g_i, \delta g'_i] \pm = 0$$

The Graviton Illusion - 8T

We have analyzed the term of Graviton as a combination of three net curvature that could be either distinct or identical. That is by the spin two trait. That spin two trait, in net curvature representation means that gravity is short ranged, as the three net curvature and the invariant three sums to be a positive number.

$$[(2N_{gravity}) + (3)] + N_{V1} + N_{V2} + N_{V3} = \left[2N_{gravity} + \frac{1}{2}\right] + \frac{1}{2} + \frac{1}{2} + \frac{1}{2} \quad (2.2)$$

$$[2N_{gravity} + 2] \rightarrow [2N_{gravity}] \quad (2.3)$$

Up until now, reader is probability familiar with every equation presented, as those are 8T thesis fundamentals. From here on out, we have a completely new paper. Since nature does not impose a restriction on the kind of particles to which are describing the term (2.2) and (2.3), it is possible to predict that there are infinite classes of Gravitons of distinct magnitudes. Alternatively, that if we take an even sum or certain sort, add a generator and three net curvature of certain magnitude, which belong to the prime ring, that combination will result a "Graviton like" particle.

$$[(2N_0) + (3)] + N_{VK1} + N_{VK2} + N_{VK3}$$

$$[N_{VK1}, N_{VK2}, N_{VK3}] \in \mathbb{P}$$

$$K1 \dots KN \in \mathbb{R}$$

Since spin two vanishes due to being an even number, the difference between each Graviton is the term that does not vanish in spin representation - $(2N_0)$. The smaller this term the stronger the Graviton should be. Therefore, if one analysis is correct, Gravitons classes are infinite in kind, and they are, in contrast to the first three interactions that are in a sense independent, is not independent and depends upon the composite elements. As previously mentioned, Gravitons are a superposition of net curvatures (equivalent or distinct is currently not known), which means that in order to sustain Graviton on quantum scale, it requires aligning three net curvatures in time and position. If one of the net curvature terms is not there, we no longer have the Graviton. The main point of this paper is to make a prediction about the nature of Graviton, and here it is the prediction:

(1) Gravitons are infinite in kind.

It is a daring statement to make given the fact that we did not detect even a single graviton to date, but the 8T is a daring theory. It also provides us a practical way to test whether Graviton like particles can be created in an artificial way. For example, for the electromagnetic coupling, we need the term in (2.4) to create a "Graviton like" particle, the Graviton is a matter of illusion, and it is everywhere and nowhere at the same time, as it is quite rare to create the term in (2.4) as far as one can see.

$$[(2N_{(2)}) + (e)] + \gamma + \gamma + \gamma \quad (2.4)$$

Suppose that we are a given the gravitation coupling as the following term:

$$[(2N_0) + (3)] + N_{VK1} + N_{VK2} + N_{VK3}$$

However, the net variation N_{VK3} suddenly vanish from the combination and being replaced by a distinct net curvature element:

$$[(2N_0) + (3)] + N_{VK1} + N_{VK2} + N_{VK4}$$

$$N_{VK3} \neq N_{VK4}$$

That seems as a trivial change, but it really is not. 8T predicts that the Gravitons are infinite in kind. That is the example of that idea. We previously mentioned gravity is different because it is a composite interaction due to the spin two trait. That is in contrast to interactions of the primordial which are not a

composite but contain one net element. That means that gravity coupling magnitude could vary over time. In particular, it means that net curvature elements can replace other elements that were part of the threefold composite. Since the total spin is invariant, i.e. two, there is not a change in the nature of gravity; the structure is the same while the composite element could be different.

$$[(2N_0) + (3)] + N_{VK1} + N_{VK2} + N_{VK3} \rightarrow 2N_0 + 2$$

$$[(2N_0) + (3)] + N_{VK1} + N_{VK2} + N_{VK4} \rightarrow 2N_0 + 2$$

Prediction: The Gravitational coupling constant is not a constant at all.

Curvature Terminators and Conservation of Energy - 8T

The point of the first 8T proof presented in pages 3-4, which was only briefly mentioned, is that nature is aspiring to eliminate the curvature. The result of the elimination is yielding the group that allowed physicists to predict the existence of the omega minus (333) in the 8T. However even if (2.12) is vanishing to zero, there is constant creation of matter. Arbitrary variation of the manifold are not obeying a time limit, they can and are created in a random fashion. So one way to put it is that **energy is not conserved**. That is because matter is constantly being created, and matter is synonymous with energy. The only way to ensure that the energy will be conserved is to present a new way of curvature terminators that is anti-matter. We allow the existence of anti-matter as the coupling magnitudes are preserved under sign reversal.

$$\left[2N_3 + \frac{1}{2}\right] + \frac{1}{2} \rightarrow \left[-2N_3 - \frac{1}{2}\right] - \frac{1}{2} \quad (1.45)$$

We analyzed the subject of anti-matter in previous paper, and in particular the subject of orthogonal curvature, which has inner product zero. So to ensure the conservation of energy, we will have to present the set of arbitrary curvature terminators, for fermions, it has two inverse elements.

$$X = [-\delta g_1, +\delta g_2] \\ \langle \delta g_i | -\delta g_i \rangle = 0 \quad (1.46)$$

So overall, there are two main stages of curvature elimination. First arbitrary variation vanish into matter, as presented in the 8T thesis and the prove above. Secondly, to ensure the conservation of energy, anti-matter terminators are presented. Whether energy is actually conserved is unknown, author tend to belief is not. That is due to the asymmetry of matter to anti-matter in the universe. If for each matter created there is also an anti-matter particle, anti-matter should be more common. We have presented the same procedure of orthogonal curvatures to leptons and Bosons. We used equation (1.46) with leptons as elements of the inner product such as the electron and positron:

$$\langle +3_i | -3_i \rangle = 0$$

In addition, with bosons, described by the term (2.12) as they were proven discrete amount of prime curvature on the matric tensor:

$$\sum_{i=1}^M \delta g_i > 0 \quad (3.13.B)$$

Same rules apply for the photon as an example:

$$\langle \gamma_i | \gamma_i^- \rangle = 0$$

Summing up, if require the conservation of energy we must present the arbitrary curvature terminators, i.e. Anti-matter. If the number of anti-matter terminators is smaller than the number of arbitrary variations which vanish into matter, which seems to be the case on our manifold, than energy is **not** conserved, as matter is constantly being created.

Direction Invariant Fermion Distributions - 8T

The sole mathematical discipline of the 8T is calculus of variations. As reader assumed familiar with it, one of the major features of this theory is the vanishing of variation.

$$\frac{\partial \mathcal{L}}{\delta q_i} = \frac{\partial \mathcal{L}}{\partial q_i} - \frac{\partial \mathcal{L}}{\partial \dot{q}_i} \left(\frac{d}{dt} \right) = 0$$

Since in our theory we have the arbitrary variation term in equation (1.48) to vanish into matter, we can represent the idea as:

$$\frac{\partial \mathcal{L}}{\delta g_i} = \frac{\partial \mathcal{L}}{\partial g_i} - \frac{\partial \mathcal{L}}{\partial \dot{g}_i} \left(\frac{d}{dt} \right) = 0 \quad (1.8)$$

If so, Bosons as net curvature isomorphic to prime numbers are interfering with the stationarity of the manifold, hence their name "Agrarian", as they cannot vanish into matter, they cause the matter clustering. For Bosons:

$$\frac{\partial \mathcal{L}}{\delta g_i} = \frac{\partial \mathcal{L}}{\partial g_i} - \frac{\partial \mathcal{L}}{\partial \dot{g}_i} \left(\frac{d}{dt} \right) > 0 \quad (1.81)$$

One final point, since the primorial coupling series is invariant to direction:

$$\mathcal{P}_0 = 8 + (1)$$

$$\mathcal{P}_N \# = \left(2^{\mathcal{M}} * \prod_{V=1}^{V=N} \mathcal{P}_V + (\mathcal{M}) \right) + \mathcal{P}_V = 30:128:850:9254.. \quad (1.2.A)$$

As presented in the idea of probability variation of the (1.2A):

$$P_A \# = \left(K * \prod_{A=3}^{A=2n+1} P(A) + \mathcal{M} \right) + P(A)$$

$$A \in \mathbb{P};$$

The matter configuration of the manifold should invariant to direction, that is precious because it is not possible to determine where the lepton is going to emit or absorb, or even which kind of bosons are in play. Put another way, because it is impossible to know where the net curvature that violate stationarity will appear, the fermion distribution across all directions of the manifold is the same, there is no special direction of any sort. That is precious the current modern picture of cosmology, the universe look everywhere the same. Same idea we presented in earlier paper of the sphere shape of starts, but now to much larger Fermion clusters.

Minimizing the Laws

The last part of this paper will revolve around a feature of nature which was mentioned briefly in previous papers, and in the 8T thesis. Lagrangian oriented theories are based upon the principle of least action, which deals with minima of certain classes, and this is the most significant feature of those theories. There is one additional minima in the 8T and in a final theory that should get our attention, as it is just as important. That is minimizing the number of laws that govern everything. In every universe, at every stage of development of the manifold, from the flattening by the packet to complete coldness, the number of laws should stand at minima. In other words, the number of equations or ideas in which we use to describe everything should be minimal, and that is a significant feature of a final theory. The minima is not only path-oriented such as in classical mechanics or QED, it is also manifested in the number of laws. 8T is that kind of theory as all we have achieved, from dark energy to the coupling series and the Quark masses series, is encompassed in just one equation and two conditions.

$$\frac{\partial \ell}{\partial s} \frac{\partial s}{\partial M} \frac{\partial M}{\partial g} \frac{\partial g}{\partial t} - \frac{\partial \ell}{\partial s'} \frac{\partial s'}{\partial M} \frac{\partial M}{\partial g'} \frac{\partial^2 g'}{\partial t^2} = 0 \quad (1)$$

In addition, all the standard model of particle physics is encompassed in just two conditions. For fermions:

$$\sum_{i=1}^N \delta g_i = 0 \quad (1.48)$$

Bosons as net curvature of discrete prime amounts:

$$\sum_{i=1}^M \delta g_i > 0 \quad (1.49)$$

Manor Ohad