

A review of the zooplankton in Singapore waters

Claire Schmoker^{1, 2*}, Mohamed-Sofiane Mahjoub¹, Albert Calbet², Shih-Hui Hsiao³, Francesca Russo¹, Ole Larsen¹, Aurore Trottet¹ & Guillaume Drillet¹

Abstract. The island of Singapore is located between 1°09'N–1°29'N and 103°38'E–104°06'E at the confluence of the Malacca Straits and the South China Sea. To date, both the marine and freshwater zooplanktons of this area are poorly studied, and availability of taxonomic identification is scarce. Moreover, most of the studies were published between the 1950s to the beginning of the 1970s. The available data are mainly qualitative, with only a few studies on zooplankton biology and ecology. Here, the literature on zooplankton communities in Singapore waters is reviewed in order to provide a baseline for future zooplankton surveys, and to better understand the aquatic ecosystems of this area. Also included are recent data obtained from a one-year plankton monitoring in 2012 from two marine stations in Singapore. The temporal variation of the plankton groups was observed in the study to be similar to what was described in some works from the 1970s. The species richness increased in these more recent studies, probably due to changes in the sampling and preservation methods. Because of these changes, comparing between data-sets is challenging; however, similarities in species richness and seasonality between a recent study and previous data-sets were evident. Finally, it is argued that continuous marine plankton monitoring would be an asset for Singapore and the region.

Key words. Singapore, South China Sea, zooplankton, species richness, diversity

INTRODUCTION

Geographic and climatic position of Singapore. The island of Singapore is located between 1°09'N–1°29'N and 103°38'E–104°06'E at the convergence of the Malacca Straits and the South China Sea. It is separated from Malaysia in the north by the Johor Strait (1.8 km wide) and from Indonesia in the south by the Singapore Strait (16 km wide) (Fig. 1). The Johor Strait is divided into the West and East Johor Strait by the Singapore Causeway. The large Malaysian city of Johor Bahru is located on the northern shoreline of the East Johor Strait (Pang & Tkalic, 2003; Mulia et al., 2013). Because of the geographical position, along important commercial routes connecting the Indian Ocean with the South China Sea, and the deep waters allowing the passage of large commercial vessels, Singapore has become one of the biggest and busiest ports in the world (Chou, 2006). The majority of Singapore's freshwater reserves are within 17 shallow, man-made reservoirs that are highly managed, in states ranging from eutrophic to hypertrophic (Low et al., 2010).

The climate of Singapore is typically equatorial, with high temperatures and large amounts of rainfall throughout the year. The general pattern of water temperature fluctuations is characterised by a minimum during December to January (~26°C) and a maximum during April to June (~32°C) (NASA, 2013). The climate is strongly influenced by the Southeast Asian Monsoon regime, which introduces seasonal variations. The seasons are divided in four periods: two main monsoon seasons, the North-East (NE) Monsoon from November to early March and the South-West (SW) Monsoon from June to September, and two inter-monsoon periods (late March to May and October to November). Heavy




Figure 1: Map of Singapore showing the location of freshwater reservoirs and the sampled stations (DHI Singapore cartography department – GIS). WQ7 and WQ8 are the stations chosen in the frame of the MadeInPlankton project and OSL is the station from the study of Wickstead (1958).

¹DHI-NTU Water & Environment Research Centre and Education Hub, CleanTech Loop 1, #03-05 CleanTech One, Singapore 637141; Email: claire_schmoker@yahoo.fr (*corresponding author)

²Institut de Ciències del Mar, CSIC. Passeig Marítim de la Barceloneta 37-49, 08003 Barcelona, Spain

³Department of Science Education, National Taipei University of Education, Taipei 106, Taiwan, R.O.C

rains and winds characterise both monsoon periods, but the SW Monsoon is usually drier than the NE Monsoon. During the NE Monsoon the general direction of the currents in the South China Sea is south-west. During the SW Monsoon, the currents bring water from the Java Sea and the Malacca Strait (Behera et al., 2013). The two inter-monsoon periods are characterised by intermittent rains and weak variable winds. The transitions between the monsoon seasons occur gradually, generally over a period of two months (Pang & Tkalic, 2003; National Environment Agency, 2009; Behera et al., 2013).

Identifying major knowledge gaps on zooplankton diversity and distribution in Singapore waters. Aquatic and terrestrial ecosystems are structured in complex food webs in which different groups of organisms interact. In the photic zone, phytoplankton cells are unicellular phototrophic organisms at the base of the planktonic trophic webs. Larger heterotrophic organisms—the micro- and mesozooplankton—actively prey on phytoplankton. Microzooplankton mostly consists of unicellular protozoans, whereas mesozooplankton comprise metazoans. Within mesozooplankton, the dominant group are crustaceans belonging to the subclass Copepoda and represent around 80% of the total mesozooplanktonic biomass (Mauchline, 1998). These organisms prey on both primary producers and microzooplankton (Calbet & Saiz, 2005; Saiz & Calbet, 2011). Additionally, they comprise the main food source for the planktonic stages of commercially important species, such as other crustaceans and fish. As elsewhere, the marine plankton distribution in Singapore waters is driven by the general oceanic patterns described above. To understand the aquatic ecosystems of Singapore it is essential to first determine the planktonic assemblages and to assess their functional diversity (how organisms interact with each other). Some studies on phytoplankton communities and their taxonomy in Singapore's coastal waters exist (Tham, 1953; Chou & Chia, 1991; Gin et al., 2000, 2003), and these have recently been reviewed in a checklist of algae species (Pham et al., 2011). To date, however, zooplanktonic organisms have been poorly studied, and their taxonomic identifications are quite scarce. Most of the available literature investigating zooplankton around Singapore was published between the 1950s and the beginning of the 1970s (e.g., Tham, 1953; Wickstead, 1958; Tham, 1973; Thia-Eng, 1973); it is likely that these studies no longer reflect the present species diversity.

Due to its geographic position and to its climate, Singapore generally is very rich in species (Pham et al., 2011). It lies in a biodiversity hotspot, which is a conservation priority (Myers et al., 2000). According to estimations of Brook et al. (2003), based on both recorded and inferred data, Singapore may have lost more than one-half of its terrestrial and freshwater taxa (including more than 80% of freshwater decapods and fish) from 1819 until 2002. In addition, due to the oceanographic features characterising the waters of Singapore, anthropogenic and natural events occurring in more distant areas (i.e., Malacca Strait, South China Sea) could affect coastal ecosystems locally.

According to Yeo et al. (2011), many factors such as the construction of dams, the causeway linking Singapore to Johor Bahru (Malaysia), shoreline modifications, and species introductions may have affected native aquatic biodiversity. Moreover, the introduction of exogenous species by ballast water may have increased as a result of the growing activity in Singapore harbours. Without knowledge of the existing baseline biodiversity waters introductions of non-native species cannot be confirmed and recorded (Pham et al., 2011). Therefore, in order to plan an efficient management of Singapore waters, a thorough knowledge of plankton diversity and ecosystem functioning is needed. In this paper, a review of the existing data on zooplankton communities in Singaporean waters is used to provide a baseline for future zooplankton surveys. Due to the scarcity of data on zooplankton from fresh waters, this review mainly focuses on marine waters.

MATERIAL AND METHODS

Literature review. Data were obtained from a literature search on zooplankton studies in the area of Singapore, including freshwater and marine studies. Initially, the words “zooplankton” and “Singapore” were searched in the Aquatic Sciences and Fisheries, Abstract electronic database, then other electronic databases such as WoS, Google Scholar or JSTOR. After, the search was extended to the older references cited in the selected papers (many of these are not incorporated into the electronic databases). Several of the studies on freshwater and marine zooplankton included also Malaysian records, but did not differentiate the origin of data; therefore, those data were presented as well.

New data on zooplankton diversity. Besides the above-mentioned literature analysis, we also present new data obtained at two stations on the west coast of Singapore (Fig. 1; WQ7 – west part; 01°17.949'N, 103°42.383'E and WQ8 – east part; 01°17.694'N, 103°43.340'E). The sampling was carried out every two weeks over the course of one year, from May 2012 to May 2013. Vertical plankton hauls were conducted with a 100 µm mesh net to collect mesozooplankton. In the laboratory, the samples were fixed with 4% formaldehyde and then divided into two subsamples; one was saved for future processing, and the other was used for the taxonomic analysis of the mesozooplankton community. Around 1000 individuals were counted per sample (Omori & Ikeda, 1984) and identified, when possible, to species level following Chen & Zhang (1965), Chen et al. (1974), Nishida (1985) and Harris et al. (2000).

RESULTS AND DISCUSSION

Historical zooplankton seasonality in Singapore waters. Investigations on copepods in the Straits of Malacca started in the early 1900s with the works of Clave (1901) and Sewell (1933). The first complete available study of zooplanktonic organisms from Singaporean seawaters is dated from 1953 (Tham, 1953; see Table 1). Tham (1953) established a “plankton calendar” for the Singapore Straits, by monitoring weather, seawater conditions, and absence

Table 1. List of zooplanktonic organisms found in Singaporean and Malaysian seawaters. (1) = reference Tham, 1953; (2) = reference Wickstead, 1958; (3) = reference Tham, 1973; (4) = reference Thia-Eng, 1973; (5) = reference Gollash et al., 2000; (6) = reference Razai et al., 2004; (7) = data from the present study, Nparks and DHLgroup; (8) = Razouls et al (2005–2012); (9) = World Register of Marine Species at <http://www.marinespecies.org/aphia.php?p=taxdetails&id=220933> on 2013-01-23.

PHYLUM	CLASS	ORDER	Reported scientific name	(1)	(2)	(3)	(4)	(5)	(6)	(7)	Authority	Currently accepted scientific name (Genus and lower) (8,9)	Authority for accepted name
Sarcomastigophora (subphylum Radiolaria) Ciliophora	Spirotrichia	Euploida	Unidentified						+				
			<i>Euploes</i> sp.						+		Müller, 1786	<i>Euploes</i>	Müller, 1786
			<i>Strombidium</i> sp.						+		Claparède & Lachmann, 1859	<i>Strombidium</i>	Claparède & Lachmann, 1859
			<i>Tontonia</i> sp						+		Fauré-Fremiet, 1914	<i>Tontonia</i>	Fauré-Fremiet, 1914
Foraminifera	Litostomatea	Choreotrichida	Unidentified Tintinnina (sub-order)	+	+					+	Kofoed & Campbell, 1929		Kofoed & Campbell, 1929
			<i>Mesodinium</i> sp.						+		von Stein	<i>Mesodinium</i>	von Stein
			Unidentified		+								
			<i>Berggrenia cf pumilio</i> <i>Streptochilus globulosus</i>						+		Parker, 1962 Cushman, 1933	<i>Berggrenia pumilio</i> <i>Streptochilus globulosus</i>	Parker, 1962 Cushman, 1933
Cnidaria	Hydrozoa	Anthomedusae	Unidentified		+								
			<i>Cladia</i> sp.						+		Lamouroux, 1812	<i>Cladia</i>	Lamouroux, 1812
			Unidentified		+								
			<i>Ceratocymba leuckarti</i> <i>Diphyes chamissonis</i> <i>Lensia</i> sp.						+		Huxley, 1859 Huxley, 1859 Totton 1932	<i>Ceratocymba leuckarti</i> <i>Diphyes chamissonis</i> <i>Lensia</i>	Huxley, 1859 Huxley, 1859 Totton 1932
			Unidentified	+	+	+	+	+					

Table 1. Cont'd.

PHYLUM	CLASS	ORDER	Reported scientific name	(1)	(2)	(3)	(4)	(5)	(6)	(7)	Authority	Currently accepted scientific name (Genus and lower) (8,9)	Authority for accepted name
Ctenophora	Cubozoa	Cubomedusae	Unidentified		+								
			Unidentified		+								
Annelida	Tentaculata	Cydippidia	<i>Pleurobrachia</i> sp.						+	Fleming, 1822	<i>Pleurobrachia</i>		Fleming, 1822
			Unidentified		+								
Platyhelminthes	Polychaeta		Unidentified larvae	+	+	+				+			
			Unidentified										
Arthropoda	Branchiopoda	Diplostraca (<i>infra order Cladocera</i>) (<i>infra order Cladocera</i>)	Unidentified					+					
			<i>Evadne</i> sp.			+				Lovén, 1836	<i>Evadne</i>		Lovén, 1836
Ostracoda		Halocyprida	<i>Penilia</i> sp.			+				Dana, 1852	<i>Penilia</i>		Dana, 1852
			Unidentified		+	+				Müller, 1890	<i>Pyrocypis</i>		Müller, 1890
Maxillopoda (Subclass Copepoda)			<i>Euconchoecia</i> sp.			+				Müller, 1890	<i>Euconchoecia</i>		Müller, 1890
			<i>Conchoecia parvidentata</i>						+	Müller, 1906	<i>Conchoecia parvidentata</i>		Müller, 1906
	Calanoida		<i>Acartia amboinensis</i>						+	Carl, 1907	<i>Acartia (Odontacartia) amboinensis</i>		Carl, 1907
			<i>Acartia erythraea</i>						+	Giesbrecht, 1889	<i>Acartia (Odontacartia) erythraea</i>		Giesbrecht, 1889
			<i>Acartia pacifica</i>						+	Steuer, 1915	<i>Acartia (Odontacartia) pacifica</i>		Steuer, 1915

Table 1. Cont'd.

PHYLUM	CLASS	ORDER	Reported scientific name	(1)	(2)	(3)	(4)	(5)	(6)	(7)	Authority	Currently accepted scientific name (Genus and lower) (8,9)	Authority for accepted name
			<i>Acartia plumosa</i>					+			Scott T., 1894	<i>Acartia (Acanthacartia) plumosa</i>	Scott T., 1894
			<i>Acartia sinijiensis</i>						+		Mori, 1940	<i>Acartia (Acanthacartia) sinijiensis</i>	Mori, 1940
			<i>Acartia</i> sp.			+					Dana, 1846	<i>Acartia</i>	Dana, 1846
			<i>Acartia</i> sp. 1		+			+			Dana, 1846	<i>Acartia</i>	Dana, 1846
			<i>Acartia</i> sp. 2					+			Dana, 1846	<i>Acartia</i>	Dana, 1846
			<i>Acartia spinicauda</i>					+			Giesbrecht, 1889	<i>Acartia (Odontacartia) spinicauda</i>	Giesbrecht, 1889
			<i>Acrocalanus</i> sp.			+					Giesbrecht, 1888	<i>Acrocalanus</i>	Giesbrecht, 1888
			<i>Acrocalanus gibber</i>					+			Giesbrecht, 1888	<i>Acrocalanus gibber</i>	Giesbrecht, 1888
			<i>Acrocalanus gracilis</i>					+			Giesbrecht, 1888	<i>Acrocalanus gracilis</i>	Giesbrecht, 1888
			<i>Acrocalanus longicornis</i>					+			Giesbrecht, 1888	<i>Acrocalanus longicornis</i>	Giesbrecht, 1888
			<i>Acrocalanus monachus</i>					+			Giesbrecht, 1888	<i>Acrocalanus monachus</i>	Giesbrecht, 1888
			<i>Bestiolina similis</i>					+			Sewell, 1914	<i>Bestiolina similis</i>	Sewell, 1914
			<i>Calanopia</i> sp.		+						Dana, 1852	<i>Calanopia</i>	Dana, 1852
			<i>Calanopia elliptica</i>					+			Dana, 1849	<i>Calanopia elliptica</i>	Dana, 1849
			<i>Calanopia minor</i>					+			Scott A., 1902	<i>Calanopia minor</i>	Scott A., 1902
			<i>Calanopia thompsoni</i>					+			Scott A., 1909	<i>Calanopia thompsoni</i>	Scott A., 1909
			<i>Calocalanus pavo</i>					+			Dana, 1852	<i>Calocalanus pavo</i>	Dana, 1852
			<i>Calocalanus styliformis</i>					+			Giesbrecht, 1888	<i>Calocalanus styliformis</i>	Giesbrecht, 1888
			<i>Calocalanus</i> sp.					+			Giesbrecht, 1888	<i>Calocalanus</i> sp.	Giesbrecht, 1888
			<i>Candacia</i> sp.			+					Dana, 1846	<i>Candacia</i>	Dana, 1846
			<i>Candacia bradyi</i>					+			Scott A., 1902	<i>Candacia bradyi</i>	Scott A., 1902
			<i>Candacia curta</i>					+			Dana, 1849	<i>Candacia curta</i>	Dana, 1849
			<i>Candacia discaudata</i>					+			Scott A., 1909	<i>Candacia discaudata</i>	Scott A., 1909
			<i>Candacia ethiopica</i>					+			Dana, 1849	<i>Candacia ethiopica</i>	Dana, 1849
			<i>Candacia pachydactyla</i>					+			Dana, 1849	<i>Candacia pachydactyla</i>	Dana, 1849
			<i>Canthocalanus</i>		+						Scott A., 1909	<i>Canthocalanus</i>	Scott A., 1909
			<i>Canthocalanus pauper</i>					+			Giesbrecht, 1888	<i>Canthocalanus pauper</i>	Giesbrecht, 1888
			<i>Centropages dorsispinatus</i>					+			Thompson I.C. & Scott A., 1903	<i>Centropages dorsispinatus</i>	Thompson I.C. & Scott A., 1903
			<i>Centropages furcatus</i>		+			+			Dana, 1849	<i>Centropages furcatus</i>	Dana, 1849
			<i>Centropages orsinii</i>					+			Giesbrecht, 1889	<i>Centropages orsinii</i>	Giesbrecht, 1889
			<i>Centropages</i> sp.			+		+			Kroyer, 1849	<i>Centropages</i>	Kroyer, 1849
			<i>Centropages tenuiremis</i>					+			Thompson I.C. & Scott A., 1903	<i>Centropages tenuiremis</i>	Thompson I.C. & Scott A., 1903

Table 1. Cont'd.

PHYLUM	CLASS	ORDER	Reported scientific name	(1)	(2)	(3)	(4)	(5)	(6)	(7)	Authority	Currently accepted scientific name (Genus and lower) (8,9)	Authority for accepted name
			<i>Clausocalanus</i> <i>arcuicornis</i>					+			Dana, 1849	<i>Clausocalanus</i> <i>arcuicornis</i>	Dana, 1849
			<i>Clausocalanus furcatus</i>					+	+		Brady, 1883	<i>Clausocalanus furcatus</i>	Brady, 1883
			<i>Clausocalanus jobei</i>					+	+		Frost & Fleminger, 1968	<i>Clausocalanus jobei</i>	Frost & Fleminger, 1968
			<i>Clausocalanus</i> <i>parapergens</i>							+	Frost & Fleminger, 1968	<i>Clausocalanus</i> <i>parapergens</i>	Frost & Fleminger, 1968
			<i>Clausocalanus pergens</i>					+			Farran, 1926	<i>Clausocalanus pergens</i>	Farran, 1926
			<i>Eucalanus attenuatus</i>					+	+		Dana, 1849	<i>Pareucalanus attenuatus</i>	Dana, 1849
			<i>Eucalanus crassus</i>					+	+		Giesbrecht, 1888	<i>Subeucalanus crassus</i>	Giesbrecht, 1888
			<i>Eucalanus pileatus</i>					+	+		Giesbrecht, 1888	<i>Subeucalanus pileatus</i>	Giesbrecht, 1888
			<i>Eucalanus</i> sp.				+		+		Dana, 1852	<i>Eucalanus</i>	Dana, 1852
			<i>Eucalanus subcrassus</i>		+			+	+		Giesbrecht, 1888	<i>Eucalanus subcrassus</i>	Giesbrecht, 1888
			<i>Eucalanus subtenius</i>			+		+	+		Giesbrecht, 1888	<i>Subeucalanus subtenius</i>	Giesbrecht, 1888
			<i>Euchaeta</i> sp.				+				Philippi, 1843	<i>Euchaeta</i>	Philippi, 1843
			<i>Euchaeta concinna</i>		+			+	+		Dana, 1849	<i>Paraeuchaeta concinna</i>	Dana, 1849
			<i>Euchaeta marinella</i>					+	+		Bradford, 1974	<i>Euchaeta marinella</i>	Bradford, 1974
			<i>Labidocera</i> sp.		+	+					Lubbock, 1853	<i>Labidocera</i>	Lubbock, 1853
			<i>Labidocera acuta</i>					+	+		Dana, 1849	<i>Labidocera acuta</i>	Dana, 1849
			<i>Labidocera bengalensis</i>					+	+		Krishnaswamy, 1952	<i>Labidocera bengalensis</i>	Krishnaswamy, 1952
			<i>Labidocera euchaeta</i>					+	+		Giesbrecht, 1889	<i>Labidocera euchaeta</i>	Giesbrecht, 1889
			<i>Labidocera kroyeri</i>					+	+		Brady, 1883	<i>Labidocera kroyeri</i>	Brady, 1883
			<i>Labidocera minuta</i>					+	+		Giesbrecht, 1889	<i>Labidocera minuta</i>	Giesbrecht, 1889
			<i>Labidocera pectinata</i>					+	+		Thompson I.C. & Scott A., 1903	<i>Labidocera pectinata</i>	Thompson I.C. & Scott A., 1903
			<i>Labidocera rotunda</i>					+	+		Mori, 1929	<i>Labidocera rotunda</i>	Mori, 1929
			<i>Labidocera</i> sp. 1					+	+		Lubbock, 1853	<i>Labidocera</i>	Lubbock, 1853
			<i>Labidocera</i> sp. 2					+	+		Lubbock, 1853	<i>Labidocera</i>	Lubbock, 1853
			<i>Lucicutia flavicornis</i>					+	+		Claus, 1863	<i>Lucicutia flavicornis</i>	Claus, 1863
			<i>Lucicutia gaussae</i>					+	+		Grice, 1963	<i>Lucicutia gaussae</i>	Grice, 1963
			<i>Metacalanus</i> sp.			+		+	+		Cleve, 1901	<i>Metacalanus</i>	Cleve, 1901
			<i>Nannocalanus minor</i>					+	+		Claus, 1863	<i>Nannocalanus minor</i>	Claus, 1863
			<i>Neocalanus gracilis</i>					+	+		Dana, 1852	<i>Neocalanus gracilis</i>	Dana, 1852
			<i>Paracalanus aculeatus</i>					+	+		Giesbrecht, 1888	<i>Paracalanus aculeatus</i>	Giesbrecht, 1888
			<i>Paracalanus denudatus</i>					+	+		Sewell, 1929	<i>Paracalanus denudatus</i>	Sewell, 1929
			<i>Paracalanus parvus</i>				+	+	+		Claus, 1863	<i>Paracalanus parvus</i>	Claus, 1863
			<i>Paracalanus nanus</i>					+	+		Sars, 1925	<i>Paracalanus nanus</i>	Sars, 1925
			<i>Paracalanus serrulus</i>					+	+		Shen & Lee, 1963	<i>Paracalanus serrulus</i>	Shen & Lee, 1963

Table 1. Cont'd.

PHYLUM	CLASS	ORDER	Reported scientific name	(1)	(2)	(3)	(4)	(5)	(6)	(7)	Authority	Currently accepted scientific name (Genus and lower) (8,9)	Authority for accepted name
			<i>Paracalanus</i> sp.			+	+		+		Boeck, 1865	<i>Paracalanus</i>	Boeck, 1865
			<i>Parvocalanus crassirostris</i>						+	+	Dahl F., 1894	<i>Parvocalanus crassirostris</i>	Dahl F., 1894
			<i>Parvocalanus elegans</i>						+		Andronov, 1972	<i>Parvocalanus elegans</i>	Andronov, 1972
			<i>Parvocalanus</i> sp.						+	+	Andronov, 1970	<i>Parvocalanus</i>	Andronov, 1970
			<i>Pontella</i> sp. 1						+		Dana, 1846	<i>Pontella</i>	Dana, 1846
			<i>Pontella</i> sp. 2						+	+	Dana, 1846	<i>Pontella</i>	Dana, 1846
			<i>Pontella</i> sp. 3						+	+	Dana, 1846	<i>Pontella</i>	Dana, 1846
			<i>Pontellina plumata</i>						+	+	Dana, 1849	<i>Pontellina plumata</i>	Dana, 1849
			<i>Pseudodiaptomus aurivilli</i>						+	+	Cleve, 1901	<i>Pseudodiaptomus aurivilli</i>	Cleve, 1901
			<i>Pseudodiaptomus marinus</i>							+	Sato, 1913	<i>Pseudodiaptomus marinus</i>	Sato, 1913
			<i>Pseudodiaptomus cf ardjuna</i>							+	Brehm, 1953	<i>Pseudodiaptomus ardjuna</i>	Brehm, 1953
			<i>Pseudodiaptomus</i> sp.						+		Herrick, 1884	<i>Pseudodiaptomus</i>	Herrick, 1884
			<i>Scolecithricella</i> sp.			+	+		+	+	Sars G.O., 1902	<i>Scolecithricella</i>	Sars G.O., 1902
			<i>Scolecithricella minor</i>							+	Brady, 1883	<i>Scolecithricella minor</i>	Brady, 1883
			<i>Subeucalanus mucronatus</i>							+	Giesbrecht, 1888	<i>Subeucalanus mucronatus</i>	Giesbrecht, 1888
			<i>Subeucalanus subcrassus</i>						+		Giesbrecht, 1888	<i>Subeucalanus subcrassus</i>	Giesbrecht, 1888
			<i>Temora</i> sp.			+	+				Baird, 1850	<i>Temora</i>	Baird, 1850
			<i>Temora discaudata</i>						+	+	Giesbrecht, 1889	<i>Temora discaudata</i>	Giesbrecht, 1889
			<i>Temora longicornis</i>					+			Müller O.F., 1785	<i>Temora longicornis</i>	Müller O.F., 1785
			<i>Temora plumosa</i>										
			<i>Temora stylifera</i>						+		Dana, 1849	<i>Temora stylifera</i>	Dana, 1849
			<i>Temora turbinata</i>						+	+	Dana, 1849	<i>Temora turbinata</i>	Dana, 1849
			<i>Tortanus</i> sp.			+					Giesbrecht in G & Schmeil, 1898	<i>Tortanus</i>	Giesbrecht in G & Schmeil, 1898
			<i>Undinula vulgaris</i>		+				+		Dana, 1849	<i>Undinula vulgaris</i>	Dana, 1849
			Unidentified (Pontellidae, Scolecithricidae)										
Cyclopoida													
			<i>Halicyclops</i> sp.							+	Norman, 1903	<i>Halicyclops</i>	Norman, 1903
			<i>Oithona</i>			+	+		+		Baird, 1843	<i>Oithona</i>	Baird, 1843
			<i>Oithona attenuata</i>						+	+	Farran, 1913	<i>Oithona attenuata</i>	Farran, 1913
			<i>Oithona brevicornis</i>						+		Giesbrecht, 1891	<i>Oithona brevicornis</i>	Giesbrecht, 1891

Table 1. Cont'd.

PHYLUM	CLASS	ORDER	Reported scientific name	(1)	(2)	(3)	(4)	(5)	(6)	(7)	Authority	Currently accepted scientific name (Genus and lower) (8,9)	Authority for accepted name
Pocillostomatoida			<i>Oithona nana</i>						+	+	Giesbrecht, 1893	<i>Oithona nana</i>	Giesbrecht, 1893
			<i>Oithona oculata</i>						+	+	Farran, 1913	<i>Oithona oculata</i>	Farran, 1913
			<i>Oithona plumifera</i>						+		Baird, 1843	<i>Oithona plumifera</i>	Baird, 1843
			<i>Oithona rigida</i>						+	+	Giesbrecht, 1896	<i>Oithona rigida</i>	Giesbrecht, 1896
			<i>Oithona similis</i>						+	+	Claus, 1866	<i>Oithona similis</i>	Claus, 1866
			<i>Oithona simplex</i>						+	+	Farran, 1913	<i>Oithona simplex</i>	Farran, 1913
			<i>Oithona vivida</i>						+	+	Farran, 1913	<i>Oithona vivida</i>	Farran, 1913
			<i>Oithona longispina</i>						+	+	Nishida, Tanaka & Omori, 1977	<i>Oithona longispina</i>	Nishida, Tanaka & Omori, 1977
			<i>Oithona fallax</i>						+	+	Farran, 1913	<i>Oithona fallax</i>	Farran, 1913
			<i>Oithona dissimilis</i>						+	+	Lindberg, 1940	<i>Oithona dissimilis</i>	Lindberg, 1940
			<i>Oithona aruensis</i>							+	Früchtl, 1923	<i>Oithona aruensis</i>	Früchtl, 1923
			<i>Copilia quadrata</i>						+		Dana, 1849	<i>Copilia quadrata</i>	Dana, 1849
			<i>Copilia mirabilis</i>						+		Dana, 1852	<i>Copilia mirabilis</i>	Dana, 1852
			<i>Corycaeus</i>			+	+				Dana, 1845	<i>Corycaeus</i>	Dana, 1845
			<i>Corycaeus affinis</i>						+	+	McMurrich, 1916	<i>Corycaeus affinis</i>	McMurrich, 1916
			<i>Corycaeus agilis</i>						+	+	Dana, 1849	<i>Onychocorycaeus agilis</i>	Dana, 1849
			<i>Corycaeus andrewsi</i>						+	+	Farran, 1911	<i>Corycaeus andrewsi</i>	Farran, 1911
			<i>Corycaeus asiaticus</i>						+	+	Dahl F., 1894	<i>Corycaeus asiaticus</i>	Dahl F., 1894
			<i>Corycaeus catus</i>						+	+	Dahl F., 1894	<i>Onychocorycaeus catus</i>	Dahl F., 1894
			<i>Corycaeus dahli</i>						+	+	Tanaka, 1957	<i>Corycaeus dahli</i>	Tanaka, 1957
			<i>Corycaeus dubius</i>						+	+	Farran, 1911	<i>Corycaeus dubius</i>	Farran, 1911
			<i>Corycaeus erythraeus</i>						+	+	Cleve, 1904	<i>Corycaeus erythraeus</i>	Cleve, 1904
			<i>Corycaeus lautus</i>						+	+	Dana, 1849	<i>Corycaeus lautus</i>	Dana, 1849
			<i>Corycaeus limbatus</i>						+	+	Brady, 1883	<i>Corycaeus limbatus</i>	Brady, 1883
			<i>Corycaeus ovalis</i>					+			Claus, 1863	<i>Corycaeus ovalis</i>	Claus, 1863
			<i>Corycaeus pacificus</i>						+	+	Dahl F., 1894	<i>Onychocorycaeus pacificus</i>	Dahl F., 1894
			<i>Corycaeus speciosus</i>						+		Dana, 1849	<i>Corycaeus speciosus</i>	Dana, 1849
			<i>Corycaeus subtilis</i>						+		Dahl M., 1912	<i>Ditrichocorycaeus subtilis</i>	M.Dahl, 1912
			<i>Corycaeus pumilus</i>							+	Dahl, 1912	<i>Onychocorycaeus pumilus</i>	Dahl, 1912
			<i>Corycaeus giesbrechti</i>							+	Dahl, 1894	<i>Corycaeus giesbrechti</i>	Dahl, 1894
			<i>Corycaeus crassiusculus</i>							+	Dana, 1849	<i>Corycaeus crassiusculus</i>	Dana, 1849
			<i>Farranula gibbula</i>						+		Giesbrecht, 1891	<i>Farranula gibbula</i>	Giesbrecht, 1891

Table 1. Cont'd.

PHYLUM	CLASS	ORDER	Reported scientific name	(1)	(2)	(3)	(4)	(5)	(6)	(7)	Authority	Currently accepted scientific name (Genus and lower) (8,9)	Authority for accepted name
Harpacticoida			<i>Farranula rostratus</i>						+		Claus, 1863	<i>Farranula rostrata</i>	Claus, 1863
			<i>Oncaea</i>								Philippi, 1843	<i>Oncaea</i>	Philippi, 1843
			<i>Oncaea clevei</i>		+				+		Früchtl, 1923	<i>Oncaea clevei</i>	Früchtl, 1923
			<i>Oncaea media</i>						+		Giesbrecht, 1891	<i>Oncaea media</i>	Giesbrecht, 1891
			<i>Oncaea paraclevei</i>						+		Böttger-Schnack, 2001	<i>Oncaea paraclevei</i>	Böttger-Schnack, 2001
			<i>Oncaea scottodicarloi</i>						+		Heron & Bradford-Grieve, 1995	<i>Oncaea scottodicarloi</i>	Heron & Bradford-Grieve, 1995
			<i>Oncaea venusta</i>					+			Philippi, 1843	<i>Oncaea venusta</i>	Philippi, 1843
			<i>Pseudomacochiron</i> sp.					+			Reddiah, 1969	<i>Pseudomacochiron</i>	Reddiah, 1969
			<i>Sapphirina angusta</i>					+			Dana, 1849	<i>Sapphirina angusta</i>	Dana, 1849
			<i>Sapphirina gastrica</i>					+			Giesbrecht, 1891	<i>Sapphirina gastrica</i>	Giesbrecht, 1891
			<i>Sapphirina intestinata</i>					+			Giesbrecht, 1891	<i>Sapphirina intestinata</i>	Giesbrecht, 1891
			<i>Sapphirina metallina</i>						+		Dana, 1849	<i>Sapphirina metallina</i>	Dana, 1849
			<i>Clytemnestra scutellata</i>						+		Dana, 1849	<i>Clytemnestra scutellata</i>	Dana, 1849
			<i>Clytemnestra</i> sp.		+						Dana, 1848	<i>Clytemnestra</i>	Dana, 1848
			<i>Distiocolus minor</i>					+			Scott T., 1894	<i>Distiocolus minor</i>	Scott T., 1894
			<i>Ectinosoma</i>		+						Boeck, 1865	<i>Ectinosoma</i>	Boeck, 1865
			<i>Euterpina</i>		+	+					Norman, 1903	<i>Euterpina</i>	Norman, 1903
Monstrilloidea			<i>Euterpina acutifrons</i>						+		Dana, 1847	<i>Euterpina acutifrons</i>	Dana, 1847
			<i>Macrosetella</i>			+							
			<i>Macrosetella gracilis</i>		+			+			Brady & Robertson, 1873	<i>Microsetella</i>	Brady & Robertson, 1873
			<i>Microsetella</i>										
			<i>Microsetella norvegica</i>					+					
			<i>Microsetella rosea</i>		+			+					
			<i>Tisbintra</i> sp.								Sewell, 1940	<i>Tisbintra</i>	Sewell, 1940
			<i>Tisbe graciloides</i>					+			Sars G.O., 1920	<i>Tisbe graciloides</i>	Sars G.O., 1920
			<i>Monstrilla longiremis</i>					+			Giesbrecht, 1893	<i>Monstrilla longiremis</i>	Giesbrecht, 1893
Euphausiacea	Subclass Thecostraca	Malacostraca		+						+			
			Cirripedia (infra class)		+								
			Mysidacea Subclass Eumalacostraca, unranked)		+		+				Haworth, 1825	<i>Mysida</i>	Haworth, 1825
			<i>Euphausia tenera</i>							+	Dana, 1852 Hansen, 1905	<i>Euphausiacea</i> <i>Euphausia tenera</i>	Dana, 1852 Hansen, 1905

Table 1. Cont'd.

PHYLUM	CLASS	ORDER	Reported scientific name	(1)	(2)	(3)	(4)	(5)	(6)	(7)	Authority	Currently accepted scientific name (Genus and lower) (8,9)	Authority for accepted name
Chaetognatha	Sagittioidea	Aphragmophora	<i>Nematoscelis</i> sp.						+		Sars, 1883	<i>Nematoscelis</i>	Sars, 1883
			<i>Sylocheiron</i> sp.						+		Sars, 1883	<i>Sylocheiron</i>	
			Un-identified		+								
			Unidentified		+						Latreille, 1817	<i>Isopoda</i>	Latreille, 1817
			Unidentified		+						Latreille, 1816	<i>Amphipoda</i>	Latreille, 1816
			Unidentified		+						Krøyer, 1846	<i>Cumacea</i>	Krøyer, 1846
			Unidentified		+								
			Gastrosaccus pelagics							+	Li, 1964	<i>Gastrosaccus pelagica</i>	Li, 1964
			<i>Siriella media</i>							+	Hansen, 1910	<i>Siriella media</i>	Hansen, 1910
			<i>Anisomysis</i> sp.							+	Hansen, 1910	<i>Anisomysis</i>	Hansen, 1910
			<i>Acetes</i> sp.										
			<i>Lucifer</i> sp.	+									
			<i>Lucifer hansenii</i>										
			<i>Lucifer intermedius</i>										
			<i>Sylocheiron</i> sp.										
			Brachyura larvae	+									
			Macrura Larvae	+									
			Unidentified (larvae)	+				+					
			Unidentified	+				+					
Chaetognatha	Sagittioidea	Aphragmophora	<i>Sagitta</i>										
			<i>Sagitta enflata</i>		+						Quoy & Gaimard, 1827	<i>Sagitta</i>	Quoy & Gaimard, 1827
			<i>Sagitta bedoti</i>		+						Grassi, 1881	<i>Flaccisagitta enflata</i>	Grassi, 1881
			<i>Sagitta hexaptera</i>								Beraneck, 1895	<i>Zonosagitta bedoti</i>	Beraneck, 1895
			<i>Sagitta lyra</i>								d'Orbigny, 1836	<i>Flaccisagitta hexaptera</i>	d'Orbigny, 1836
			<i>Sagitta minima</i>								Krohn, 1853	<i>Pseudosagitta lyra</i>	Krohn, 1853
			<i>Sagitta pacifica</i>								Grassi, 1881	<i>Mesosagitta minima</i>	Grassi, 1881
			<i>Sagitta regularis</i>								Tokioka, 1940	<i>Aidanosagitta regularis</i>	Aida, 1897
			<i>Sagitta pseudoserratodentata</i>								Aida, 1897	<i>Serratiasagitta</i>	Aida, 1897
											Tokioka, 1939	<i>pseudoserratodentata</i>	Tokioka, 1939
			<i>Sagitta crassa</i>								Tokioka, 1938	<i>Aidanosagitta crassa</i>	Tokioka, 1938
			<i>Krohnittia pacifica</i>			+					Aida, 1897	<i>Krohnittia pacifica</i>	Aida, 1897

Table 1. Cont'd.

PHYLUM	CLASS	ORDER	Reported scientific name	(1)	(2)	(3)	(4)	(5)	(6)	(7)	Authority	Currently accepted scientific name (Genus and lower) (8,9)	Authority for accepted name
Mollusca	Gastropoda	Pteropoda (Super Order)	Unidentified		+			+					
			Unidentified		+			+					
		Littorinimorpha	<i>Carinaria</i> sp.		+					+	Lamarck, 1801	<i>Carinaria</i>	Lamarck, 1801
			Superfamily Pterotracheoidea (Heteropoda)		+						Rafinesque 1814	<i>Pterotracheoidea</i>	Rafinesque, 1814
Echinodermata	Bivalvia	Mytiloidea	Unidentified (larvae)	+				+					
			<i>Mytilus viridis</i> (larvae)				+				Linnaeus, 1758	<i>Perna viridis</i>	Linnaeus, 1758
			<i>Modiolus penelegans</i> (larvae)				+				Iredale, 1939	<i>Modiolus modulaides</i>	Röding, 1798
			Unidentified		+								
Hemichordata	Echinoidea	Appendicularia	Unidentified					+					
			Unidentified						+				
			Unidentified					+					
Chordata	Pisces (Super class)	Copelata	<i>Oikopleura cornutogastra</i>							+	Gegenbaur, 1855	<i>Oikopleura (Coecaria) fusiformis cornutogastra</i>	Gegenbaur, 1855
			<i>Oikopleura</i> sp.	+	+					+	Mertens, 1830	<i>Oikopleura</i>	Mertens, 1830
			Unidentified					+					
			Unidentified					+					

or presence of the different plankton groups (Table 1). He found that copepods were present or “very common” during the whole year. Loricated ciliates, tintinnids, were observed only during January, June, November, and December. The dinoflagellates *Ceratium* sp. and *Noctiluca* sp. were very common in February. The organisms classified as larger zooplankton, such as decapod larvae, siphonophores and chaetognaths were also described during the entire year studied by the author.

Wickstead (1958) published a study on the larger zooplankton of Singapore waters (Table 1). One station located at the Outer Shoal Light Buoy (1°15'N; 103°52'E; see Fig. 1) was sampled during one year for plankton enumeration. Climatic aspects, water temperature and salinity were also described. In the protozoa group, foraminifera and cystoflagellata (currently referred to as dinoflagellates) were found. Due to the sampling method, a “25 mesh per inch net” (i.e., 700 µm pore-size mesh), Tintinnidae were seldom found. Hydrozoa, Scyphozoa and Ctenophora were recorded in the group of Coelenterata. The polychaete annelids occurred occasionally throughout the year. The crustaceans were mainly represented by copepods in the groups of Gymnoplea, Podoplea, Harpacticoida and more rarely monstriloida. Cladocerans, ostracods, cirripedes, mysids, isopods, amphipods, euphausiids, decapods were much more uncommon (Wickstead, 1958). In this last work, the author showed a seasonal variation with almost all the planktonic groups being more abundant in the first half of the year and towards the end of the year, with the exception of copepods that only showed a peak of abundance in February/March. The author, however, points that due to the large pore size net used, copepods were likely to be underestimated. It is noteworthy that there are high numbers of Siphonophora found in the study, with this group being the most numerous during the second half of the year, as they likely preyed upon copepods.

Tham (1973) provided the general seasonal pattern of phyto- and zooplankton (Table 1). The author showed that zooplankton abundance was higher between March and November, and that the group was dominated by copepods, accounting for 40–70% of the total abundance. In his study, copepod peaked at April/May, July, and October/November. These findings partially contrast with those of Wickstead (1958), who described the highest biomass of copepods only at the beginning of the year (February/March). The little representation of copepods during the rest of the year found in Wickstead's (1958) survey could be due either to particular conditions during the sampling year or to a high predatory control of blooming siphonophores during this period, or even to a mere underestimation of the group as the author admits in his work. The other groups found were appendicularians, cirripedes, molluscs, decapod larvae, siphonophores and chaetognaths.

In that same year, Thia-Eng (1973) published an ecological study of the Ponggol Estuary (Table 1; see Fig. 1 for location of the estuary), a short, narrow and shallow mesohaline to polyhaline estuary, now transformed in a freshwater reservoir. In this particular environment, which is not comparable to

the literature discussed so far, dinoflagellates, copepods and molluscs, accounted for 80% of the total zooplankton biomass.

Over the last decade, only a few studies on zooplankton from Singapore waters have been published. Rezai et al. (2004) described the spatial and temporal distribution of 117 copepod species belonging to 37 genera in the Strait of Malacca (Table 1). Among these species, 13 accounted for 70% of the total copepod abundance. No consistent seasonal patterns of individual species were observed, but the authors found differences in abundance between the northern, the central, and the southern parts of the Strait. Nakajima et al. (2008) published a study on the zooplankton community of a coral reef, which was located at the Malaysia peninsula coast (Redang Island, North East Malaysia). The authors size-fractionated the organisms and found that the 100–200 µm fraction was composed of 51.3% adult copepods and copepodites and the rest (47.1%) of copepod nauplii. In the following fraction (200–335 µm), adult copepods and copepodites were also the more abundant (74.3%) of the total biomass but they also found other crustaceans, larvae and chaetognaths. They found that the larger organisms (>335 µm) were more diverse. In addition to adult copepods and copepodites, they found chaetognaths, larvae, hydrozoans, ostracods and other crustaceans.

In the study conducted as part of this paper, the station situated in the west part of the channel was observed to vary in the total abundance of copepods (orders Calanoida, Harpacticoid, and Cyclopoida, including copepodid larva) throughout the year. Two peaks in abundance in November 2012 and in March 2013 were observed. These highest abundances found in March and November and are in accordance with the work of Tham (1973). However, it was not clearly observed whether these changes in abundances occurred in the east part of the channel. Indeed, the total abundance of copepods did not draw any temporal pattern. It was only observed that an increase of the order Calanoida (which accounted for most of the total abundance of copepods) occurred in December 2012.

Different zooplankton groups of freshwater bodies from Singapore and Malaysia have also been described (Table 2). The intensive sampling was carried out during 1966–1967, and gave rise to several publications in the early eighties. The samples were collected in all types of freshwater habitats available, like ponds, lakes, marshes, reservoirs, pools, rice fields, rivers and streams in West Malaysia (a total of 503 zooplankton samples) and Singapore (a total of 38 samples). Lai & Fernando (1978) focused only on calanoid copepods, and Fernando & Ponzi (1981) studied the free-living freshwater cyclopoid copepods of Malaysia and Singapore (Table 2). In spite of extensive sampling, Fernando & Ponzi (1981) found a low number of species, which could be the result of the high human impact on the aquatic fauna. Another work, which came out of this extensive sampling, was that of Fernando & Zankai (1981) on Rotifera (Table 2). They described 165 species belonging mainly to Ploimida and Gnesiotrocha. The authors argued that the low record

of Bdelloida was due to an issue in sampling methodology. Most of the species found in this study were only observed in some samples. They observed that the diversity was higher in ponds and rice fields than in open waters of reservoirs and streams. The last work is from Idris & Fernando (1981) on Cladocera (Table 2). They described 63 cladoceran species from which, 26 were recorded for the first time in that region and they provided their detailed morphological description. Similar to Idris & Fernando (1981), the highest diversity of cladocerans was found in ponds and rice fields. They found that the species composition was typical of tropical region; that *Daphnia* was extremely rare, and that families Polyphemidae, Leptodoridae, and Holopedidae were absent. More recently, Low et al. (2010) described trophic links between phytoplankton and zooplankton organisms in freshwater reservoirs from Singapore. They based their study on an historical data set over 15 years, between January 1992 and December 2006, of a monthly sampling regime in 12 shallow reservoirs. They categorised zooplanktonic organisms in six groups. Four of them, calanoids, cyclopoids, cladocerans and rotifers, were present in more than 98% of the samples. However, two groups, harpacticoids and ostracods, were rare. Their study did not provide a species-level list for zooplankton, nor their geographical and seasonal distribution.

Species richness as descriptor of diversity. For a better understanding of planktonic historical trends, the use of quantitative diversity descriptors would have been ideal; unfortunately, most of the available data are only qualitative. However, a comparison of the species richness among can be undertaken for the different studies. For freshwater habitats, unfortunately, this is not possible because the disparity of target groups amongst studies; as, each study focused on one group. Regarding marine systems, even if the data are limited it is evident that Rezai et al. (2004) found more species than all the other studies (a total of 112 species versus 8–32 species for the rest of studies). Unfortunately, the work of Rezai et al. (2004) is limited to the description of copepods, and does not include other zooplankters. This exemplifies the lack of broad-based studies considering all zooplanktonic groups at high taxonomic resolution.

The present study included an extensive plankton investigation, conducted once every two weeks during one year (Table 1). We found differences in the species richness of some groups compared to that of previous works. For instance, the species richness for cnidarians and copepods was higher than that found in studies from the fifties, but it was lower than the one described by Rezai (Rezai et al., 2004; Table 1). However, we are closer in total number of zooplankton species to the work of Rezai et al. (2004), as we found, a total of 91 species including species from all groups (Table 1). These discrepancies between the previous work and the present study may simply be result of the different geographic locations of the sampling sites. In the present study, the west coast of Singapore was studied (e.g., in the area of Jurong Island), while Rezai (Rezai et al., 2004) studied the Straits of Malacca, which covers a wide geographic area, and Nakajima et al. (2008) undertook their survey in the north-east of Malaysia. Our sampling sites

have a lower water circulation rate and are more affected by human activities, which should affect the planktonic community. Nevertheless, considering the particularities of the sampling stations, it is rather surprising that the total number of zooplanktonic species found in our study is similar, or even higher, to those found in previous reports.

Among the 57 species of copepods found, 32 belong to the order Calanoida, 12 to the order Cyclopoida, eight to the Poecilostomatoida, and five to the order Harpacticoida. Some copepods from our study have been described only from this region. *Acartia pacifica* Steuer, 1915, *Acartia sinijiensis* Mori, 1940 and *Bestiolina similis* (Sewell, 1914) are found only in the Pacific. Other species have a more broad distribution, such as *Acrocalanus gibber* Giesbrecht, 1888, and *Acrocalanus gracilis* Giesbrecht, 1888, which are found mainly in the Indian Ocean, but also in Pacific and Atlantic oceans. Some are cosmopolitan (found in all oceans except polar zones), like *Calocalanus styliformis* (Giesbrecht, 1888), and *Clausocalanus furcatus* (Brady, 1883). In all cases, the calanoid copepods found in the present survey all have a tropical to temperate distribution. However, cyclopoid copepods such as *Oithona similis* Claus, 1866, are cosmopolitan, including polar zones. Chaetognaths, also important components of the zooplankton in our study, have a species-dependent distribution: four of our species are found in all tropical to temperate oceans, e.g., *Flaccisagitta enflata* (Grassi, 1881), and five of our species are limited to the Indian and Pacific oceans, e.g., *Sagitta bedoti* (Beraneck, 1895). Their temporal variation showed a similar trend to that of copepods at both stations; they were present the whole year in the study of Tham (1953).

Final remarks on the need of long term plankton-monitoring programs. Data on zooplankton diversity and distribution in the area of Singapore (including combined works with Malaysia) have been mostly obtained at the beginning and the middle of the last century. It is only recently, at the beginning of the 2000s, that investigations in this field have restarted in Malaysia and only in 2013 in Singapore. This review was therefore necessary to update the information available to detect possible trends associated with climatic or anthropogenic events, and to inform an efficient water management plan for Singapore. However, because of the changes in sampling techniques and methods, the heterogeneity of sampling stations and groups studied, the comparisons are not obvious.

From a conservation biology point of view, it is important to compare the state of actual ecosystems to their description in the past. Without a proper description of biodiversity and its functioning over time, it is difficult to ensure appropriate ecosystem management. Many factors, such as the construction of dams, the causeway linking Singapore to Johor Bahru (Malaysia), shoreline modifications, and species introductions (e.g., from ballast water or aquaculture) may have affected native aquatic biodiversity. For instance, it has been shown that ballast waters are responsible for ca. 30% of bio-invasions, the risk of dispersal of exogenous species increasing yearly by approximately 3% (Gollasch, 2007;

Table 2. List of zooplanktonic organisms found in the available literature on Singaporean and Malaysian freshwaters. (1) = Fernando & Pongyi, 1981; (2) = Fernando & Zankai, 1981; (3) = Idris & Fernando, 1981; (4) = Lai & Fernando, 1978; (5) = Low et al., 2010; (6) = World Register of Marine Species at <http://www.marinespecies.org/aphia.php?p=taxdetails&id=220933> on 2013-01-23.

Reported scientific name	(1)	(2)	(3)	(4)	(5)	Authority	Currently accepted scientific name (Genus and lower) (6)	Authority for accepted name
Phylum Arthropoda								
Order Cladocera					+			
Family Silidae								
Genus <i>Diaphanosoma</i>						Fisher, 1850		
<i>Diaphanosoma modigliani</i>			+			Richard, 1895		
<i>Diaphanosoma excisum</i>			+			Sars, 1885	<i>Diaphanosoma excisum</i>	Sars, 1885
<i>Diaphanosoma sarsi</i>			+			Richard, 1895		
<i>Diaphanosoma aspinosum</i>			+			Chiang, 1956		
Genus <i>Pseudosida</i>						Herrick, 1884		
<i>Pseudosida bideniata</i>			+			Herrick, 1884		
Genus <i>Latonopsis</i>						Sars, 1888		
<i>Latonopsis australis</i>			+			Sars, 1888		
Family Daphniidae								
Genus <i>Daphnia</i>						O.F. Muller, 1785		
<i>Daphnia similis</i>			+			Claus, 1876		
Genus <i>Ceriodaphnia</i>						Dana, 1853		
<i>Ceriodaphnia cornuta</i>			+			Sars, 1888	<i>Ceriodaphnia cornuta</i>	Sars, 1885
Genus <i>Scapholeberis</i>						Schoedler, 1853		
<i>Scapholeberis kingi</i>			+			Sars, 1903		
Genus <i>Simocephalus</i> S						Schoedler, 1858		
<i>Simocephalus serrulatus</i>			+			(Koch, 1841)	<i>Simocephalus serrulatus</i>	(Koch, 1841)
<i>Simocephalus latirostris</i>			+			Stingelin, 1906		
Family Moinidae								
Genus <i>Moinodaphnia</i>						Herrick, 1887		
<i>Moinodaphnia macleayi</i>			+			(King, 1853)		
Genus <i>Moina</i>						Baird, 1850		
<i>Moina micrura</i>			+			Kurz, 1874	<i>Moina micrura</i>	Kurz, 1874
Family Bosminidae								
Genus <i>Bosminopsis</i>						Richard, 1895		
<i>Bosminopsis deitersi</i>			+			Richard, 1895		
Family Macrothricidae								
Genus <i>Ilyocryptus</i>						Sars, 1862		
<i>Ilyocryptus spinifer</i>			+			Herrick, 1882		
Genus <i>Grimaldina</i>						Richard, 1892		
<i>Grimaldina brazzai</i>			+			Richard, 1892		
Genus <i>Guernella</i>						Richard, 1892		

Table 2. Cont'd.

Reported scientific name	(1)	(2)	(3)	(4)	(5)	Authority	Currently accepted scientific name (Genus and lower) (6)	Authority for accepted name
<i>Guernella raphalis</i>		+				Richard, 1892		
Genus <i>Streblocercus</i>						Sars, 1862		
<i>Streblocercus pygmaeus</i>		+				Sars, 1901		
Genus <i>Macrothrix</i>						Baird, 1843		
<i>Macrothrix spinosa</i>		+				King, 1852		
<i>Macrothrix capensis-monodi</i>		+				Gauthier, 1930		
<i>Macrothrix triserialis</i>		+				Brady, 1886		
Family Chydoridae								
Genus <i>Pleuroxus</i>						Baird, 1843		
Pleuroxus laevis		+				Sars, 1862		
Genus <i>Disparalona</i>						Fryer, 1968		
<i>Disparalona rostrata</i>		+				(Koch, 1841)	<i>Disparalona rostrata</i>	(Koch, 1841)
Genus <i>Alonella</i>						Sars, 1862		
<i>Alonella nana</i>		+				(Baird, 1850)		
<i>Alonella excisa</i>		+				(Fisher, 1854)		
<i>Alonella hamulatus</i>		+				(Birge, 1910)		
Genus <i>Chydorus</i>						Leach, 1816		
<i>Chydorus barroisi</i>		+				Richard, 1894		
<i>Chydorus ventricosus</i>		+				Daday, 1898		
<i>Chydorus parvus</i>		+				Daday, 1898		
<i>Chydorus eurynotus</i>		+				Sars, 1901		
<i>Chydorus</i> cf. <i>pubescens</i>		+				Sars, 1901		
<i>Chydorus reticulatus</i>		+				Daday, 1898		
<i>Chydorus faviformis</i>		+				Birge, 1893		
Genus <i>Pseudochydorus</i>						Fryer, 1968		
<i>Pseudochydorus globosus</i>		+				(Baird, 1893)		
Genus <i>Dunhevedia</i>						King, 1853		
<i>Dunhevedia crassa</i>		+				King, 1853		
<i>Dunhevedia serrata</i>		+				Daday, 1898		
Genus <i>Dadaya</i>						Sars, 1901		
<i>Dadaya macrops</i>		+				(Daday, 1898)		
Genus <i>Leydigia</i>						Kurz, 1875		
<i>Leydigia acanthocercoides</i>		+				(Fisher, 1854)	<i>Leydigia acanthocercoides</i>	(Fisher, 1854)
Genus <i>Graptoleberis</i>						Sars, 1862		
<i>Graptoleberis testudinaria</i>		+				(Fisher, 1851)	<i>Graptoleberis testudinaria</i>	(Fisher, 1848)
Genus <i>Alona</i>						Baird, 1843		
<i>Alona quadrangularis</i>		+				(O.F. Mueller, 1785)	<i>Alona quadrangularis</i>	(O.F. Mueller, 1776)

Table 2. Cont'd.

Reported scientific name	(1)	(2)	(3)	(4)	(5)	Authority	Currently accepted scientific name (Genus and lower) (6)	Authority for accepted name
<i>Alona affinis</i>			+			Leydig, 1860	<i>Alona affinis</i>	Leydig, 1860
<i>Alona eximia</i>			+			Kiser, 1948		
<i>Alona intermedia</i>			+			Sars, 1862		
<i>Alona cf. dentifera</i>			+			Sars, 1901		
<i>Alona macronyx</i>			+			Daday, 1898		
<i>Alona davidi</i>			+			Richard, 1895		
<i>Alona cf. sarasinorum</i>			+			Stingelin, 1900		
<i>Alona costata</i>			+			Sars, 1862	<i>Alona costata</i>	Sars, 1862
<i>Alona guttata</i>			+			Sars, 1862	<i>Alona guttata</i>	Sars, 1862
<i>Alona cf. karelica</i>			+			Stenroos		
<i>Alona cf. pulchella</i>			+			King, 1853		
<i>Alona karua</i>			+			(King, 1853)		
<i>Alona monacantha</i>			+			Sars, 1901		
<i>Alona rectangula</i>			+			Sars, 1862	<i>Alona rectangula</i>	Sars, 1861
<i>Alona verrucosa</i>			+			Sars, 1901		
<i>Alona freyi</i>			+			Idris & Fernando, 1980		
Genus <i>Kurzia</i>						Dybowski & Growchowski, 1894		
<i>Kurzia longirostris</i>			+			(Daday, 1898)		
Genus <i>Acroperus</i>						Baird, 1843		
<i>Acroperus harpae</i>			+			(Baird, 1834)	<i>Acroperus harpae</i>	(Baird, 1834)
Genus <i>Oxyurella</i>						Dybowski & Growchowski, 1894		
<i>Oxyurella sinhalensis</i>			+			(Daday, 1898)		
Genus <i>Euryalona</i>						Sars, 1901		
<i>Euryalona orientalis</i>			+			Baird, 1843		
Genus <i>Camptocercus</i>						Baird, 1843		
<i>Camptocercus cf. australis</i>			+			Sars, 1896		
Genus <i>Indialona</i>						Petkovski, 1966		
<i>Indialona globulosa</i>						(Daday, 1898)		
Order Calanoid				+				
<i>Neodiaptomus handeli</i>				+		Brehm, 1921	<i>Neodiaptomus schmackeri</i>	(Brehm, 1921)
<i>Neodiaptomus blachei</i>				+		Brehm, 1951	<i>Vietodiaptomus blachei</i>	Brehm, 1951
<i>Neodiaptomus botulifer</i>				+		Kiefer, 1974	<i>Mongolodiaptomus botulifer</i>	Kiefer, 1974
<i>Neodiaptomus laii</i>				+		Kiefer, 1974	<i>Neodiaptomus laii</i>	Kiefer, 1974
<i>Neodiaptomus malaindosenensis</i>				+		Lai & Fernando, 1981	<i>Mongolodiaptomus botulifer</i>	Lai & Fernando, 1977
<i>Neodiaptomus meggitti</i>				+		Kiefer, 1932	<i>Neodiaptomus meggitti</i>	Kiefer, 1932
<i>Neodiaptomus mephistopheles</i>				+		Brehm, 1933	<i>Mongolodiaptomus mephistopheles</i>	Brehm, 1933
<i>Pseudodiaptomus danglishi</i>				+		Sewell, 1932	<i>Pseudodiaptomus danglishi</i>	Sewell, 1932

Table 2. Cont'd.

Reported scientific name	(1)	(2)	(3)	(4)	(5)	Authority	Currently accepted scientific name (Genus and lower) (6)	Authority for accepted name
<i>Pseudodiaptomus tollingerae</i>				+		Sewell, 1924	<i>Pseudodiaptomus tollingerae</i>	Sewell, 1919
<i>Tropodiaptomus</i> spp.				+			<i>Tropodiaptomus</i>	Kiefer, 1932
Order Cyclopoid					+			
<i>Cryptocyclops bicolor</i>	+					(Sars, 1863)	<i>Microcyclops bicolor</i>	(Sars, 1863)
<i>Eutocyclops phaleratus</i>	+					(Koch, 1838)		
<i>Eucyclops serrulatus</i>	+					(Fisher, 1851)	<i>Eucyclops serrulatus</i>	(Fisher, 1851)
<i>Macrocyclus albidus</i>	+					(Jurine, 1820)	<i>Macrocyclus albidus</i>	(Jurine, 1820)
<i>Macrocyclus distinctus</i>	+					(Richard, 1887)	<i>Macrocyclus distinctus</i>	(Richard, 1887)
<i>Macrocyclus fusus</i>	+					(Jurine, 1820)	<i>Macrocyclus fusus</i>	(Jurine, 1820)
<i>Mesocyclops leuckarti</i>	+					(Claus, 1857)	<i>Mesocyclops leuckarti</i>	(Claus, 1857)
<i>Metacyclops minutus</i>	+					(Claus, 1863)	<i>Metacyclops minutus</i>	(Claus, 1863)
<i>Microcyclus dengizicus</i>	+					(Lepeschkin, 1900)	<i>Microcyclus dengizicus</i>	(Lepeschkin, 1900)
<i>Microcyclus varicans</i>	+					(Sars, 1863)	<i>Microcyclus varicans</i>	(Sars, 1863)
<i>Paracyclops affinis</i>	+					(Sars, 1863)	<i>Paracyclops affinis</i>	(Sars, 1863)
<i>Paracyclops fimbriatus</i>	+					(Fisher, 1853)	<i>Paracyclops fimbriatus</i>	(Fisher, 1853)
<i>Thermocyclops crassus</i>	+					(Fisher, 1853)	<i>Thermocyclops crassus</i>	(Fisher, 1853)
<i>Thermocyclops</i> cf. <i>schmeili</i>	+					(Poppe & Mrázek, 1895)	<i>Thermocyclops</i> cf. <i>schmeili</i>	(Poppe & Mrázek, 1895)
<i>Tropocyclops prasinus</i>	+					(Fisher, 1860)	<i>Tropocyclops prasinus</i>	(Fisher, 1860)
Phylum Rotifera					+			
Order Ploimida								
Family Ephiphanidae								
<i>Epiphanes clavulata</i>		+				(Ehrenberg, 1832)		
<i>Epiphanes macrourus</i>		+				(Barrois & Daday, 1894)	<i>Epiphanes macrourus</i>	(Barrois & Daday, 1894)
Family Brachionidae								
<i>Anuraeopsis fissa fissa</i>		+				(Gosse, 1851)		
<i>Anuraeopsis navicula coelata</i>		+				De Beauchamp, 1832		
<i>Brachionus angularis angularis</i>		+				Gosse, 1851	<i>Brachionus angularis angularis</i>	Gosse, 1851
<i>Brachionus angularis</i> f. <i>bidens</i>		+				(Plate, 1886)		
<i>Brachionus bidentata bidentata</i>		+				Anderson, 1889		
<i>Brachionus budapestinensis budapestinensis</i>		+				(Daday, 1885)		
<i>Brachionus calyciflorus calyciflorus</i>		+				Pallas, 1766	<i>Brachionus calyciflorus calyciflorus</i>	Pallas, 1776
<i>Brachionus calyciflorus</i> f. <i>anuraeiformis</i>		+				(Brehm, 1909)		
<i>Brachionus calyciflorus</i> f. <i>amphiceros</i>		+				(Ehrenberg, 1838)	<i>Brachionus calyciflorus calyciflorus</i>	Pallas, 1776
<i>Brachionus caudatus</i>		+				Barrois & Daday, 1894		
<i>Brachionus caudatus</i> f. <i>apsteini</i>		+				Fedeev, 1925		

Table 2. Cont'd.

Reported scientific name	(1)	(2)	(3)	(4)	(5)	Authority	Currently accepted scientific name (Genus and lower) (6)	Authority for accepted name
<i>Brachionus caudatus</i> f. <i>vulgatus</i>		+				Ahlstrom, 1940	<i>Brachionus diversicornis</i>	(Daday, 1883)
<i>Brachionus diversicornis diversicornis</i>		+				(Daday, 1883)		
<i>Brachionus falcatus falcatus</i>		+				Zacharias, 1898		
<i>Brachionus forficula</i>		+				Wierzejski, 1891		
<i>Brachionus forficula</i> f. <i>minor</i>		+				(Voronkov, 1913)		
<i>Brachionus leydigi</i> var. <i>rotundus</i>		+				(Rousselet, 1907)		
<i>Brachionus patulus patulus</i>		+				(O.F. Mueller, 1786)		
<i>Brachionus patulus</i> var. <i>macracanthus</i>		+				(Daday, 1905)		
<i>Brachionus plicatilis plicatilis</i>		+				(O.F. Mueller, 1786)		
<i>Brachionus plicatilis</i> f. <i>rotundiformis</i>		+				(Tschugunoff, 1921)		
<i>Brachionus plicatilis</i> f. <i>longicornis</i>		+				(Fedeev, 1925)		
<i>Brachionus quadridentatus quadridentatus</i>		+				(Hermann, 1783)		
<i>Brachionus quadridentatus</i> f. <i>melheni</i>		+				(Barrois & Daday, 1894)		
<i>Brachionus quadridentatus</i> f. <i>brevispinus</i>		+				(Ehrenberg, 1832)		
<i>Brachionus quadridentatus</i> f. <i>rhenanus</i>		+				(Lauterborn, 1893)		
<i>Brachionus quadridentatus</i> var. <i>ancylognathus</i>		+				(Schmarda, 1859)		
<i>Brachionus quadridentatus</i> var. <i>cluniorbicularis</i>		+				(Skorikov, 1894)	<i>Brachionus quadridentatus quadridentatus</i>	Hermann, 1783
<i>Brachionus quadridentatus mirabilis</i>		+				(Daday, 1897)		
<i>Brachionus quadridentatus zernovi</i>		+				Voronkov, 1907		
<i>Brachionus urceolaris</i>		+				(O.F. Mueller, 1773)		
<i>Brachionus urceolaris</i> var. <i>rubens</i>		+				(Ehrenberg, 1838)		
<i>Brachionus urceolaris</i> var. <i>sessilis</i>		+				(Varga, 1951)		
<i>Brachionus urceolaris nilsoni</i>		+				(Ahlstrom, 1940)		
<i>Brachionus variabilis</i>		+				Hempel, 1896		
<i>Keratella cochlearis cochlearis</i>		+				(Gosse, 1851)		
<i>Keratella cochlearis</i> var. <i>tecta</i>		+				(Lauterborn, 1900)	<i>Keratella cochlearis cochlearis</i>	(Gosse, 1851)
<i>Keratella javana</i>		+				Hauer, 1932		
<i>Keratella lenzi lenzi</i>		+				(Hauer, 1953)		
<i>Keratella procurva procurva</i>		+				(Thorpe, 1891)		
<i>Keratella tropica tropica</i>		+				(Apstein, 1907)		
<i>Platygaster quadricornis</i>		+				(Ehrenberg, 1832)		
Family Eucalanidae								
<i>Beauchampella eudactylota eudactylota</i>		+				(Gosse, 1886)		
<i>Dipleuchlanis propatula propatula</i>		+				(Gosse, 1886)		
<i>Dipleuchlanis propatula</i> f. <i>macroductyla</i>		+				(Hauer, 1965)		
							<i>Keratella tropica</i>	(Apstein, 1907)
							<i>Platygaster quadricornis</i>	(Ehrenberg, 1832)

Table 2. Cont'd.

Reported scientific name	(1)	(2)	(3)	(4)	(5)	Authority	Currently accepted scientific name (Genus and lower) (6)	Authority for accepted name
<i>Euchlanis alata</i>		+				Voronkov, 1912		
<i>Euchlanis callysta</i>		+				Myers, 1930		
<i>Euchlanis contorta</i>		+				Wulfert, 1939		
<i>Euchlanis deflexa deflexa</i>		+				(Gosse, 1851)		
<i>Euchlanis dilatata dilatata</i>		+				Ehrenberg, 1832		
<i>Euchlanis dilatata</i> f. <i>luckstiana</i>		+				Hauer, 1930		
<i>Euchlanis incisa incisa</i>		+				Carlin, 1939		
<i>Euchlanis oropha</i>		+				Gosse, 1887		
Family Mytilinidae								
<i>Mytilina acanthophora</i>		+				Hauer, 1938		
<i>Mytilina bisulcata</i>		+				(Luks, 1912)		
<i>Mytilina ventralis ventralis</i>		+				(Ehrenberg, 1832)		
Family Trichotridae								
<i>Macrochaetus collinsi collinsi</i>		+				(Gosse, 1867)		
<i>Macrochaetus sericus</i>		+				(Thorpe, 1893)		
<i>Trichotria pocillum</i>		+				(O.F. Mueller, 1776)	<i>Trichotria pocillum</i>	(Müller, 1766)
<i>Trichotria terractis</i> var. <i>similis</i>		+				(Stenroos, 1898)		
Family Colurellidae								
<i>Colurella uncinata</i> f. <i>bicuspidata</i>		+				(Ehrenberg, 1832)		
<i>Lepadella apsicora</i>		+				Myers, 1934		
<i>Lepadella monodactyla</i> f. <i>caudata</i>		+				(Koste, 1972)		
<i>Lepadella ovalis</i>		+				O.F. Mueller, 1786	<i>Lepadella ovalis</i>	(O.F. Muller, 1896)
<i>Lepadella patella patella</i>		+				O.F. Mueller, 1776		
<i>Lepadella rhomboides rhomboides</i>		+				Gosse, 1886		
Family Lecanidae								
<i>Lecane (Lecane) arcuata</i>		+				Harring, 1914	<i>Lecane (Lecane) arcuata</i>	(Bryce, 1891)
<i>Lecane (L.) aspasia</i>		+				Myers, 1917		
<i>Lecane (L.) crepida crepida</i>		+				Harring, 1914		
<i>Lecane (L.) curvicornis curvicornis</i>		+				(Murray, 1913)		
<i>Lecane (L.) curvicornis nitida</i>		+				(Murray, 1913)		
<i>Lecane (L.) doryssa</i>		+				Harring, 1914		
<i>Lecane (L.) hastata</i>		+				(Murray, 1913)	<i>Lecane (L.) hamata</i>	(Stokes, 1896)
<i>Lecane (L.) hornemanni</i>		+				(Ehrenberg, 1834)		
<i>Lecane (L.) inermis</i>		+				(Bryce, 1892)	<i>Lecane (L.) inermis</i>	(Bryce, 1892)
<i>Lecane (L.) leontina</i>		+				(Turner, 1893)		
<i>Lecane (L.) ludwigi ludwigi</i>		+				(Eckstein, 1893)		

Table 2. Cont'd.

Reported scientific name	(1)	(2)	(3)	(4)	(5)	Authority	Currently accepted scientific name (Genus and lower) (6)	Authority for accepted name
<i>Lecane (L.) luna luna</i>		+				(O.F. Mueller, 1776)		
<i>Lecane (L.) ohioensis</i>		+				(Herrick, 1885)		
<i>Lecane (L.) papuana</i>		+				(Murray, 1913)		
<i>Lecane (L.) pusilla</i>		+				Harring & Myers, 1914		
<i>Lecane (L.) saginata</i>		+				Harring & Myers, 1926		
<i>Lecane (L.) signifera ploenensis</i>		+				(Voigt, 1902)		
<i>Lecane (L.) cingulata ungulata</i>		+				(Gosse, 1887)		
<i>Lecane (L.) verecunda</i>		+				Harring & Myers, 1926		
<i>Lecane (Hemimonestyla) inopinata f. sympoda</i>		+				(Hauer, 1929)		
<i>Lecane (Monostyla) arcuata</i>		+				(Bryce, 1891)	<i>Lecane (Monostyla) arcuata</i>	(Bryce, 1891)
<i>Lecane (M.) bulla bulla</i>		+				(Gosse, 1886)		
<i>Lecane (M.) bulla diabolica</i>		+				(Hauer, 1936)		
<i>Lecane (M.) closteroerca closteroerca</i>		+				(Schmarda, 1859)		
<i>Lecane (M.) decipiens</i>		+				(Daday, 1913)	<i>Lecane (M.) decipiens</i>	(Murray, 1913)
<i>Lecane (M.) elachis</i>		+				Harring & Myers, 1926		
<i>Lecane (M.) lamellata thalera</i>		+				(Harring & Myers, 1926)		
<i>Lecane (M.) lunaris crenorta</i>		+				(Harring, 1913)		
<i>Lecane (M.) lunaris lunaris</i>		+				(Ehrenberg, 1832)		
<i>Lecane (M.) monostyla</i>		+				(Daday, 1897)		
<i>Lecane (M.) pyriformis</i>		+				(Daday, 1905)		
<i>Lecane (M.) stenroosi</i>		+				(Meissner, 1908)		
<i>Lecane (M.) unquitata unquitata</i>		+				(Fadeew, 1925)		
Family Proalidae								
<i>Proales</i> sp.		+				similar to Hauer's (1938)	<i>Proales</i> sp.	Gosse, 1886
Family Notommatidae								
<i>Cephalodella gibba gibba</i>		+				(Ehrenberg, 1838)		
<i>Cephalodella tenuior tenuior</i>		+				(Gosse, 1886)		
<i>Eosphora anthadis</i>		+				(Harring & Myers, 1922)		
<i>Eosphora elongata</i>		+				(Ehrenberg, 1832)		
<i>Itura myersi</i>		+				Wulfert, 1935		
<i>Monommata aequalis</i>		+				(Ehrenberg, 1832)		
<i>Monommata maculata</i>		+				Harring & Myers, 1924		
<i>Monommata grandis</i>		+				Tessin, 1890		
<i>Notommata allantois</i>		+				Wulfert, 1935		
<i>Notommata copeus</i>		+				Ehrenberg, 1834		
<i>Notommata endoxa</i>		+				Myers, 1933		

Table 2. Cont'd.

Reported scientific name	(1)	(2)	(3)	(4)	(5)	Authority	Currently accepted scientific name (Genus and lower) (6)	Authority for accepted name
<i>Notommata saccigera</i>		+				Ehrenberg, 1832		
<i>Notommata tripus</i>		+				Ehrenberg, 1838		
<i>Scaridium longicaudum</i>		+				(O.F. Mueller, 1786)		
Family Trichocercidae								
<i>Trichocerca bicristata bicristata</i>		+				(Gosse, 1887)		
<i>Trichocerca cylindrica chattoni</i>		+				(De Beauchamp, 1907)		
<i>Trichocerca dixon-nuttali</i>		+				(Jennings, 1903)		
<i>Trichocerca elongata</i>		+				(Gosse, 1886)		
<i>Trichocerca elongata braziliensis</i>		+				(Murray, 1913)		
<i>Trichocerca flagella</i>		+				Hauer, 1937		
<i>Trichocerca jenningsi</i>		+				Voigt, 1957		
<i>Trichocerca myersi</i>		+				(Hauer, 1931)		
<i>Trichocerca pusilla</i>		+				(Lauterborn, 1898)		
<i>Trichocerca rattus</i>		+				(O.F. Mueller, 1776)	<i>Trichocerca pusilla</i>	(Jennings, 1903)
<i>Trichocerca similis similis</i>		+				(Wierzejski, 1893)	<i>Trichocerca rattus</i>	(Müller, 1776)
<i>Trichocerca stylata</i>		+				(Gosse, 1851)	<i>Trichocerca stylata</i>	(Gosse, 1851)
<i>Trichocerca tigris</i>		+				(O.F. Mueller, 1786)		
Family Synchaetidae								
<i>Ploesoma lenticulare</i>		+				Herrick, 1885		
<i>Polyarthra vulgaris vulgaris</i>		+				Carlin, 1943		
<i>Synchaeta pectinata</i>		+				(Ehrenberg, 1832)	<i>Synchaeta pectinata</i>	Ehrenberg, 1832
Family Asplanchnidae								
<i>Asplanchna brightwelli</i>		+				Gosse, 1850	<i>Asplanchna brightwellii</i>	Gosse, 1850
<i>Asplanchna girodi</i>		+				(De Guerne, 1888)	<i>Asplanchna girodi</i>	De Guerne, 1888
<i>Asplanchna sieboldi</i>		+				(Leydig, 1856)		
<i>Asplanchna multiceps</i>		+				Schrank, 1793		
<i>Harringia rousseleti</i>		+				Herrick, 1885		
Family Dicranophoridae								
<i>Dicranophorus caudatus caudatus</i>		+				(Ehrenberg, 1834)		
<i>Dicranophorus claviger</i>		+				(Hauer, 1965)		
<i>Dicranophorus epicharis</i>		+				Harring & Myers, 1928		
<i>Dicranophorus robustus</i>		+				Harring & Myers, 1928		
<i>Encentrum diglandula</i>		+				(Zawodowski, 1926)		
Family Testudinellidae								
<i>Pompholyx complanata</i>		+				Gosse, 1851		
<i>Pompholyx sulcata</i>		+				Hudson, 1885	<i>Pompholyx sulcata</i>	Hudson, 1885

Table 2. Cont'd.

Reported scientific name	(1)	(2)	(3)	(4)	(5)	Authority	Currently accepted scientific name (Genus and lower) (6)	Authority for accepted name
<i>Testudinella parva</i>		+				(Ternetz, 1892)		
<i>Testudinella patina patina</i>		+				(Hermann, 1783)		
<i>Testudinella patina tribolata</i>		+				(Anderson & Shepard, 1982)		
<i>Testudinella tridentata curvata</i>		+				(Wulfert, 1965)		
Family Floscularidae								
<i>Conochilus dossuarius dossuarius</i>		+				(Hudson, 1875)		
<i>Conochilus dossuarius dossuarius</i> var. <i>coenobasis</i>		+				(Skorikov, 1914)		
<i>Hexarthra intermedia intermedia</i>		+				Wisniewski, 1929		
<i>Hexarthra mira</i>		+				(Hudson, 1871)		
<i>Limnias melicerta melicerta</i>		+				Weisse, 1848		
<i>Sinantherina spinosa</i>		+				(Thorpe, 1893)		
<i>Sinantherina procera</i>		+				(Thorpe, 1893)		
Family Filinidae								
<i>Filinia longiseta longiseta</i>		+				(Ehrenberg, 1834)		
<i>Filinia pejeri</i>		+				Hutchinson, 1965		
<i>Filinia terminalis</i>		+				(Plate, 1886)		
<i>Filinia opoliensis opoliensis</i>		+				(Zacharias, 1898)		
Bdelloidea								
<i>Rotaria rotaria</i>		+				(Pallas, 1766)		
<i>Rotaria neptunia</i>		+				(Ehrenberg, 1832)		
<i>Philodina</i> sp.		+						
<i>Dissotrocha</i> sp.		+						

UNCTAD, 2012). We thus argue that a continuous monitoring of plankton communities in Singapore would be a great asset for the region. These long-term monitoring programs exist in other areas of the world and allow for continuous information on ecosystem response to environmental changes, which is especially needed in zones heavily industrialised and very sensitive to climate change events, such as Singapore.

ACKNOWLEDGEMENTS

The authors would like to thank the National Parks of Singapore, the DHI-NTU Research Centre for the financial support of project MadeInPlankton, where the present study is framed. The work was also supported by Elite Forsk grants nb 10-093759 and 10-094773 from the Danish Agency for Science Technology and Innovation to GD, and by project PROTOS (CTM2009-08783) from the Spanish Ministry of Science and Innovation to AC. Also, N. Chan is thanked for linguistic improvements of the manuscript.

LITERATURE CITED

- Behera MR, Chun C, Palani S & Tkalic P (2013) Temporal variability and climatology of hydrodynamic, water property and water quality parameters in the West Johor Strait of Singapore. *Marine Pollution Bulletin*, 77(1–2): 380–395.
- Beranek E (1895) Les chétognathes de la baie d'Amboine (voyage de MM. M. Bedot et C. Pictet dans l'archipel Malais). *Revue Suisse de Zoologie et Annales du Musée d'Histoire naturelle de Genève*, 3: 137–159.
- Brady GS (1883) Report on the Copepoda collected by H.M.S. Challenger during the years 1873–1876. Report of the Scientific Results of the Voyage of H.M.S. Challenger (Zoology), 8: 1–142.
- Brook BW, Sodhi NS & Ng PKL (2003) Catastrophic extinctions follow deforestation in Singapore. *Nature*, 424: 420–426.
- Calbet A & Saiz E (2005) The ciliate-copepod link in marine ecosystems. *Aquatic Microbial Ecology*, 38: 157–167.
- Chen & Zhang (1965) The planktonic copepods of the Yellow Sea and the East China Sea. I. Calanoida. *Studia Marina Sinica*, 7: 20–131, 53 pls [In Mandarin with English summary].
- Chou LM & Chia LS (1991) The marine environment. In: Chia LS, Rahman A & Tay BH (eds.) *The Biophysical Environment of Singapore*. Singapore University Press for the Geography Teachers' Association of Singapore, Singapore. Pp. 155–184.
- Chou LM (2006) Marine habitats in one of the world's busiest harbours. In: Wolanski E (ed.) *The Environment in Asia Pacific Harbours*, Springer, New York. Pp. 377–391.
- Claus C (1866) Die Copepoden-Fauna von Nizza. Ein Beitrag zur Charakteristik der Formen und deren Abänderungen "im Sinne Darwin's". *Schriften der Gesellschaft zur Beförderung der gesamten Naturwissenschaften zu Marburg*, Supplement 1: 1–34.
- Fernando & Ponzi (1981) The freeliving freshwater cyclopoid copepoda (crustacea) of Malaysia and Singapore. *Hydrobiologia* 78, 113–123.
- Fernando CH & Zankai NP (1981) The Rotifera of Malaysia and Singapore with remarks on some species. *Hydrobiologia*, 78: 205–219.
- Giesbrecht W (1893) Systematik und Faunistik der pelagischen Copepoden des Golfes von Neapel. *Fauna Flora Golfes Neapel angrenzenden Meeresabschnitte*, 19: 1–831.
- Gin KYH, Lin X & Zhang S (2000) Dynamics and size structure of phytoplankton in the coastal waters of Singapore. *Journal of Plankton Research*, 22: 1465–1484.
- Gin KYH, Zhang S & Lee YK (2003) Phytoplankton community structure in Singapore's coastal waters using HPLC pigment analysis and flow cytometry. *Journal of Plankton Research*, 25: 1507–1519.
- Gollasch S, Lenz J, Dammer M & Andre H-G (2000) Survival of tropical ballast water organisms during a cruise from the Indian Ocean to the North Sea. *Journal of Plankton Research*, 22: 923–937.
- Grassi GB (1881) I Chetognati. *Fauna Flora Golfo Neapel Monografie V*, 1–126.
- Harris RP, Irigoien X, Head RN, Rey C, Hygum BH, Hansen BW, Niehoff B, Meyer-Harms B & Carloti F (2000) Feeding growth and reproduction in the genus *Calanus*. *ICES Journal of Marine Science*, 57: 1708–1726.
- Idris BAG & Fernando CH (1981) Cladocera of Malaysia and Singapore with records, redescrptions, and remarks on some species. *Hydrobiologia*, 77: 233–256.
- Lai HC & Fernando CH (1978) The freshwater calanoida (Crustacean: Copepod) of Singapore and Peninsular Malaysia. *Hydrobiologia*, 61: 113–127.
- Low EW, Clews E, Todd PA, Tai YC & Ng PKL (2010) Top-down control of phytoplankton by zooplankton in tropical reservoirs in Singapore? *The Raffles Bulletin of Zoology* 58: 311–322.
- Mauchline J (1998) The biology of calanoid copepods. *Advances in Marine Biology*, 33: 1–710.
- Mori T (1940) Two new copepods from Japanese waters. *Zoological Magazine (Dobutsugaku Zasshi)*, Tokyo, 52(8): 328–330, figs. 1–12.
- Mulia IE, Tay H, Roopsekhar K & Tkalic P (2013) Hybrid ANN-GA model for predicting turbidity and chlorophyll-a concentrations. *Journal of Hydro-environment Research*, 7(4): 279–299.
- Myers N, Mittermeier RA, Mittermeier CG, da Fonseca GAB & Kent J (2000) Biodiversity hotspots for conservation priorities. *Nature*, 403: 853–858.
- Nakajima R, Yoshida T, Othman BHR & Toda T (2008) Diel variation in abundance, biomass and size composition of zooplankton community over a coral-reef in Redang Island, Malaysia. *Plankton & Benthos Research*, 3: 216–226.
- NEA (National Environment Agency) (2009) WEATHERWise Singapore, National Environment Agency, Singapore. <http://app2.nea.gov.sg/data/cmsresource/20090721544571208250.pdf> (Accessed 2011).
- NASA (2013) Ocean Motion, Washington, USA. <http://oceanmotion.org/> (Accessed 23 April 2013).
- Nishida (1985) Taxonomy and distribution of the Family Oithonidae (Copepoda, Cyclopoida) in the Pacific and Indian Oceans. *Bulletin of the Ocean Research Institute of the University of Tokyo*, 20: 11–67.
- Omori & Ikeda, 1984: *Methods in Marine Zooplankton Ecology*. John Wiley & Sons, New York, 332 pp.
- Pang WC & Tkalic P (2003) Modeling tidal and monsoon driven currents in the Singapore Strait. *Singapore Maritime & Port Journal*, 2003: 151–162.
- Pham MN, Tan HTW, Mitrovic S & Yeo HHT (2011) A Checklist of the Algae of Singapore. *Raffles Museum of Biodiversity Research, National University of Singapore, Singapore*, 100 pp.
- Razouls C., de Bovée F., Kouwenberg J. et Desreumaux N., 2005–2012. – Diversity and Geographic Distribution of Marine Planktonic Copepods. Available at <http://copepodes.obs-banyuls.fr/en> [Accessed 23 January 2013]
- Rezai, H., F. M. Yusoff, A. Arshad, Kawamura A, Nishida S & Othman Bin HJ Ross (2004) Spatial and temporal distribution of Copepods in the Straits of Malacca. *Zoological Studies*, 43: 486–497.

- Saiz E & Calbet A (2011) Copepod feeding in the ocean: scaling patterns, composition of their diet and the bias of estimates due to microzooplankton grazing during incubations. *Hydrobiologia*, 666: 181–196.
- Sewell SRB (1914) Notes on the surface Copepoda of the Gulf of Mannar. *Spolia Zeylanica*, 9: 191–262.
- Sewell SRB (1933) Note on a small collection of marine copepods from the Malay States. *Bulletin of the Raffles Museum*, Singapore, 8: 25–31.
- Steuer A (1915) Revision der Gattung *Acartia* Dana. *Zoologischer Anzeiger*, 45: 392–397.
- Tham AK (1953) The plankton calendar of Singapore Straits with suggestions for a simplified methodology. *Marine Biological Association of India*, Special Issue: 60–73.
- Tham AK (1973) Seasonal distribution of the plankton in Singapore Straits. In: *Marine Biological Association of India (MBAI) Special Publication dedicated to Dr. N K Panikkar (1)*, Kochi, India. Pp. 60–73.
- Thia-Eng C (1973) An ecological study of the Punggol estuary in Singapore. *Hydrobiologia*, 43: 505–533.
- UNCTAD (2012) United Nation Conference on Trade and Development. *Review of Maritime Transport*. United Nation Publication, Geneva, Switzerland, 176 pp.
- Wickstead JH (1958) A survey of the larger zooplankton of Singapore Straits. *Ices Journal of Marine Science*, 23(3): 340–353.
- Yeo DCJ, Ng PKL, Corlett RT & Tan HTW (2011) Threats to Singapore Biodiversity. In: Ng PKL, Corlett R & Tan HTW (eds.) *Singapore Biodiversity: An Encyclopedia of the Natural Environment and Sustainable Development*, Editions Didier Millet, Singapore. Pp 96–105.