

# Taxonomy of Piscian Cestodes of Freshwater fish: A cladistic model of Morphotaxonomic Study of Rare Tapeworm from the Dehradun Region (U.K.), India



Amar Jeet Singh, Aditya Narayan, Sarvjeet Singh

**Abstract:** *The current experimental study has been conducted to analyze cladistic taxonomic aspects of the piscian cestode parasites including its morphological characters. Freshwater fishes caught on the spot at selected sampling stations with the help of local fishermen and collected for tapeworm investigations. Freshwater catfish Clarias batrachus (Linnaeus, 1758)<sup>[3]</sup> collected from Dehradun sampling stations and examined for tapeworm parasites as per experimental design. The collected fishes were brought to the Parasitology laboratory for experimental study and morphological as well as anatomical facts investigation. Parasites obtained from infected host fishes kept intact in 5% formalin for further investigation. Ten cestode (tapeworm) parasites were found from its infested Intestine. The morphological study of tapeworms indicates that parasitic diversity is accessible within the Ganga River at Dehradun (U.K.) region. Morphotaxonomic characterization reported as - H, butterfly flap shaped ovary, testes numerous and well developed, vitelline follicle - cortical and medullary. This study provides an overview of the piscian tapeworm parasites of some freshwater fishes inhabiting the Ganga River.*

**Keywords:** Morphotaxonomic, Morphometry, Tapeworm, Cestode, Ganga River, Dehradun (U.K.).

## I. INTRODUCTION

Fish provides the essential nutrients for the human body mainly in the form of micronutrients as well as macronutrients as complex proteins, lipids, vitamins, and minerals. Fish matter an important source of income and economic issues for such people who are related in fisheries, aquaculture and fish trade. The government of India conducts many research projects and training schemes from time to time, to improve fish quality and growing supply with fulfilled nutritional rich fishes [1]. Edible fishes are found

infested by tapeworm parasites which reduce the nutritional and economic value of infected fish because tapeworm parasites consume required nutrition from the host and secrete harmful substances. Due to this, the physical weight and size of fishes reduce and ultimately lose their market value [2]. Keeping all these facts in focus the morphotaxonomic study of tapeworm parasites is an urgent necessity today. Very little exploration has been done on the taxonomy of tapeworm parasites hence more experimental work is required. Much research has shown that the generic helminth parasites are found as cestode (tapeworm) parasites in freshwater edible fishes. The curiosity of the authors to develop understanding about the tapeworm parasites found in freshwater fishes lead them to undertake this project at different sampling stations. In the inclusion of this research study, the authors have limited restricted themselves to the taxonomy of cestode parasites of infected host fishes from various sampling stations of Ganga riverside at the Dehradun region. An analysis has been carried out incorporating all the taxonomic aspects and statistical parameters involved in the research (Table I, II).

## II. MATERIAL & METHOD

The methodology has been descended in this study with these following steps as an experimental protocol –

1. Sampling (Field Work)
  - a) Sampling from water bodies (Including local ponds, reservoirs, rivers and water tanks, etc.)
  - b) Sampling from local fish markets (Including village and township based street market)
2. Collection of fishes and rearing in water tanks/pots
3. Dissection
  - a) Virtual Dissection (Computer-based practice)
  - b) Handmade dissection at spot/at the sampling site (Experimental framework)
4. Removal of parasites from host fish (Collection of parasites)
5. Preservation of parasites
6. Permanent slides preparation in parasitology laboratory
7. Micrographs/Camera Lucida of prepared slides
8. Identification and observation of parasites (with morphological, anatomical and morphometrically aspects)

Revised Manuscript Received on April 20, 2020.

\* Correspondence Author

**Amar Jeet Singh**, Department of Zoology, Institute of Basic Science, Bundelkhand University, Jhansi (U.P.), India. Email: amarjeet7321@gmail.com


**Aditya Narayan**, Department of Zoology, Institute of Basic Science, Bundelkhand University, Jhansi (U.P.), India. Email: adityanarayan21@gmail.com

**Sarvjeet Singh**, Department of NST, DTU, Delhi, India. Email: sarvjeetsingh779@gmail.com


© The Authors. Published by Blue Eyes Intelligence Engineering and Sciences Publication (BEIESP). This is an [open access](http://creativecommons.org/licenses/by-nc-nd/4.0/) article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

The freshwater catfish namely *Clarias batrachus* (Linnaeus, 1758) [3] collected from various sampling stations of Ganga riverside at the Dehradun region of Uttarakhand state for the morphotaxonomic study of piscian tapeworms.

Freshwater local fishes angled to caught on the spot at sampling stations with the help of local fishermen, collected and brought to the lab for tapeworms investigation. Fishes were dissected to check out internal organs, the alimentary canal and various organs of the infected host were removed and cut open in 0.9% (w/v) aqueous isotonic saline water in the petri dish. It was shaken several times and many of the associated waste materials were removed. The intestine, gills, stomach and body cavity of parasites were examined thoroughly under a dissecting microscope to ensure that no parasites are left behind. During the autopsy, scolex of parasites were attached deeply embedded in the fish body, which was taken out by scrapping the mucosa of the intestine. It requires a sharp-edged needle and scalpel to remove it carefully. After this, the segment of the mucosa and tissues attached to the parasite was removed by shaking the body of worm in saline water. The stretched and torsion worms were replaced in lukewarm saline water till changing in normal status. In the case of longer worms, these were lifted and repeated several times with the help of needles against the edges of the Petri dish and later on fixed in 5% formalin. Harris' hematoxylin solution and eosin were used to stain whole worm specimen, finally cleared in xylene and mounted in Canada balsam as per experimental protocol. Camera lucida drawings were made using stage micrometer with an ocular micrometer. All the measurements (L×W) were measured in millimeters. For calculation of bio-statistical parameters as X1, X2, Y1 and Y2 values, Margolis equations adopted [4].



**Fig. 1. (a) The anterior part of the worm with smooth scolex and neck [100x] (b) Middle and posterior part of the worm [100x] (c) Eggs [400x] (d) Eggs (bilobed & ovular) [100x]**



**Fig. 2. Map showing of Ganga river the study area at Rishikesh (Dehradun) region.**

The following statistics equations adopted to calculate the various bio-statistical parameters of the parasite (Table I) [4].

$$\text{Incidence of infection (X1)} = \frac{\text{No. of infected hosts (b)}}{\text{No. of examined hosts (a)}}$$

$$\text{Intensity of infection (X2)} = \frac{\text{No. of collected parasites (c)}}{\text{No. of infected hosts (b)}}$$

$$\text{Density of infection (Y1)} = \frac{\text{No. of collected parasites (c)}}{\text{No. of examined hosts (a)}}$$

$$\text{Index of infection (Y2)} =$$

$$\frac{\text{No. of infected hosts (b)} \times \text{No. of collected parasites (c)}}{\text{No. of examined hosts (a}^2\text{)}}$$

**Table- I: Showing statistical analysis of the population of host fishes at sampling stations.**

Sampling station District – Dehradun (Tehsil - Rishikesh) U.K.								
Sr. No.	Sampling stations	No. of examined hosts (a)	No. of infected hosts (b)	No. of collected parasites (c)	Incidence of infection (X1)	Intensity of infection (X2)	Density of infection (Y1)	Index of infection (Y2)
1.	Birpur Khurd near Ganga river line, Rishikesh	30	6	11	0.2	1.833	0.366	0.073
2.	Bishan Garh near Ganga river line, Rishikesh	30	4	7	0.133	1.75	0.233	0.031
3.	Ram Jhula near river Ramganga, Rishikesh	30	5	9	0.166	1.8	0.3	0.05
4.	Fish reservoirs, Mal Devta farm, Dehradun	30	5	10	0.166	2	0.333	0.055
5.	Fish tanks, Malsi Dear Park, Dehradun	30	3	6	0.1	2	0.2	0.02
6.	Water bodies in Aglar Valley, hill town of Mussoorie	30	7	13	0.233	1.857	0.433	0.101
<b>Total</b>		180	30	56	0.998	11.24	1.865	0.33

**Table- II: Biometric characteristic (weight and total length) of freshwater host fishes with Prevalence of Cestode/ Nematode/ Trematode/ Acanthocephala parasites.**

Sr. No. (Sampling stations)	No. of examined hosts	No. of infected hosts	Host fishes	Infected site (Habitat)	Weight (gm)	Total Length (cm)	No. of collected Parasites	Parasites			
								C	N	T	A
1.	30	6	<i>Clarias batrachus</i>	Intestine	185-352	12.3-19.6	11	1	5	4	1
2.	30	4	<i>Clarias batrachus</i>	Intestine	189-325	16.5-22.5	7	1	4	2	0
3.	30	5	<i>Clarias batrachus</i> , <i>Channa punctatus</i>	Intestine, Body cavity	240-372	14.5-18.6	9	2	4	2	1
4.	30	5	<i>Clarias batrachus</i>	Intestine	205-342	19.4-31.2	10	2	4	3	1
5.	30	3	<i>Clarias batrachus</i> , <i>Channa punctatus</i>	Intestine, Gills	315-389	14.8-21.6	6	1	3	2	0
6.	30	7	<i>Clarias batrachus</i>	Intestine, Body cavity	257-412	17.4-23.9	13	3	5	3	2

Table- III. Showing morphometric measurement L×W (in mm) and cladistic taxonomical characters of Genus *Lytocestus* Cohn<sup>[17]</sup>, 1908 tapeworms under generic description

Sr. No.	Characters	<i>Lytocestus indicus</i> (Re-described)	<i>Lytocestus osmanabensis</i> (Bhure et. al., 2010) <sup>[5]</sup>	<i>Lytocestus marathwadensis</i> (Shinde et. al., 1988) <sup>[6]</sup>	<i>Lytocestus follicularae</i> (Bhure et. al., 2010) <sup>[5,7]</sup>	<i>Lytocestus puranensis</i> (Kasar et. al., 2010) <sup>[8]</sup>	<i>Lytocestus clariae</i> (Tandon et. al., 2005) <sup>[9]</sup>	<i>Lytocestus longicollis</i> (Devi PR et. al., 1973) <sup>[10]</sup>	<i>Lytocestus parvulus</i> (Kasar et. al., 2010) <sup>[8]</sup>	<i>Lytocestus clariae</i> (Re-described) (Singh et. al., 2020) <sup>[11]</sup>	<i>Lytocestus birmanicus</i> (Lynsdale, 1956) <sup>[12]</sup>	<i>Lytocestus filiformis</i> (Woodland, 1923) <sup>[13]</sup> and Fuhrman et. al., 1925) <sup>[14]</sup>	<i>Lytocestus attenuatus</i> Tandon et. al., 2005) <sup>[9]</sup>	<i>Lytocestus heteropneustii</i> (Tandon et. al., 2005) <sup>[9]</sup>
1.	Length of the worm	13	11.52	12.22	12.26	14.69	12.10	11.76	12.46	12.06	11.58	12.67	13.20	11.80
2.	Maximum breadth of the worm	0.61	0.82	0.90	0.70	0.95	0.79	0.75	0.80	0.69	0.85	0.96	1.05	0.62
3.	Neck	0.48×0.55	0.61×0.22	0.53×0.23	0.46×0.18	0.49×0.24	0.69×0.19	0.58×0.24	0.68×0.19	0.69×0.16	0.62×0.20	0.50×0.23	0.65×0.31	0.60×0.26
4.	Ovary: shape	H and butterfly flap shaped	H shaped	Butterfly shaped	H shaped	Inverted U shaped	Butterfly shaped	Inverted A-shaped	X shaped	H shaped like a butterfly	H shaped	Inverted U shaped	Butterfly shaped	Inverted A-shaped
5.	Genital aperture (pore)	Present	Present	Present	Present	Present	Present	Present	Present	Present	Present	Present	Present	Present
6.	Vitelline follicles	0.098×0.052	0.6×0.13	0.5×0.12	0.5×0.12	0.6×0.12	0.7×0.11	0.5×0.9	0.05×0.12	0.05×0.13	0.6×0.12	0.6×0.13	0.4×0.10	0.6×0.9
7.	Cirrus Pouch	well-developed oval to round	Round to oval	Round	Symmetrical oval	Round to oval	Oval to round	Mostly oval	Round to oval	Well developed, oval to round	Round to oval	Round to oval	Round	Mostly oval
8.	Eggs	0.037×0.022	0.04×0.02	0.04×0.01	0.05×0.03	0.04×0.03	0.05×0.02	0.05×0.02	0.06×0.03	0.05×0.02	0.04×0.02	0.04×0.04	0.05×0.01	0.05×0.03
9.	Excretory pore	0.016×0.035	0.03×0.07	0.04×0.07	0.03×0.06	0.05×0.08	0.03×0.06	0.05×0.08	0.04×0.9	0.03×0.08	0.03×0.08	0.05×0.09	0.04×0.7	0.05×0.08




Fig. 3. Graphical presentation of sampling stations.

A detailed account of rare tapeworm parasite of freshwater catfish (*Clarias batrachus* Linnaeus, 1758)<sup>[3]</sup> is described. Shape, size, morpho-micrometry, infection site, etc. are observed under a compound microscope and camera lucida figures are drawn (Fig.1, Table II). All morphometrical measurements are measured in millimeters using a stage micrometer and an ocular micrometer (Table III).

Tapeworms measure 12-14×0.35-0.87 (13×0.61). Scolex is medium-sized, flat, smooth, blunt without any major groove. Narrow at the top measures 0.22-0.45×0.35-0.53 (0.33×0.44). Neck present with measures 0.43-0.53×0.48-0.62 (0.48×0.55). Proglottids absent. Numerous testes are present in the medullary region, numerable, oval to round shaped Measures 0.05-0.14×0.07-0.09 (0.09×0.11). Vas deferens are visible. Vitelline Follicle present in the cortical and medullary region measures 0.090-0.106×0.045-0.060 (0.098×0.052). Some VF surrounds touches the ovarian lobes. Uterus extends the posterior region to the ovary. Uterus long, non-glandular, coiled medullary, situated posterior and anterior to the ovarian isthmus. Cirrus pouch well-developed oval to round with internal seminal vesicle. Median measures 0.04-0.05×0.03-0.02 (0.04×0.02). Genital aperture present. Ovary H shaped like a butterfly flap, both arms even, lateral lobes of ovary situated in cortical and medullary regions measures 0.45-0.43×0.39-0.27 (0.42×0.33) behind the cirrus pouch. Genital pore present at the posterior region of the worm. Isthmus present and curved in the medullary region. Anterior uterus measures 0.03-0.04×0.12-0.14 (0.03×0.13). Excretory pore measures 0.015-0.018×0.030-0.040 (0.016×0.035). Eggs are some oval and bilobed, broader than length, some coffee bean-like shape and operculate structure measures 0.045-0.030×0.030-0.015 (0.037×0.022).

### III. RESULT AND DISCUSSION

This parasite species comes closer to genera *Lytocestus*. This species differs from *Pseudolytocestus* in having

well-developed scolex, external seminal vesicle absent, internal seminal vesicle present, ovary H shaped like butterfly and uterus extend up to post ovarian region [15].

- 1) The present form differs from *Lytocestus osmanabadensis* in having a well-developed Scolex with neck, ovary H shaped, and bilobed eggs [5,7].
- 2) The present form differs from *Lytocestus marathwadensis* in having medium size worm with well-developed scolex, H shaped ovary, uterus extends up to post ovarian region and bilobed, operculated eggs [6].
- 3) The Present form differs from *Lytocestus follicularae* having mediums size of worm, H butterfly-shaped ovary in absence of Mehlis gland [7].
- 4) The present form differs from *Lytocestus puranensis* in having a well-developed Scolex with neck, internal seminal vesicle present, and H shaped ovary [8].
- 5) This form differs from *Lytocestus clariae* in having medium-sized worms, presence of neck (under segmented), presence of internal seminal vesicle and butterfly-shaped ovary [9].
- 6) The present form differs from *Lytocestus longicollis* in having internal seminal vesicle, H shaped ovary and bilobed eggs [10].
- 7) The present form differs from *Lytocestus parvulus* in having an H shaped ovary and bilobed, operculate eggs [8].
- 8) The present form differs from *Lytocestus birmanicus* in having butterfly flap shaped ovary, longer than broader eggs [12].
- 9) The present form differs from *Lytocestus filiformis* in having H shaped ovary, round to oval eggs, Internal seminal vesicle present [13].

- 10) The present form differs from *Lytocestus attenuates* in having long Isthmus, butterfly-shaped ovary, Internal seminal vesicle present, broader than longer ovular eggs [9].
- 11) The present form differs from *Lytocestus heteropneustii* in having a well-developed ovary, uterus, internal seminal vesicle, excretory pore [9].

On the basis of taxonomic comparison with morphometrical characters present tapeworm holds a specific classified hierarchy in the Lytocestidae family.

Taxonomic Summary	
Kingdom	Animalia
Phylum	Platyhelminthes
Class	Cestoda
Order	Caryophyllidea
Family	Lytocestidae Wardle et McLeod, 1952 <sup>[16]</sup>
Genus	<i>Lytocestus</i> Cohn, 1908 <sup>[17]</sup>
Species	<i>indicus</i> (Redescribed)
Host	<i>Claris batrachus</i> (Linn. 1758) <sup>[3]</sup>
Habitat	Intestine
Locality	Ganga riverside at Dehradun district, Rishikesh tehsil (U.K.)
Number of Parasites	06/10
NCBI Taxonomy ID	647077
ITIS TSN	1060037
Global Biodiversity Information Facility ID	7010074
IRMNG ID	11469553

#### IV. CONCLUSION

The *Lytocestus indicus* is considered as uncommon Caryophyllidea cestode of *Clarias batrachus* in terms of cladistic morphotaxonomy. The morphological demographic diversity of its population has observed in listed sampling stations of the Dehradun region. The anatomical structure and morphotaxonomic studies are particularly needed to prevent cestode parasite infection in edible fishes. In case of heavy infestation of cestode parasites in host fishes, it is observed as serious injuries in intestine, gills and body mass by worm's scolex and hooks penetration. To get more information about cestode and observing the cladistic morphotaxonomy and morphometry this type of research is necessary.

#### ACKNOWLEDGMENT

The authors are thankful to the Head, Department of Zoology, Institute of Basic Science, Bundelkhand University Campus, Jhansi (U.P.) for providing the laboratory facilities and valuable suggestions.


#### REFERENCES

1. National portal of India.
2. <https://www.india.gov.in/topics/agriculture/fisheries>
3. Case studies on fish loss assessment of small-scale fisheries in Indonesia, FIAM/C1129 (En), FAO Fisheries and Aquaculture Circular, ISSN 2070-6065.
4. C. Linnaeus, Systema Naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. Editio Decima, Reformata. Tomus I Laurentii salvii Stockholm 1758, 824 pp. DOI: 10.5962/bhl.title.542 bhl
5. L. Margolis, G. W. Esch, J. C. Holmes, A. M. Kuris and G. A. Schad, "The use of ecological terms in parasitology" (report of an ad hoc committee of the American Society of Parasitologists). *Journal of Parasitology* 1982, 68 pp. 131-133.
6. D. B. Bhure, S. B. Waghmare, C. R. Kasar and K. M. Shaikh, "Taxonomic Observation of the Caryophyllidea tapeworm *Lytocestus* Cohn, 1908 from *Clarias batrachus* (Linnaeus, 1758)". *Journal of Ecology and Environmental Sciences* 2010, 1(1) pp. 01-06.
7. G. B. Shinde and A. N. Phad, "On a new Cestode *Lytocestus marathwadensis* from freshwater fish". *Riv Di Para* 1988, 47(2) pp. 295-298.
8. D. B. Bhure and S. S. Nanware, "Faunestic Diversity and Distribution of Cestode Parasites of Piscean, Avian and Mammalian Host". *American Journal of Biology and Life Sciences* 2015, 3(6) pp. 241-245 (<http://www.openscienceonline.com/journal/ajbls>)
9. C. R. Kasar, D. B. Bhure, S. S. Nanware and M. B. Sonune, "New species of the caryophyllidean tapeworm *Lytocestus* Cohn, 1908 from *Clarius batrachus* (Linnaeus, 1758)". *The Asian Journal of Animal Science* 2010, Vol. 5(2) pp. 219-222.
10. V. Tandon, R. Chakravarty and B. Das, "Four new species of the genus *Lytocestus* (Caryophyllidea, Lytocestidae) from Edible Catfishes in Assam and Meghalaya, India". *Journal of Parasitic Diseases* 2005, Vol. 29(2) pp. 131.
11. P. R. Devi, "*Lytocestus longicollis* sp. nov. (Cestoidea: Caryophyllidea) from catfish *Clarias batrachus* (L.) in India". *Journal of Helminthology* 1973, Vol. 47 pp. 415-420.
12. A. J. Singh, A. Narayan and S. Singh, "Morphotaxonomic Cladistic Analysis of Rare Cestode Parasites from the Prayagraj Region (U.P.), India". *International Journal of Basic Sciences and Applied Computing (IJBSAC)* 2020 ISSN: 2394-367X, Volume-2 Issue-10, pp 16-19.
13. J. A. Lynsdale, "On two new species of *Lytocestus* from Burma and the Sudan respectively". *J Helminthol.* 1956, 30(2-3) pp. 87-96.
14. W. N. F. Woodland, "On some remarkable new forms of Caryophyllaeidae from the Anglo-Egyptian Sudan, and a revision of the families of the Cestodaria". *Q J Microsc Sci (New Series)* 1923, Vol. 67 pp. 435-472.
15. O. Fuhrmann, J. G. Baer, "Zoological results of the third Tanganyika expedition conducted by Dr. W. A. Cunningham, 1904-1905". Report on the Cestodes. In: *Proceedings of the Zoological Society of London* 1925, pp. 79-100.
16. A. Pathak and A. K. Srivastav, "Morphotaxonomical study of a new cestode, *Pseudoclarialis pandei* n.g., n.sp from catfish, *Clarias batrachus* Linn.," *Flora and Fauna* 2015, Vol. 21(1) pp. 15-20, ISSN 0971 6920.
17. R. A. Wardle and J. A. McLeod, *The Zoology of tapeworms.* University of Minnesota Press, Minneapolis 1952, pp. 780.
18. L. Cohn, Die Anatomie eines neuen Fischcestoden. *Centralblatt für Bakteriologie, Parasitenkunde, Infektion-shrank-heiten und Hygiene. Abteilung L Originale* 1908; Vol. 46 pp. 134-139.

#### AUTHORS PROFILE



**Amar Jeet Singh**, Research Scholar (Zoology), Parasitology Laboratory, Department of Zoology, Institute of Basic Science, Bundelkhand University Campus, Jhansi (U.P.) India. Present working project: Taxonomy of piscian cestodes of fresh water fish. Specialization: Parasitology & Immunology Email - [amarjeet7321@gmail.com](mailto:amarjeet7321@gmail.com)





**Aditya Narayan**, Assistant Professor, Department of Zoology, Institute of Basic Science, Bundelkhand University, Jhansi (U.P.) India. Email - [adityanarayan21@gmail.com](mailto:adityanarayan21@gmail.com)



**Sarvjeet Singh**, Research Scholar (NST) Department of NST, Science Block, DTU, Delhi Email - [sarvjeetsingh779@gmail.com](mailto:sarvjeetsingh779@gmail.com)