


**Supplementary Figure 1. Controls for the separation of EVs from *Fusarium graminearum* (Fgr) by SEC.** (A) An Fgr culture was processed for EV separation and mixed with DPBS instead of FM5-95 before SEC. Fluorescence of all fractions was recorded (red line) and their particle number determined by NTA (blue line) (n=1). (B) Fgr was grown for 5 d, then the mycelia were heat-treated before being transferred to fresh medium that was incubated for 5 d. NTA detected  $4.1 \times 10^{10}$  particles/L in the 0.45- $\mu$ m filtrate of the heat-treated Fgr culture,  $4.8 \times 10^{10}$  particles/L in the untreated 0.45  $\mu$ m filtrate, and  $1.9 \times 10^{10}$  particles/L in the sterile YNB+ (n=1).

[illegible]

**Supplementary Figure 2. The superoxide dismutase [Cu-Zn] (SOD1) from *Fusarium graminearum* (*Fgr*) contains a diacidic amino acid motif implicated in unconventional secretion.** The computational prediction of putative effectors from *Fgr* returned two candidates. One was a SOD1 without a predicted signal peptide, that was interrogated to find links to unconventional protein secretion. Its sequence (*Fgr* SOD1) was aligned with the *S. cerevisiae* (yeast) and human SOD1 sequences. The *Fgr* SOD1 contains a diacidic Asp-Glu motif (highlighted in red) previously implicated in non-Golgi secretion of SOD1 in *S. cerevisiae* (Cruz-Garcia et al., 2017).





**Supplementary Figure 4. Computational prediction of effector candidates detected in EV samples from *Fusarium graminearum* (Fgr).** LFQ-based proteomics revealed 647 proteins in the EV samples from *Fgr*. All proteins were processed with EffectorP 2.0 to predict effector activity, PrediSi, Uniprot annotation, SignalP 5, and Phobius to predict signal peptide (SP), SecretomeP 2.0 to predict unconventional secretion, PredGPI to detect GPI anchoring, ApoplastP 1.0, WolfPSORT, and DeepLoc 1.0 to predict cellular location (ex: extracellular, cyt: cytoplasmic, mito: mitochondrial, nucl: nuclear). The percentage cysteine sequence content was calculated manually. Housekeeping, ribosomal and transmembrane (TM) proteins were omitted.