

Supplementary information to “Toward better phenology analyses based on bud traits and energy budget”

Marc Peaucelle, Josep Penuelas and Hans Verbeeck

2022-01-24

Contents

| | |
|----------------------------|---|
| Overview | 1 |
| IP information | 1 |
| Approach | 2 |
| Process sequence | 3 |
| References | 5 |

Overview

Article acceptance date: submitted

This R-Markdown document provides the instructions for executing the analysis and producing figures presented in the paper *Toward better phenology analyses based on bud traits and energy budget* by Peaucelle, Penuelas and Verbeeck (submitted). All data used for the analyses are freely available after registration on the respective website:

- PEP data: <http://www.pep725.eu/>
- CRU-JRA data: <https://catalogue.ceda.ac.uk/uuid/13f3635174794bb98cf8ac4b0ee8f4ed>
- FLUXNET data: <https://fluxnet.fluxdata.org/data/fluxnet2015-dataset/>
- ERA5 data: Downloaded from <https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-land?tab=form> with the KrigR package

These data are under specific licence, this is the reason why we did not provide raw data but only processed data in the **database** folder, and file templates in the **templates** folder. Further information about data processing and creating the data file is given below (section ‘Data processing’). Using RMarkdown and open access code available through github (<https://github.com/mpeaucelle/Tbud>) this is supposed to allow for full reproducibility of published results - from publicly accessible data files to published figures.

IP information

All code in this repository is free software. It may be redistributed and/or modified under the terms of the GNU General Public License as published by the Free Software Foundation, version 3. Present code is distributed in the hope that it will be useful, but WITHOUT ANY WARRANTY; without even the implied warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE. See the GNU General Public License (file `./LICENSE`) for more details.

In case you use or modify this code, please cite the original work as: Peaucelle, Penuelas and Verbeeck (2022), Toward better phenology analyses based on bud traits and energy budget, *under review*

Copyright (C) 2020 Marc Peaucelle

Approach

Bud energy budget

We quantify the effective temperature of buds by applying a simple energy budget model based on (Landsberg, Butler, and Thorpe 1974), (Hamer 1985) and equations from (Muir 2019). For an isolated bud, the amount of absorbed incoming radiation (R_{abs}) is balanced by the thermal infrared radiation loss (LW_{bud}) and the energy lost by conduction and convection, generally called the sensible heat flux (H). To simulate bud temperature, we considered that the energy balance is close to equilibrium at a time scale of a few minutes, consistent with temporal resolution of meteorological observations from FLUXNET sites (30 min) and CRU-JRA (6 h):

$$R_{\text{abs}} = H + LW_{\text{bud}}$$

The amount of absorbed energy by buds is the sum of incoming shortwave (SW, visible and near-infrared) and longwave (LW, infrared) radiations:

$$R_{\text{abs}} = \alpha_{\text{SW}}(1 + r)SW + \alpha_{\text{LW}}(LW_{\text{sky}} + LW_{\text{gnd}})$$

where α_{SW} is the bud absorptivity to SW; r is the fraction of SW reflected by the ground and α_{LW} is the bud absorptivity to LW, which is here defined as the average of LW coming from the atmosphere LW_{sky} and coming from the ground LW_{gnd} . LW emitted by surrounding objects such as branches were considered negligible. We set α_{LW} to 0.97, which corresponds to the average absorptivity for wood and leaves Howard and Stull (2013). Since no data were available in the literature, we tested two different values of α_{SW} , 0.5 and 0.8, corresponding to values commonly used for leaves and needles, respectively. We set r to 0.2, which corresponds to a reasonable value for the fraction of SW reflected by grasses. Of course, r will strongly vary according to the albedo of the ground (e.g. understory/grass, forest litter, snow, etc.), which was simplified here for our perspective.

In our example, we simplified the estimation of LW_{gnd} by considering that ground temperature equals air temperature:

$$LW_{\text{gnd}} = \sigma \varepsilon T_{\text{air}}^4$$

where σ is the Stefan-Boltzmann constant and ε is the bud emissivity (which is equal to its absorptivity) to longwave radiations. This is an important oversimplification to keep in mind since ground temperature will strongly depend on soil type, vegetation and snow cover, soil humidity, but also the fact that soil heat capacity is generally higher than air.

Buds lose thermal infrared radiation proportionally to their temperature as:

$$LW_{\text{bud}} = 2\sigma \varepsilon T_{\text{bud}}^4$$

where σ is the Stefan-Boltzmann constant and ε is the bud emissivity (which is equal to its absorptivity) to longwave radiations.

Finally, the sensible heat flux depends on the air to bud temperature gradient and is formulated as in Muir (2019):

$$H = \rho_a C_p g_b (T_{\text{bud}} - T_{\text{air}})$$

where ρ_a is the density of dry air; C_p is the specific heat capacity of air at constant pressure and g_b is the boundary-layer conductance to heat.

In our example, we simplified bud's shape as a spherical object with a diameter of 5 mm to compute the boundary-layer conductance to heat. Please refer to Hamer (1985) for a detailed discussion of this




Figure 1: *Illustration of the bud energy budget.*

assumption, as well as empirical formulations for the convective heat transfer of apple buds. Condition for laminar and turbulent flows, as well as constants a , b , c and d for a spherical object can be found in Monteith and Unsworth (2013). In addition, we performed two simulations varying ground albedo from 0.1 (~wet bare soil) to 0.9 (~snow) and bud diameter from 5 to 13 mm (Supplementary Figure 5 and 6) The bud energy model simulates bud temperature in idealized conditions, without distinction between species and sites and several simplifications as described above. Note that a proper calibration for species and site properties is needed to use this model for robust predictions.

Bud temperature estimation

Bud temperature was estimated by solving for the equilibrium of the energy budget at each time-step (30 min for FLUXNET, 6 h for CRU-JRA) and for each site.

!!! Important Note: This model is simplified to represent idealized conditions. To have robust simulations of bud temperature and to use this approach to model phenology, a proper calibration of model parameters, including bud traits (e.g. solar absorptivity, size and shape), but also the surrounding environment (e.g. ground albedo and soil temperature) Running the model with micrometeorological data should also produce better results for bud temperature

Process sequence

The following provides a documentation and instruction to reproduce results and figures presented in the paper. Data processing (`0_create_database_Tbud.R` and `2_Figures_Fluxnet.R`) requires raw data to run properly. Please download raw data from the dedicated website and format the data according to the template files in the `templates` folder.

Figures and analyses with PEP data (`1_Figures_PEP_CRUJRA.R`) can be directly reproduced with processed data available in the `database` folder for each studied species.

1. Requirements and environment setup

The execution of the scripts provided here requires the following R packages only for data preprocessing. Analysis can be performed with the base package. Refers to the following script `0_create_database_Tbud.R`.

```
list.of.packages <- c("RNetCDF", "raster", "xts", "ggplot2")
new.packages <- list.of.packages[!(list.of.packages %in% installed.packages()[,"Package"])]
if( length(new.packages) ) install.packages(new.packages)
```

```
## Installation des packages dans '/home/marc/R/x86_64-pc-linux-gnu-library/4.1'
## (car 'lib' n'est pas spécifié)
```

```
## installation des dépendances 'colorspace', 'farver', 'labeling', 'munsell', 'RColorBrewer', 'viridis'
```

```
## Warning in install.packages(new.packages): installation of package 'terra' had
## non-zero exit status
```

```
## Warning in install.packages(new.packages): installation of package 'raster' had
## non-zero exit status
```

Define working directory. Define the path variable used throughout multiple scripts and functions.

```
workingdir <- getwd()
print( paste("working directory directory:", workingdir ) )
```

```
## [1] "working directory directory: /media/marc/Data/CAVELab/Tbud/R_src"
```

```
workdir <- "/AddWorkdirHere"
```

Get additional utility functions.

```
source("Tbud.R") # Estimate bud temperature. Call Ebalance.R
source("Ebalance.R") # Estimate the energy balance of bud
source("functions.R") # Several functions for averaging, etc..
```

2. Data processing

FLUXNET data were directly used without any particular pre-processing. Only the quality and continuity of data was checked for the example provided in the paper. PEP data were processed with the script `0_create_database_Tbud.R`.

PEP and CRU-JRA raw data were processed following these steps for each species:

1. Load PEP and CRU-JRA data
2. Select PEP data corresponding to leaf unfolding observations (BBCH = 11)
3. Retrieve corresponding pixel of CRU-JRA (0.25° resolution) for each site
4. Extract meteorological data for each pixel and each site
5. Computes pre-season (3 months preceding budburst) meteorological conditions, as well as forcing and chilling, for each observation.
6. Merge data and save the new dataset in .rds

1 Note that CRU-JRA are based on a no-leap calendar. Leap years were added manually to match phenological observations by duplicating the last day (i.e. XXXX/12/31) of the corresponding year.

2 Note that CRU-JRA data were linearized to improve computing time. Meteorological variables can be easily extracted from raw 2D CRU-JRA data using the ‘raster’ package and site longitude-latitude, however the script `0_create_database_Tbud.R` will have to be adapted in accordance.

3. Data analysis

PEP data (Supplementary Figures 2, 3 and 4) Processed .rds data files in `./database/` contain for each site all the data used for analyses and for plotting the figures presented in the paper.

The script `1_Figures_PEP_CRUJRA.R` plots Supplementary Figures 2, 3 and 4. To generate the figures simply run the script as:

```
source("1_Figures_PEP_CRUJRA.R")
```

This script executes the following tasks:

1. Load .rds data for each species and merge them all in a unique variable with all species and sites together
2. Select sites with at least 20 years of observations over 1990-2015
3. Computes mean yearly values (and standard deviation) with all sites and species pooled together
4. Generate Figures

FLUXNET data (Figure 2a and 2b, Supplementary Figures 1, 5 and 6) The script `2_Figures_Fluxnet.R` plots Figures 2a and 2b, as well as Supplementary Figures 1, 5 and 6 from FLUXNET data. As mentioned above, FLUXNET data are not provided in the repository, but are freely accessible after registration on the dedicated website.

To generate the figures, please use raw FLUXNET data and simply source the script with appropriate working directories

```
source("2_Figures_Fluxnet.R")
```

This script executes the following tasks:

1. Load Fluxnet data
2. Computes Bud temperature with two different solar absorptivity by calling the function `Tbud` in `Tbud.R`
3. Computes Chilling and forcing requirement of buds from air temperature and bud temperature
4. Generate analysis by varying bud diameter and ground albedo
5. Generate Figures

References

- Dupleix, Anna, Domingos De Sousa Meneses, Mark Hughes, and Rémy Marchal. 2013. “Mid-Infrared Absorption Properties of Green Wood.” *Wood Science and Technology* 47 (6): 1231–41.
- Hamer, PJC. 1985. “The Heat Balance of Apple Buds and Blossoms. Part i. Heat Transfer in the Outdoor Environment.” *Agricultural and Forest Meteorology* 35 (1-4): 339–52.

- Howard, Rosie, and Roland Stull. 2013. "IR Radiation from Trees to a Ski Run: A Case Study." *Journal of Applied Meteorology and Climatology* 52 (7): 1525–39.
- Jones, Hamlyn G. 2013. *Plants and Microclimate: A Quantitative Approach to Environmental Plant Physiology*. Cambridge university press.
- Landsberg, JJ, DR Butler, and MR Thorpe. 1974. "Apple Bud and Blossom Temperatures." *Journal of Horticultural Science* 49 (3): 227–39.
- Monteith, John, and Mike Unsworth. 2013. *Principles of Environmental Physics: Plants, Animals, and the Atmosphere*. Academic Press.
- Muir, Christopher D. 2019. "Tealeaves: An r Package for Modelling Leaf Temperature Using Energy Budgets." *AoB Plants* 11 (6): plz054.