

Derivation of Special Relativity Transformation Formulas Without Time Slowing and Length Shrinking

——Modification of Relativistic Aberration Formula

Author: Sheng Qin

E-Mail: qinsheng@smail.swufe.edu.cn

Southwestern University of Finance and Economics, China, 610000

Abstract: In this paper, we try to use a new way to deduce the transformation formula of special relativity. On the premise of not assuming that time slows down and length shrinks, we use the experimentally verified Planck energy formula, momentum formula and mass-energy equation as the basis, namely, The Doppler frequency shift formula of the special theory of relativity, the moving mass formula and the velocity transformation formula in the special theory of relativity can be obtained. At the same time, the biggest innovation of this paper is: (1) The preconditions of the Doppler frequency shift formula are supplemented, and it is pointed out that the Doppler frequency shift formula The relative motion speed of the object in the frequency shift formula of Doppler relativity is actually relative to the speed after the object emits photons, not relative to the speed before emission, which makes up for the deficiency of the frequency shift formula of Doppler relativity; (2) Prove and The traditional relativistic aberration formula is revised, and the correct relativistic aberration formula is introduced. The correction can be used for the calculation of laser absorption and reflection of moving objects, and the calculation of radiation and absorption angle of high-speed moving particles; (3) This paper also proves that in the special theory of relativity, the two conclusions of time slowing down and length shrinking are superfluous. Then our conclusion in another article may be correct: simultaneity is possible in different inertial frames, and the space is relatively independent. In this way, we can make more reasonable explanations for phenomena such as quantum entanglement at a distance and cosmic inflation.

Key words: Time slows down; length shrinks; Doppler frequency shift formula; moving mass formula; relativistic velocity transformation formula.

0 Introduction

In the past, many physicists have proposed many novel and simple methods to deduce the conclusions of special relativity. For example, Feng Shimeng proposed a characteristic deduction method of the relationship between mass and velocity^[1], but retained the time and coordinate

transformation functions; Wang Yonggang, Cao Xuecheng^[2], etc. Trying to use space-time algebra to deduce the velocity transformation formula of relativity theory is more of a formal transformation; Dai Youshan and Song Xiaotong^[3], in their article "New Derivation of Special Relativity Without Assumption of Invariant Speed of Light", based on the mechanical model A method that does not require the assumption of constant speed of light, but still requires the assumption of time transformation. There are also some scholars who use the relativistic time and length transformation relationship to deduce the Doppler frequency shift effect, such as Lu Junling, Wang Rongbao^[4], Gao Bingkun, and Wang Fenglin^[5].

However, the inference methods in the above articles still retain the time transformation and length transformation formulas of relativity. The new method used in this paper does not need to rely on the relativistic time transformation and length transformation, but only relies on the traditional conservation of momentum and energy, and the following transformation formula of the special theory of relativity can be deduced: The relativistic Doppler frequency shift formula, relativistic dynamic mass formula, relativistic velocity transformation formula.

And more importantly, this paper proves that the relative motion speed of the object in the Doppler relativistic frequency shift formula is actually relative to the speed of the object after the photon is emitted, not the speed before the emission. In the calculations of past experimental studies, this point has been ignored^[6], which makes up for the deficiency of the relativistic Doppler frequency shift formula.

This paper finds a new way to derive the special relativity transformation formula, which does not need to rely on the time transformation and length transformation formulas. The author believes that only based on a rigorous experimental basis is a solid foundation for constructing a physical theory.

The biggest difference between this article and the special theory of relativity:

(1) There is no need to assume that length shrinks, time becomes slower, or even the speed of light remains unchanged. This paper is based on the same physical postulates as the laws of physics in the inertial system.

(2) The basis of the theory is based on the mass-energy equation which has been strictly proved by experiments $E = mc^2$ ^{[7][11]}. Energy Formula of Planck Photon $E = h\omega$ and momentum

formula $P = \frac{h\omega}{c}$ (Checked by the Compton scattering experiment^[8]).

(3) The relativistic Doppler frequency shift effect is a direct conclusion of Planck's formula and mass-energy equation, not a new assumption.

1 Proof of the Transformation Formula of Special Relativity

1.1 Proof of Relativistic Doppler Frequency Shift Formula

First, based on the Planck formula $E = h\omega$ and $P = h\omega / c$ (E is the energy, P is the momentum, h is Planck's constant, ω is the photon frequency). We prove two theorems:

Theorem 1: The red-shift or blue-shift Doppler effect of photons emitted from each other between two relatively moving objects A and B, the relative velocity of the two objects A and

B in the corresponding photon frequency transformation formula, refers to the object The relative motion speed of A and B after A or B emits a photon.

Prove:

As shown in Fig. 1, we assume that a black box is stationary in free cosmic space, we emit a photon α with frequency ω from one side of the black box, and it is absorbed on the other side of the black box. We know that the black box as a whole will maintain momentum and energy conservation and will eventually continue to remain at rest.




Fig. 1 Photon emission and absorption in a stationary black box

Then we can reason: due to the conservation of momentum between the black box and the photon, after the black box emits photon α and before the photon α is absorbed by the other side, the black box will acquire a velocity v (relative to the opposite direction of the photon motion), then It means that the black box is no longer in a static state when the photon α is absorbed, but has a relative velocity v .

We assume that the Doppler blueshift formula is: $\omega \cdot \omega_{blue}(|v|)$.

where $\omega_{blue}(|v|) \geq 1$, $\omega \cdot \omega_{blue}(|v|)$ is the frequency after the blue shift of the photon; We assume that the Doppler redshift formula is: $\omega \cdot \omega_{red}(|v|)$

where $\omega_{red}(|v|) \leq 1$, $\omega \cdot \omega_{red}(|v|)$ is the frequency after photon redshift.

And: $\omega_{blue}(0) = \omega_{red}(0) = 1$

If the photon frequency ω is relative to the static state before the black box is emitted, then when the black box absorbs the photon α , a blue-shift effect will be obtained because the black box has a relative velocity v , and the frequency ω of the photon α will be blue-shifted as:

$$\omega \cdot \omega_{blue}(|v|).$$

Then, according to Planck's formula, the black box obtains a momentum relative to the photon when it is emitted: $P_1 = h\omega / c$; when it absorbs the photon, it obtains another momentum due

to the Doppler blue shift effect: $P_2 = -h\omega \cdot \omega_{blue}(|v|) / c$, obviously $P_2 > P_1$, the

momentum of the black box is not conserved before and after the photon α is emitted, which Clearly violates the law of conservation of momentum.

Therefore, only assuming that the frequency ω of the photon α is actually the state after the photon is emitted relative to the black box and the speed v is obtained, then for the photon α , when it is absorbed by the other side of the black box, it is equivalent to being absorbed by a The emitting object is absorbed by the object in a relatively static state. At this time, the frequency of the photon α when it is absorbed is still ω (due to $\omega_{blue}(0) = \omega_{red}(0) = 1$, the relative motion speed between the emitting object and the absorbing object is 0). At this time, the black box will maintain momentum before and after the absorption. and energy conservation.

Therefore, when we calculate the formula of Doppler red shift or blue shift, $\omega \cdot \omega_{blue}(|v|)$ or $\omega \cdot \omega_{red}(|v|)$ the V in actually refers to the relative velocity of A and B after the relative motion objects A and B emit photons.

Rather than the relative velocity of A and B before the emission of the photon.

Therefore, if we use high-speed moving particles to radiate photons to test the correctness of the Doppler frequency shift formula, then the velocity v used in our calculation should be the speed after the particles radiate photons, not the speed before high-speed particles radiate photons. Movement speed.

Prove it.

This result is different from that in Einstein's special theory of relativity^[9], which makes up for the deficiency of the Doppler shift formula in the special theory of relativity and answers the premise of the Doppler shift formula more rigorously.

Theorem 2: Two objects move toward or away from each other at the same speed v , and the Doppler blue-shift and red-shift effects are reciprocal. Suppose objects A and B move towards each other at speed v , and emit photons of frequency ω from A to B, and assume that their blue-shifted frequency is $\omega \cdot \omega_{blue}(|v|)$; then when A and B move away at speed v ,

their red-shift frequency will be $\omega \cdot \omega_{red}(|v|) = \omega \cdot \frac{1}{\omega_{blue}(|v|)}$, (where $\omega_{blue}(|v|) \geq 1$, is a

velocity-dependent Doppler shift function).

Proof: Figure 2 below




Fig. 2 Doppler red shift and Doppler blue shift

Suppose an object A moves in the direction of the X-axis at a speed v relative to the ground

observer, and A first emits photons of frequency ω to the ground P_1 in front of the movement. At this time, because of the blue shift effect, it is assumed that the frequency of the photons received by the ground observer at P_1 is for $\omega \cdot \omega_{blue}(|v_1|)$. According to the conclusion of the previous "Theorem 1", we know that v_1 at this time refers to the movement speed of A relative to the ground after A emits photons.

After receiving the photon at the ground P_1 in front, the ground observer emits a photon with the same frequency $\omega \cdot \omega_{blue}(|v_1|)$ from the rear P_2 to the rear of A, and is then absorbed by A. At this time, since object A moves away from the direction of photons emitted by P_2 on the ground behind with v_1 , a red-shift effect will occur, and the frequency of photons absorbed by A will become: $\omega \cdot \omega_{blue}(|v_1|) \cdot \omega_{red}(|v_1|)$.

Assuming that the ground is large enough, we can ignore the speed change of the ground itself before and after the ground receives and emits photons. Then for a ground observer, after receiving a photon of frequency $\omega \cdot \omega_{blue}(|v_1|)$ and transmitting a photon of frequency $\omega \cdot \omega_{blue}(|v_1|)$, the momentum and energy of the ground as a whole will remain unchanged (because the ground observer absorbs and emits the same frequency $\omega \cdot \omega_{blue}(|v_1|)$ photons in opposite directions).

Then at the same time, before the object A emits photons of frequency ω and absorbs photons of frequency $\omega \cdot \omega_{blue}(|v_1|) \cdot \omega_{red}(|v_1|)$, the velocity v of object A relative to the ground and the mass (overall energy) of A will remain unchanged, that is, the momentum of A and the The energy will remain the same.

Therefore, from the perspective of A, it is necessary to keep the momentum and energy before and after the emission and absorption of photons unchanged: it must be that a photon of frequency ω is first emitted to the front P_1 , and then absorbed from the rear P_2 of the movement. Only in this way can the photon with frequency ω maintain the conservation of its own momentum and energy before and after. We assume that its redshift formula is $\omega_{red}(|v_1|)$, then there must be the following relationship:

$$\omega \cdot \omega_{blue}(|v_1|) \cdot \omega_{red}(|v_1|) = \omega$$

Therefore, we get: $\omega_{blue}(|v_1|) = \frac{1}{\omega_{red}(|v_1|)}$

Prove it.

Theorem 3: The relativistic Doppler redshift formula is: $\omega_{red}(|v|) = \sqrt{\frac{c - v}{c + v}}$; **The blue**

shift formula is: $\omega_{blue}(|v|) = \sqrt{\frac{c + v}{c - v}}$.

Prove:

We continue to do the following thought experiment, as shown in Figure 3 below: Suppose an object A moves along the X-axis with a velocity v relative to the observation system S , and at a certain moment A simultaneously emits two equivalent photons to the front and back of its motion, assuming that both photons have frequencies ω .




Fig. 3 Proof diagram of Doppler frequency shift formula

The two photons are observed by the observation system S where the ground observer is located. For A, the velocity v relative to the ground before and after the emission is unchanged (the total momentum of the photons emitted before and after is $P=0$), the only decrease is The overall mass of A (we can assume that these two photons are a pair of symmetric photons produced by the annihilation of a pair of positrons and electrons moving together with A, and the only thing that decreases for A before and after emission is the mass, the velocity v of object A relative to the ground before and after launch remains unchanged). The same seems to be the case for ground observers of the S series, the relative velocity v of A remains unchanged before and after launch.

Then at P_1 in front of the movement, the photons observed by the ground observer have a blue shift. Assuming that the frequency of the detected photon is $\omega \cdot \omega_{blue}(|v|)$, then according to the conclusion of our previous "Theorem 2", the red light detected at P_2 behind the ground observer

is The shifted photon frequency is $\omega \cdot \frac{1}{\omega_{blue}(|v|)}$.

Then from the point of view of the ground observer of the S system, assuming that the mass reduced by the object A is $2\Delta m$, it will be equal to the sum of the energies of the photons observed by the ground observer (note that $2\Delta m$ is a moving object from the ground observer's point of view here). mass), which is derived from Planck's formula:

$$2\Delta m = \frac{h \left(\omega \cdot \omega_{blue}(|v|) + \omega \cdot \frac{1}{\omega_{blue}(|v|)} \right)}{c^2}$$

The reduced momentum ΔP of A observed by the S-series ground observer is:

$$\Delta P = 2\Delta m \cdot v = \frac{h \left(\omega \cdot \omega_{blue}(|v|) + \omega \cdot \frac{1}{\omega_{blue}(|v|)} \right)}{c^2} \cdot v \quad (1)$$

At the same time, according to the Planck formula of photon momentum, the momentum is equal to the sum of the momentums of the two front and rear photons observed by the ground observer (because the total momentum obtained by the ground observer is composed of only the sum of the momentums of the two front and rear photons), Since the two photons are in opposite directions, the relationship is as follows:

$$\Delta P = 2\Delta m \cdot v = \frac{h \left(\omega \cdot \omega_{blue}(|v|) - \omega \cdot \frac{1}{\omega_{blue}(|v|)} \right)}{c} \quad (2)$$

There is only one unknown quantity $\omega_{blue}(|v|)$ in the above equation system formed by equations (1) and (2), we get the following equation:

$$\frac{h \left(\omega \cdot \omega_{blue}(|v|) + \omega \cdot \frac{1}{\omega_{blue}(|v|)} \right)}{c^2} \cdot v = \frac{h \left(\omega \cdot \omega_{blue}(|v|) - \omega \cdot \frac{1}{\omega_{blue}(|v|)} \right)}{c} \quad (3)$$

We obtain the Doppler blue shift formula by solving equation (3) as:

$$\omega_{blue}(|v|) = \sqrt{\frac{c + v}{c - v}}$$

According to the conclusion of the previous "Theorem 2", we get the formula of Doppler redshift:

$$\omega_{red}(|v|) = \frac{1}{\omega_{blue}(|v|)} = \sqrt{\frac{c - v}{c + v}}$$

Thus, we have the relativistic Doppler shift formula.

Prove it.

1.2 Doppler shift in other directions

As shown in Figure 4 below:




Fig. 4 Doppler frequency at other angles

If the two photons with frequency ω are emitted in the direction of the A system and the angle between the X axis is θ , it is assumed that the photon pair is emitted symmetrically, and it is assumed that the angle at which the photon is detected by the S system in front of the motion is θ_1 , and the photon frequency is ω_1 ; The angle at which the photon is detected is θ_2 and the photon frequency is ω_2 (assuming θ 、 θ_1 、 θ_2 are in the range $\left[0, \frac{\pi}{2}\right]$).

Since the photons emitted by object A are symmetrical, the velocity v of A before and after the emission is unchanged, and the momentum in the Y-axis direction is also unchanged.

Then photons ω_1 and ω_2 will satisfy the following momentum and energy conservation relations:

$$h\omega_1 + h\omega_2 = 2h\omega \cdot \sqrt{\frac{c^2}{c^2 - v^2}} \quad (4)$$

$$\frac{h\omega_1}{c} \cdot \sin \theta_1 - \frac{h\omega_2}{c} \cdot \sin \theta_2 = 0 \quad (5)$$

$$\frac{h\omega_1}{c} \cdot \cos \theta_1 - \frac{h\omega_2}{c} \cdot \cos \theta_2 = \frac{2h\omega}{c^2} \cdot \sqrt{\frac{c^2}{c^2 - v^2}} \cdot v \quad (6)$$

$$\frac{h\omega_1}{c} \cdot \sin \theta_1 = \frac{h\omega}{c} \cdot \sin \theta \quad (7)$$

$$\frac{h\omega_2}{c} \cdot \sin \theta_2 = \frac{h\omega}{c} \cdot \sin \theta \quad (8)$$

Where, Equation (4) refers to the conservation of energy from the perspective of the S system; Equation (5) refers to the conservation of momentum in the Y-axis direction from the perspective of the S-system, and the total momentum in the Y-axis direction is 0; Equation (6) It refers to the conservation of momentum in the X-axis direction from the perspective of the S system.

Both the aberration formula and the Doppler frequency shift formula in the theory of relativity will be obtained from the solution of the above formula: (4) (5) (6) (7) (8). Among them, equations (7) and (8) refer to the conclusions of the transformation relationship of the special theory of relativity.^[10]

Solving equations (4) (5) (6) (7) (8) is easy to get:

$$\omega_1 = \omega \cdot \frac{\left(1 + \frac{v}{c} \cos \theta\right)}{\sqrt{1 - \frac{v^2}{c^2}}}$$

$$\omega_2 = \omega \cdot \frac{\left(1 - \frac{v}{c} \cos \theta\right)}{\sqrt{1 - \frac{v^2}{c^2}}}$$

Therefore, the correct aberration formula is:

$$\frac{\sin \theta}{\sin \theta_1} = \frac{\omega_1}{\omega} = \frac{\left(1 + \frac{v}{c} \cos \theta\right)}{\sqrt{1 - \frac{v^2}{c^2}}}$$

Or:

$$\frac{\sin \theta}{\sin \theta_2} = \frac{\omega_2}{\omega} = \frac{\left(1 - \frac{v}{c} \cos \theta\right)}{\sqrt{1 - \frac{v^2}{c^2}}}$$

Compared with the traditional relativistic aberration formula:

$$\tan \theta = \frac{\sin \theta_1}{\sqrt{1 - \frac{v^2}{c^2} \left(\cos \theta_1 - \frac{v}{c}\right)}} \Rightarrow \frac{\sin \theta}{\sin \theta_1} = \frac{\cos \theta}{\sqrt{1 - \frac{v^2}{c^2} \left(\cos \theta_1 - \frac{v}{c}\right)}}$$

The correct aberration formula we obtained is significantly different from the traditional aberration formula, which is conducive to correct calculation in observation and experiment.

1.3 Proof of the dynamic mass formula of relativistic moving objects

Reasoning: From the Doppler frequency shift formula, we can further infer that the

dynamic mass formula $m(v)$ of the moving object is: $m(v) = m_0 \sqrt{\frac{c^2}{c^2 - v^2}}$, where m_0 is

the static mass of the object, and v is the moving speed of the object.

Prove:

Assume that two positive and anti-matter masses of m_0 are moving relative to the ground S observation system in the direction of the X-axis of the S system at a speed v , and the positive and

anti-matter annihilate each other to generate two photons of frequency ω , which are respectively directed towards the X-axis of the S system. It is emitted in two directions, where the photon frequency ω is obtained from the mass-energy equation $m_0 \cdot c^2 = h\omega$. Assume that the

dynamic mass of matter and antimatter is $m_0(v)$ in the view of the S-system ground observer.

After the annihilation of positive and negative matter, two photons with frequency ω are respectively emitted towards the front and rear directions of the motion. According to the conclusion of "Theorem 3" above, the total photon energy E received by the ground observer is:

$$E = h \left(\omega \cdot \omega_{blue}(v) + \omega \cdot \frac{1}{\omega_{blue}(v)} \right) = h \left(\omega \cdot \sqrt{\frac{c+v}{c-v}} + \omega \cdot \sqrt{\frac{c-v}{c+v}} \right)$$

Because $E = 2m_0(v) \cdot c^2$, there are:

$$2m_0(v) \cdot c^2 = h \left(\omega \cdot \sqrt{\frac{c+v}{c-v}} + \omega \cdot \sqrt{\frac{c-v}{c+v}} \right) \quad (9)$$

Solving equation (9) yields:

$$m_0(v) = h\omega \cdot \sqrt{\frac{c^2}{c^2 - v^2}} = m_0 \cdot \sqrt{\frac{1}{1 - \frac{v^2}{c^2}}}$$

This is the dynamic mass formula in the chivalrous theory of relativity.

Prove it.

1.4 Proof of the relativistic velocity transformation formula

1.4.1 The first way of proof




Fig. 5 Relativistic velocity transformation formula of Doppler effect

As shown in Figure 5 above, in the view of the ground observer S system, both objects A and B move along the X-axis direction of the S system, where the speed of B relative to the S system is v , and the speed of A relative to the S system is u (u moves in the negative direction of the X axis, so it has a negative sign).

Now we ask for the relativistic velocity transformation formula: What is the speed of motion of A in the view of the B system?

We assume that B emits a photon of frequency ω to A, which is absorbed by an object P in

the S system between A and B, and assume that the photon redshifted frequency is ω_1 when the photon absorbed by the S system. Then the S-system object P emits a photon of the same frequency ω_1 to A from the other side. We assume that the frequency of the photon observed when A absorbs the photon is ω_2 (we assume that the masses of A and B are relatively high relative to the photon. is large, and the difference in motion speed before and after emitting photons and absorbing photons can be ignored).

From the point of view of the S system, the energy of the absorbed and emitted photons of frequency ω_1 is the same and in opposite directions, so both the momentum and the energy of the S system remain unchanged.

At this point, the whole process is equivalent to directly emitting a photon of energy ω from B and absorbed by the A object. According to our previous conclusion of "Theorem 3", the photon frequency ω_2 received by A is:

$$\omega_2 = \omega_1 \cdot \sqrt{\frac{c+u}{c-u}} = \omega \cdot \sqrt{\frac{c-v}{c+v}} \cdot \sqrt{\frac{c+u}{c-u}} \quad (10)$$

(Note: Since the velocity u is moving in the negative direction of the X axis, its sign is negative.)

If we assume that in the view of B, the speed of A's movement is u' (the direction of movement of u' is the negative direction of the X-axis of B), then the above formula (10) will become:

$$\omega \cdot \sqrt{\frac{c-v}{c+v}} \cdot \sqrt{\frac{c+u}{c-u}} = \omega \cdot \sqrt{\frac{c+u'}{c-u'}}$$

Solving the above equation yields:

$$u' = \frac{u-v}{1 - \frac{uv}{c^2}}$$

This is the velocity transformation formula in the special theory of relativity. The proofs in other motion situations can refer to the above proofs.

Prove it.

Of course, we can derive it in another way as follows and obtain more useful conclusions.

1.4.2 The second way of proof

1.4.2.1 Let's do the following thought experiment:

Assuming that the static mass of the object A is m_A , A moves in the X-axis direction with a velocity v_A relative to the S-series ground observer (the X-axis direction of the A-series is

parallel to the ground X-axis). There are two bodies B and C with static mass $m_B = m_C = m_0$. Their velocity relative to the ground S system is unknown, but their velocity relative to object A is v and $-v$, respectively. Now, what is the speed of the objects B and C in the view of the ground observer S system ? (we assume that their velocities are v_1 and v_2 respectively in the view of the S system), as shown in Figure 6 below, we can also call the ground reference system For the S system, the reference frame of the object A is the S' system or the A system:




Fig. 6 Transformation of object motion along x-axis

There are two potential hypotheses here, which we will conversely prove correct through the speed-shifting argument:

1: the static mass of any object is consistent in any inertial reference frame, for example, we observe the mass of electron e in any inertial frame, and it will have the same properties.

2: Assuming that the objects in the two reference frames do not feel any acceleration, their relative velocity relative to the other frame of reference will remain unchanged (we will not discuss the explanation of the equivalence principle of gravitational fields for now, which will be explained in the next article).

(1) We assume that the moving directions of objects B and C are both in the direction of the X-axis. From the perspective of the A system, B and C collide with velocities v and $-v$ respectively and merge together after the collision. Then from the point of view of system A, since B and C have equal masses and opposite speeds, B and C will remain stationary relative to system A after collision. At this time, in the view of the ground observer S, B and C will move along the X-axis with a velocity v_A relative to the ground after the collision.

Then in the view of the ground observer S system, due to the energy conservation before and after the collision of B and C, the total mass before the collision is:

$$m_0 \sqrt{\frac{c^2}{c^2 - v_1^2}} + m_0 \sqrt{\frac{c^2}{c^2 - v_2^2}} \quad (11)$$

From the point of view of the A system, B and C are stationary relative to the A system after the collision, and the static mass after the B and C collision is the sum of the dynamic masses before

the collision seen by the A system: $2m_0 \sqrt{\frac{c^2}{c^2 - v^2}}$.

The static mass is also applicable to the ground observer S system, so from the ground observer S system, the total mass of B and C after collision is (dynamic mass):

$$2m_0 \sqrt{\frac{c^2}{c^2 - v^2}} \cdot \sqrt{\frac{c^2}{c^2 - v_A^2}} \quad (12)$$

The above equations (11) and (12) are equal, so we obtain the first equation:

Equation 1:

$$m_0 \sqrt{\frac{c^2}{c^2 - v_1^2}} + m_0 \sqrt{\frac{c^2}{c^2 - v_2^2}} = 2m_0 \sqrt{\frac{c^2}{c^2 - v^2}} \cdot \sqrt{\frac{c^2}{c^2 - v_A^2}}$$

Similarly, we can get a momentum conservation equation:

Equation 2:

$$\vec{v}_1 \cdot m_0 \sqrt{\frac{c^2}{c^2 - v_1^2}} + \vec{v}_2 \cdot m_0 \sqrt{\frac{c^2}{c^2 - v_2^2}} = \vec{v}_A \cdot 2m_0 \sqrt{\frac{c^2}{c^2 - v^2}} \cdot \sqrt{\frac{c^2}{c^2 - v_A^2}}$$

Since we assume that the motion directions of v_1 、 v_2 、 v 、 v_A are all along the X-axis direction of the S system and the A system, then we directly substitute the following relativistic velocity transformation formula for verification.

According to the velocity transformation formula of special relativity^[3], the X-axis velocity component of B is:

$$v_1 = v_{1x} = \frac{v_x + v_A}{1 + \frac{v_x \cdot v_A}{c^2}} = \frac{v + v_A}{1 + \frac{v \cdot v_A}{c^2}} \quad (13)$$

The X-axis velocity direction of C is:

$$v_2 = v_{2x} = \frac{-v_x + v_A}{1 - \frac{v_x \cdot v_A}{c^2}} = \frac{-v + v_A}{1 - \frac{v \cdot v_A}{c^2}} \quad (14)$$

It is easy to verify that the two equations (13) and (14) are the solutions of "Equation 1" and "Equation 2" by substituting (13) and (14) into the above "Equations 1" and "Equations 2".

Conclusion: The velocity transformation formula of special relativity is valid in the direction of X axis.

1.4.2.2 Discussion of the more general case

As shown in Figure 7 below:




Fig. 7 Speed transformation in general case

Objects B and C have the same static mass, and B and C collide and combine together. In the view of system A, the velocities of objects B and C are v and v' respectively. The three-direction velocity components of the x, y, and z axes of B and C are (v_x, v_y, v_z) and $(v_x, -v_y, -v_z)$, respectively.

That is to say, from the point of view of the A system, the movement speed of the two objects after the collision will become v_x and move along the X axis. Since v_x remains unchanged before and after the collision, obviously for the S system, the velocity component of the X-axis direction before and after the collision of objects B and C is also unchanged.

Assuming that the moving velocities of the two objects B and C are v_1 and v_2 respectively in the S system, we assume that their velocity components in the directions of the three coordinate axes are: (v_{1x}, v_{1y}, v_{1z}) and (v_{2x}, v_{2y}, v_{2z}) , then there are: $v_{1x} = v_{2x}$, $v_{1y} = -v_{2y}$, $v_{1z} = -v_{2z}$.

Since we can always rotate the coordinate axis while keeping the direction of the X axis unchanged, the velocity components of the objects B and C in the Y and Z axes become only moving along the Y' axis in the new coordinate system. (The axes of the A system and the S system are always kept parallel), so in fact we only need to consider the speed transformation in the direction of the Y axis, so we can simply assume $v_{1z} = -v_{2z} = 0$, $v_z = -v_z = 0$, so there are:

$$v_1^2 = v_2^2 = v_{1x}^2 + v_{1y}^2 = v_{2x}^2 + v_{2y}^2$$

Assuming that the static mass before the collision between B and C is m_0 , and the static mass after the collision is m_1 , according to the conservation of energy before and after the collision, in the view of the A system, there are:

$$2m_0 \cdot \sqrt{\frac{c^2}{c^2 - (v_x^2 + v_y^2)}} = 2m_1 \cdot \sqrt{\frac{c^2}{c^2 - v_x^2}}$$

where m_1 is the static mass of objects B and C after collision, so the static mass m_1 of objects B and C after collision is obtained as:

$$m_1 = m_0 \cdot \sqrt{\frac{c^2}{c^2 - (v_x^2 + v_y^2)}} \cdot \sqrt{\frac{c^2 - v_x^2}{c^2}} \quad (15)$$

According to our previous conclusion on the velocity transformation formula in the X-axis direction, in the S system, there are:

$$v_{1x} = v_{2x} = \frac{v_x + v_A}{1 + \frac{v_x \cdot v_A}{c^2}} \quad (16)$$

Due to the conservation of energy before and after the collision, in the S system, there are:

$$2m_0 \cdot \sqrt{\frac{c^2}{c^2 - (v_{1x}^2 + v_{1y}^2)}} = 2m_1 \cdot \sqrt{\frac{c^2}{c^2 - v_{1x}^2}} \quad (17)$$

$$\text{where, } v_1^2 = v_2^2 = v_{1x}^2 + v_{1y}^2 = v_{2x}^2 + v_{2y}^2$$

Substituting the above formula (15) into the formula (17), we get:

$$2m_0 \cdot \sqrt{\frac{c^2}{c^2 - (v_{1x}^2 + v_{1y}^2)}} = 2m_0 \cdot \sqrt{\frac{c^2}{c^2 - (v_x^2 + v_y^2)}} \cdot \sqrt{\frac{c^2 - v_x^2}{c^2}} \cdot \sqrt{\frac{c^2}{c^2 - v_{1x}^2}}$$

Then substitute equation (16) into the above equation, and solve the equation after expansion to obtain:

$$v_{1y} = \pm \frac{v_y}{1 + \frac{v_x v_A}{c^2}} \cdot \sqrt{1 - \frac{v_A^2}{c^2}}$$

Taking into account the consistency of the directions of v_y and v_{1y} , the positive sign is taken, which is the component of Einstein's special relativity velocity transformation formula in the Y-axis and Z-axis directions.

Prove it.

Therefore, we have proved the relativistic velocity transformation formula through the above thought experiment; at the same time, we have also proved the correctness of the previous two assumptions: the static mass of the object in the inertial frame is unique, otherwise the velocity transformation formula will not hold; at the same time, we also The premise of proving that the relative speed of motion between two objects changes is the force.

1.4.3 Explanation of the above derivation

The derivation of the relativistic transformation formula in this paper is based on the assumption of inertial system, mass-energy equation, energy conservation, momentum conservation, Planck's energy-frequency formula, etc. The core essence of the transformation formula is actually implied by the inertial system. The transformation relationship between the conservation of energy and momentum. We did not use the time transformation relationship and length transformation relationship of relativity, and also deduced the entire relativistic transformation formula.

This proves that the core of the relativistic transformation formula is actually the observance and transformation of momentum conservation and energy conservation.

2 Conclusion

In this paper, based on the mass-energy equation and Planck's energy and momentum formula, without the hypothesis of time slowing down and length contraction, only based on the conservation of momentum and energy, the Doppler red shift and blue shift formula, the kinetic mass equation and the velocity transformation formula of relativity are strictly deduced and proved.

In addition, this paper also supplements a defect of special relativity in demonstrating the Doppler frequency shift formula through rigorous derivation: we have proved that the relative velocities of two objects A and B in the photon frequency conversion formula actually refer to The relative velocity of A and B after object A or B emits photons. In this way, the Doppler frequency shift formula can be measured and calculated more rigorously in the experiment.

At the same time, this paper also revises the traditional relativistic aberration formula, which is also one of the most important conclusions of this paper. This can be used in high-speed particle radiation experiments, astronomical observations, or laser experiments on moving objects.

This paper also proves that in the special theory of relativity, the two conclusions of time slowing down and length shrinking are superfluous. Then our conclusion in another article may be correct: simultaneity is possible in different inertial frames, and the space is relatively independent. In this way, we can make more reasonable explanations for phenomena such as quantum entanglement at a distance and cosmic inflation^[12].

References:

- [1] Feng Shimeng. The characteristic derivation method of the relationship between mass and velocity in the special theory of relativity. University Physics, 2021, Vol.40, No.6: 32-35.
- [2] Wang Yonggang, Cao Xuecheng, etc., Deriving the Arbitrary Direction Acceleration Transformation of Special Relativity Using Space-Time Algebra. University Physics, 2019, Vol.38, No.4: 8-12.
- [3] Dai Youshan, Song Xiaotong, New derivation of special relativity without the assumption of constant speed of light. Journal of Zhejiang University, 2016, Vol.43, No.2: 247-251.
- [4] Lu Junling, Wang Rongbao, Simple derivation of Doppler effect formula. University Physics, 2005, Vol.24, No.8:25-29.

-
- [5] Gao Bingkun, Wang Fenglin, Simple derivation of the Doppler effect of relativistic theory. University Physics, 2003, Vol.22, No.8:15-16.
- [6] Hirsch I. Mandelberg; Louis Witten, Journal of the Optical Society of America, Volume 52, Issue 5.1962.PP:529-535
- [7] National Institute of Technology, USA. Einstein's special theory of relativity confirmed by precise experiments in the United States. Bulletin of Physics. 2006, Issue 1: P64
- [8] A. H. Compton, Phys. Rev., Vol.22 (1923), p409
- [9] Einstein, translated by Yang Zi. Theory of Relativity. Beijing: Taihai Publishing House, 2017.
- [10] Zhao Kaihua, Chen Ximou. Electromagnetics. Beijing: Higher Education Press, 2021.
- [11] Simon Rainville, James K. Thompson et al, A direct test of $E=mc^2$, Nature, 2005, 438:1096 – 1097.
- [12] Qin, Sheng (2022): A possible explanation for the twin paradox and action at a distance.docx. figshare. Thesis. <https://doi.org/10.6084/m9.figshare.19807654.v2>