

DEER analysis report on dataset DEER4p_Q_50K_PTBP1_IRES_WT_109-475 _18us_166sc



**DEERNet Spinach SVN Rev 5662 and DeerLab
0.9.1 Tikhonov regularization**

ComparativeDEERAnalyzer version 2.0

see: S. G. Worswick et al., DOI: 10.1126/sciadv.aat5218, L. Fabregas Ibanez et al., DOI: 10.5194/
mr-1-209-2020


23-Feb-2022 18:15:20

1. Distance distributions





2. Fits of time-domain data

DEERNet fits and background fits



Tikhonov fit





3. Experimental and processing parameters

Modulation depth: 0.211

Signal-to-noise ratio: 146.6 (w.r.t. modulation)

Noise estimates normalized to maximum signal

From imaginary part: 0.00165

From DEERNet fit: 0.00144

From Tikhonov fit: 0.00138

Zero time: 0 ns

Maximum time: 18000 ns

Time increment: 40 ns

Phase: -11.9 degree

Ensemble of 32 neural networks

Background separation by neural network

Background dimension: 3

Regularization parameter by best overlap with neural network solution

Regularization parameter used: 5.60

Reg. par. initial estimate by L-curve corner: 15.85

Overlap between DEERNet and regularization solutions: 0.931

Predicted overlap of consensus solution with ground truth: 0.80...0.97

Mean distance: 78.3 Å

Distance standard deviation: 11.6 Å

Full data set in Matlab format:

G:\projects\Christoph_Gmeiner\modelling\master_shot\Deer\DEER4p_Q_50K_PTBP1_IRES_WT_109-475_18us_166sc_comparative_DEER_analysis.mat

Distance distributions in text format:

G:\projects\Christoph_Gmeiner\modelling\master_shot\Deer\DEER4p_Q_50K_PTBP1_IRES_WT_109-475_18us_166sc_consensus_DEER_distribution.csv

Fit and background in text format:

G:\projects\Christoph_Gmeiner\modelling\master_shot\Deer\DEER4p_Q_50K_PTBP1_IRES_WT_109-475_18us_166sc_consensus_DEER_fit.csv

3. Experimental and processing parameters

Metadata:

G:\projects\Christoph_Gmeiner\modelling\master_shot\Deer\DEER4p_Q_50K_PTBP1_IRES_
WT_109-475_18us_166sc_comparative_DEER_meta_data.csv