

# W2W: A Python package that injects WUDAPT's Local Climate Zone information in WRF

Matthias Demuzere<sup>\*1</sup>, Daniel Argüeso<sup>2</sup>, Andrea Zonato<sup>3</sup>, and Jonas Kittner<sup>1</sup>

<sup>1</sup> Urban Climatology Group, Department of Geography, Ruhr-University Bochum, Bochum, Germany <sup>2</sup> Physics Department, University of the Balearic Islands, Palma, Spain <sup>3</sup> Atmospheric Physics Group, Department of Civil, Environmental and Mechanical Engineering, University of Trento, Trento, Italy

DOI: [10.21105/joss.0XXXX](https://doi.org/10.21105/joss.0XXXX)

## Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Editor: [Editor Name](#) ↗

Submitted: 01 January XXXX

Published: 01 January XXXX

## License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

## Summary

The Python-based WUDAPT-to-WRF (W2W) package is developed to translate Local Climate Zone (LCZ) maps into urban canopy parameters readable by WRF, the community “Weather Research and Forecasting” model (Skamarock et al., 2021). It is the successor of the Fortran-based W2W package developed by Brousse et al. (2016) and Martilli et al. (2016), and provides an improved, simpler, and more efficient procedure to use LCZ information in WRF. Some important changes include direct manipulation of the geogrid files without the creation of temporary files, and the use of average LCZ-based urban morphological parameters instead of assigning them to the modal LCZ class.

This development of this package is in line with the objectives of WUDAPT, the World Urban Database and Access Portals Tools community project, that aims to 1) acquire and make accessible coherent and consistent information on the form and function of urban morphology relevant to climate weather and environmental studies, and 2) provide tools that extract relevant urban parameters and properties for models and model applications at appropriate scales for various climate, weather, environment, and urban planning purposes (Ching et al., 2018).

## Statement of need

Since the pioneering work of Brousse et al. (2016) and Martilli et al. (2016), the level-0 WUDAPT information, the Local Climate Zone maps, have been used increasingly in WRF.

We expect this trend to continue because of three recent developments: 1) the creation of city-wide LCZ maps is now easier than ever with the launch of the LCZ Generator web application (Demuzere et al., 2021), 2) the availability of a global LCZ map (Demuzere et al., 2022), and 3) WRF versions > 4.3 (Skamarock et al., 2021) can ingest 10 or 11 built classes (corresponding to WUDAPT's LCZs) by default, whereas previous WRF versions required manual code changes (see Martilli et al. (2016), Zonato & Chen (2021) and Zonato et al. (2021) for more information).

Because of these developments, an improved, Python-based, WUDAPT-to-WRF (W2W) routine is presented here to translate LCZ-based parameters better and simpler. It differs from its Fortran-based predecessor mainly by 1) using a more up-to-date LCZ-based urban extent, 2)

<sup>\*</sup>corresponding author

38 aggregating the morphological parameters instead of using modal values, and 3) the fact that  
39 all processing is done with one (automated) tool, whereas the Fortran-based version required  
40 multiple pre-processing steps and manual namelist changes, described in more detail by Martilli  
41 et al. (2016).

## 42 Initial data requirements

43 To use the tool, two input files are required:

- 44 1. A `geo_em.dXX` (.nc) file for the inner WRF model domain in which one would like to  
45 use the LCZ-based information. This file can be produced by WRF's `geogrid.exe` tool  
46 as part of the WRF Preprocessing System (WPS), without additional modifications of  
47 the standard procedure.
- 48 2. A **Local Climate Zone map** (.tif) file that is slightly bigger than the domain extent of  
49 the `geo_em.dXX.nc` file. There are several ways to obtain an LCZ map for your region  
50 of interest (ROI):
  - 51 ■ Extract your ROI from the global LCZ map (Demuzere et al., 2022), or the continental-  
52 scale LCZ maps for Europe (Demuzere et al., 2019) or the United States (Demuzere et  
53 al., 2020) (see [here](#) for more info).
  - 54 ■ Check if your ROI is already covered by the many LCZ maps available in the [submission](#)  
55 [table](#) of the LCZ Generator.
  - 56 ■ Use the [LCZ Generator](#) to make your LCZ map for your ROI. See [here](#) for more informa-  
57 tion. When using LCZ maps produced with the LCZ Generator, by default the Gaussian  
58 filtered LCZ map is used (corresponding to argument `--lcz-band = 1`).

59 **Note:** When using LCZ information from any of the large-scale LCZ maps, please make sure  
60 to crop your domain of interest first to avoid memory issues.

## 61 Workflow

62 The goal of the Python-based W2W tool is to obtain an inner WRF domain file (`geo_em.dX`  
63 `X.nc`) that contains the built LCZ classes and their corresponding urban canopy parameters  
64 relevant for all urban parameterizations embedded in WRF: the single-layer urban canopy  
65 model (Noah/SLUCM, Kusaka et al. (2001)), the Building Environment Parameterization  
66 (BEP, Martilli et al. (2002)), and BEP+BEM (Building Energy Model, Salamanca et al.  
67 (2010)).

68 To get to that point, a number of sequential steps are required:

### 69 Step 1: Remove the default urban land cover

70 The default urban land cover from MODIS is replaced with the dominant surrounding veg-  
71 etation category, as done in Li et al. (2020). This procedure affects WRF's parameters  
72 `LU_INDEX`, `LANDUSEF`, and `GREENFRAC`. First, an initial number of neighboring pixels  
73 (corresponding argument `--npix-area`, default = `--npix-nlc ** 2`) are selected using  
74 `scipy`'s k-d tree algorithm (Maneewongvatana & Mount, 1999), assuming a spherical Earth,  
75 and using the Euclidean distance along the great circle arc (i.e., section of the Earth that di-  
76 vides the sphere into two equal parts) to find the nearest pixels. Afterwards, the `LU_INDEX`

is set by sampling the dominant category from the corresponding argument `--npix-nlc` (default = 45) nearest initial grid points (excluding ocean, urban, and lakes). GREENFRAC is calculated as the mean over all grid points with that dominant vegetation category among the `--npix-nlc` nearest points. For each grid point, if LANDUSEF had any percentage of urban, it is set to zero and the percentage is added to the dominant vegetation category assigned to that grid point.

Resulting output: `geo_em.dXX_NoUrban.nc`

## Step 2: Define the LCZ-based urban extent

LCZ-based impervious fraction values (FRC\_URB2D, available from `LCZ_UCP_default.csv`) are assigned to the original 100 m resolution LCZ map, and are aggregated to the WRF resolution. Areas with `FRC_URB2D < 0.2` (corresponding to argument `--frc-threshold`) are currently considered non-urban. This choice has been made to avoid the use of the urban schemes in areas where the majority of the land use is vegetated since the impact of the impervious surfaces is low. The FRC\_URB2D field is also used to mask all other urban parameter fields so that their extent is consistent.

Resulting output: `geo_em.dXX_LCZ_extent.nc`

## Step 3: Introduce modal built LCZ classes

For each WRF grid cell, the mode of the underlying built LCZ classes is added to LU\_INDEX (numbered from 31-41). See [here](#) for more info. Note that the W2W routine by default considers LCZ classes 1-10 as built classes (corresponding to argument `--built-lcz`). In some cases, also LCZ E (or 15 - Bare rock or paved) can be considered as a built LCZ class, as it might reflect large asphalt surfaces such as big parking lots or airstrips. In that case, the user must make sure the `--built-lcz` argument is set appropriately.

## Step 4: Assign urban canopy parameters

Two procedures are followed when assigning the various urban canopy parameters to the LCZ map and translating this information onto WRF's grid:

**Procedure 1: Morphological parameters** are assigned directly to the high-resolution LCZ map, and are afterward aggregated to the lower-resolution WRF grid. As a result, the method produces a unique urban morphology parameter value for each WRF grid cell. This was found to be more efficient in reproducing urban boundary layer features, especially in the outskirts of the city (Zonato et al., 2020), and is in line with the [WUDAPT-to-COSMO](#) routine (Varentsov et al., 2020).

Morphological urban canopy parameter values are provided in `LCZ_UCP_default.csv` and are generally based on values provided in Stewart & Oke (2012) and Stewart et al. (2014). Note however that the values of MH\_URB2D\_MIN, MH\_URB2D, and MH\_URB2D\_MAX for LCZ 7 are set to 4, 5, and 6 m instead of 2, 3, and 4 m because the minimum building height that can be assigned to BEP-BEM is 5m if `dz_u = 5m` (standard value) is used.

In addition:

- While `URBPARM_LCZ.TBL` (stored in WRF's `run/` folder) has values on street width (SW), W2W derives street width from the mean building height (MH\_URB2D) and the Height-to-Width ratio (H2W), to have these fields consistent.
- Building width (BW), is derived from `(BLDFR_URB2D / (FRC_URB2D - BLDFR_URB2D)) * SW`, these values are available in the look-up table `LCZ_UCP_default.csv`.

- 120 ■ Plan (LP\_URB2D), frontal (LF\_URB2D), and total (LB\_URB2D) area indices are
- 121 based on formulas in Zonato et al. (2020).
- 122 ■ HI\_URB2D is obtained by fitting a bounded normal distribution to the minimum
- 123 (MH\_URB2D\_MIN), mean (MH\_URB2D), and maximum (MH\_URB2D\_MAX)
- 124 building height, as provided in LCZ\_UCP\_default.csv. The building height
- 125 standard deviation is also required and is approximated as  $(MH\_URB2D\_MAX -$
- 126  $MH\_URB2D\_MIN) / 4$ .
- 127 ■ For computational efficiency, HI\_URB2D values lower than 5% were set to 0 after
- 128 resampling, and the remaining HI\_URB2D percentages are re-scaled to 100%.

129 **Procedure 2:** In line with the former Fortran-based W2W procedure, **radiative and thermal**  
 130 **parameters** are assigned to the modal LCZ class that is assigned to each WRF grid cell (see  
 131 *Step 3*). These parameter values are not stored in the NetCDF output but are read from  
 132 URBPARAM\_LCZ.TBL and assigned automatically to the modal LCZ class when running the  
 133 model.

## 134 Step 5: Adjust global attributes


135 In a final step, some global attributes are adjusted in the resulting NetCDF files:

- 136 ■ NBUI\_MAX is added as a global attribute, reflecting the maximum amount of
- 137 HI\_URB2D classes that are not 0 across the model domain. This parameter can be
- 138 used when compiling WRF to optimize memory storage.
- 139 ■ NUM\_LAND\_CAT is set to 41, to reflect the addition of 10 or 11 built LCZ classes.
- 140 This is not only done for the highest resolution domain file (e.g. d04), but also for **all of**
- 141 **its lower-resolution parent domain files (e.g. d01, d02, d03)**. As such, make sure
- 142 these files are also available in the input data directory. In case the parent domain files
- 143 have  $NUM\_CAT\_LAND \neq 41$ , new parent domain files will be written to your drive
- 144 with the extension \_41.

145 Resulting output: `geo_em.dXX_LCZ_params.nc` (and `geo_em.dXX_41.nc`)

## 146 Integration in WRF's preprocessing

147 The current tool is designed to work with the `geo_em.dXX` files produced by `geogrid.exe`,  
 148 which is available in the WRF Preprocessing System (WPS). WPS needs to be at a version  
 149  $>3.8$ , to incorporate the urban geometrical parameters in the URB\_PARAM matrix (Glotfelty  
 150 et al., 2013). The user should run `geogrid.exe` using its default settings, which will provide  
 151 the various `geo_em.dXX.nc` files containing the static data fields. No additional variables  
 152 are required, neither in the `namelist.wps` nor within the `GEOGRID.TBL` table. The W2W  
 153 tool ([Figure 1](#)) reads the standard `geo_em.dXX.nc` files (for all the domains) and produces  
 154 the aforementioned `geo_em.dXX_LCZ_params.nc` files. The user should then rename these  
 155 files to the standard name for each of the domains (e.g. rename `geo_em.d01_41.nc` to  
 156 `geo_em.d01.nc` and `geo_em.d04_LCZ_params.nc` to `geo_em.d04.nc`), which will serve as  
 157 input to the `metgrid.exe` module ([Figure 1](#)).



**Figure 1:** Modified workflow to set up and run a WRF simulation including urban parameters derived from LCZs using W2W.

## Potential use cases

The files provided as output by W2W allow a wide range of applications, including - but not limited to - addressing the impact of:

- urbanization, by running WRF with the default `geo_em.dXX.nc` and the `geo_em.dXX_NoUrban.nc` files (see for example Li et al. (2020) and Hirsch et al. (2021)).
- an improved urban land cover extent description, by running WRF with the default `geo_em.dXX.nc` and the `geo_em.dXX_LCZ_extent.nc` files (similar to for example Bhati & Mohan (2018) and Mallard et al. (2018)).
- a more detailed (LCZ-based) urban description, by running WRF with the default `geo_em.dXX.nc` and the `geo_em.dXX_LCZ_params.nc` files (see for example Brousse et al. (2016), Hammerberg et al. (2018), Molnár et al. (2019), Wong et al. (2019), Patel et al. (2020), Zonato et al. (2020), Ribeiro et al. (2021), Hirsch et al. (2021) and Patel et al. (2022)).

## Important notes

- Make sure to set `use_wudapt_lcz=1` (default is 0) and `num_land_cat=41` (default is 21) in WRF's `namelist.input` when using the LCZ-based urban canopy parameters.
- The LCZ-based urban canopy parameter values provided in `LCZ_UCP_default.csv` and `URBPARAM_LCZ.TBL` are universal and generic, and might not be appropriate for your ROI. If available, please adjust the urban canopy parameters values according to the characteristics of your ROI. A custom csv file can be specified using the `--lcz-ucp path/to/custom_file.csv` flag.
- It is advised to use this tool with urban parameterization options BEP or BEP+BEM (`sf_urban_physics = 2` or `3`, respectively). In case you use this tool with the SLUCM model (`sf_urban_physics = 1`), make sure your lowest model level is above the highest building height. If not, `real.exe` will provide the following error message:

183 ZDC + ZOC + 2m is larger than the 1st WRF level - Stop in subroutine  
 184 urban - change ZDC and ZOC.  
 185 ■ At the end of running W2W, a note is displayed that indicates the nbui\_max value, e.g. for  
 186 the sample data: Set nbui\_max to 5 during compilation, to optimize memory  
 187 storage. This is especially relevant for users that work with the BEP or BEP+BEM  
 188 urban parameterization schemes (sf\_urban\_physics = 2 or 3, respectively). See  
 189 num\_urban\_nbui in [WRF's README.namelist](#) for more info.  
 190 ■ It is advised to use WRF versions > 4.3, that can ingest 10 or 11 built classes (corresponding  
 191 to WUDAPT's LCZs) by default (Skamarock et al., 2021), and WPS versions  
 192 > 3.8, to incorporate the urban geometrical parameters in the URB\_PARAM matrix (Glotfelty  
 193 et al., 2013).

## 194 Acknowledgements

195 We acknowledge contributions and support from Alberto Martilli, Alejandro Rodriguez  
 196 Sanchez, and Oscar Brousse.

## 197 References

- 198 Bhati, S., & Mohan, M. (2018). WRF-urban canopy model evaluation for the assessment of  
 199 heat island and thermal comfort over an urban airshed in India under varying land use/land  
 200 cover conditions. *Geoscience Letters*, 5(1). <https://doi.org/10.1186/s40562-018-0126-7>
- 201 Brousse, O., Martilli, A., Foley, M., Mills, G., & Bechtel, B. (2016). WUDAPT, an efficient  
 202 land use producing data tool for mesoscale models? Integration of urban LCZ in WRF  
 203 over madrid. *Urban Climate*, 17, 116–134. <https://doi.org/10.1016/j.uclim.2016.04.001>
- 204 Ching, J., Mills, G., Bechtel, B., See, L., Feddema, J., Wang, X., Ren, C., Brorousse, O.,  
 205 Martilli, A., Neophytou, M., Mouzourides, P., Stewart, I., Hanna, A., Ng, E., Foley,  
 206 M., Alexander, P., Aliaga, D., Niyogi, D., Shreevastava, A., ... Theeuwesits, N. (2018).  
 207 WUDAPT: An urban weather, climate, and environmental modeling infrastructure for  
 208 the anthropocene. *Bulletin of the American Meteorological Society*, 99(9), 1907–1924.  
 209 <https://doi.org/10.1175/BAMS-D-16-0236.1>
- 210 Demuzere, M., Bechtel, B., Middel, A., & Mills, G. (2019). Mapping Europe into local climate  
 211 zones. *PLOS ONE*, 14(4), e0214474. <https://doi.org/10.1371/journal.pone.0214474>
- 212 Demuzere, M., Hankey, S., Mills, G., Zhang, W., Lu, T., & Bechtel, B. (2020). Combining expert  
 213 and crowd-sourced training data to map urban form and functions for the continental  
 214 US. *Scientific Data*, 7(1), 264. <https://doi.org/10.1038/s41597-020-00605-z>
- 215 Demuzere, M., Kittner, J., & Bechtel, B. (2021). LCZ Generator: A Web Application to  
 216 Create Local Climate Zone Maps. *Frontiers in Environmental Science*, 9(April). <https://doi.org/10.3389/fenvs.2021.637455>
- 217  
 218 Demuzere, M., Kittner, J., Martilli, A., Mills, G., Moede, C., Stewart, I. D., Vliet, J. van,  
 219 & Bechtel, B. (2022). A global map of Local Climate Zones to support earth system  
 220 modelling and urban scale environmental science. *Earth System Science Data Discussions*.  
 221 <https://doi.org/10.5194/essd-2022-92>
- 222 Glotfelty, T., Tewari, M., Sampson, K., Duda, M., Chen, F., & Ching, J. (2013). *NUDAPT 44:*  
 223 *How to use NUDAPT dataset in WRF/SLUCM/MLUCM models* (p. 9). National Center  
 224 for Atmospheric Research. <https://www.yumpu.com/en/document/read/26871494/how-to-use-nudapt-dataset-in-wrf-slucm-mlucm-models>  
 225



- 226 Hammerberg, K., Brousse, O., Martilli, A., & Mahdavi, A. (2018). Implications of employing  
227 detailed urban canopy parameters for mesoscale climate modelling: a comparison between  
228 WUDAPT and GIS databases over Vienna, Austria. *International Journal of Climatology*,  
229 38, e1241–e1257. <https://doi.org/10.1002/joc.5447>
- 230 Hirsch, A. L., Evans, J. P., Thomas, C., Conroy, B., Hart, M. A., Lipson, M., & Ertler,  
231 W. (2021). Resolving the influence of local flows on urban heat amplification during  
232 heatwaves. *Environmental Research Letters*, 16(6), 064066. <https://doi.org/10.1088/1748-9326/ac0377>  
233
- 234 Kusaka, H., Kondo, H., Kikegawa, Y., & Kimura, F. (2001). A Simple Single-Layer Ur-  
235 ban Canopy Model For Atmospheric Models: Comparison With Multi-Layer And Slab  
236 Models. *Boundary-Layer Meteorology*, 101(3), 329–358. <https://doi.org/10.1023/A:1019207923078>  
237
- 238 Li, Y., Fowler, H. J., Argüeso, D., Blenkinsop, S., Evans, J. P., Lenderink, G., Yan, X.,  
239 Guerreiro, S. B., Lewis, E., & Li, X. F. (2020). Strong Intensification of Hourly Rainfall  
240 Extremes by Urbanization. *Geophysical Research Letters*, 47(14), 1–8. <https://doi.org/10.1029/2020GL088758>  
241
- 242 Mallard, M. S., Spero, T. L., & Taylor, S. M. (2018). Examining WRF’s sensitivity to con-  
243 temporary land-use datasets across the contiguous united states using dynamical down-  
244 scaling. *Journal of Applied Meteorology and Climatology*, 57(11), 2561–2583. <https://doi.org/10.1175/JAMC-D-17-0328.1>  
245
- 246 Maneewongvatana, S., & Mount, D. M. (1999). Analysis of Approximate Nearest Neighbor  
247 Searching with Clustered Point Sets. *Computer Science - ArXiv*. <https://doi.org/10.48550/arXiv.cs/9901013>  
248
- 249 Martilli, A., Brousse, O., & Ching, J. (2016). *Urbanized WRF modeling using WU-*  
250 *DAPT*. <http://www.wudapt.org/wudapt-to-wrf/>. <https://www.wudapt.org/wp-content/uploads/2016/05/Urbanized-WRF-modeling-using-WUDAPT-web-version-March2016.pdf>  
251  
252
- 253 Martilli, A., Clappier, A., & Rotach, M. W. (2002). An urban surface exchange parame-  
254 terisation for mesoscale models. *Boundary-Layer Meteorology*, 104(2), 261–304. <https://doi.org/10.1023/A:1016099921195>  
255
- 256 Molnár, G., Gyöngyösi, A. Z., & Gál, T. (2019). Integration of an LCZ-based classification into  
257 WRF to assess the intra-urban temperature pattern under a heatwave period in Szeged,  
258 Hungary. *Theoretical and Applied Climatology*, 138(1-2), 1139–1158. <https://doi.org/10.1007/s00704-019-02881-1>  
259
- 260 Patel, P., Jamshidi, S., Nadimpalli, R., Aliaga, D. G., Mills, G., Chen, F., Demuzere, M.,  
261 & Niyogi, D. (2022). Modelling Large-Scale Heatwave by Incorporating Enhanced Urban  
262 Representation. *Journal of Geophysical Research : Atmospheres*, 127, 1–33. <https://doi.org/10.1029/2021JD035316>  
263
- 264 Patel, P., Karmakar, S., Ghosh, S., & Niyogi, D. (2020). Improved simulation of very heavy  
265 rainfall events by incorporating WUDAPT urban land use/land cover in WRF. *Urban*  
266 *Climate*, 32(July 2019), 100616. <https://doi.org/10.1016/j.uclim.2020.100616>
- 267 Ribeiro, I., Martilli, A., Falls, M., Zonato, A., & Villalba, G. (2021). Highly resolved  
268 WRF-BEP/BEM simulations over Barcelona urban area with LCZ. *Atmospheric Research*,  
269 248(August 2020), 105220. <https://doi.org/10.1016/j.atmosres.2020.105220>
- 270 Salamanca, F., Krpo, A., Martilli, A., & Clappier, A. (2010). A new building energy model  
271 coupled with an urban canopy parameterization for urban climate simulations—part I.  
272 formulation, verification, and sensitivity analysis of the model. *Theoretical and Applied*  
273 *Climatology*, 99(3-4), 331–344. <https://doi.org/10.1007/s00704-009-0142-9>

- 274 Skamarock, W. C., Klemp, J. B., Dudhia, J. B., Gill, D. O., Liu, Z., Berner, J., Wang, W.,  
 275 Powers, J. G., Duda, M. G., Barker, D. M., & Huang, X.-Y. (2021). *A Description of*  
 276 *the Advanced Research WRF Model Version 4.3* (July). National Center for Atmospheric  
 277 Research. <https://doi.org/10.5065/1dfh-6p97>
- 278 Stewart, I. D., & Oke, T. R. (2012). Local Climate Zones for Urban Temperature Studies.  
 279 *Bulletin of the American Meteorological Society*, 93(12), 1879–1900. <https://doi.org/10.1175/BAMS-D-11-00019.1>  
 280
- 281 Stewart, I. D., Oke, T. R., & Krayenhoff, E. S. (2014). Evaluation of the 'local climate  
 282 zone' scheme using temperature observations and model simulations [Journal Article].  
 283 *International Journal of Climatology*, 34(4), 1062–1080. <https://doi.org/10.1002/joc.3746>  
 284
- 285 Varentsov, M., Samsonov, T., & Demuzere, M. (2020). Impact of Urban Canopy Parameters  
 286 on a Megacity's Modelled Thermal Environment. *Atmosphere*, 11(12), 1349. <https://doi.org/10.3390/atmos11121349>  
 287
- 288 Wong, M. M. F., Fung, J. C. H., Ching, J., Yeung, P. P. S., Tse, J. W. P., Ren, C., Wang,  
 289 R., & Cai, M. (2019). Evaluation of uWRF performance and modeling guidance based on  
 290 WUDAPT and NUDAPT UCP datasets for Hong Kong. *Urban Climate*, 28(June 2018),  
 291 100460. <https://doi.org/10.1016/j.uclim.2019.100460>
- 292 Zonato, A., & Chen, F. (2021). *Updates of WRF-urban in WRF 4.3: Local Climate Zones,*  
 293 *Mitigation Strategies, building materials permeability and new buildings drag coefficient.*  
 294 <http://www.wudapt.org/wudapt-to-wrf/>. [https://ral.ucar.edu/sites/default/files/public/](https://ral.ucar.edu/sites/default/files/public/product-tool/urban-canopy-model/WRF_urban_update_Readme_file_WRF4.3.pdf)  
 295 [product-tool/urban-canopy-model/WRF\\_urban\\_update\\_Readme\\_file\\_WRF4.3.pdf](https://ral.ucar.edu/sites/default/files/public/product-tool/urban-canopy-model/WRF_urban_update_Readme_file_WRF4.3.pdf)
- 296 Zonato, A., Martilli, A., Di Sabatino, S., Zardi, D., & Giovannini, L. (2020). Evaluating the  
 297 performance of a novel WUDAPT averaging technique to define urban morphology with  
 298 mesoscale models. *Urban Climate*, 31(May 2019), 100584. <https://doi.org/10.1016/j.uclim.2020.100584>  
 299
- 300 Zonato, A., Martilli, A., Gutierrez, E., Chen, F., He, C., Barlage, M., Zardi, D., & Giovannini,  
 301 L. (2021). Exploring the effects of rooftop mitigation strategies on urban temperatures  
 302 and energy consumption. *Journal of Geophysical Research: Atmospheres*, 1–30. <https://doi.org/10.1029/2021JD035002>  
 303