



CODEN [USA]: IAJPBB

ISSN : 2349-7750

INDO AMERICAN JOURNAL OF PHARMACEUTICAL SCIENCES

SJIF Impact Factor: 7.187

Available online at: <http://www.iajps.com>

Research Article

FREE RADICAL SCAVENGING PROPERTIES OF COLEUS VETTEROIDES LEAF EXTRACTS AND DNA PROTECTION AGAINST HYDROGEN PEROXIDE RADICAL DAMAGE: A PRELIMINARY ANALYSIS OF ENZYMATIC AND NON- ENZYMATIC ANTIOXIDANTS

Nagesh Ramya ¹, Kiruthika Balasubramanian ^{2*}, Jayshree Srinivas ³¹Junior Research Fellow, Department of Community Medicine, Sri Devaraj Urs Academy of Higher Education and Research, Kolar, Karnataka, India^{2*}Secretary, New Jersey Academy of Science, Kean University, New Jersey, USA³Post Graduate, Department of Biochemistry, MS Ramaiah College of Arts, Science and Commerce, Bangalore, Karnataka, India.

Article Received: September 2022 Accepted: September 2022 Published: October 2022

Abstract:

Background: For thousands of years, nature provides medicinal agents, and an incredible number of therapeutic drugs have been derived from natural sources, many based on their usage in traditional medicine. **Objective:** The goal of this study was to examine the antioxidant properties of *Coleus vetteroides* leaves. **Methodology:** Enzymic antioxidants-POD, CAT, and GST, Non-enzymic antioxidants- ascorbic acid, flavonoids, tocopherol, chlorophyll, phenols, carotenoids, and lycopene. The free radical scavenging effects and chelating properties were observed by extracting the leaves with three different solvents (Hexane, Ethylacetate, and Isopropanol). Agarose gel electrophoresis was used to test the DNA protective activity of various leaf extracts against radicals. **Result:** The plant leaves exhibited strong antioxidant and radical scavenging properties. **Conclusion:** The leaves of *Coleus vetteroides* have a potential content of both enzymatic and non-enzymatic antioxidants that could protect against oxidants and free radical damage. The phytochemical screening revealed the presence of flavonoids, alkaloids, and phenols.

Corresponding author:

Dr. Kiruthika Balasubramanian,
email id- medijoywithbk@gmail.com

QR code



Please cite this article in Kiruthika Balasubramanian et al, Free Radical Scavenging Properties Of *Coleus vetteroides* Leaf Extracts And DNA Protection Against Hydrogen Peroxide Radical Damage: A Preliminary Analysis Of Enzymatic And Non-Enzymatic Antioxidants., Indo Am. J. P. Sci, 2022; 09(10).

INTRODUCTION:

The demand for medicinal plants has grown exponentially as a result of their significance and has been used to treat illnesses from the earliest times. India is well known as the “EMPORIUM OF MEDICINAL PLANTS”⁽¹⁻³⁾. Unlike humans plants and other animals, lack a distinct immune system. Plants use a mix of chemicals to protect themselves from predators, and parasites attack plants with similar pathways that attack humans. As a result, substances used by plants to defend themselves may be useful to humans in the treatment of diseases. Popular perceptions of medicinal plant use and efficacy play an important role in the revelation of their therapeutic capabilities, thus they are regularly prescribed, even if their chemical ingredients are not always fully understood⁽⁴⁻⁷⁾.

Antioxidants are also referred to as phytochemicals, however, not all phytochemicals are antioxidants. Phytochemical consumption has been linked to a variety of health advantages. Non-infectious disorders such as cardiovascular disease, cancer, and diabetes are leading causes of death, however, phytochemicals in the diet may help to reduce the risk of these conditions.⁽⁸⁻⁹⁾.

A single free radical can harm millions of cells in the human body, preventing it from functioning properly⁽¹⁰⁾. Antioxidants are essential chemicals that protect the body from damage caused by oxidative stress generated by free radicals⁽¹¹⁾. Many enzyme systems in the human body produce reactive oxygen species (ROS) such as hydroxyl radicals, and hydrogen peroxide as a result of oxygen consumption⁽¹²⁾. These ROS are useful as signal transducers and growth regulators at modest levels⁽¹³⁾. A large range of plants has been shown to have potent antioxidant and free radical scavenging properties all around the world⁽¹⁴⁾. Antimicrobial activity testing of phytochemicals has revealed that higher plants could be a source of novel antibiotic prototypes⁽¹⁵⁾.

The purpose of this study was to look into the antioxidant capabilities, and free radical scavenging activity of *Coleus vetiveroides* leaves. It is a small and succulent herb, with pubescent leaves and fibrous roots, which are strongly aromatic. The plant's leaves can be utilized in a variety of ways, including juice mixed with honey to alleviate stomach distress and crushed leaves to treat headaches⁽¹⁶⁾.

MATERIALS AND METHODS:

The goal of the study was to analyze the antioxidant levels in the *Coleus vetiveroides* leaves. For

Enzymatic, Non-enzymatic, Radical Scavenging, Chelating, and Phytochemical screening, leaves were collected from the University of Agricultural Science, GKVK, Karnataka.

For each parameter, fresh leaves were procured, washed in running water to remove any surface contaminants, and blotted dry between the folds of soft tissue papers.

Enzymatic Antioxidants

The enzymatic antioxidants analyzed were Catalase, Peroxidase, and Glutathione S-transferase. Catalase was analyzed by the method of Luck (1974)⁽¹⁷⁾, Peroxidase by the method of Reddy et al. (1985)⁽¹⁸⁾, and Glutathione-S-Transferase by Habig et al. (1974)⁽¹⁹⁾ methods proposed respectively.

Non-Enzymatic antioxidants

Ascorbic acid, Tocopherol, Total Carotenoids and Lycopene, Total Phenols, Flavonoids, Reduced Glutathione, and Chlorophyll were analyzed using the spectrophotometric methods of Roe and Keuther (1943)⁽²⁰⁾, Rosenberg (1992)⁽²¹⁾, Zakaria *et al.* (1979)⁽²²⁾, Mallick and Singh (1980)⁽²³⁾, Cameron *et al.* (1943)⁽²⁴⁾, Moron *et al.* (1979)⁽²⁵⁾ and Witham *et al.* (1971)⁽²⁶⁾ respectively.

Radical Scavenging effects

The free radical scavenging effects of the leaves were analyzed against Hydrogen peroxide and Hydroxyl radicals

Preparation of plant extract:

Fresh *Coleus vetiveroides* leaves were collected, washed in water to remove surface pollutants, and air-dried for 10 days in the shade. The dried leaves were ground into a fine powder, which was then used to make three distinct solvent extracts using a sequential extraction process.

Sequential extraction/Successive extraction

A shaker incubator was used to dissolve 10g of powder in 100ml of hexane for 24 hours. The solvent was then filtered, and the resulting filtrate was used to create the hexane extract, which was then stored in an airtight container. The residue was air-dried, weighed, and dissolved in 100mL Ethyl acetate solvent before being incubated in a shaker for 24 hours. Ethyl acetate extract was created by filtering the solvent and storing it. The leftover residue was air-dried and weighed before being dissolved in Isopropanol solvent and the same procedure was repeated. After drying all three solvent extracts (hexane, ethyl acetate, and isopropanol), 100mg was utilized for free radical scavenging and phytochemical assays.

The H₂O₂ scavenging effect of the leaf extracts was estimated according to the method of Ruch *et al.* (1989)⁽²⁷⁾. The scavenging capacity for hydroxyl radical was measured according to the method of Elizabeth and Rao (1990)⁽²⁸⁾ and the chelating property of *Coleus vetiveroides* leaf extracts was determined based on the method of Brown *et al.* (1998)⁽²⁹⁾.

Agarose gel electrophoresis for assessing the DNA protective effect of *Coleus vetiveroides* :

Sample preparation

The procedure of Marmur (1961)⁽³⁰⁾ was used to isolate goat liver DNA: 2 g liver tissue was weighed and homogenized in 8 ml SSC solution. The pellet was separated by centrifuging the suspensions at 300 rpm for 10 minutes. The pellet was re-centrifuged at 10,000rpm for 10 minutes after adding 4ml of 0.2M NaCl. Cold ethanol was added to the supernatant twice its capacity. The precipitated DNA pellet was dissolved in TE buffer and used as part of the experiment.

1% agarose gel was prepared, and samples were loaded in the wells by mixing 2ul of 6x gel loading dye with 15ul of control DNA. 15 ul of different plant extracts were combined with 15ul of DNA and loaded in individual wells with 6x gel loading dye. DNA was damaged by adding hydrogen peroxide to generate free radicals, which were then loaded into the appropriate wells. The DNA was treated with extracts and hydrogen peroxide before being loaded into the migration pattern well. 100 volts were used for the gel electrophoresis.

Preliminary phytochemical screening of *Coleus vetiveroides* leaves

The extracts of *Coleus vetiveroides* leaves were tested for the presence of various phytochemicals using three different tests (A) Mayer's test: An aliquot of the extract was treated with Mayer's reagent (1.36g of mercuric chloride and 5g of potassium iodide in 100ml of distilled water) and were observed for the formation of cream coloured precipitate. (B) Dragendroff's test: A fraction of the extract was treated with Dragendroff's reagent and observed for the formation of a reddish-orange coloured precipitate. (C) Wagner's test: A small fraction of the extract was treated with Wagner's reagent (1.27g of iodine and 2g of KI in 100ml of distilled water) and observed for the formation of a reddish-brown coloured precipitate.

Phenolics detection was carried out using (D) Ferric chloride test: An aliquot of the extract was treated with

5% FeCl₃ reagent and observed for the formation of a deep blue-black colour and (E) Lead acetate test: A fraction of the extract was treated with 10% lead acetate solution and observed for the formation of white precipitate.

Flavonoids were detected by using (F) aqueous sodium hydroxide test: A fraction of the extract was treated with 1N aqueous sodium hydroxide and observed for the formation of yellow-orange colouration and (G) Sulphuric acid test: A fraction of the extract was treated with concentrated sulphuric acid and observed for the formation of orange colour.

RESULTS:

Enzymatic antioxidants

Coleus vetiveroides, the plant used for the study, has been shown to have high enzymatic antioxidant content. Table 1 shows the enzymatic activities that were obtained. Catalase, peroxidase, and glutathione-S-transferase activity were found in abundance in the leaves. These three enzymatic antioxidants act as an innate defensive mechanism in plants, helping them to withstand oxidative damage⁽³²⁾.

Non-Enzymatic antioxidants

Non-enzymic, low-molecular-weight antioxidants also contribute to the antioxidant spectrum. Table 2 shows the amounts of non-enzymic antioxidants found in *Coleus vetiveroides* leaves. Ascorbic acid and alpha tocopherol were found in significant amounts. Total carotenoids, reduced glutathione, total phenols, flavonoids, and chlorophyll were all associated with the greatest amounts in the leaves.

Radical Scavenging activity

For radical scavenging activity, the extracted leaf samples yielded the following findings. Isopropanol and Ethyl acetate extracts of *Coleus vetiveroides* leaves had significant hydrogen peroxide scavenging capacity, followed by Hexane extract, and Isopropanol extracts of *Coleus vetiveroides* leaves had a strong hydroxyl radical scavenging ability, followed by Ethyl acetate and Hexane extracts. Figures 1 and 2 depict the findings of hydrogen peroxide and hydroxyl scavenging effects, respectively.

Chelating property of *Coleus vetiveroides*

Plants that have iron-chelating activity are the most effective at preventing lipid peroxidation, and so play an important role in medicine. *Coleus vetiveroides* extracts in Isopropanol and Ethyl acetate showed the most chelating activity, followed by Hexane extract. Figure 3 illustrates the chelating properties of three distinct extracts.

DNA protective action of *Coleus vetiveroides* against radical damage

Extracts of *Coleus vetiveroides* leaves in Hexane, Ethyl acetate, and Isopropanol showed potential in protecting DNA from free radicals created by hydrogen peroxide. The migratory pattern of goat liver DNA treated with hydrogen peroxide in the presence and absence of three distinct leaf extracts is represented in Figure 4.

The presence of alkaloids, flavonoids, and phenols was observed in the phytochemical analysis of *Coleus vetiveroides* leaves of different extracts.

DISCUSSION:

Antioxidants function effectively as radical scavengers protecting cells from degenerative disease. The human body endogenously releases antioxidants to neutralize the detrimental effects of free radicals. A constant supply of external sources of antioxidants maintains the balance and achieves significant effects. Peroxides and free radicals produced by disturbances in the normal redox state of cells damage cell components leading to cytotoxicity.

The objective of this study was to discover the antioxidant and scavenging activities of *Coleus vetiveroides* leaves which have been shown to have high antioxidant properties. Previous research on the roots of *coleus vetiveroides* exhibited significant antibacterial activity. Plants produce a large number of bioactive molecules making them rich sources of different types of medicines.

The leaves of *Coleus vetiveroides* showed a considerable activity of enzymic antioxidants; catalase, peroxidase, and glutathione-s-transferase. Catalase plays a significant role in providing antioxidant defenses to an organism. The catalase activity of *Plectranthus aromaticus* and *Catharanthus* under salinity was boosted after treatment with triazole⁽³³⁾. Paclobutrazol treatment improved it, even more, allowing it to effectively scavenge H₂O₂⁽³⁴⁾. In oxidatively challenged plants, Jaleel et al.⁽³⁵⁾ found that the H₂O₂ scavenging mechanism represented by CAT and APX is more essential than SOD in conferring tolerance.

Rhinaocanthus nasutus leaves have been reported to be the richest source of both enzymic and non-enzymic antioxidants among three underexploited medicinal plants⁽³⁶⁾. Glutathione-s-transferase is substantially more efficient on a molar basis than other

enzymes⁽³⁷⁾. According to Zimmermann and Zentgraf (2005)⁽³⁸⁾, the components of both the enzymic and non-enzymic antioxidant systems correlate well with oxidative stress during plant development and senescence.

The leaves of *Coleus vetiveroides* contain significant amounts of non-enzymic antioxidants. Many plants have been tested for non-enzymic antioxidants and the results have been published. In comparison to the leaf, stem, and roots of *Moringa oleifera*, total levels of ascorbic acid were found to be highest in the pods⁽³⁹⁾. These antioxidant activities were also observed to be dependent on the sample concentration⁽⁴⁰⁾. In a comparative study of a few plants, the highest α -tocopherol content was reported in *Sauropus androgynus* leaves, followed by *Citrus hystrix* leaves etc.,⁽⁴¹⁾

Carotenoids are a good non-enzymatic antioxidant that protects against free radical damage by quenching singlet oxygen. According to a study, lycopene levels in the leaves and β -Carotene levels in the tubers of *Planctranthus aromaticus* were much higher than in other parts of the plant.

Antioxidant, antimutagenic, and anticancer effects are all prevalent in phenols. As a result, their presence in *Coleus vetiveroides* leaves is credited with the plant's remarkable antioxidant qualities. In the *Plectranthus aromaticus* plant, the total phenol concentration was found to be substantially higher in the tubers than in the roots and stem⁽⁴²⁾

Amorphophallus commutatus, for example, was shown to have a significant amount of reduced glutathione in developed leaves, followed by juvenile leaves⁽⁴³⁾. The *Coleus vetiveroides* leaves show a high level of reduced glutathione. According to Karthika et al., 2014⁽⁴⁴⁾, the chlorophyll concentration of healthy and tender papaya leaves was higher than that of dried leaves. Our findings support the findings of earlier research and imply that *Coleus vetiveroides* leaves are a rich source of both enzymic and non-enzymic antioxidants.

Although hydrogen peroxide is not particularly reactive, it can occasionally be hazardous to cells because it can produce hydroxyl radicals in the cells. By reducing unsaturated bonds in biomolecules, the most physiologically reactive hydroxyl radical produces hydroxylation (Lipinshi *et al.*, 2011)⁽⁴⁵⁾. Thus, eliminating H₂O₂ from cells or dietary systems is critical for antioxidant defense.

The water and ethanol extracts of *Crataegus monogyna* exhibited hydrogen peroxide scavenging

ability⁽⁴⁶⁾. All of the extracts of *Pouzolzia zeylanica* leaves had good hydroxyl radical scavenging activity, Cold ethyl extract showed the most effective hydroxyl radical scavenger⁽⁴⁷⁾. About the above literature, the Hexane and Ethyl acetate extracts of *Coleus vetiveroides* leaves exhibited strong radical scavenging effects against Hydrogen peroxide and hydroxyl radicals.

Plants that have iron-chelating activity are the most effective at preventing lipid peroxidation and thus play an important role in medicine. Isopropanol and Ethyl acetate extracts of *Coleus vetiveroides* leaves exhibited high levels of the chelating property followed by Hexane extract. In the study of *Coriander sativum* and *Petroselinum crispum* (stem and leaves) Iron chelating activity of *Coriander sativum* extracts were reported to be higher than *Petroselinum crispum* plant⁽⁴⁸⁾.

The rhizome extract of *Dioscorea alata* possessed radical scavenging activity and showed a protective effect on calf thymus DNA and plasmid DNA as evaluated by EtBr (Wang *et al.*, 2004)⁽⁴⁹⁾. The alcohol: water (1:1) extract of curry leaves (*Murraya koenigii* L.) showed the highest antioxidant as reflected by the

inhibition of ferrous sulphate:ascorbate-induced fragmentation and sugar oxidation of calf thymus DNA (Ningappa *et al.*, 2008)⁽⁵⁰⁾. The leaf extracts of *Coleus vetiveroides* showed potential in protecting DNA from hydrogen peroxide-induced radicals.

Studies with leaves of *Adathoda vasica* and *Carissa carandas* linn plants⁽⁵¹⁻⁵⁴⁾, along with *Coleus vetiveroides*, revealed high quantities of enzymatic and non-enzymatic antioxidants, as well as free radical scavenging activity against hydroxyl and hydrogen peroxide radicals in the leaf extracts. Both plants have a high level of DNA protection against hydrogen peroxide radicals.

CONCLUSION:

The medicinal efficacy of the *Coleus* genus, which is widely utilized in Indian medicine, has been proven by modern testing and evaluation in a variety of illness states. These findings make this indigenous medication a promising candidate for bioprospecting and therapeutic development for wound, microbial infection, and liver disorders, among other conditions. The medical benefits of these plants, as well as a variety of research opportunities, are yet relatively unexplored aspects of their function.

Table 1: Enzymatic antioxidant levels in the leaves of *Coleus vetiveroides*

Parameters	Amount of enzymatic antioxidants present in <i>Coleus vetiveroides</i> leaf
CAT (U [*] /g)	17.70 ± 0.196
POD (U [#] /g)	1.548 ± 0.062
GST(U ^{\$} /g)	0.16 ± 5.88

The values are Mean ± SD of triplicates

*1 Unit = Amount of enzyme required to decrease the absorbance at 240 nm by 0.05units/minute


1 Unit = Change in absorbance at 430 nm /minute

\$1 Unit = nmoles of CDNB conjugated / minute

Table-2: Levels of Non-enzymatic antioxidants in the *Coleus vetiveroides* leaves

Parameters	Non-enzymatic antioxidant levels <i>Coleus vetiveroides</i> leaves
Ascorbic acid	4.476 ± 0.025 mg/gram
Tocopherol	227.3 ± 1.12 µg/gram
Total carotenoides	120.6 ± 0.763 mg/gram
Total lycopene	2.34 ± 0.0472 mg/gram
Reduced glutathione	102.5 ± 0.05 nM/gram
Total phenol	1.44 ± 0.040 mg/gram
Total chlorophyll	103.5 ± 1.01 mg/gram

Values are mean ± SD of triplets

**Figure 1: Hydrogen peroxide scavenging effect of *Coleus vetiveroides* leaf extracts**




Figure 2: Hydroxyl radical scavenging effect of *Coleus vetiveroides* leaf extracts




Figure 3: Chelating property of *Coleus vetiveroides* leaf extracts




Figure 4: Migration pattern of goat liver DNA treated with H₂O₂ with and without the leaf extracts (Starting from left side of the figure) Lane1: DNA + H₂O₂, Lane 2: DNA, Lane 3: DNA + Hexane extract, Lane 4: DNA + Ethyl acetate extract, Lane 5: DNA + Isopropanol extract, Lane 6: DNA + H₂O₂ + Hexane extract, Lane 7: DNA + H₂O₂ + Ethyl acetate extract, Lane 8: DNA + H₂O₂ + Isopropanol extract

REFERENCES:

1. Paargavi SA, Aanandaroopan A, Priya TS, Merish S (2015) Review of Medicinal Plants Common to Traditional Chinese and Indian Medicines with Relevance to Non-Communicable Diseases. *Current Traditional Medicine*, 1:116-121
2. Frie, B. and Carr, C. (1993) vitamin C in human health and disease. *J. Nutri Med.*, 8(13):14-16.
3. Silva and Fernandes A. (2010) Biological properties of medicinal plants: a review of their antimicrobial activity, *The Journal of Venomous Animals and Toxins including tropical Diseases*, 16(3), 402-413.
4. Sarma, A.S, Mallick, A.R. and Ghosh, A.K. (2010) Free radicals and their role in different clinical conditions: An overview, *International Journal of Pharma Sciences and Research*, 1(3):185-192.
5. Saranraj, P. and Sivasakthi, S. (2014) Medicinal Plants and its Antimicrobial Properties: A Review, *Global Journal of Pharmacology* 8 (3): 316-327.
6. Sindhe, A., Guru, J., and Naik, P., (2012) Effect of free radicals and antioxidants on oxidative stress: A review, *Journal of Dental and Allied science*; 1(2):63-66.
7. Hemalatha, M., Thirumalai, T., Saranya, R., Elumalai, E.K., and David, E. (2013) A Review On Antimicrobial Efficacy Of Some Traditional Medicinal Plants In Tamilnadu, *Journal Of Acute Disease*, 99-105.
8. Pryor, W.A. (2000) Dietary reference intake for vitamin C, vitamin E, selenium, and carotenoids. *J. of Free Radic Chem.*, 9(12):201-205.
9. Dhalla, N.S. Gupta, K.C. Sastry, M.S. and Malhotra (2000) The effect of stobadine on the copper induced peroxidation and its effects on cardiovascular diseases., *J. Rev.* 3.1152-1153.
10. Rahaman, I. The role of oxidative stress in pathogenesis and treatment of inflammatory bowel diseases. *J. Naunyn Schmiedebergs Arch pharmacol.*, 387(7):605-620.
11. Rajmohan, S.B. and Gupta, N.T. (2008) Transduction with antioxidant enzyme catalase protects human T cells against oxidative stress. *J. Immunol.*, 181(12):8382-8390.
12. Finkel, T. (1998) Oxygen radicals and signaling. *Current Opinion in Cell Biology*, 10, 248-253
13. Redza-Dutordoir M, Averill-Bates DA. Activation of apoptosis signalling pathways by reactive oxygen species. *Biochim Biophys Acta*. 2016 Dec;1863(12):2977-2992.
14. Pandey, K. B., & Rizvi, S. I. (2009). Plant polyphenols as dietary antioxidants in human health and disease. *Oxidative medicine and cellular longevity*, 2(5), 270-278.
15. Naz, R., & Bano, A. (2013). Phytochemical screening, antioxidants and antimicrobial potential of *Lantana camara* in different solvents. *Asian Pacific Journal of Tropical Disease*, 3(6), 480-486.

16. Saraswathy A, Lavanya MS (2013) Chemical Composition and Antibacterial Activity of the Essential Oil from the Roots of *Coleus vettiveroides* K.C. Jacob. *International Journal of Pharmacy and Biological Sciences*, 3(3)212-217
17. Luck, H. (1974), In: *Methods in enzymatic analysis.*, Ber. Acad. N. Y., 885.
18. Reddy, K. P, Subhani, S. M., Khan, P. A. and Kumar (1995), Effect of light and benzyl adenine and dark- treated graving rice (*Oryza sativa*) leaves II. Changes in peroxidases activity, *Plant. Cell. Physiol.*, 26, 987- 994.
19. Habig, W. H, Pabst, M. J. and Jokoby, W. (1974), The first enzymatic step in mercapturic acid III formation, *J. Biol. Chem.*, 249, 7130- 7139.
20. Roe, J. H. and Keuther, C. A. (1943), The determination of ascorbic acid in whole blood and urine through 2, 4-dinitro phenyl hydrazine derivative of dehydro ascorbic acid, *J. Biol. Chem.*, 147, 399- 407.
21. Rosenberg, H.R. (1992), *Chemistry and physiology of the vitamins*, Intersci. Pub, N.Y., 452- 453
22. Zakaria, H., Simpson, K., Brown, P. R. and Krotulovic, A. (1979), Use of reversed phase HPLC analysis for the determination of provitamin A carotenes in tomatoes, *J. Chrom.*, 176, 109- 117.
23. Mallick, C. P. and Singh, M. B. (1980), *Plant enzymology and plant histoenzymology*, Kalaya. Pub., New. Delhi., 281- 286.
24. Cameron, G.R., Mitton, R.F. and Allan, J.W. (1943), *Lancet*, 179.
25. Moron, M.S., Depierre, J.N. and Mannervik, Y.C. (1979), Levels of glutathione, glutathione reductase and glutathione S-transferase activities in rat lung and liver, *Biochem. Biophys. Acta.*, 582, 67-68.
26. Witham, F.H., Blaydes, B.F. and Devlin, R.M. (1971), *Experiments in plant physiology*, Van Nostrand-Reinhold. 245.
27. Ruch, R.J., Cheng, S. and Klaunig, J.E. (1989) Prevention of cytotoxicity and inhibition of intercellular communication by antioxidant catechins isolated from Chinese green tea, *Carcinogenesis*, 10, 1003-1008.
28. Elizabeth, K. and Rao, H.N.A. (1990) Oxygen radical scavenging activity of curcumin, *Int. J. Pharm.*, 58, 237-240.
29. Brown, D.J. and Goodman, J. (1998), A review of vitamin A, C and E and their realationship to cardiovascular disease, *Clin. Excell. Nurse Pract.*, 2, 10-12.
30. Marmur, J. (1961) A procedure for the isolation of deoxyribonucleic acid from microorganisms, *journal of Molecular Biology*, 3(2), 208-218.
31. Khandelwal, K.R. (2002) *Practical pharmacognosy-techniques and experiments*, IX edition, NiraliPrakashan Publishers, Pune, pp. 149-157.
32. Verma, S. and Dubey, R.S. (2003) Lead Toxicity Induces Lipid Peroxidation and Alters the Activities of Antioxidant Enzymes in Growing Rice Plants. *Plant Science*, 164, 645-655
33. Gomathinayagam M, Jaleel CA, Lakshmanan GM, Panneerselvam R. (2007) Changes in carbohydrate metabolism by triazole growth regulators in cassava (*Manihot esculenta* Crantz); effects on tuber production and quality. *C R Biol.* 330(9):644-55.
34. Sankhla, N., Upadhyaya, A., Davis, T.D. et al. (1992) Hydrogen peroxide-scavenging enzymes and antioxidants in *Echinochloa frumentacea* as affected by triazole growth regulators. *Plant Growth Regulations*, 11, 441–443
35. Jaleel, C.A., Riadh, K., Gopi, R. et al. (2009) Antioxidant defense responses: physiological plasticity in higher plants under abiotic constraints. *Acta Physiol Plant* 31, 427–436
36. Nirmaladevi R, Padma PR (2008) A study on the antioxidant potential of three under-exploited plants of medicinal value. *Plant Archives.* 8(1):339-341
37. Starlin T, Gopal Krishna VR (2013) Enzymatic and non-enzymatic antioxidant properties of *Tylophora pauciflora* Wight and Arn.—An in vitro study. *Asian Journal of Pharmaceutical and Clinical Research.* 6(4): 68-71
38. Zimmermann P, Zentgraf U (2005) The correlation between oxidative stress and leaf senescence during plant development. *Cellular and Molecular Biology Letters.*;10(3):515-34
39. Shih, M. C., Chang, C. M., Kang, S. M., & Tsai, M. L. (2011). Effect of different parts (leaf, stem and stalk) and seasons (summer and winter) on the chemical compositions and antioxidant activity of *Moringa oleifera*. *International journal of molecular sciences*, 12(9), 6077–6088
40. Serteser A, Kargioğlu M, Gök V, Bağcı Y, Özcan MM, Arslan D (2008) Determination of antioxidant effects of some plant species wild growing in Turkey. *International Journal of Food Sciences and Nutrition.* 59(7-8):643-51
41. Ching LS, Mohamed S. (2001) Alpha-tocopherol content in 62 edible tropical plants. *Journal of Agricultural and Food Chemistry.* 49(6):3101-5.

42. Velioglu YS, Mazza G, Gao L, Oomah BD (1998) Antioxidant Activity and Total Phenolics in Selected Fruits, Vegetables, and Grain Products. *Journal of Agricultural and Food Chemistry* 46(10):4113-4117
43. Krishna RK, Krishnakumari S, Chandrakala S (2012) Evaluation of antioxidant properties of different parts of *Amorphophallus commutatus*, an endemic aroid of Western Ghats, South India. *International Journal of Pharma and Bio Sciences* 3(3):B443-B455
44. Karthika P, Hema Karthika R, Jayachandran N, Joice Rathinam E (2014) Estimation of Chlorophyll Content in Papaya Leaf using Mathematical Operations. *International Journal of Computer Applications* 6-11
45. Lipinski B (2011) Hydroxyl Radical and Its Scavengers in Health and Disease, *Oxidative Medicine and Cellular Longevity* (5):809696
46. Keser S, Celik S, Turkoglu S, Yilmaaz O (2012) Hydrogen Peroxide Radical Scavenging and Total Antioxidant Activity of Hawthorn, *Chemistry Journal* 2(1):9-11
47. Peiyuan Li, Lini Huo, Wei Su, Rumei Lu (2011) Free radical-scavenging capacity, antioxidant activity and phenolic content of *Pouzolzia zeylanica*, *Journal of the Serbian Chemical Society* 76(5):709-717
48. Peter Y.Y. Wong, David D Kitts (2006) Studies on the dual antioxidant and antibacterial properties of Parsley (*Petroselinum crispum*) and Cilantro (*Coriandrum sativum*) Extracts, *Food Chemistry* 97(3):505-515
49. Wang TS, Liang SJ, Kueii Li C, Yie Liu S (2004) Protective Effect of Water Yam (*Dioscorea alata* L.) Extract on the Copper-driven Fenton Reaction and X-ray Induced DNA Damage in vitro, *Phytotherapy Research* 18(4):325-8
50. Ningappa MB, Ramdas D, Srinivas L (2008) Antioxidant and free radical Scavenging activities of polyphenol-enriched curry leaf (*Murrayakoenigii* L.) extracts, *Food Chemistry* 106(2)
51. Ramya N, Nagarathna A, Kiruthika B (2020) Antioxidant Potential of *Adathoda vacia* leaves, *World Journal of Pharmacy and Pharmaceutical Science* 9(4): 1426-34
52. Ramya N, Nagarathna A, Kiruthika B (2020) Phytochemical Screening and Protective Effect of *Adhatoda Vasica* Leaf Extracts against Free Radical Damage, *Indo American Journal of Pharmaceutical Sciences* 7(9): 691-7
53. Swathi KA, Nagarathna A, Kiruthika B (2020) Antioxidant analysis of *Carissa carandas* linn plant leaves, , *Indo American Journal of Pharmaceutical Sciences* 7(4):980-985
54. Swathi KA, Nagarathna A, Kiruthika B (2020) Radical Scavenging, Preliminary Phytochemical Screening and DNA Protective effect of *Carissa Carandas* Linn Plant Leaves, *International Journal of Pharma and Bio Sciences* 12(1):1-8