



Contents lists available at:  
<https://journals.irapa.org/index.php/BCS/issue/view/7>

## Biomedicine and Chemical Sciences

Journal homepage: <https://journals.irapa.org/index.php/BCS>



# Synthesis of Some New 1,3,4- Oxadiazole Derivatives and Thiazolidine Derived from Cysteine and Evaluation their Anticancer (MCF7) Activity

Ahmed A. Majed<sup>a\*</sup>, Dawood S. Abid<sup>b</sup>

<sup>a</sup> Education Directorate of Thi-Qar, Ministry of Education, Thi-Qar – Iraq

<sup>b</sup> Department of Chemistry, College of Education, University of Basrah, Basrah – Iraq

### ARTICLE INFO

#### Article history:

Received on: December 21, 2021

Revised on: January 05, 2021

Accepted on: February 5, 2022

Published on: April 01, 2022

#### Keywords:

Anticancer MCF-7

L-cysteine

Oxadiazole

Thiazolidines

### ABSTRACT

The study focused on the preparation of oxadiazole derivatives containing thiazolidine. Thiazolidine firstly was prepared from the reaction of benzaldehyde with L-cysteine with a good yield and then it was reacted with acetic anhydride to prepare acetyl thiazolidine, then with ethanol in the presence of H<sub>2</sub>SO<sub>4</sub>, then steps were taken to prepare a thiazolidine hydrazide (A<sub>3</sub>), which was reacted with aromatic carboxylic acid in presence POCl<sub>3</sub> or carbon disulphide and base KOH to obtain oxadiazole derivatives A<sub>4-8</sub>. These compounds characterized using FT-IR, NMR and Mass (EI) were diagnosed and the synthesized compound were validated. The activity of oxadiazole derivatives were studied against breast cancer cells, the two compound A<sub>8</sub>, A<sub>4</sub> showed good activity against the cells as for the compounds A<sub>5</sub>, A<sub>6</sub>, A<sub>7</sub> it was showed little activity against these cells and the value IC<sub>50</sub> was calculated.

Copyright © 2022 Biomedicine and Chemical Sciences. Published by International Research and Publishing Academy – Pakistan, Co-published by Al-Furat Al-Awsat Technical University – Iraq. This is an open access article licensed under CC BY:

(<https://creativecommons.org/licenses/by/4.0>)

## 1. Introduction

Heterocyclic compounds play an effective role in many fields, where the compound of oxadiazole, triazole, thiazole (Guessas, et al., 2007) and thiazolidine have received great attention in medicinal, pharmaceutical and industrial chemistry (Leod & Ames, 1987). Thiazolidine is an aliphatic cyclic compound found in natural produced (Busacca, et al., 1996), foodstuffs and antibiotics, which is produced from the interaction of cysteine or cysteamine (penicillamine) with aliphatic and aromatic aldehydes or ketones (Tawfiq, 2016), thiazolidine plays a distinctive character in natural products similar to antibiotics including penicillin and cevasporine, one of the most important thiazolidine derivatives is thiazolidine-4-carboxylic acid (Subr & Ulbrich, 2006), which is considered the basis for the basis for the production of thiazolidine drugs (El-Sharkawy, 2011), as it has against

cancer, bacteria (Pavin, et al., 2011), fungi, inflammation and diabetes (Da Silva, et al., 2015). Oxadiazole is considered an effective cyclic aromatic compound and is include in the composition of many derivatives of biologically active drugs (Şahin, et al., 2002), which include anti-inflammatory (Guessas, et al., 2007), antifungal, HIV (Wang, et al., 2012) creating pharmaceutical scaffolds, antimicrobials (Bondock, et al., 2012), anticancer, antihypertensive and cardio-tonic. The most famous method used to prepare oxadiazole is a reaction of hydrazide with carboxylic acids in the presence of POCl<sub>3</sub> (Gilani, et al., 2010) or it is condensation with carbon disulfide and base (KOH) (Zhang, et al., 2013). The aim of the study was to prepare and study thiazolidine and oxadiazole ring as anti-cancer of type MCF-7.

## 2. Materials and Methods

Melting points, FT-IR spectra, and <sup>1</sup>H, <sup>13</sup>C-NMR spectrum were recorded on Bruker 400MHz in DMSO and CDCl<sub>3</sub> solvent. For abbreviation used: s :singlet, d:doublet, t: triplet, dd :doublet of doublets and m: multiblasty. Mass(EI) spectra and spectrophotometry.

2.1. Synthesis the Compound 2-Phenyl thiazolidine-4-carboxylic acid (A).

\*Corresponding author: Ahmed A. Majed, Education Directorate of Thi-Qar, Ministry of Education, Thi-Qar – Iraq

E-mail: [eduppg.ahmed.majed@uobasrah.edu.iq](mailto:eduppg.ahmed.majed@uobasrah.edu.iq)


#### How to cite:

Majed, A. A. (2022). Synthesis of some 1,3,4- oxadiazole, thiazolidine derivatives resulting from cysteine and their study as an anticancer of breast cancer. Biomedicine and Chemical Sciences, 1(2), 83-87.

DOI: <https://doi.org/10.48112/bcs.v1i2.109>

A mix 0.01 mole (1.02 ml) of Benzaldehyde and 0.01 mole (1.21 g) of L-cysteine were mixed in (10 ml) water and (50 ml) ethanol, it was stirred at room temperature for (12 h), the resulting white precipitate formed was filtered, collected, washed, with ether, dried and recrystallized with ethanol; water (1:3) (Nawar, et al., 2020). Structure was showed in Scheme1. This product as a mix of diastereomers, cis-(2R,4R) and trans-(2S,4R) as shown in Equation 1. **(2R, 4R) 2-Phenyl thiazolidine-4-carboxylic acid (A)** **61%Cis Isomer (Trans39%)**

Yield: 80%, mp:159-161°C. FT-IR(KBr): 3100-2700(ZwitterionNH<sub>2</sub><sup>+</sup>), 1573s (COO<sup>-</sup>). <sup>1</sup>H NMR(400 MHz, CDCl<sub>3</sub>): 82.9 dd, 1H, J=10.28, 8.7 Hz (2.99dd, J=10.51, 5.58Hz) (**H5a**), 83.24dd, 1H, J=10.33, 7.25Hz (3.14 dd, J=10.51, 7.51Hz) (**H5b**), 83.77 dd, 1H, J=8.36, 7.32Hz (3.96t, J=5.91) (**H4**), 85.29 s, 1H (5.56s) (**H2**), 7.00-7.27(5H) (**Har**).



**Equation 1:** Cis and Trans diastereomers

## 2.2. Synthesis the Compound 3-Acetyl-2-Phenyl thiazolidine-4-carboxylic acid (A<sub>1</sub>).

0.01 mole (2.09 g) was dissolved of (A) in (40 ml) of (6 %) Na<sub>2</sub>CO<sub>3</sub> cool in ice bath, followed by the addition 0.04 mole (3.77 ml) of acetic anhydride in the form drops. The mixture was stirred for 1 h and it is acidified by HCl, the mixture was extracted using with ethyl acetate and evaporation leaving the solid white, then washed by water, dried and recrystallized with ethyl alcohol (Majed & Abid, 2015) to give (A<sub>2</sub>). **(2R, 4R)-N-Acetyl-2-Phenyl thiazolidine-4-carboxylic acid (A<sub>1</sub>) 93%Cis Isomer (Trans7%)**

Yield:85%, mp:147-149°C. FT-IR(KBr): 3327s (OH), 1731m, 1680m(C=O). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): 81.98s, 3H (2.19s) (**H7**), 3.32dd, 1H, J=12.12, 6.66Hz (3.42d, J=6.4Hz) (**H5a**), 83.36dd, 1H, J=12.07, 6.98Hz (3.42, d, J=6.4Hz) (**H5b**), 85.06t, 1H, J=6.8Hz (4.8s) (**H4**), 86.05s, 1H (6.39s) (**H2**), 7.27-7.35(5H) (**Har**), 811.13s, 1H (11.13s) (**H6**)(OH).

## 2.3. Synthesis the Compound Ethyl 3-Acetyl-2-Phenyl thiazolidine-4-carboxylate(A<sub>2</sub>).

A 0.01 mole of (A<sub>1</sub>) was mixed with 30ml ethanol and drops of concentrated sulphuric acid. The mixture was refluxed for 6 h, then filtered and evaporated to give oily yellow product (Tawfiq, 2016).

## 2.4. Synthesis the 3-Acetyl-2-Phenyl thiazolidine-4-carbohydrazone(A<sub>3</sub>).

Compound A<sub>2</sub> was dissolved in (30 ml) ethanol then (5 ml) of (80%) hydrazine hydrate added. The solutions were refluxed for 24 h, and the solvent evaporated to leave a white precipitate which was recrystallized with water: ethanol (2:8) (Al-Badrany, et al., 2019), to give (A<sub>3</sub>) m.p=123-125°C.

## 2.5. 3-Acetyl-2-Phenyl thiazolidine-4-carbohydrazone (A<sub>3</sub>) 75%Trans Isomer (Cis25%)


mp:122-125°C FT-IR(KBr): 3317, 3182s (NH<sub>2</sub>), 3226w (NH), 1700m, 1662m (C=O). <sup>1</sup>HNMR (400MHz, DMSO-d<sub>6</sub>): 81.86s, 3H (2.08s) (**H7**), 3.1 dd, 1H, J=10.4Hz (3.3m,) (**H5a**), 83.46 m, 1H, J=4 Hz (3.37m) (**H5b**), 84.21 m, 1H, J=8, 4Hz (4.21s) (**H4**), 84.78 d, 2H, J=4 (4.79s) (**H5**), 85.3 d, 1H, J=4 (5.3) (**H6**) 86.43 s, 1H (6.22s) (**H2**) 7.22-7.71(5H) (**Har**). Mass(EI): 265.1 M.Wt, 185.1 peas beak.

## 2.6. General method for the prepare 2-(2-Phenyl thiazolidine- 4-yl)-5-(aryl)-1,3,4-Oxadiazole (A<sub>4-7</sub>)

Totally compounds are synthesize the same procedure (Husain, et al., 2012). A solution 0.01 mole of compound (A<sub>3</sub>) and 0.01 mole of aromatic carboxylic acid in POCl<sub>3</sub> (5 ml) was reflux to (90 °C) for 6-8 h on water bath. The reaction was followed using the TLC (ethanol ethyl: acetate 2:8), then evaporated, poured into crushed ice, neutralized using NaHCO<sub>3</sub> (10 %) and left to form a precipitate, the result was filtered, washed many step with water, dried and recrystallized from the mixture chloroform: Hexane or ethanol: water. Structures and symbols of prepared compounds are illustrated in Scheme2. Trans and Cis diastereomers of oxadiazole showed in Equation 2 (Jafari, et al., 2017).

## 2.7. 2-((2R,4S)-N-Acetyl-2-phenylthiazolidine-4-yl)-5-(8-hiazoli-4-yl)-1,3,4-oxadiazole (A<sub>4</sub>) 85%Trans Isomer (Cis15%)

Yield:20%, mp:187-189°C, FT-IR(KBr disk): 1701s (C=O), 1627m (C=N). <sup>1</sup>HNMR(400 MHz, DMSO-d<sub>6</sub>): 81.81s, 3H (2.08s) (**H7**), 3.13t, 1H, J=8Hz (3.32m, J=4Hz) (**H5a**), 83.46m, 1H, J=8 Hz (3.40m, J=4Hz) (**H5b**), 84.46t, 1H, J=8Hz (4.45s) (**H4**), 86.36 s, 1H (6.16s) (**H2**), 7.30-8.24(9H) (**Har**). <sup>13</sup>CNMR: 8 22.42(**CH**<sub>3</sub>), 30.44(**CH**<sub>2</sub>)**C**<sub>5</sub>, 64.9(**CH**)**C**<sub>4</sub>, 73.3(**CH**) **C**<sub>2</sub>, 121-150 (**CAr**), 159, 163 (**2C=N**), 170.55 (**C=O**) Mass(EI): 352.2 M.Wt, 180.2 peas beak.



**Equation 2:** Cis and Trans diastereomers of oxadiazole A4-7

## 2.8. 2-((2R,4S)-N-Acetyl-2-phenylthiazolidine-4-yl)-5-Phenyl-1,3,4-oxadiazole (A<sub>5</sub>) 73%Trans Isomer (Cis27%)

Yield:24%, mp:164-167°C). FT-IR(KBr): 1685m (C=O), 1598m (C=N). <sup>1</sup>H NMR(400 MHz, DMSO-d<sub>6</sub>): 81.83s, 3H (2.06s) (**H6**), 83.1dd, 1H, J=10.4Hz (3.3m) (**H5a**), 83.47 m, 1H, J=4Hz (3.41d) (**H5b**), 84.21 m, 1H, J=4Hz (4.21m) (**H4**), 86.42 s, 1H (6.2s) (**H2**), 7.23-7.81(10H) (**Har**). <sup>13</sup>CNMR: 8 23.37(**CH**<sub>3</sub>), 34.13(**CH**<sub>2</sub>) **C**<sub>5</sub>, 67.7(**CH**)**C**<sub>4</sub>, 74.15(**CH**)**C**<sub>2</sub>, 120-140 (**CAr**), 159, 162(**2C=N**), 170 (**C=O**). Mass(EI): 351.2 M.Wt, 180.2 peas beak.

## 2.9. 2-((2R,4S)-N-Acetyl-2-phenylthiazolidine-4-yl)-5-(4-nitrophenyl)-1,3,4-oxadiazole (A<sub>6</sub>) 91% Trans Isomer (Cis9%)

Yield:22%, mp:191-194°C, FT-IR(KBr disk): 1689s (C=O), 1635m (C=N), 1346, 1540 (NO<sub>2</sub>). <sup>1</sup>HNMR(400 MHz, DMSO-d<sub>6</sub>): 81.81s, 3H (1.78s) (**H7**), 3.1m, 1H, J=4Hz (3.3m) (**H5a**), 83.37m, 1H, J=8 Hz (3.7m) (**H5b**), 84.27m, 1H, J=8Hz

(4.27m) (**H4**),  $\delta$ 6.38 s, 1H (6.34s) (**H2**), 7.2-8.1(9H) (**Har**). Mass(EI): 396 M.Wt, 43.2 peas beak.

2.10. 2-((2R,4S)-N-Acetyl-2-phenylthiazolidine-4-yl)-5-(4-tolyl)-1,3,4-oxadiazole (**A7**) 78%Trans Isomer (Cis22%)

Yield:16%, mp:177-179°C, FT-IR(KBr disk): 1741s (C=O), 1627m (C=N).  $^1\text{H}$ NMR(400 MHz, DMSO- $d_6$ ):  $\delta$ 1.85s, 3H (2.05s) (**H7**), 3.13dd, 1H, J=8.4Hz (3.3m) (**H5a**),  $\delta$ 3.45m, 1H, J=8 Hz (3.38m) (**H5b**),  $\delta$ 4.28m, 1H, J=8Hz (4.4m) (**H4**),  $\delta$ 6.34 s, 1H (6.14s) (**H2**), 7.2-8.1(9H) (**Har**). Mass(EI): 362.2 M.Wt, 65.2 peas beak.

2.11. Synthesis the 2-(3-Acetyl-2-Phenyl thiazolidine)-5-mercapto-1,3, 4-oxadiazole (**A8**)

A mixture of 0.01mole (**A3**) in ethanol, 0.015mole KOH and (5 ml) CS<sub>2</sub> in 0 °C was refluxed for 8 h and followed using TLC(ethanol ethyl: acetate 2:8). The mixtures concentrated and poured in (100 ml) water ice, later acidified by 10 % HCl. The product was filtered, washed, dried (**Pitasse-Santos, et al., 2018**), recrystallized from ethyl alcohol to give (**A7**). M.p=163-165°C.

2.12. 2-((2R,4S)-N-Acetyl-2-phenylthiazolidine-4-yl)-5-mercapto-1,3,4-oxadiazole (**A8**) 55%Trans Isomer (Trans45%)

Yield:30%, mp:200-202°C. FT-IR(KBr): 3122w (NH), 2626w (SH), 1720 (C=O), 1612s (C=N).  $^1\text{H}$  NMR(400 MHz, DMSO- $d_6$ ):  $\delta$ 1.78s, 3H (1.96s) (**H7**),  $\delta$ 2.89t, 1H, J=12Hz (3.1m, J=4) (**H5a**),  $\delta$ 3.4m, 1H, J=4 Hz (3.34) (**H5b**),  $\delta$ 4.24t, 1H, J=4Hz (4.55t) (**H4**),  $\delta$ 6.42s, 1H (6.54s) (**H2**),  $\delta$ 6.2s, 1H (**SH**) 7.34-7.84(5H) (**Har**). Mass(EI):307.1M.Wt, 237.2 peas beak.

2.13. Anticancer Study

Cell growth and cell viability were quantified using the MTT (3-(4, 5-dimethylthiazol-2-yl)-2, 5-diphenyltetrazolium Bromide) assay. In brief, for monolayer culture, cells (MCF7) were digested with trypsin, harvested, adjusted to a density of  $1.4 \times 10^4$  cells/well and seeded to 96-well plates filled with 200  $\mu$ l fresh medium per well for 24 h. When cells formed a monolayer, they were treated with 500-15.62  $\mu$ g/ml of the compounds for 24 h at 37 °C in 5% CO<sub>2</sub>. At the end of the treatment (24 h), while the monolayer culture was left untouched in the original plate, the supernatant was removed and 200  $\mu$ l/well of MTT solution (0.5 mg/ml in phosphate-buffered saline (PBS)) was added and the plate was incubated at 37 °C for an additional 4 h. MTT solution (the supernatant of cells was removed and dimethyl sulfoxide was added (100  $\mu$ l per well)). Cells were incubated on a shaker at 37 °C until crystals were completely dissolved. Cell viability were quantified by measuring absorbance at 570 nm using an ELISA reader (Model wave xs2, Biotek, USA). The concentration of the compounds that resulted in 50% of cell death (IC<sub>50</sub>) was determined from respective dose-response curves.


MCF7 cancer cells were cultured in a dish containing 96 vacuoles filled with (200  $\mu$ l) of fresh medium for each holes for (24 h) in optimal conditions (37°C and atmosphere of carbon dioxide at 5% concentration and in a humid atmosphere). After that, serum media (10% FBS was + Antibiotics- penicillin and 100  $\mu$ g/ml streptomycin) and wash the cells twice with buffer phosphate solution (PBS), Fresh culture media containing dilute concentrations of the compounds to be tested were used, and the cells were then incubated for 48, 72 and 24 hours. Then the prepared

compounds were studied in a range of concentrations of 500-15.62 micrograms per liter. Gaps in the plate were analyzed for each concentration by adding 10  $\mu$ l of 5% of freshly prepared mtt (methanol) in PBS in 100 ml of DMSO, then stirring the plates to facilitate dissolution of the crystals. The absorbance was measured at a wavelength of 570 nm. A Biotek device was used and the half inhibitory concentration was calculated (Ozyazici, 2021) .

### 3. Results & Discussion

#### 3.1. Organic Synthesis

The amino acid (L-cysteine) attacks the carbonyl group of the benzaldehyde via the NH<sub>2</sub>, SH group to product thiazolidine-4-carboxylic acid (**A**), thiazolidines were easily attained in yields of 80-85%, this product as a mix of diastereomers, Cis-(2R,4R) and Trans-(2S,4R), the couldn't be separated. An equilibrium resulting from epimerization at C(2) occurs between two isomers. The percentage Cis/Trans was strongly dependent on a nature of the solvent. In CDCl<sub>3</sub> the major isomer was the Cis isomer while in DMSO- $d_6$  major isomer was trans isomer. Then protect the amine group will react with acetic anhydride to form 3-Acetyl-2-Phenylthiazolidine-4-carboxylic acid (**A1**), the compound (**A1**) reacts with ethyl alcohol in the presence H<sub>2</sub>SO<sub>4</sub> to synthesis (**A2**) (Mohammed, et al., 2021) and before it was condensed with (80%)hydrazine in a presence ethanol solvent to obtain 3-Acetyl-2-Phenylthiazolidine-4-carbohydrazide (**A3**), the compound (**A3**) goes in two ways, either it was reacted with carboxylic acid in the presence of phosphoryl chloride (POCl<sub>3</sub>) to obtain 1,3,4-oxadiazole derivatives (**A4-7**), or condensing it with carbon disulfide and potassium hydroxide KOH to prepare 2-(3-Acetyl-2-Phenylthiazolidine)-5-mercapto-1,3,4-Oxadiazole (**A8**), as shown in (Taha, et al., 2016) (scheme 1).



**Scheme 1.** Compounds prepared in the study **A1-8**


#### 3.2. Anti-cancer Activity(MCF-7)

The prepared oxadiazole derivatives were studied on breast cancer cells type MCF-7 using five concentrations of each compound, the inhibition activity and IC<sub>50</sub> value were calculated according to the following relationship:

$$\text{Viability \%} = \frac{\text{mean of OD sample}}{\text{mean of OD controle}} \times 100$$

The Values of the scavenging activity of the synthesis compounds ( $A_4$ ,  $A_5$ ,  $A_6$ ,  $A_7$  and  $A_8$ ) at concentration (500, 250, 125, 63, 31.5 and 15.5  $\mu\text{g/ml}$ ) were measured by the absorbance at 570nm. The inhibition activity was showed that all compounds possess an inhibitory ability at high concentration and gradually decreasing concentration, as compound  $A_8$  showed inhibitory activity towards cancer cells and this may be due to the compound's ability to form hydrogen bonds with the amino acid present in the active center of the cancer cells protein. Through the results  $\text{IC}_{50}$  shown in Fig.(1), it was observed that the arrangement of a studied compounds according to their ability to inhibit cancer cells is as follows (Ahsan, 2018, Ozyazici, et al., 2021):

$$A_8 > A_4 > A_5 > A_7 .$$



**Fig. 1.** The relationship of the percentage of inhibition with the concentration and calculate the ( $\text{IC}_{50}$ ) for the studied compounds

#### 4. Conclusions

Thiazolidines were obtained easily in yield of 80-85% from the react of L-cysteine and benzaldehyde under slightly conditions. 1,3,4-oxadiazoles were obtained difficult in low yields 15-30% from the condensate of thiazolidine-4-hydrazides and aromatic carboxylic acid in the presence of  $\text{POCl}_3$  or reacted with  $\text{CS}_2$  and  $\text{KOH}$  under a little conditions. From the product we conclude that the all compounds oxadiazole and thiazolidine have a biological activities against cancer cells MCF-7, therefore, we recommend the preparation of many thiazolidine and oxadiazole derivatives, and the possibility of studying them on different cancer cells and using them as treatment (drug).

#### References

Ahsan, M. J. (2018). Synthesis and cytotoxicity evaluation of [(2, 4-dichlorophenoxy) methyl]-5-aryl-1, 3, 4-oxadiazole/4 $\beta$  H  $\beta$ -1, 2, 4-triazole analogues. Turkish Journal of Chemistry, 42(5), 1334-1343.

Al-Badrany, K. A., Mohammed, A. S., & Alasadi, Y. K. (2019). Synthesis of some new 1, 3, 4-oxadiazole

compounds derived from 1H-imidazole and study their biological activity. Eurasian Journal of Biosciences, 13(1), 501-507.

Bondock, S., Adel, S., Etman, H. A., & Badria, F. A. (2012). Synthesis and antitumor evaluation of some new 1, 3, 4-oxadiazole-based heterocycles. European Journal of Medicinal Chemistry, 48, 192-199. <https://doi.org/10.1016/j.ejmech.2011.12.013>

Busacca, C. A., Dong, Y., & Spinelli, E. M. (1996). A one step synthesis of thiazolines from esters. Tetrahedron letters, 37(17), 2935-2938. [https://doi.org/10.1016/0040-4039\(96\)00451-0](https://doi.org/10.1016/0040-4039(96)00451-0)

Da Silva, T. L., Miolo, L. M. F., Sousa, F. S., Brod, L. M., Savegnago, L., & Schneider, P. H. (2015). New thioureas based on thiazolidines with antioxidant potential. Tetrahedron Letters, 56(48), 6674-6680. <https://doi.org/10.1016/j.tetlet.2015.10.037>

El-Sharkawy, K. A. (2011). Synthesis and Antimicrobial Activity of 2-Substituted-3-Acetyl Thiazolidine-4-Carbonyl-Amino acid Derivatives. Journal of Pharmaceutical Sciences and Research, 3(1), 1005-1014.

Gilani, S. J., Khan, S. A., & Siddiqui, N. (2010). Synthesis and pharmacological evaluation of condensed heterocyclic 6-substituted 1, 2, 4-triazolo-[3, 4-b]-1, 3, 4-thiadiazole and 1, 3, 4-oxadiazole derivatives of isoniazid. Bioorganic & Medicinal Chemistry Letters, 20(16), 4762-4765. <https://doi.org/10.1016/j.bmcl.2010.06.125>

Guessas, B., Othman, A. A., & Khiati, Z. (2007). Synthesis and antibacterial activity of 1, 3, 4-oxadiazole and 1, 2, 4-triazole derivatives of salicylic acid and its synthetic intermediates. South African Journal of Chemistry, 60(1), 20-24. <https://hdl.handle.net/10520/EJC24423>

Husain, A., Rashid, M., Mishra, R., Parveen, S., Shin, D. S., & Kumar, D. (2012). Benzimidazole bearing oxadiazole and triazolo-thiadiazoles nucleus: Design and synthesis as anticancer agents. Bioorganic & medicinal chemistry letters, 22(17), 5438-5444. <https://doi.org/10.1016/j.bmcl.2012.07.038>

Jafari, E., Mohammadi, T., Jahanian-Najafabadi, A., & Hassanzadeh, F. (2017). Synthesis and antimicrobial evaluation of some 2, 5 disubstituted 1, 3, 4-oxadiazole derivatives. Research in pharmaceutical sciences, 12(4), 330. <https://dx.doi.org/10.4103%2F1735-5362.212051>

Leod, G. M., & Ames, J. M. (1987). Effect of water on the production of cooked beef aroma compounds. Journal of Food Science, 52(1), 42-45. <https://doi.org/10.1111/j.1365-2621.1987.tb13968.x>

Majed, A. A., & Abid, D. S. (2015). Synthesis, Characterization and Biological Activity Study of some New Thiazolidine Derivatives. Basrah Journal of Science (C), 33(1), 101-117.

Majed, A. A., & Abid, D. S. (2021). Synthesis, Characterization and Biological Activity and Anti-Oxidant Study of Some New Thiazolidine Derivatives

- Containing Oxadiazole. *Annals of the Romanian Society for Cell Biology*, 11935-11945.
- Mohammed, M. A., Abid, D. S., & Sayyah, S. G. (2021). Preparation, Characterization of some Thiazolidine Derived from Penicillamine and their Antioxidants Activity. *Annals of the Romanian Society for Cell Biology*, 4653-4663.
- Nawar, F., Al-Asadi, R., & Abid, D. (2020). Synthesis, Antibacterial Activity and DFT Calculations of Some Thiazolidine-4-Carboxylic acid Derivatives and Their Complexes with Cu (II), Fe (II) and VO (II). *Egyptian Journal of Chemistry*, 63(1), 349-362. <https://dx.doi.org/10.21608/ejchem.2019.16096.1986>
- Özyazici, T., Şahin, F., & Akkoç, M. K. (2021). Synthesis, spectral characterization, and biological studies of 3, 5-disubstituted-1, 3, 4-oxadiazole-2 (3H)-thione derivatives. *Turkish Journal of Chemistry*, 45(3), 749-760.
- Pavin, N. F., Donato, F., Cibin, F. W., Jesse, C. R., Schneider, P. H., de Salles, H. D., ... & Savegnago, L. (2011). Antinociceptive and anti-hypernociceptive effects of Se-phenyl thiazolidine-4-carboselenoate in mice. *European journal of pharmacology*, 668(1-2), 169-176. <https://doi.org/10.1016/j.ejphar.2011.06.038>
- Pitasse-Santos, P., Sueth-Santiago, V., & Lima, M. E. (2018). 1, 2, 4-and 1, 3, 4-Oxadiazoles as Scaffolds in the Development of Antiparasitic Agents. *Journal of the Brazilian Chemical Society*, 29, 435-456. <https://doi.org/10.21577/0103-5053.20170208>
- Şahin, G., Palaska, E., Ekizoğlu, M., & Özalp, M. (2002). Synthesis and antimicrobial activity of some 1, 3, 4-oxadiazole derivatives. *Il Farmaco*, 57(7), 539-542. [https://doi.org/10.1016/S0014-827X\(02\)01245-4](https://doi.org/10.1016/S0014-827X(02)01245-4)
- Šubr, V., & Ulbrich, K. (2006). Synthesis and properties of new N-(2-hydroxypropyl) methacrylamide copolymers containing thiazolidine-2-thione reactive groups. *Reactive and Functional Polymers*, 66(12), 1525-1538. <https://doi.org/10.1016/j.reactfunctpolym.2006.05.002>
- Taha, M., Ismail, N. H., Imran, S., Wadood, A., Rahim, F., Saad, S. M., ... & Nasir, A. (2016). Synthesis, molecular docking and α-glucosidase inhibition of 5-aryl-2-(6'-nitrobenzofuran-2'-yl)-1, 3, 4-oxadiazoles. *Bioorganic chemistry*, 66, 117-123. <https://doi.org/10.1016/j.bioorg.2016.04.006>
- Tawfiq, M. T. (2016). Synthesis and Characterization of Some New 4-Oxothiazolidine-2-Carboxylic Acid Derivatives With the Evaluation of Their Biological Activity. *Diyala J. For PureSci*, 12, 43-64.
- Wang, Z., Wang, M., Yao, X., Li, Y., Qiao, W., Geng, Y., ... & Wang, Q. (2012). Hydroxyl may not be indispensable for raltegravir: Design, synthesis and SAR studies of raltegravir derivatives as HIV-1 inhibitors. *European journal of medicinal chemistry*, 50, 361-369. <https://doi.org/10.1016/j.ejmech.2012.02.015>
- Zhang, M. Z., Mulholland, N., Beattie, D., Irwin, D., Gu, Y. C., Chen, Q., ... & Clough, J. (2013). Synthesis and antifungal activity of 3-(1, 3, 4-oxadiazol-5-yl)-indoles and 3-(1, 3, 4-oxadiazol-5-yl) methyl-indoles. *European Journal of Medicinal Chemistry*, 63, 22-32. <https://doi.org/10.1016/j.ejmech.2013.01.038>