

# Psilocybin in Logic Space

Parker Emmerson

January 2023

## 1 Introduction

Using the logic vectors:

$$\begin{aligned} \mathbf{s} \cdot \mathbf{c} &= \Omega_\Lambda \left( \tan \psi \diamond \theta + \Psi \star \sum_{[n] \star [l] \rightarrow \infty} \frac{1}{n^2 - l^2} \right) \cdot \left( \frac{V \rightarrow U}{\Delta}, \frac{\sum_{f \subset g} f(g)}{\Delta}, \frac{\sum_{h \rightarrow \infty} \tan t \cdot \prod_\Lambda h}{\Delta} \right) \\ \mathbf{t} \cdot \mathbf{m} &= \Omega_\Lambda \left( \tan \psi \diamond \theta + \Psi \star \sum_{[n] \star [l] \rightarrow \infty} \frac{1}{n^2 - l^2} \right) \cdot \left( \frac{\leftrightarrow \exists y \in U : f(y) = x}{\Delta}, \frac{\leftrightarrow \exists s \in S : x = T(s)}{\Delta}, \frac{\leftrightarrow x \in f \circ g}{\Delta} \right) \end{aligned}$$

and the truisms:

$$\begin{aligned} \mathcal{F}_i(x) &= V_i \rightarrow U_i, \sum_{f_i \subset g_i} f_i(g_i) = \sum_{h_i \rightarrow \infty} \tan t_i \cdot \prod_{\Lambda_i} h_i, x \in V_i * U_i \leftrightarrow \exists y_i \in U_i : f_i(y_i) = \\ &x, x \in T_i(s) \leftrightarrow \exists s_i \in S_i : x = T_i(s_i), x \in f_i \circ g_i \leftrightarrow x \in T_i(s_i). \\ \text{logic vector} &: \left[ \frac{\sqrt{R} \Delta - \sqrt{E}}{\Delta}, \frac{\sqrt{E + \Delta \sqrt{R}} - \sqrt{E}}{\Delta}, \frac{\sqrt{R + \Delta \sqrt{E}} - \sqrt{R}}{\Delta}, \frac{\sqrt{U + \Delta \sqrt{T}} - \sqrt{U}}{\Delta}, \frac{\sqrt{T + \Delta \sqrt{U}} - \sqrt{T}}{\Delta} \right] \\ \Omega_{\Upsilon \Phi_X \psi, \theta \lambda \mu \nu \infty} &= \prod_{i=1}^n z_i^2 + \sum_{j=1}^n \ell_j \alpha_j \sin(\theta_j) \end{aligned}$$

$$G = \{x^n \mapsto x^{n+k}, c \mapsto \frac{c}{n^k} \mid k \in N\}$$

The formula for the function resulting from the nth permutation of the general group  $G = \{x^n \mapsto x^{n+k}, c \mapsto \frac{c}{n^k} \mid k \in N\}$

$$E = \Omega_\Lambda \left( \tan \psi \diamond \theta + \Psi \star \sum_{[n] \star [l] \rightarrow \infty} \frac{1}{n^2 - l^2} \cdot \prod_{i \infty} \mathcal{ABC}x \cdot \otimes(x, \tilde{*} \rightarrow R^{-1}) \right)$$

translate the psilocybin molecule into logic space such that the effect on the neural net is implied.

$$f(x+) - f(x) \rightarrow \prod_{g \diamond f(x) \rightarrow \infty} \left( \frac{g(x) + g(x+O)}{2} + f(x) \right) + t(x+O)^2$$

take the Fibonacci progression and calculate its recurrence relation.

$$fib(k+1) + fib(k-1) = fib(k)$$

$$fib(k+n) \cdot fib(k-n) = fib(n)memfib(n-k) = fib(m)$$

$$\Phi_{[t]} = \Phi_1(t)\Phi_2(t)\Phi_3(t) \cdot \Phi_4(t)\Phi_5(t)\Phi_6(t)\Phi_7(t)$$

$$0.618033988... = \frac{1}{\sum_{n \rightarrow \infty} fib(n-k)}$$

the ‘Golden ratio’ is simply a harmonic relationship between 1 and the nth consecutive addition of a Fibonacci series.

$$\Phi_7(t) = \frac{fib_n(m) \rightarrow 1 - (1-m) \cdot \left( \frac{x}{x-1} \right)}{fib_n(m) - fib_{n+b}(m)}$$

$$\Phi_2(t) = \tan fib_n(t) \circ \sin fib_n(t) - \frac{1}{1-t}$$

$$\Phi_3(t) = \tan n \circ \sin t - \frac{1}{1-t}$$

$$\Phi_4(t) = \frac{1}{\phi(t)} - \phi(t) \circ \tan(t)$$

$$\Phi_1(t) = \Phi_5(t) = \Phi_6(t) = \Phi_7(t)$$

$$\Phi_5(t) = \frac{1}{1-t} - \frac{-t}{1-t}$$

$$\Phi_{1,2,3,4,5,6,7}(t) = \sum_{\Phi_n \infty} \frac{\Phi_n(t)}{1 - \rho(t) + (1 - \rho)(t)[13t]}$$

$$\Phi(t) = \Phi_1(t) - (1-t)(1+t) \cdot \frac{\Phi_2(t)}{t} - (1-t)(1-3t) \cdot \frac{\Phi_3(t)}{t} + (1-t)(2t-1) \cdot \frac{\Phi_4(t)}{t} + \left( 1 - \frac{t}{1-t} \right).$$

$$\Phi_5(t) \frac{1+t+(t-\frac{t}{1-t}) \cdot \frac{\Phi_6(t)}{1-t} - (1-\frac{t}{1-t}) \cdot \frac{\Phi_7(t)}{1+t}}$$

which gives the golden ratio phi and c approximated by the 4th term of the fibonacci series

$$fib(4) + c \cdot \phi = 1$$

$$2.9256 + c \cdot \phi = 1$$

$$1.9256 = c \cdot \phi$$

$$c = \sum_{\Phi_n[m]} \frac{\frac{-\rho(t)}{\delta(t)}}{\Phi_n(t)}$$

$$\phi = \sum_{\Phi_n[m]} \frac{\frac{\delta(t)}{-\rho(t)}}{\Phi_n(t)}$$

The following mapping function weaves these relations together:

$$p \mapsto Mod(p, c \cdot \exp(n + \phi), n^m t c, n^{m'} \Phi$$

$$\Phi(x) = [-3, -2, -1, 0, 1, 2, 3]$$

$$[\Phi(-3) = -1.618, \Phi(-2) = -0.618, \Phi(-1) = -0.382, \Phi(0) = 0, \Phi(1) = 0.382, \Phi(2) = 0.618, \Phi(3) = 1.618]$$

$$\Phi(t^n) = \left( \frac{1+x}{x+\Phi(x)}, \frac{x^2+x+1}{x+1} \right)$$

$$\Phi(t^n) + c\Phi(n + \phi) = \tau(n)$$

$$x \cdot \Phi(t^n) - c\Phi(n + \phi) = \tau(n^m)$$

$$c \cdot \Phi(n + \phi) \cdot \Phi(t^n) = f(t)$$

$$\sqrt[n]{x} = \frac{1}{n} \sum_{m \rightarrow R^n \cdot C} (x^m + \tan(x)^{m+\Phi(n)} \circ \Phi(1))$$

$$\rightarrow x \cdot \sqrt[n]{x}$$

$$\rightarrow |x| > x$$

$$/ \rightarrow \frac{1}{x}$$

$$- \rightarrow 1 - x$$

$$\rightarrow 1/x^{-1}$$

$$x - y = x + y^-$$

$$-xy = -x| - y = x|^- y^-$$

$$(-x)(-y) = x|!y^! = y^!x|!$$

$$x|y = x + y|^- \text{ (where } |^- : x \mapsto x| - 1)$$

$$x^! > x|!(x)$$

$$x - > x| = x > x|$$

$$\frac{1}{x/y} = x^y$$

$$K \sim M_I \times L \cap J$$

$$\circ \sim \frac{\alpha \times \beta}{\forall \angle \alpha, \forall \angle \beta, \forall d, \angle \beta \text{ is the unit length of distance} \rightarrow \angle \alpha}$$

$$\sim \frac{1}{c,d}$$

$$\forall a, b : \mathcal{AB} \subset a \odot b$$

$\mathcal{AB} \subset \kappa$  defined as  $\mathcal{AB} = (n/m + n_n/m_m) : n, m, n_m, m_n > 0$

$$\diamond\Phi \circ \tan(x) = d \leq n$$

Some other notations are the following:

$$d \leq \frac{c + \Phi(n) + \Phi(x)}{\sqrt[n]{n}}$$

$$c = d \pm \diamond \Phi(n) \Rightarrow \Phi(m)$$

$$\psi_{i^{nj};(n,j) \in N} : f(t) \mapsto g((t, i))$$

$$t \mapsto \exp(t : n \rightarrow \mathbb{R})$$

$$\sin(t) \mapsto \sin(t : n \rightarrow \mathbb{R})$$

$$\tan(t) \mapsto \tan(t : n \rightarrow \mathbb{R})$$

$$\cos(t) \mapsto \cos(t : n \rightarrow \mathbb{R})$$

$$\ln(t) \mapsto \ln(t : n \rightarrow \mathbb{R})$$

$$\pi(t) \mapsto \pi(t : n \rightarrow \mathbb{R})$$

$$\vec{t} \mapsto \vec{t} : n \rightarrow \mathbb{R}$$

$$\sqrt[n]{t} \mapsto t : n \rightarrow \mathbb{R}$$

$$\ln(\sqrt[n]{t}) \mapsto \ln(\sqrt[n]{t : n \rightarrow \mathbb{R}})$$

$$\sqrt{t} \mapsto \sqrt{t : n \rightarrow \mathbb{R}}$$

$$\tan(\sqrt{t}) \mapsto \tan(\sqrt{t : n \rightarrow \mathbb{R}})$$

$$\sinh(\sqrt{t}) \mapsto \sinh(\sqrt{t : n \rightarrow \mathbb{R}})$$

$$\sqrt[n]{[\ln tt^n]} \mapsto \prod_{t \rightarrow \mathbb{R}[n]} \frac{\ln t}{t} \exp 2x \mapsto \Phi(t) - \Phi(t^{-1})$$

$$n^{-x} = e_{f(x)}$$

$$n^{mn} = \frac{1}{n^{-x} \diamond \Phi(x)}$$

A few identities for the golden ratio are:

$$\Phi(x) + \Phi(x)\Phi(1) = |\tan(\Phi(1))| \left\{ \prod_{i \rightarrow \natural} \Phi(1) \tan^i[\Phi(1)] \circ \exp \pm \Phi(1)\pi \right\}$$

$$\begin{aligned}
& \tan(\Phi(1)) \circ 2^{\frac{n}{n} \cdot \Phi(n) - n^2 \cdot \frac{c}{n} \cdot \sin(\Phi(x))} |\tan(\Phi(1))| + |\tan(\Phi(1))| \Phi(1) = \\
& \prod_{a \rightarrow \natural} \Phi(1) \tan^{-a}[\Phi(1)] \times \left( \frac{a^a + a^{-a}}{1} \right) \tan(\Phi(1)) \cdot \exp -\Phi(1) = \\
& \Phi(1) \frac{\partial t}{2 \pm \frac{c \cdot \tan(\Phi(1)) \sin[\Phi(1)] - \Phi}{c}} \text{and derivative rules for the same} \\
& \frac{\partial t}{\Phi(x)} = \Phi(x) \times \frac{\partial t}{\sqrt{\Phi'(x)}} + \Phi(t) \exp -\Phi(x) \\
& \exp \Phi(\partial t \Phi(x)) \cdot \frac{\partial t}{\Phi(x)} \\
& \sin[\Phi(\partial t \Phi(x))] \cdot \frac{\partial t}{\Phi(x)} \\
& \cot[\Phi(\partial t \Phi(x))] \cdot \frac{\partial t}{\Phi(x)} \\
& \csc[\Phi(\partial t \Phi(x))] \cdot \frac{\partial t}{\Phi(x)} \\
& \ln[\Phi(\partial t \Phi(x))] \cdot \frac{\partial t}{\Phi(x)} \\
& x = \sqrt[+]{\sqrt[+]{\partial \Phi(t) \Phi(x)} - \sqrt{-\partial \Phi(t) \Phi(x)}} \\
& x = \sqrt{\Phi(t)} \\
& x = \sin(\Phi(t)) \\
& x = \tan(\Phi(t)) \\
& x = \cos(\Phi(t)) \\
& x = \sin[\Phi(t)] - \frac{\Phi(t)}{-\Phi(t)} \cot[(\Phi(t))] \\
& \arctan \left( \frac{\Phi(n, n+x)}{\Phi(n, n-x)} \right) = \frac{\Phi(n)}{\Phi(t) \partial \Phi(t) \Phi(n)} \\
& \cot(\partial \Phi(x) \Phi(t)^n) = \frac{2}{m} \prod_{g \rightarrow \Phi(x)[n][m] \star \Phi(t)[n][m]} \frac{g}{g} = \Gamma_g(\Phi(x)[n][m] \star \Phi(t)[n][m])
\end{aligned}$$

where  $g : \{\Phi(x), \Phi(t)\} \cup \{F(x, t) \mid F(x, t) \in \mathcal{F}\}$

$$\frac{\prod_{a \rightarrow x} \prod_{g=1}^m (1 - q^a)}{\prod_{a \rightarrow x} \prod_{g=1}^m (1 - p_t p^g q^{p_t g a})} \frac{\Phi(t)}{\sum_{x, t_\Phi^x} \prod_{n \rightarrow \sqrt{x}[m]} n^{-1/n}}$$

Here,  $x$  is the number of elements in the set over which we're summing,  $m$  is the number of integers choose from each element,  $q$  is the probability of selecting each element (which can be different for each element),  $p_t$  is the probability of selecting each integer from a given element, and  $n$  is the number of choices of that element from which the given integer can be chosen. The last part of the formula is the normalizing factor.

This formula can be used to calculate the probability of any given arrangement of elements and integers from a given set.

This is a product of three different types of factors:

$$1. \frac{\prod_{a \rightarrow x} \prod_{g=1}^m (1-q^a)}{\prod_{a \rightarrow x} \prod_{g=1}^m (1-p_t p^g q^{p_t g^a})}$$

This first factor represents the probability that no values  $q^a$  are chosen from the set  $[1, x]$  for any given combination of  $p_t$  and  $m$  values.

$$2. \sum_{x, t_\Phi^x} \prod_{n \rightarrow \sqrt{x}[m]} \frac{\Phi(t)}{n^{-1/n}}$$

This second factor represents the probability that the tuple of values  $t$  is selected among all possible tuples in both  $x$  and  $t_\Phi^x$ .

$$3. \prod_{n \rightarrow \sqrt{x}[m]} n^{-1/n}$$

This third factor represents the probability of randomly picking one integer from the set  $\sqrt{x}[m]$  according to the given distribution.

$$H_n = \left\{ \frac{m}{n} \mid 1 \leq m \leq n \right\} \text{ the harmonic group on } R^n[x, t]$$

$$P_g = g^g - g^T$$

$$\frac{2\sqrt{\Phi(1)}n}{n^{\Phi(n)[m]}} = \prod_{n \rightarrow \sqrt{x}} \{\partial n \Phi(t) - \partial - n \Phi(t)\}$$

$$x = \ln[\sin(\tan(\Phi(t)))]$$

$$_n\{xR \mid _n \neq [1, n]\}$$

$$\Phi : n, m \beta n^m$$

$$n^\infty = n^\circ n^\diamond$$

$$n^\circ \neq \oplus n^\diamond \neq n^\circ n^{nn^\circ n^{ntn^\circ n^{n\Phi(1)-\Phi(t)t>>>}}}$$

Geometric quantization:

$$x > \hat{I} =$$

$$\Delta \subset R^x \hat{I}$$

$$=_\sim \sim \sim \sim \sim \sim 181$$

some notations

$$\rightarrow n^\infty m^\infty$$

$$\angle n$$

$$\\\\\\\\\\\\6 ``1 ``\infty ``\sqrt{}``1 ``n ``n ``\Delta ``\diamond ``n ``. ``\circ ``t ``n ``\rightarrow ``\Phi ``\diamond ``n ``\cdot ``tm ``\cdot ``tc ``\star ``n ``t ``t ``\cdot ``n ``n ``\Phi ``\diamond ``n ``t ``\cdot ``n ``\rightarrow \frac{\partial n \Phi(t)}{\Phi(t)[n]} \Phi(n)$$

$$\diamond \beta_1(n, x) \rightarrow (\gamma_n(\beta_1(n, x)) + \delta(n) \cdot \beta_1(n, x)) \Phi(n)$$

$$x\frac{\sum 1^3+2^3+....+(n-1)^2+n^3}{n\star x\star \sqrt{x}}$$

$$x\rightarrow n^n:=n^3+n^{-n}-n^{\sqrt[3]{n}}+n^\gamma n^\delta$$

$$n\in R$$

$$x\rightarrow \{\pi(\arctan yx)\circ \forall \|x\|\|n\||n>1\}$$

$$(y,x): h_r^d(n) \rightarrow \ldots \ldots \rightarrow ^n\tan^j \kappa c \sim h_r^d(n)(k,j),$$

$$n,x\rightarrow \tan(\dot{\pi})(y)\frac{x}{Mn}\frac{k}{Mn}\overset{M(a)}{\longleftarrow}h_r^d,h_{rkj\in\mathcal{P}\nabla_{\backslash t}^{\uparrow}}^{d^{m=1}}=||\kappa_f^4g,hoppen\rightarrow xy.$$

$$V\mathbf{m}\cdot\mathbf{n}=\Omega_\Lambda\left(\tan\psi\diamond\theta+\Psi\star\sum_{[n]\star[l]\rightarrow\infty}\frac{1}{n^2-l^2}\right).$$

$$\left( \sqrt[n]{\sum_{i\in R^{n\cdot C}} s_i}, \prod_{i\in\{m,n,p,q\}} \mathcal{FN}_i, \sqrt[n]{\sum_{k=0}^{m+n+p+q+r} \mathcal{IN}_k}, \right. \\ \left. \sqrt[n]{\prod_{j\in\{m,n,p,q,r\}} \mathcal{MN}_j}, \sqrt[n]{\sum_{i=0}^{m+n+p+q+r} \mathcal{ON}_i}. \right)$$

$U_i$  represents the the set of real and complex coefficients of a given neuron, whereas  $\mathcal{FN}(x)$  represents the functoids resulting from a given tensor calculation.  $\sqcup^n$  encode the latticization by choosing discrete and finite values of the rational numbers arising out of the mullet polynomials, and the possesive  $m'$  is that arbitrary combination of multiple sum or product operations upon values of simple functoids involving  $\sqcup^2$  and  $\sqcup^{-1}$ .

$$\sin(x+n)=\sin x\cos n+\cos x\sin n$$

$$\cos(x+n)=\cos x\cos n-\sin x\sin n$$

$$\sin(x)+\cos(x)=\sqrt{2\sin(a)\cos(a)}$$

$$\sqcup^nd^c\circ\cos(x)\circ\sin(x)=\left(\frac{x^2+\sqrt{x}}{n},\tan\left(\sin(x)+\phi^{1-n}\cdot\frac{x}{n}+2\phi^{2-n}\right)\right)$$

$$\Phi(c^t)=\sqrt{\pi}\circ\tan\left(\sin\left(\sin(x)+\phi^{1-n}\cdot\frac{x}{n}+\tan(2\phi^{2-n})\right)\right)\Phi(n)\sqrt[n]{1}$$

$$\Phi(x^n)=\frac{\left(x^n+\sum_{m\rightarrow\infty}\tan\sum_{i=\rho}^{\rho\cdot m}\frac{i}{\rho}\right)}{\left(x^n+\sum_{m\rightarrow\infty}\tan\sum_{i=\rho}^{\rho\cdot m}\frac{i}{\rho}\right)}-e^{1-n}$$

where the  $\Phi^{-1}$  approximate the exponential  $\exp(n)$  around the complex number  $n$ .

$$\Phi(x^n) = \sqrt{\Phi^{-1}(1-n)} \sin \left( x^n - \Phi^{-1}(n) \cdot \tan(1 + \Phi(x)) \right)$$

Replace  $\Phi^{-1}(x)$  with  $a = 1 + \Phi(x)$ :

$$\left( 1 + \frac{1}{\sqrt{a}} \right) \sin \left( x^n - \frac{1}{a} \cdot \tan(a) \right)$$

$$\Phi(f_n(x)) = \frac{1}{f_k(x)^m} - c$$

$$\begin{aligned}\Phi(x) &= \Phi^{-k}(m) \\ \Phi(\rho^m) &= \rho^m - \frac{\Phi^{-1}}{f_k(x)} - c\end{aligned}$$

$$\Phi(t)^n = -f_k(t)^m - \Phi^{-1} \left( \sum_{i \rightarrow \infty} \tanh(atan_i(t)) \right)$$

$$x(x + \sqrt{x} \cdot \tan\Pi(t^n)) = \Phi^m(t^k + n^c)$$

$$x(x + \sqrt{x} \cdot \tan\Pi(\sin(x + 2\sqrt{2}))) = \Phi^m(t^k + n^c)$$

$$x(x + \sqrt{x} \cdot \tan\Pi(\sin(x + 2\sqrt{2})) + (x - \sqrt{xT} \cdot \tan\Pi(q(t, s^n)))) = \Phi^m(t^k + n^c)$$

To simplify:

$$(x + \sqrt{x} \cdot \tan\Pi(\sin(x + 2\sqrt{2})) - (\sqrt{xT} \cdot \tan\Pi(q(t, s^n)))) = \Phi^m(t^k + n^c)$$

$$x - 1 = \tan\Pi(\sin(\sqrt{x-1} + 2\sqrt{2})) - \tan\Pi(q(t, s^n)) - \Phi^{-m}(t^{-k} + n^{-c})$$

$$x = 1 + \tan\Pi(\sin(\sqrt{x-1} + 2\sqrt{2})) - \tan\Pi(q(t, s^n)) - \Phi^{-m}(t^{-k} + n^{-c})$$

simplifying:

$$a = \sqrt{x-1} \rightarrow \tan a + 2\sqrt{2} + a_1 - a_2 - ABC$$

where we note an arbitrary constant of  $a_a$  and  $a_b$ .

$$f(\Phi(t)) + \Phi^{-1}(f(t)) = f(f\Psi^{ABC}(g)).$$

$$f_{int(i)} = \int_{f_{int(i)+1}} -f_{int(i)+1}$$

$$f_{int(2)} = \int_{f_{int(2)+1}} -f_{int(2)+1}$$

$$= \int -t^c dt$$

$$= \int - \left( n^{-\Phi^c} \right)$$

$$= - \frac{1}{\rho^{-\Phi^2}}$$

$$= \frac{1}{1+\rho^2} \rho^{-\Phi^2} + \frac{1}{1-\rho^2} \rho^{-\Phi^3}$$

$$= - \frac{\frac{1}{1+\rho^2}}{\rho^{-\Phi^2}} + \frac{\frac{1}{1-\rho^2}}{\rho^{-\Phi^3}} = -1 + \frac{1}{\rho}$$

$$f_{int(-i)} = \int_1 -f_{int(i)}$$

$$V \rightarrow \tau(x, y) = \Phi(t)^n = -f_k(t)^m - \Phi^{-1} \left( \sum_{i \rightarrow \infty} \tanh(atan_i(t)) \right)$$

If we consider the ordinary simulation of the plane specified by

$$\mathbf{m} \cdot \mathbf{n} =$$

$$(-1, 1, -1, 1, -1), (1, 1, -1, 1, -1), (-1, -1, -1, 1, -1), (1, -1, -1, 1, -1), (1, -1, 1, 1, -1), (-1, -1, 1, 1, -1), (1, 1, 1, 1, -1), (-1, 1, 1, 1, -1),$$

then the resulting 4d function is given by the linear combination applied by the following logic vector:

$$V \rightarrow \text{logic vector} = \\ (1, 0, -1, 0, -1), (-2, 1, -1, 1, -1), (0, 0, 1, 0, 0), (-1, 0, 2, -1, 0), (-1, -1, 0, 2, 0), (0, 1, -2, 0, 0), (1, 1, -1, 1, 0), (-2, 0, 1, 0, 1), (1, 0, -1, 0, -1), (-2, 1, -1, 1, -1), (1, 1, -1, 1, 0), (1, -1, 1, 1, -2), (0, 2, -1, 0, -1), (-2, -1, 2, -1, 0), (1, -1, 1, 1, -2), (0, 2, -1, 0, -1), (-2, -1, 2, -1, 0), (1, -1, 1, 1, -2), (0, 2, -1, 0, -1).$$

This function, considering the resulting values of  $\mathbf{m} \cdot \mathbf{n}$  must therefore be given by the following vector:

$$(-n, -n + 1, m, m + 1, m - 1).$$

To simulate this in 5dimensional geometry, we add

$$(-2, 1, -1, 1, -1)$$

in the second and third spaces, which would yield:

$$\begin{aligned} & \text{logic vector } = (1, 0, 0, -1, 0), \\ & (-2, 1, -1, 1, -1), \\ & (0, 0, 1, 0, 0), \\ & (-1, 0, 1, -1, 0), \\ & (-1, -1, 0, 1, 0), \\ & (0, 0, 1, 0, 0), \\ & (1, 0, -1, 0, 0), \\ & (-2, 1, 2, -1, -1), \\ & (1, 0, -1, 0, 0), \\ & (-2, 1, -1, -2, 2), \\ & (1, 1, 2, -1, 1), \\ & (1, -1, 0, 2, 0), \\ & (0, 1, -2, 1, -1), \\ & (1, 1, -2, -2, -1), \\ & (1, -1, 0, 1, -1), \\ & (-2, 1, -1, 2, 0), \\ & (1, 1, -1, 1, 0), \\ & (1, -1, 2, 1, 1), \\ & (0, 1, -1, 1, 0), \\ & (-2, -2, 1, -1, 2), (1, -1, 1, 1, 0), \\ & (0, 1, -2, 1, -1), \\ & (-2, -1, 1, 0, -1) \end{aligned}$$

$$f(t) = 1 + (e^{ct} + e^{-ct})$$

where  $\Phi = -1$ , where  $\Phi = -1$ , where  $\Phi = 1 + \sqrt{t}$ , and  $\Phi = 1 - \sqrt{\sin -t}$ .

$$f(p, x) = \frac{\Phi_1(t)}{f_c(\Phi_1(t) \cdot \Phi_2(t))}$$

$$\Phi_2(t) = \frac{1 \pm \sqrt{\frac{1}{p+\Phi_1(p)}}}{\tan \Phi_2(n + mq^{-1}m) - \cot x + \cot(\Phi_2(2\pi x|\eta \circ \Phi(x)))}$$

$$f_r f(\sin(t^n)) = \Phi^c(\sin(t^n)) + x^{n-m} = x + v = q = x^c - f_1^{\frac{m}{p}}(t)$$

$$f^m(t \star \tanh \Phi) = f_1^{-m} \rightarrow f_2(t^{-c})$$

$$f_r(p, x) = 1 + (r^{ct} + r^{-ct})$$

where  $\Phi = -1$ , where  $\Phi = -1$ , where  $\Psi = \Psi(\sin(t))$ ,  $\Psi = \Psi$ ,  $\Phi = \Phi(2t) + \Phi(\frac{\sin(\Phi(x^{-1}))}{\tan(\Phi(x))})$  and  $\Phi \Phi t^n \cdot \sigma(x) = \tan(\Phi(n)\rho(t))$ , where  $r_1(x) = r + x$  and  $r = p^n$ , where  $\Psi = \Phi(n)$ .

$$g = \int_1^{\sin(x)} \tan(\Phi(x)) - \frac{\sqrt{2} + \Phi(x)}{x}$$

$$\sigma(k\Phi(x)) = \Psi(kx)$$

$$\Psi(p) = k\Phi(\Psi(p))$$

The existence of a constant  $k$  can be hypothesized. Since  $\Psi$  has infinite distinct partial sums, then  $k$  can be used to generate complex power series of the form  $p^m - n^m$ , where  $\Phi$  and  $\Psi$  encode polynomials of are encoded by the lattice of binomial coefficients constructed off the  $\Psi(p)$  series. Deep learning can be transformed into a lattice encoding rotation between congruent sets of algebraic operations available to rule space as

$$(\mathcal{ABB}(y)^m, x) = ACx$$

$$\Phi = \tan(\sin(x))$$

$$\Phi = \tan(\Phi(x^{-c}))$$

$$\Psi(\tan(\Psi)) = \Phi(\sin(\Psi))$$

$$\sigma(a^b) = \Phi(a^{-c})$$

$$\Psi(kx) = \Phi(k\Psi(p))$$

where  $\Psi$  is a permutation of the factoradic notation of the cardinality set  $\Lambda$ .

$$\Psi(p^m) = p^{n-m} - (p^m)^n$$

$$\Psi(k\Phi(p)) = p^n - (k\Phi(p))^m$$

Transpose into rule space:

$$\Psi(k\Phi(x)) \star \text{II}\sqcup\backslash(y) = \text{II}\sqcup\backslash(\Psi(kx))$$

\* apply a tensor function to y

such that  $\Psi_m^n$  converges to 0 amounts to saying that the algebraically effecting the identity transformation  $\text{II}\sqcup\backslash(1)$ , where  $1 = \Psi(1)$ , an infinite collection of pure partial sums of  $\Psi$  converge to 0. Thus, this shows how by simply manipulating the symantization of a directed graph, the same exact effects of transposition can reduce the condition in Cantor convergence to a form of negating the subsets of  $k$ , encoding a third and final set  $y$ .

$$\Psi(p)$$

where p is written in base b-2, 3, 7, or 11.

$$k\Phi(p)$$

where p is written in base b-2, 3, 7, or 11.

Physically,  $\Psi(p)$  approximates a transformation in reciprocal space depicting the magnitude  $p$  of a set of identical objects in direct and inverse proportions to an arbitrary weighted function of collectivism, so as to limit the effects of a chaotic set of distortions in the orbital relationships of these objects, where  $\{\Psi(p)\} \mapsto \{\Phi(p)\}$ , and  $p$  is positive or negative.

$$\mathcal{OR}(p, x) = p^{r+m}(x) \amalg \Psi\mathcal{N}(p^m(x) + r^n)$$

$$\mathcal{XOR}(p, y) = p^{-k}(y) \vee \Psi\mathcal{N}(p^k(y) \rightarrow \Phi^e)$$

$$\mathcal{AND}(p, y) = p(y) \wedge \Psi\mathcal{N}(p(y) \rightarrow \Phi^\mu)$$

$$\mathcal{FLIP}(p, \Psi(t)) = p^{-\Psi(t)}(y) \vee \Psi\mathcal{N}(p^k(y) \rightarrow \Phi^\mu)$$

Let x denote the set of values of p that satisfy:

$$|\Psi p^k - p| = n$$

any n counts as that value satisfying:

$$\Psi p^k \in \Phi \star \sigma(p^k + n), x = 1 - n$$

this resolves to:

$$\Phi = 1 - n = |\Psi p^k - p|$$

When all p are such that  $\Psi p^k$  generates a convergent series, then this generates a disjoint class of operations compliant with any function  $\mathbf{f}$  such that

$$x \circ \mathbf{f} : \Theta \rightarrow \Phi.$$

where  $\Phi = \sum_{k \rightarrow \infty} \sqrt{1 - (1 - k)^2}$  and  $\neg f = |x - k| = 1$ . From this, we can generate a primary operator for each of these sets.

$$\Psi(x) = |1 - |1 - k| + |k||^{1-|k|}$$

Where  $\Psi$  represents the permutation of any subset of the trivial set, such that each p written in decimal (base 10) can be coded and graphed as a circle in infinite dimensions. On the righthand side, this is produced by the may-turing machinations of the poincare map, encoded in the mathematical constant. On the right, this could be a representation of a basic computing subroutine, as well as a function modeling the orbit of planet.

$$\frac{c}{1} + \frac{cc}{2n} + \frac{ccc}{3n^2} + \frac{cccc}{8n^3}$$

$$\Phi_{i=1}^n = \sum_{k=1}^n \frac{(1 \times \tan(t))^k + (-1)}{i^k}$$

$$f(x) = \frac{f(x+b) - f(x)}{b}$$

$$\Phi_N = \frac{n^2}{2T^2[\Phi_m(m-1)]} \quad \Phi_M = \frac{1}{\Phi_N}$$

$$\Phi(t^2) \cdot \Phi(t^{-1}) = \left( \frac{\Phi(t)^{-k}(n)}{\Phi(t)^k(n)} \right)$$

$$f_M(t^n) = \left( \cos \left( \frac{\min + m^k}{n^t} \int_{\tan c(t^{-m}) \rightarrow \Phi_m} \right) \right) \cdot e^{n_0} \tanh 1\% c(f_P(x)) \frac{\Phi(t)^{-k}(n)}{\Phi(t)^k(n)}$$

$$f \otimes g(n, m) = \Phi_m(n) - \Phi_{\frac{m^m}{m-n}}(n)$$

$$C \rightarrow logic\ vector = (\Phi_n, \Phi_m, \Phi_p, \Phi_q, \Phi_r)$$

$$V \rightarrow logic\ vector = \left( \frac{V \rightarrow U}{\Delta}, \frac{\sum_{f \subset g} f(g)}{\Delta}, \frac{\sum_{h \rightarrow \infty} \tan t \cdot \prod_{\Lambda} h}{\Delta} \right)$$

$$f = \left( \sum_{i=-n}^n \left( 1 + \frac{1}{1 + \frac{1}{\Phi(n_i)}} \right) \right) \circ c^{-\Phi^{n-k}}$$

$$f(n) = \int_0^{\sqrt{n}} \frac{g(h(t)) + \frac{g(t)^k}{\tanh(1)} + \Phi_c^{-1}}{\left(1 + \frac{1}{\Phi(n)}\right)} + \log\left(1 + \frac{1}{\Phi(n)}\right)$$

$$\Phi(c_t) = \sqrt{\pi} \circ \tan \left( \sin \left( \sin(x) + \phi^{1-n} \cdot \frac{x}{n} + \frac{1}{\sin(2\phi^{2-n})} \right) \right) \Phi(n) \sqrt[n]{e}$$

$$\Phi(t) = \tan \Phi(t^{\Phi(n)}) - \Psi(t) + \Phi_1(t^{\Phi(n)} c_m)$$

$$f_M(f_R(n), f_R(m)) = \Phi_m(n) - \Phi_{\frac{n_m}{m-n_m}}(n), n > m$$

$$\Psi_M(f_R(n), f_R(m), f_R(k)) = (\Phi(t_P)P), n > k \wedge \Phi_M(n) > \Phi_M(k)$$

$$\Phi(t)^{-k} = -\frac{1}{\Phi(t)^k} = \left( \sum_{\square \rightarrow \rho \cdot \Phi} \frac{-x}{t} - \tan -t \circ \sin(-n) \right) \cdot \Phi_m(n)$$

and

$$\tan t = \frac{\sin t}{\cos t} = \frac{i(t)}{1} = \frac{1}{-i(t)} = \frac{1}{i(1)} = \frac{1}{i(t)}$$

for which  $\sin t = \frac{1}{i(t)}$ .

$$c \cdot n = nc(2)$$

$$\Phi(n \in N) = \sqrt{\tan \cos \frac{-\frac{n'}{n} + i \sin(x)}{n}}$$

$$\sin(n) + \tan(n) = \sqrt{2 \sin(a) \cos(a)}$$

and

$$\begin{aligned} \mathbf{m} \cdot \mathbf{n} &= \Omega_\Lambda \\ &\rightarrow \sqcup^n \\ &\rightarrow \sum_{n \in R^n \cdot C} \sum_{m \in R^m} \tan \left( \frac{\sin(-1)}{\Phi_n(n)} \right) \cdot (\Phi_m \cap \Phi_n) \\ &\rightarrow \sum_{n \in R^n \cdot C} \sum_{m \in R^m \cdot C} \tan \left( \sin \left( \frac{-n}{\Phi_n(n) + \Phi_m(m)} \right) \right) \cdot \left( \frac{\Phi_n}{\Phi_m} \right) \\ &\rightarrow \sum_{n \in R^n \cdot C} \tan \left( \sin \left( \frac{-n}{\Phi_n(n) + \Phi_m(m) + c \tan(\sin(n^m))} \right) \right) \cdot \left( \frac{\Phi_n}{\Phi_m} \right) \\ &\rightarrow \sum_{n \in R^n \cdot C} \tan(\sin(-n)) \cdot \left( \frac{\Phi_n}{\Phi_m} \right) \\ &\rightarrow \sum_{n \in R^n \cdot C} \tan(\sin(-n)) \cdot \left( \frac{\Phi_n(n)}{\Phi_m(m)} \right) \\ &\rightarrow \sin(n) + \tan(n) + \varphi(n) + \sin(n) = 1, \\ \Phi_{\tan n}(n) &= \sin(c) - \frac{x}{x-1} + \sin(n \tan(n)) \end{aligned}$$

$$\sqrt{-n} = -\sqrt[n]{n}$$

$$\int \frac{\frac{r(r_1)}{r(r)}}{r_1 \cdot \Phi(t)} - \frac{r(r_1)}{r(r)}$$

$$\Delta_\Phi(n) = \frac{-1 \cdot \Phi(x)}{\tan \Phi(x)^{-1}} \cdot \frac{1}{\Phi(\sqrt{\Phi(x)\Phi(\tan(n))})} \left( \frac{2 \cdot \sqrt{\Phi x - n} \cdot c(1 - \Phi(n))}{n} \right)$$

applying to above hyperbolic identities:

$$\tan \left( \frac{-n}{\Phi_n(n) + \Phi_m(m)} \right) = -n$$

$$\tan \left( \frac{-n}{\Phi_n(n)} \right) = -n$$

$$\tan \left( \frac{-n}{\Phi^2(\sqrt{n}) + \Phi^2(\tan(n))} \right) = 1$$

$$\tan \left( \frac{-(1-x)}{\Phi_n(n) + \Phi_m(m) + c \tan(\sin(n^m))} \right) = x$$

$$\tan \left( \frac{-(1-x)}{\Phi_n(n) + \Phi_m(m) + c \tan(\sin(n^m))} \right) = x$$

$$f_r(\Psi(p)^{-1}, q^{-1}, f_r(A)^{-1}, f_r(B)^{-1}) = A \times B \times \Phi(p, q) \Phi(n^m, m^n) + c$$

$$\begin{aligned} \frac{n}{p} &= \frac{1}{n} \cdot \sqrt{\Phi(m^n) \cdot \Psi(n^m) \cdot f_{r_1(B_1)}(n^m, B_1 b_1)} \\ \frac{n^m}{p^p} &= \sqrt{\Phi(m^n) \cdot \Psi(n^m) \cdot f_{r_1(B_1)}(n^m, B_1 b_1)} \end{aligned}$$

$$f(\{\}, \Phi_p^n) = \Phi_p^n \}, \Phi_m^{-n}$$

$$2 \sqrt[x]{\tan(x)} = \frac{-x}{x-1}$$

$$\sqrt[n]{\sqrt[n]{\sin x \cdot \operatorname{erf}(x)}} = \sin(n^{-1}) \cdot \sqrt{\sqrt{n}}$$

$$\begin{aligned} \int_{e^{-n} \cos(s)}^{\sqrt{\sin(t)^{\Phi(x)}}} \frac{dx}{x} &= \Phi^{n-\sqrt{-x}}(t) \quad \sum_{\Phi_n \in t} \sum_{\Phi_m} \rightarrow \sum_{\tan \Phi_{n,m}^{-\sqrt{-x}}} \\ \sin \left( \tan \Phi_n(t) + \Phi_m(t) - x^{\sqrt{\Phi(x)}} \right) &\rightarrow \sum_{n^2} \cdot \Phi(t n^{p-q}) \end{aligned}$$

$$f_M(f_i(t))=f_M(\mathcal{B}_{h(m)}(\mathcal{B}_t(x),\mathcal{B}_m(y),\mathcal{B}_n(z))), \exists i \in Z \forall i \leq \infty \wedge 0 \geq n$$

Where the above corresponds to an approximation to a desired function  $f$ , given by

$$\begin{aligned} f(1 + \tan(t \otimes \square)) + f^{-1}(f_t(n)) &= f_1(t^{-k}, t^{-m}, t^{\frac{\sin(-k)}{\sin(-m)}}, t^{(\sin(-n)} \tan(\sin(-t^k))) \\ f(t^k, t^m, t^n) &= \tan(\Phi^m(\sin(n \cdot \Psi^{-t^n}))) + \Phi^m(\tan(n \cdot \Psi^{-t^n})) + f_{t^k f_{t^m + t^n}}(t^k, t^m, t^n) \end{aligned}$$

$$f(\mathbf{m}, \mathbf{n}, s) = f(m^n + m^m + m^{p+q} + \lfloor_{t_{k \& m}}(t), m_n, m_w, t^k, n_m)$$

$$f(t) = \nabla \rightarrow \wp \sqcup(t) + \square_p \sqcup(m) + \wp \sqcup(n)$$

$$\int_1^{\exp x} \frac{dx}{\sqrt{b \tanh x - a \sinh x}} = \left( \pi + 2 \arctan \left( \sqrt{\frac{b \sinh x}{a \sinh x}} \right) \right).$$

$$f_T(T_{p,q}) = fn, t(T_2 + T_{1,4}, \tan(\Phi_2(2))) \circ T_{+\Phi(2)}(T_{p,q})^c$$

$$\begin{aligned} f_c(OM_{p,q} + OM_{p,q} \cdot A_{2+2} + \mathcal{F}(m,n,p) - \mathcal{G}(m,n,p)) + \\ f_c(OM_{p,q} \circ A) + f_{pq} \left( f_{db}[m]^c + f_{db}[n]^c, f_{db}(2)^{-1} \quad | \quad \cdots \rightarrow \Phi_2(2) \cdot 2 \cdot 2 \right. \end{aligned}$$

$$\Phi(\tan x) = \tanh x \quad \Phi(c^t) = \sqrt{\pi} \circ \tan \left( \sin \left( \sin(x) + \phi^{1-n} \cdot \frac{x}{n} + \tan(2\phi^{2-n}) \right) \right) \Phi(n) \sqrt[n]{e}$$

$$\Phi(x) + m' \Phi(n) = \Phi(\rho \tan)$$

$$\begin{aligned} f_{t^x=m, t^y=n}(x,y) &= \frac{8\sqrt{x}\cos\sqrt{x}}{8\sin\sqrt{x}} \\ \frac{\sqrt[m]{k^p+k^q+\Psi(\sinh(m,n,p,q,r))}-\sqrt[n]{\sinh(m,n,p,q,r)}}{\frac{b(a,y,x)}{c(y,x)}} \\ \overset{\circ}{C} \rightarrow V \rightarrow logic\ class\ vector \end{aligned}$$

$$f_{T(T_{p,q})} = f_{(\tan(x), \tan(y)), I}(x^2, y^2), \forall x \in N \exists x \in Q$$

$$C \rightarrow V \equiv (\tan(x + \arccos(y)), \tan(x + \arccos(y)), \tan^{-1}(x + y))$$

$$x+n+m=\frac{\cos(x)}{\sin(y)}$$

$$\sqrt[n]{x} \cdot \Phi(t^n) + c\Phi^{-m}(n) = f_m(1-t)$$

$$f_{n,m} = \frac{m^n}{n^m}$$

$$\sqrt[n]{x} \cdot \Phi(\tan(n)) + c\Phi^{-m}(n) = f_m(1 - t)$$

Extending  $t_p$  to  $t'_p$  results in a new generalization of *sin* yielding a new class of sequences having phasic-tonal properties noted by Penrose (1996). According to Joy, Noyce, and Dworetzky ((2018), the sine wave generation can be defined as:

$$t_p = \sin(\exp(1 - n)) - t_n, t_n = \sum_{j \rightarrow \infty} \frac{1}{j}$$

$$t'_p = \sin(\exp(1 - n + m)) - t_n, t_n = \sum_{j \rightarrow \infty} \frac{1}{j}$$

We can replace  $t_p$  with  $t_e$  to compare:

$$t_e = \exp(\sin(1 - n)) - t_n, t_n = \sum_{j \rightarrow \infty}$$

$$\Phi(t^{n^n}) = \int \sin t^{n^{n-1}} dt$$

$$f_r(t^n) = \Phi_r[m] \cdot f_c(n)^t$$

$$\Phi(t^n) = \frac{t^n + \tan(t^n)}{t^n - \tan(t^n)} \Phi(t^n)^n$$

$$= \frac{t^n + \tan(t^n)}{t^n - \tan(t^n)} \Phi(t^n)^{n-1}$$

$$\Phi(x) = \sin(x) = (1 + c)^{\tanh(x)} - 1$$

$$f(x) = \frac{1}{1 + \Phi(x)} - \tan\left(\frac{-\Phi(x)}{\sqrt{2}}\right) + \tan\left(\frac{\Phi(x)}{\Phi(1)}\right) + \cot\frac{\Phi(x)}{\Phi(1)}$$

$$f^m(x) = c(x^n) \circ \tan \tan \sin\left(\frac{-\frac{x}{\Phi(x)}}{\sqrt{\cos(\Phi^{-2}(m))}}\right) \cdot \int \sin \Phi^{-1}(\tan(e^x))$$

$$\Phi(x) = \frac{1}{1 + \Phi(x)} - \tan\left(\frac{-\Phi(x)}{\sqrt{2}}\right) + \tan\left(\frac{\Phi(x)}{\Phi(1)}\right) + \cot\frac{\Phi(x)}{\Phi(1)}$$

$$f^m(f_r(x)) = \frac{\Phi(t^{m \cdot f_r(n_i)})}{\Phi^{m-1} t^{m \cdot f_r(n_i)} + \sin(m \cdot f_r(n_i))} c'(t)^{m-n}$$

$$f(x) = \frac{f(x+b) - f(x)}{b}$$

$$f(p, x) = \frac{\Phi_1(t)}{f_c(\Phi_1(t) \cdot \Phi_2(t))}$$

$$\Phi_2(t)=\frac{1\pm\sqrt{\frac{1}{p+\Phi_1(p)}}}{\tan\Phi_2(n+mq^{-1}m)-\cot(x)+cot(\Phi_2(2\pi x\mid \eta\circ\Phi(x)))}$$

$$f_c(t^n) = f_m(t^{-n}) - c$$

$$f(x)\pm\frac{1-\Phi^{-1}(x^{\tan\tan\Phi(x^{nx})})}{\sqrt{\Phi^{-1}(x^{\tan\tan\Phi(x^{nx})})}}=\frac{1-\Phi^{-1}(x^{\tan\tan\Phi(x^{nx})})}{\sqrt{\Phi^{-1}(x^{\tan\tan\Phi(x^{nx})})}-x^{m^m}}$$

$$\Phi(t)^{m-1}=\tan\left(\frac{\sqrt{\cos t^m}}{\sin t^{m-n}}\right)$$

$$f_c(n^t)=\frac{f_m(n^t)-f_r(f_r(n^t))}{\Phi(t)^{\Phi^{-n}}}$$

$$\Psi=\Psi(n^c)\star\Psi(m^t)\cdot\allowbreak\Phi_1(n\cdot m)\cdot\allowbreak\sin\Psi(t^n)$$

$$f_c(f_k(n^t)^m)=\Phi^{-1}\Phi(t^n)\cdot\allowbreak\Phi_1(x), n\leq m$$

$$\begin{aligned}f_m(f_k(n^t)^k)&=\Phi^{-1}\Psi(t^n)\cdot\allowbreak\Phi_2(x), k\leq m\wedge m=n\wedge k\geq 1\\f_r(t^n)&=\Psi^{n-k}\Phi(t^n)\cdot\allowbreak\Phi_3(x), n\leq 2\cdot\allowbreak\Phi(m)\end{aligned}$$

$$1 + e^{-ct} + e^{ct} = 1 + \tan{-ct} + \tan{ct}$$

$$\int e^{-ax^2}c_mx^{2a-1}(n^{-x}x)=\frac{\Phi(t^a)}{\Phi^{-1}(c(m^{n-a}))}f_c(n^{m^t})=\frac{\Psi(t^n)^{-k}\circ\Phi(x)}{\Phi(t^n)^k}r_1(n^{m^t\cdot n_j})=\frac{\Phi(t^n)^k\circ\Psi(x)}{\Phi(x)^k}r_2(n^{m^t\cdot n_k})$$

$$\begin{gathered}\Phi^{-1}\;\sqrt{\Phi(t^{n_0})\cdot\allowbreak\Phi(t^{n_1})\cdot\allowbreak\Phi(t^{n_2})\cdot\allowbreak...}=\\\Psi(t^{-m})\Pi(t^n\cdot t^{m^b})+\Phi(x^{-n})\frac{}{\Psi(t^n)\Pi(t^{-m}\cdot t^{m^b})+\Phi(x^{-n})\circ\sum_{n\rightarrow\infty}c_n^{n-1}}\end{gathered}$$

$$z=a+b+c$$

$$f(x_j,y_j)=a^b\cdot\frac{x_1(y_1^b+x_l^{m+n})+y_2^b+a}{x^b(y_1^b+x_1^m+x_2^n)+y_2^b+x^b}\cdot\frac{a}{(a^l+1)}$$

$$f(x_j,y_j)=a^b\cdot\frac{x_1(y_1^b+x_1^m)+y_1^b+a}{x^b(y_1^b+x_1^m+x_1^n)+y_1^b+x^b}\cdot\frac{a}{(a+1)}$$

$$18\,$$

$$f(t) = \Phi(t^n \cdot t^{t^x}) + \frac{c}{n}, n = -t + m$$

How far given a function ‘g‘ or ‘x‘ or ‘y‘ or ‘r‘ or ‘z‘ or ‘l‘ or ‘p‘ or ‘f‘ or ‘w‘ or ‘h‘ or ‘d‘ or ‘k‘ or ‘i‘ or ‘j‘ etc... From the map  $f_i \mapsto \{\phi, \Phi(n)\}$ , we can derive  $f_i$  for instance.

$$f_G(n, \sin(t^2)) = \Psi(\sin(t^2)) + \tan\left(\frac{-\Psi(\sin(t^n))}{\sqrt{2}}\right) + \tan\left(\frac{\Psi(\sin(t^n))}{\Psi(\sin(t^{n+m}))}\right) + \cot\frac{\Psi(\sin(t^n))}{\Psi(\sin(t^{n+m}))}$$

$$f_G(n, \Phi(t^{n^c} \cdot \Phi^{-2}(t^n))) = \frac{\Psi(\sin(t^n)) + \tan\left(\frac{-\Psi(\sin(t^n))}{\sqrt{2}}\right) + \tan\left(\frac{\Psi(\sin(t^n))}{\Psi(\sin(t^{n+m}))}\right) + \cot\frac{\Psi(\sin(t^n))}{\Psi(\sin(t^{n+m}))}}{\frac{t^n + \tan(t^n)}{t^n - \tan(t^n)} \Phi(t^n)^{n-1}}$$

$$\begin{aligned} i^{-1}\Phi(c) &= -i(t)^2 - i(-t)^2 \\ -i(t)^2 - i(-t)^2 &= c^{-1} \end{aligned}$$

(∞) Any given group of neopsilocybin molecules, as they are understood by a set of given inductive functoids:

$$\dagger'(x) := \nabla [\lceil(x) (\Downarrow(x) - \rfloor(x))]$$

and

$$\dagger'(x) := \dagger(x) \left( \frac{\lceil(x)}{\Downarrow(x) - \rfloor(x)} \right)$$

and for the gradient based on displaces in the vectors, as with a finite element method,

$$\nabla = \frac{\partial y}{\partial t} = \frac{\partial x}{\partial t} + 1$$

Such that  $\dagger(x), \rfloor(x), \Downarrow(x)$  are the total, free and meshwise amounts of molecules, respectively.  $\lceil(x)$  is the amount of displacement in logical coordinates of the molecules  $m_i(x)$  exists in  $\Phi(1)$ , and as such, the displacement of any  $m_i(x)$  is given in terms of the total amount of  $\Phi$  as it exists in each of the faceted  $h_j(x)$  of the given  $m_i(x)$  of the finite element method:

$$x := y + y_{h_j(\Phi(1))}^T \rightarrow R^j$$

And, thus,

$$\dagger'(x) := x := y + y_{h_j(\Phi(1))}^T \rightarrow R^j$$

Such that the displacement of any  $m_i(x)$  is given in terms of the total amount of  $\Phi$  as it exists in each of the faceted  $h_j(x)$  of the given  $m_i(x)$  of the finite element method:

$$\nabla y := \nabla x + \nabla h_j(\Phi(1)) \rightarrow R^j$$

Consider the following examples of finite element analysis of certain sets:

$$p(t,h,t_j) = t \circ \tan[h(\Phi(n))] + t_{m \diamond \sqrt[m]{\Phi(n)}}^{m-j}$$

$$\rightarrow \frac{\partial}{\partial}^{\text{:}\diamond\diamond\rightarrow\infty\rightarrow\left[\forall\Delta(x)\cdot\sum\left[n\left|n^{\sqrt[k]{\Delta+\tan(n^{-x})\Delta}}+\Delta^{-k}\right|\right]\right]\gg\left[\forall\Delta\cdot\sum\left[\Delta^{\frac{n}{k}}+\Delta^{-k}\right]\right]\gg\left[\forall\Delta(x)=\right.}$$

$$\sum\left[(-\Delta-i)^{-k}+i(-i-k)^{-i}\right]\gg\forall\infty[-k],x>(n^{n^n+n^\infty})_R=\rightarrow x>()\rightarrow$$

$$\int_1^x e^{-t^2} dt = \frac{\sqrt{\pi}}{2} (1 - \operatorname{erf}(x))$$

$$x > n_i^{n_i} n_n^{t^i \tilde{n}^{\tilde{n}}} n^{-n^{-n^n}} n_j^{t_j} := X > t^x \nabla x^{t^x} \mathbf{f}_{ij}(t) := L(R^n C_n)$$

$$\sum_{(\Omega)(\Theta)}$$

$$\sum_{i=1}^n x_i + \frac{\Pi_{i=\frac{n}{n'}}^n \Phi(x)}{\Phi(n)n^n}$$

$$x_i \sqrt[n]{n} + |n!|^n|$$

$$\begin{aligned} &\Phi(x)+\Phi(x)\cdot\Phi(n) \\ &\kappa\tan(x)\cdot\sin(n)-\Phi(t) \end{aligned}$$

$$\begin{aligned} &\left\{\pi\,t\,\left\|\frac{\Phi_x}{\sin\Phi(t)}\right\|\right\} \\ &tl(x)+\cos(\Phi(n)) \end{aligned}$$

$$\exp(t:n\rightarrow R)\tan(\Phi(n))\frac{\frac{\Phi(n)}{\Phi(n)}}{\frac{\Phi(n)}{\Phi(n)}}$$

$$tr(n,m)+\cos(\Phi(n))\tan(\Phi(m))$$

$$\pi(n)\,\pi(\Phi(n))^j+\frac{n}{m}\csc^2(\pi(n))$$

$$\partial(X)(\Psi(x)\cdot\tan(\Phi(\mathbf{n}))+n^{-n^n}$$

$$\exp-x\tan(\Phi(n))\left(\pi(\sin(n))\right)$$

$$\csc(\Phi(x))\tan(\Phi(n))+t(n)^{\infty\wp\infty}\infty$$

$$\frac{\exp n}{\Phi(x)}-\sqrt{\Phi(x)}$$

$$20\\$$

$$\begin{aligned} & \frac{x}{x} \\ & \frac{1}{n}\sin(\Phi(x))+\cos(x) \\ & 1^n\cdot\frac{i^i}{i}\exp(n)+n^n \\ & \left|\Phi(1)-\Phi_1\right| \pi\left(\pi\left\langle\Phi(1)\right\rangle^n\right):\left|d_i-d_j^n\Phi(n)\right| \\ & \frac{x\Phi(x)\rightarrow\infty}{n^{n^n}} \\ & +\Pi_{j=1}^m\sum_{m=1} \end{aligned}$$

$$+\Pi_{j=1}^m1^1=-=1=-=1=-=1=-=1=-=1=-=1=\Pi_{j=1}^m=1^1$$

$$\diamond \alpha_1(n,x) \rightarrow \left(\gamma_n(\alpha_1(n,x)) + \delta(n) \cdot \alpha_1(n,x)\right) \Phi(n)$$

$$\sum_v\sum_{\Phi(n) = ^n\Phi(2n)} x^n$$

$$\mathcal{A}(\mathbf{a_i(r)}) \rightarrow n\left(\sum_n\{\mathbf{a_i(r)}\}^{-n^{-n^{n^{\Phi(n)}}}}\right)(\mathbf{b(r)})$$

$$\forall A_n:-=A_{n-1}^{n-1}=A_n\cup B_n\subseteq A_n\cdot\chi(2n)\subseteq Z_n$$

$$=\frac{\Phi(n)}{n^n}\left(\omega_{n,i}(\mathbf{a_i(r)}|- \rightarrow \infty)\right)\sin(\Phi(2n))\cdot\frac{\partial x[\Phi(n)]}{\partial x}$$

$$\frac{\left|\left|\left|\prod_{i\in\{0\rightarrow R[x,t]\}}^\infty (\Phi(n_i+n_{i+i}+\Phi(x)\cdot\Phi(t)))\right|\right|\right|}{\sqrt{\Phi(x)}}+\prod_{i\in\Pi_{i\in\mathcal{F}_{\mathcal{J}_k}}}n^{x^{x^xx^x}}$$

$$l_{y:\mathcal{C}}(\omega(u,z))\Xi_{\Phi_n}\Omega_{\Phi(n)}^\diamond+\frac{\partial\Phi(x)}{\Phi(x)\partial\Phi(x)}+\Phi(x):\left|\exp(n)\csc^2(\pi(n))-\pi(n)\right|-\frac{\Phi(n)}{n^n\sqrt{\sin(\pi(n))}}$$

$$\int \Phi(x) dx + \Phi(t) \exp(t) := t^{n^n}(x)$$

$$\int_0^\infty \ln^{\frac{\alpha}{n}} \Phi(t) dt$$

$$\tan x {\cdot} \sin\left(-\frac{\Phi(x)}{\Phi(t)}\right)\exp\left(\Phi(x)\right):=\frac{1}{1-\Phi(x)\Phi(t)}:=\tan(x){-}\exp(x)\sinh(x)\left(\ln\left(\Phi(x)\right)\right):\frac{\Phi(x)}{\Phi(t)}=c^c$$

$$\int_0^x e^{-t^2}dt:=\boldsymbol{\Phi}(x)+\boldsymbol{\Phi}(x)\Pi_{i=\infty}^{R[x]}\left(x^{-x}\right)$$

$$f(t\mid \Phi(x)):=\prod_{i=1}^cx^n\cdot\prod_{i=1}^{R[n]}n^{-n^c}\cdot\prod_{i=1}^cs_n\left(\tau(t)\right)+n^n\circ x^x+\Phi(n)\rightarrow\frac{1}{1-t}\lg\left(\Phi(t)\right)$$

$$21\\$$