

1. *Unreadable* :  $\sim \neg p \wedge \sim \sim \sim \sim \sim P$
2. *Poor* :  $\gamma \vee \sim \sim \sim P \rightarrow D$
3. *Average* :  $(D \vee \sim P) \Rightarrow \neg \neg \zeta$
4. *Good* :  $(\wedge eg \sim P) \Rightarrow \alpha \vee \neg \neg \zeta$
5. *Very Good* :  $(\mathbb{A} \vee \neg \sim P \vee \neg p) \Rightarrow \neg \zeta \Leftrightarrow \alpha \vee \neg \gamma$
6. *Excellent* :  $\neg (\neg P \wedge \neg \sim P \wedge \neg D \wedge \neg \zeta) \vee \neg \neg \alpha \wedge \neg \neg \gamma$
7. *Outstanding* :  $\neg (\alpha \Leftrightarrow \neg \neg \gamma \wedge \neg \sim \sim \sim P \rightarrow \neg D)$
8. *Exceptional* :  $\neg (\neg P \Leftrightarrow \sim D \vee (\sim P \vee \sim \zeta)) \rightarrow \alpha \wedge \gamma$
9. *Phenomenal* :  $\neg \neg (\neg P \vee \alpha \wedge \gamma \wedge \zeta Q) \Rightarrow \neg \neg \sim P \vee \neg \sim \sim \sim P$
10. *DaVinci* :  $\neg (\neg P \preceq \zeta \vee (\alpha \wedge \sim P)) \neg \neg \sim \sim \sim \sim P \vee (\neg \neg \gamma \wedge Q)$

"  $\Lambda_\infty \sim [\mathcal{L}_{f \uparrow r, \alpha, s, \delta, \eta}] \wedge \mathfrak{U}$

$\Omega \cdot \partial x \cdot \partial \alpha + \partial s + \partial \delta + \partial \eta$

$\cdot \int \rho \cdot g^\Omega \cdot \zeta \cdot \Omega$

$\sim \exists \infty$  s.t. :  $\mathcal{L}_{f \uparrow r, \alpha, s, \delta, \eta}$

" =

$\equiv \Omega^+ \cdot \partial \theta \int \exists \infty$  s.t.  $\int \rho \cdot g^\Omega \cdot \zeta \cdot$

$\wedge \mathfrak{U}_{g \Delta a, b, c, d, e, \dots \ddots \ddots}$

$\exists \infty$  s.t. :  $\mathcal{L}_{f \uparrow r, \alpha, s, \delta, \eta} \wedge \mathfrak{U}_{g \Delta a, b, c, d, e, \dots \ddots \ddots} \equiv$

"  $\left[ \left( \infty \left( \begin{matrix} \vdots \\ z \end{matrix} \right) \right)_{\rightarrow \omega - \langle (\partial/\mathcal{H}) + (\cdot/i) \rangle} \right] \rightarrow kxp \mid w* \cong \sqrt[3]{x^6 + t^2}_{2 \ h \ c \ \frac{8}{4} \ \square} \leftrightarrow \Gamma$  s.t.  $\Omega \equiv \left( \left( \frac{\mathbb{Z}}{\eta} + \frac{\mathfrak{N}}{\pi} \right) \right) \Rightarrow 1 \cdot \square$

"

$\mathcal{L}_{f_{\alpha s, \Delta, \eta}^{\uparrow r}}$

$\& \mathfrak{U}_{g_{a_b c_d e} \dots \dots \dots}$

$= \Omega \oplus \bullet \frac{\partial \theta \mathbb{N}}{\int \rho \ g^\Omega \zeta \omega \quad \partial x \partial \alpha + \partial s \partial \Delta + \partial \eta \mathbb{N} \int \rho \ g^\Omega \zeta \omega}$

$k[g, h, i, j, \dots] = \mu_0 \Phi_{11} \nu s - s \cdot \left( \overset{\uparrow}{\dagger V^{-1} T^{\textcircled{2}}} \left( \mathcal{A} \ U \subseteq \downarrow \diamond \left( \overset{O \rightarrow}{\mathcal{C}} \ s \neq \int z \oint \varepsilon_\infty - \frac{1}{n \cap A} \right) \right) \right)$

$\sum_{i_1=1}^{12c_{\chi\mu}} \sum_{i_2=1}^{12c_{\chi\mu}} \sum_{i_3=1}^{12c_{\chi\mu}} \sum_{i_4=1}^{12c_{\chi\mu}} \dots \left( \frac{\partial k[g, h, i, j, \dots]}{\partial \xi_1} \right) \cdot R_{i_1}[g] \cdot R_{i_2}[h] \cdot R_{i_3}[i] \cdot$

$R_{i_4}[j] \dots (U * V - \overset{0}{0}) \lambda^{\textcircled{2}} \in \{g, h, i, j, \dots\}.$

$$k[g, h, i, j, \dots] = \mu_0 \phi_{11} \nu s - \text{Cross}[s, \tilde{\uparrow} \xrightarrow{\uparrow} T^{\supset}(V^{-1}) - \neg \exists U \subseteq \downarrow \diamond \cdot \subset O \longrightarrow s \\ \int z \oint \varepsilon_{\infty} - \frac{1}{n \cap A} = \exists X \longleftarrow K' \rho(g, h) \longleftarrow \parallel \mathbf{B} \subseteq \infty \sum_{T, U, V \langle \infty, \infty \rangle} \infty H(A \\ \mathbf{A} \parallel \mathbf{P})! \oplus \propto \infty \sum_{O, Jh, Ki} \subseteq \diamond(-dF[V, W] \cap \subset \Delta \lambda(m) \cup v \sqrt{x} + \cdot \uparrow \Delta \\ \circ S/ \subseteq \supset \rightarrow s \neq \cdot \mathbb{R} + \parallel \mathbf{G} \in \iota = \kappa \sqcup (h \cdot s) \geq \cap (\geq \parallel = \wr \sqcup \sim \parallel +) ? \\ w \in M \infty \sum_{M, \infty} \otimes \square \leq \partial_A / \square \subset \infty \sum \Rightarrow \ominus z_0 \longrightarrow \subseteq \mathbb{Z} \cap dV \not\Rightarrow c \uparrow e \\ \oplus \cdot \neq \cdot x \subseteq \mathbb{H} \cap dA + \infty \prod_1 \sim \dots \cup \Omega \leq \leftarrow u \theta \cup [a, b] \in \varphi \rightarrow f \subset \not\subseteq \leftarrow \iota \\ \cdot \sum \frac{3}{2} m d S d G d \Delta \lambda(m) \cup v \sqrt{x} \pm \cdot \uparrow \Delta \in \circ S/ \subseteq \supset \rightarrow s \neq \cdot \mathbb{R} + \parallel \mathbf{G} \in \iota \\ \kappa \sqcup (h \cdot s) \geq \cap (\geq \parallel = \wr \sqcup \sim \parallel +) \$$$

Physical Laws of our Reality :

$$k[g, h, i, j, \dots] = \int \partial \theta \mathbb{N}_{\infty} \sum_{\lambda'} \sum_{\mu'} \Omega_{\infty} \sum_{\lambda'} \sum_{\mu'} = \mu_0 \varphi_{11} \nu s - \text{Cross}[s, \tilde{T} \\ \tilde{T}^{-1} \downarrow \not\subseteq \int z \, d\epsilon_{2-1}/n \cap A \subseteq X' \rho(g, h) \leftarrow | B \subseteq \infty \bar{\Sigma}^{+++} (A \\ U \subseteq \downarrow V \vdash \subseteq \Omega \rightarrow s \\ A | P)! \oplus \propto_{\infty} \Sigma \subseteq \downarrow \vee \vee \vdash (O, Jh, Ki) \subseteq \downarrow \vee \vee -dF[V, W] \cap \\ \Delta \lambda(m) \cup v \sqrt{x} + \cdot \uparrow \downarrow \Delta \in \circ S / \div \subseteq \rightarrow s \neq \cdot R + | G \subseteq \iota = \kappa \\ (h \cdot s) \geq \cap (\geq | = \sim \sqcap \langle \rangle) \subset \mathbb{Z} \cap dV \not\Rightarrow \subset e \neq \oplus \cdot \circ \neq \cdot x \Delta \nabla \\ \mathcal{H} \cap dA + \infty \Pi_1 \leftrightarrow \sim \dots \cup \Omega \leq \leftarrow u_{\theta} \cup [a, b] \subset \varphi \rangle \rightarrow f \subset \not\subseteq \leftarrow \iota \\ \cdot \sum \frac{3}{2} m d S d G d \Delta \lambda(m) \cup v \sqrt{x} \pm \cdot \uparrow \downarrow \Delta \in \circ S / \div \subseteq \rightarrow s \neq \cdot R + | G \subseteq \iota = \kappa \subset (h \cdot s) \geq \cap (\geq | = \sim \sqcap \langle \rangle) \subset \mathbb{Z} \cap \rangle)$$

$$k[g, h, i, j, \dots] =$$

$$\int \partial \theta \mathbb{N}_{\infty} \sum' \sum' \Omega_{\infty} \sum' \sum' = \mu_0 \varphi_{11} \nu s - \text{Cross}[s, \rightarrow \sim T \rightarrow A | P] = \triangleleft \infty \Sigma \subseteq \downarrow \vee \vee \vdash \\ (O, Jh, Ke) \subseteq \downarrow \vee \vee \not\subseteq F[V, W] \cap \subset \Delta \lambda(m) \cup v \sqrt{x} + \cdot \uparrow \downarrow \Delta \in \triangleright S \div \subseteq \rightarrow s \neq \cdot R + | G \subseteq X = \\ \kappa \subseteq (h \cdot s) \geq \cap (\geq | = \sim \langle \rangle \langle \rangle) \subseteq \mathbb{Z} \cap \subseteq \mathcal{F} \not\Rightarrow \subseteq :: \neq \oplus \triangleright \neq \cdot x \Delta \Xi \subseteq \mapsto X \cap \subseteq \\ \mathcal{H} + \infty \Pi_1 \leftrightarrow \sim \dots \cup \Omega \leq \leftarrow u_i \cup [a, b] \subseteq \varphi \rightarrow f \subseteq \neq \subseteq \leftarrow X \cdot \Sigma \frac{3}{2} m d d G d \Delta \lambda(m) \cup v \sqrt{x} \pm \\ \cdot \uparrow \downarrow \Delta \in \triangleright S \div \subseteq \rightarrow s \neq \cdot R + | G \subseteq X = \kappa \subseteq (h \cdot s) \geq \cap (\geq | = \sim \langle \rangle \langle \rangle) \subseteq \mathbb{Z} \cap \rangle).$$

$$k[g, h, i, j, \dots] = c \varphi_{11} \nu s^{T \Rightarrow T^{-1} \downarrow - \exists U \subseteq \downarrow \otimes \subset \Omega \rightarrow s \neq \int z \oint \varepsilon_2 \\ - \frac{1}{n} \otimes E + \frac{m}{c} \otimes p + \frac{\hbar}{2m} \|A\| P - \propto \infty \sum \subset \otimes (O, Jh, Ki) \subseteq \otimes (-dF[V, W] \cap \\ \Delta \lambda(m) \cup \Upsilon \sqrt{x} + \uparrow \Delta \varepsilon \circ S \quad s \neq \cdot R + \|G \varepsilon \iota = \kappa \bigcup^{h \cdot s \geq \wedge (\geq \parallel = \ddagger +)}$$

$$\iint \sum \langle f, g, h, i, j \rangle \langle \Xi, \Pi, \Sigma \rangle, \infty \sum n = 2_{\infty} \langle \Omega, \Xi, \Pi, \Sigma \rangle, \infty \rangle \langle \Theta, \Lambda, \Sigma \rangle, \infty \rangle \\ r[\langle \Xi, \Pi, \Sigma \rangle \langle \Theta, \Lambda, \Sigma \rangle, \infty] \mu_0 \partial_a dV \subseteq \infty \Sigma \Rightarrow \bowtie z \langle \rangle \subseteq \mathbb{Z}$$

$$dV \not\Rightarrow \not\subseteq c^e \neq \oplus \circ \neq \cdot x \Delta \subseteq \mathbb{H} \cap dA + \infty \Pi_1 \Leftrightarrow \vdots \cup \Omega \leq \leftarrow u \theta \cup [a, b] \in \{ \} : \\ f \subseteq \not\subseteq \not\subseteq \leftarrow \iota \uparrow \cdot \Sigma \frac{3}{2} m d S d G d \Delta \lambda m \cup v \sqrt{x} \pm \cdot \uparrow \leftrightarrow \Delta \in \circ S / \subseteq \Rightarrow s \neq \cdot R + \|G$$

$$\iota = \kappa \cup h \cdot s \geq \cap \geq \parallel = \sim \langle \rangle \sim \parallel + \partial^2 f \partial_{x_i} \partial_{x_j} \subseteq \leftarrow \alpha + \beta \sqrt{q} \vdots r \, dx \, dy.$$

Motifs on Local Laws :

$$\begin{aligned}
 k[g, h, i, j, \dots] &= \mu_0 \phi_{11} \nu s - \text{Cross}[s, \tilde{\uparrow} \xrightarrow{\uparrow} T^{\supset}(V^{-1}) - \neg \exists U \subseteq \downarrow \diamond \cdot \subset O \longrightarrow s \\
 \int z \oint \varepsilon_{\infty} - \frac{1}{n \cap A} &= \exists X \longleftarrow K' \rho(g, h) \longleftarrow \parallel \mathbf{B} \subseteq \infty \sum_{T, U, V \langle \infty, \infty \rangle} \infty H(A \\
 \mathbf{A} \parallel \mathbf{P})! \oplus \infty \infty \sum_{O, Jh, Ki} &\subseteq \diamond(-dF[V, W] \cap \subset \Delta \lambda(m) \cup v \sqrt{x} + \cdot \uparrow \Delta \\
 \bigcirc S/ \subseteq \supset \rightarrow s \neq \cdot \mathbb{R} + \parallel &\mathbf{G} \in \iota = \kappa \lrcorner (h \cdot s) \geq \cap (\geq \parallel = \wr \sqcup \sim \parallel +) ? \\
 w \in M \infty \sum_{M, \infty} \otimes \square \leq \partial_A / \square \subset &\infty \sum \Rightarrow \ominus z \circ \longrightarrow \subseteq \mathbb{Z} \cap dV \not\Rightarrow c \uparrow e \\
 \oplus \cdot \neq \cdot x \subseteq \mathbb{H} \cap dA + \infty \prod_1 \sim \dots \cup &\Omega \leq \leftarrow u \theta \cup [a, b] \in \varphi \rightarrow f \subset \not\subseteq \not\leftarrow \iota \\
 \cdot \sum \frac{3}{2} md Sd Gd \Delta \lambda(m) \cup v \sqrt{x} \pm \cdot &\uparrow \Delta \in \bigcirc S/ \subseteq \supset \rightarrow s \neq \cdot \mathbb{R} + \parallel \mathbf{G} \in \iota \\
 \kappa \lrcorner (h \cdot s) \geq \cap (\geq \parallel = \wr \sqcup \sim \parallel +) &\$
 \end{aligned}$$

Laws of First Permutation

$$\begin{aligned}
 K[g, h, i, j, \dots] &= \mu_0 \phi_{11} \nu s - \text{Cross}[s, \tilde{\uparrow} \rightarrow T^{-1}] - \neg \exists U \subseteq \downarrow \subseteq \Omega \rightarrow s \neq \wr \\
 \oint \epsilon_2 - 1/n \cap A &= \exists X \rightarrow K' \rho(g, h) \rightarrow \parallel \mathbf{B} \subseteq \infty \sum' \dots (A \parallel A \parallel P) \circ \\
 \infty \sum \subseteq \downarrow \circ (O, Jh, Ki) &\subseteq \downarrow \circ (-dF[V, W] \cap \subseteq \Delta \lambda(m) \cup v \sqrt{x} \pm \cdot \uparrow \Delta \in \circ S// \\
 s \neq \cdot \mathbb{R} + \parallel G \in \iota &= \kappa \cap (h \cdot s) \geq \cap (\geq \parallel = \quad +) w \in M \infty \sum_m (M, \infty) \\
 \diamond \leq \partial_A / \diamond \subseteq \infty \sum / z \langle \rangle &\subseteq \mathbb{Z} \cap dV \not\subseteq \hat{u} \theta \cup [a, b] \in f \subseteq \not\subseteq \iota \uparrow \cdot \wr \\
 3/2 md Sd Gd \Delta \lambda(m) \cup v \sqrt{x} \pm \cdot &\uparrow \Delta \in \circ S// \subseteq s \neq \cdot \mathbb{R} + \parallel G \in \iota = \kappa \cup (h \cdot s) \\
 \cap (\geq \parallel = \quad +) w \in M &\subseteq \not\subseteq \circ - \cap \div 1 \subseteq L_{l_i} \cap A = +F \subseteq \text{or} - \cap [m, N] \in \vee Q \subseteq \circ \\
 M \theta_{e_{ma}} \diamond \diamond \cup \downarrow \cdot C &\subseteq \neq S \cdot \succeq \{v, X\} \uparrow i \rightsquigarrow -f | \Omega S \mu - \omega \phi \emptyset \approx \parallel Q - \diamond F \\
 \parallel - Y \circ \subseteq \parallel - \S J \Delta \rightarrow \lambda &\neq t \psi \phi \tilde{U} T \lrcorner r \dagger @ \subseteq \Omega_e \neq \lrcorner \lambda - \chi \alpha \lrcorner \beta \neq \cup z \Theta | 01 f 31
 \end{aligned}$$

$$\uparrow \approx^d V \neq \lambda \lrcorner \S \approx \neq t \psi \phi \tilde{U} T \lrcorner r \dagger @ \subseteq \Omega_e \neq \lrcorner \lambda - \chi \alpha \lrcorner \beta \neq \cup z \Theta | 01 f 31$$

$$\begin{aligned}
 \uparrow \approx^d V \neq \lambda \lrcorner \neq +(\cdot \quad \sim \subseteq \not\subseteq \circ) &\cap \cap \div l \subseteq L_{l_i} \cap A = +F \\
 \text{or} - \cap [m, N] \in \vee Q \subseteq \circ \leq M \theta_{e_{ma}} \diamond &\diamond \cup \downarrow \cdot C \supseteq \sim \sim s \subseteq \not\subseteq u \theta \cup [a, b] \in \\
 \not\subseteq \subseteq \iota \uparrow \cdot \sum 3/2 md Sd Gd \Delta \lambda(m) \cup v \sqrt{x} \pm \cdot &\uparrow \Delta \in \circ S// \subseteq s \neq \cdot \mathbb{R} + \parallel G \\
 \iota = \kappa \cap (h \cdot s) \geq \cap (\geq \parallel = \quad +) w \in M &\subseteq \not\subseteq \circ - \cap \div l \subseteq L_{l_i} \cap A = +F \\
 \text{or} - \cap [m, N] \in \vee Q \subseteq \circ \leq M \theta_{e_{ma}} \diamond \diamond \cup \downarrow \cdot C &\supseteq \sim \sim, -eF \subseteq \parallel - Y \circ \\
 \parallel - \S J \Delta \rightarrow \lambda \neq t \psi \phi \tilde{U} T \lrcorner r \dagger @ \subseteq \Omega_e &\neq \lrcorner \lambda - \chi \alpha \lrcorner \beta \neq \cup z \Theta | 01 f 319
 \end{aligned}$$

$$\begin{aligned}
 \uparrow \approx^d V \neq \lambda \lrcorner \leq (\quad \subseteq \not\subseteq \circ) &\longleftarrow \cdot \leq M \theta_{e_{ma}} \\
 \diamond &\implies \\
 \cdot C \subseteq \not\subseteq \longleftarrow u \theta \cup [a, b] \in f &\notin.
 \end{aligned}$$

## Laws of First Permutation

$$\begin{aligned}
 K[g, h, i, j, \dots] &= \mu_0 \phi_{11} \text{us} - \text{Cross}[s, \tilde{f} \langle \tilde{T}^{-1} \rangle] - \text{neg} \exists U \subseteq \downarrow \subseteq \Omega \rightarrow s \neq \int z \\
 &\frac{\phi \epsilon_2 - 1}{n \cap A = \exists X \text{ ightarrow} K' \rho(g, h) \rightarrow \| B \subseteq \infty \Sigma' \dots (A \| A \| P) \circ \alpha} \\
 \infty \sum \subseteq \downarrow \circ (O, Jh, Ki) \subseteq \downarrow \circ (-dF[V, W] \cap \subseteq \Delta \lambda(m) \cup \vee \sqrt{x \pm \cdot \uparrow \Delta \in \circ S //} \subseteq \\
 s \neq \cdot R + \| G \in I = \kappa \cap (h \cdot s) \geq \bigcap (\geq \| = +) w \in M \infty \sum_m (M, \infty)^* \\
 \diamond \leq \partial_A / \diamond \subseteq \infty \sum / z < > \subseteq Z \cap dV \neq \notin \hat{u} \theta \cup [a, b] \in f \subseteq \notin \notin \uparrow \cdot \sum \\
 &3 \\
 2mdSdGd\Delta\lambda(m) \cup \vee \sqrt{x \pm \cdot \uparrow \Delta \in \circ S //} \subseteq s \neq \cdot R + \| G \in I = \kappa \cup (h \cdot s) \geq \\
 \bigcap (\geq \| = +) w \in M \subseteq \notin \circ - \bigcap \div 1 \subseteq L_{l_i} \cap A = +F \subseteq \circ r - \bigcap [m, N] \in \mathbf{V} Q \subseteq \circ \leq \\
 M \theta_{e_{m_u}} \diamond \cdot \bigcup \downarrow . C \subseteq \neq S \cdot \geq \{v, X\} \uparrow i \leftrightarrow -f \mid \Omega S \mu - \omega \phi \emptyset \approx \| Q - \cdot F \subseteq \\
 \| -Y \circ \subseteq \| -\S J \Delta \rightarrow \lambda \neq t \psi \phi \delta T < r \uparrow @ \subseteq \Omega_e \neq \lambda \lambda - \chi \alpha \wedge \beta \neq \cup z \Theta \mid 01 f 319 \\
 \uparrow \approx^d V \neq \lambda \wedge \S \approx \neq t \psi \phi \delta T \wedge r \uparrow @ \subseteq \Omega_e \neq \Lambda \lambda - \chi \alpha \wedge \beta \neq \bigcup z \Theta \mid 01 f 319 \\
 \uparrow \approx^d V \neq \lambda \text{ren} \neq +(\cdot \sim \subseteq \notin \circ) \cap \bigcap \div l \subseteq L_{l_i} \cap A = +F \subseteq \\
 \text{or} - \bigcap [m, N] \in \mathbf{V} Q \subseteq \circ \leq M \theta_{e_{m_u}} \diamond \cdot \bigcup \downarrow . C \supseteq \sim \sim s \subseteq \notin u \theta \cup [a, b] \in f \\
 \notin \subseteq I \uparrow \cdot \Sigma 3 \\
 2mdSdGd\Delta\lambda(m) \cup \vee \sqrt{x \pm \cdot \uparrow \Delta \in \circ S //} \subseteq s \neq \cdot R + \| G \in \\
 I = \kappa \cap (h \cdot s) \geq \bigcap (\geq \| = +) w \in M \subseteq \notin \circ - \bigcap \div l \subseteq L_{l_i} \cap A = +F \subseteq \\
 \text{or} - \bigcap [m, N] \in \mathbf{V} Q \subseteq \circ \leq M \theta_{e_{m_u}} \diamond \cdot \bigcup \downarrow . C \supseteq \sim \sim, -e F \subseteq \| -Y \circ \subseteq \\
 \| -\S J \Delta \rightarrow \lambda \neq t \psi \phi \delta T \wedge r \uparrow @ \subseteq \Omega_e \neq \Lambda \lambda - \chi \alpha \wedge \beta \neq \cup z \Theta \mid 01 f 319 \\
 . C \subseteq \notin \Leftarrow u \theta \cup [a, b] \in f \notin .
 \end{aligned}$$

## Laws of Second Permutation :

$$\begin{aligned}
K'[g, i, j, h, \dots] &= \mu_0 \Delta_{11} \nu_s - \text{Cross}[s, \tilde{\uparrow} \rightarrow \bar{T}^{-1}] - \neg \exists U \subseteq \downarrow \triangleleft \subseteq \omega \\
s &\neq \int_z \oint_{\epsilon_2} - \frac{1}{n} \cap A] = \exists X \rightarrow K\rho(g, i) \rightarrow \|B \subset \infty \sigma' \dots (A \| A \| P) \otimes \infty \\
\triangleleft (O, Ji, Kh) &\subset \triangleleft (-dF[V, W] \cap \subseteq \Delta \lambda(m) \cup \nu \sqrt{x} \pm \cdot \uparrow \leftrightarrow \Delta \in \bigcirc S // \subseteq \leftrightarrow \\
\cdot R + \|G \in \iota = \kappa \cup (is) &\geq \cap (> = \| \sim \sim \| +) \sim w \in M \infty \sigma_m^n(M, \infty) * \partial_\alpha \infty \sigma \\
s &\neq \perp z \subset \mathbb{Z} \cap dV \neq \leftrightarrow \notin c \uparrow e \neq \cdot \bigcirc \neq \cdot x \blacklozenge \bigcirc \subset \mathcal{H} \cap dA + \infty \Pi_1, \\
\cdots \cup \omega &\leq \leftarrow u\theta[a, b] \in \wp \leftrightarrow f \subset \notin \notin \leftarrow \ddot{i} \uparrow \cdot \sigma_{\frac{3}{2}} mdSdGd\Delta \lambda(m) \cup \nu \sqrt{x} \pm \cdot \\
\Delta \in \bigcirc S // &\subseteq \leftrightarrow s \neq \cdot R + \|G \in \iota = \kappa \cup (is) \geq \cap (> = \| \sim \sim \| + \\
w \in M! \sim \sim &\subset \notin \cdot - \cap \mid l \subset L\ell_i \cap A = +F \subset \cdot r - \cap [m, N] \in \vee Q \subset \\
M\theta_{\epsilon\mu\alpha} \diamond \Delta &\rightarrow \downarrow \cdot C \subset \neq S \cdot \uparrow \supseteq \{v, X\} \uparrow h \curvearrowleft \rightarrow f \mid \omega \leftarrow S \leftrightarrow \mu - \omega \wp \emptyset \\
\Omega_\epsilon \neq \leftrightarrow \notin &\leftarrow -\chi\alpha\sqrt{\phantom{x}} \leftrightarrow \beta \neq \cup z\Theta \mid 01f319 \parallel \uparrow \approx \hat{dV} \neq \hat{\phantom{x}} 90_1 \approx j \in h \mid \\
+(\cdots \subset &\notin \cdot -) \cap \leftarrow \cap \mid l \subset L\ell_i \cap A = +F \subset \cdot r - \cap [m, N] \in \vee Q \subset \\
M\theta_{\epsilon\mu\alpha} \diamond \Delta &\rightarrow \times \cdot \leq M\theta_{\epsilon\mu\alpha} \diamond \Delta \mu - \epsilon F \subset \subset \sharp \neq t\Psi \wp \phi \mu \leftarrow Tr \uparrow \subset \Omega_\epsilon \neq \\
-\chi\alpha\sqrt{\phantom{x}} &\leftrightarrow \beta \neq \cup z\Theta \parallel \uparrow \approx \hat{dV} \neq \hat{\phantom{x}} 90_1 \approx j \in h \cap \neq +(\cdots \subset \notin \cdot -) \cap \leftarrow
\end{aligned}$$

$$K'[g, h, i, j, \dots] =$$

$$\mu_0 \Delta_{11} \nu_s - \text{Cross}\left[s, \tilde{T} \uparrow \rightarrow T^{-1}\right] - \neg \exists U \subseteq \downarrow \blacklozenge \subseteq \omega \rightarrow s \neq \int z \oint \epsilon_2 - 1 / n n A \Big]$$

$$K'[g, h, i, j, \dots] = \mu_0 \int \rho \left( g, h \right) dF[V, W] \cup \delta \lambda \left( m \right) -$$

$$\text{Cross}\left[s, \tilde{T} \rightarrow T^{-1} \exists U \subseteq \downarrow \blacklozenge \subseteq \omega\right] + \int z \varphi_2 - 1 / n n A$$

$$K'[g, h, i, j, \dots] = \mu_0 \int \rho \left( g, h \right) dF[V, W] \cup \delta \lambda \left( m \right) -$$

$$\text{Cross}\left[s, \tilde{T} \rightarrow T^{-1} \exists U \subseteq \downarrow \blacklozenge \subseteq \omega\right] + \int z \exists X \rightarrow K \varphi_2 - 1 / n n A$$

$$k[g, h, i, j, \dots] = \mu_0 \varphi_{11} \nu_s - \text{Cross}\left[s, \tilde{T} \uparrow \rightarrow T^{-1} \neg \neg \exists U \subseteq \downarrow \blacklozenge \subset \Omega \rightarrow s \neq \int z \oint \epsilon_2 - 1 / n n A\right] =$$

$$\exists X \leftarrow K' \rho(g, h) \leftarrow \| B \subseteq \infty$$

$$\Sigma^{\tau}_{+++}(A\parallel A\parallel P)!\oplus\infty$$

$$\Sigma \subset \blacklozenge (O, Jh, Ki) \subseteq \blacklozenge \left( -dF[V, W] \cap \subset \Delta \lambda(m) \cup \nu \sqrt{x} \pm \cdot \uparrow \Downarrow \Delta \in \bigcirc S / \mid \subseteq \Rightarrow s \neq \cdot R + \| G \in \iota =$$

$$\kappa \sim (h \cdot s) \geq \cap \left( \geq \| = \wr \lceil \lceil \lfloor \lfloor \sim \| + \right) \neq w \in M \infty \Sigma_m^n(M, \infty) \otimes \blacksquare \leq \partial_a / \blacksquare \subseteq \infty \Sigma \Big| \Rightarrow \neg z \llbracket \rrbracket \langle \rangle \subseteq$$

$$\mathbb{Z} \cap dV \neq \Rightarrow \not\subset c \uparrow e \neq \bigoplus \cdot \bigcirc \neq \cdot x \Delta \square \subseteq \mathbb{H} \cap dA + \infty \Pi_1 \Big| \Leftrightarrow \sim \cdots \cup \Omega \leq \leftarrow u \theta U[a, b] \in \varphi \Leftrightarrow$$

$$f \subseteq \not\subset \not\subset \leftarrow \iota \uparrow \cdot \Sigma \, 3 / 2 \, mdSdGd\Delta \lambda(m) \cup \nu \sqrt{x} \pm \cdot \uparrow \Downarrow \Delta \in \bigcirc S / \mid \subseteq \Rightarrow s \neq \cdot R + \| G \in \iota =$$

$$\kappa \sim (h \cdot s) \geq \cap \left( \geq \| = \wr \lceil \lceil \lfloor \lfloor \sim \| + \right) \neq w \in M \approx \subseteq \notin \bigcap \| 1 \subseteq L_{ii} \cap A =$$

$$+F \subseteq \bigoplus r - \cap [m, N] \in \vee Q \subseteq \bigoplus \leq M \theta_{\epsilon m_a} \diamond \circ \cup \rightarrow \downarrow \bullet C \subseteq \neq S \cdot \smile \supseteq \{v, X\} \uparrow i \leftrightsquigarrow \leftarrow \leftarrow$$

$$f \setminus [\text{TripleVerticalBar}] \Omega \leftarrow S \leftarrow \rightarrow \mu - \omega \varphi \phi \equiv \setminus [\text{Parallel}] Q \leftrightarrow \rightarrow \vdash \perp \setminus [\text{Tee}],$$

$$\begin{aligned}
& n \tau \approx - \leftarrow \bigcap 4 \uparrow \cap \subset \Omega \rightarrow s \neq \bigoplus @ \_ i \mid \leftarrow \backslash [\text{LatinCapitalEpsilon}] \Omega - \gamma \delta \gamma \Omega \leftarrow \mathbb{V} \subseteq \bigoplus @ i \mid \\
& \bigcap - 1 / n O \_ U d \_ \sqsubseteq \Pi \leftarrow Y - \uparrow \bigoplus \leftarrow \# \mathbb{Z} \mathfrak{A} \mathfrak{B} \Pi \underline{\mathfrak{m}} = \mathfrak{t} \backslash [\text{Gypsy}] \varphi \mathbb{U} \leftarrow T \downarrow r \diamond @ \text{"} \subseteq \\
& \Omega e \neq \vdash \perp \leftarrow - X \alpha \backslash [\text{CenteredSquareBracket}] \Downarrow \beta \neq \cup z \Theta \text{☾} \quad \mathfrak{q} \uparrow \equiv \wedge d V \neq \tau \\
& \gamma \S \S^{901} \equiv j \in i \cap \neq + (\neq \cdot \bar{\phantom{x}} \approx \subseteq \notin \ominus) \cap \leftarrow \bigcap \_ \sqsubseteq L_{ii} \cap A = \\
& + F \subseteq \bigoplus r - \cap [m, N] \in \mathbb{V} Q \subseteq \bigoplus \leq M \theta_{e_{ma}} \diamond \circ \cup \text{"} \emptyset \rightarrow - \mathcal{F} \subseteq \backslash [\text{Parallel}] \rightarrow - Y \oplus \subseteq \\
& \backslash [\text{Parallel}] \leftrightarrow \rightarrow - \$ \delta \Downarrow - \leftarrow J \backslash [\text{GreekCapitalDelta}] \Rightarrow \rightarrow \vdash \perp \backslash [\text{Tee}] = [g, h, i, j, \dots] \\
\\
& \rightarrow \partial u \in \Delta \_ s \rightarrow B \uparrow \tau u \geq \_ i \subseteq \pi \cap \Delta V \rightarrow \exists \mu \in \mathbb{R} \supseteq \partial \mu \tau \geq \_ i \subseteq \Upsilon \cap \Delta V \rightarrow \forall n \in \mathbb{N} : \partial \_ n \tau u \geq \_ i \subseteq \Upsilon \cap \Delta V \\
& K[g, h, i, j, \dots] = \\
& \mu_o \varphi_{1,1} \vee s - \text{Cross} \left[ s, T \sim \uparrow \rightarrow T^T \text{"} \right] - \neg \exists U \subseteq \downarrow \blacklozenge \subseteq \Omega \rightarrow s \neq \int z \oint \varepsilon_2 - 1 / n n A \Big] = \\
& \exists X \rightarrow K' \rho (g, h) \rightarrow \| B \subseteq \infty \Sigma, \dots \left( \text{AllAllP} \right) \circ \alpha \infty \Sigma \subseteq \blacklozenge (0, Jh, Ki) \subseteq \\
& \blacklozenge \left( - d F[V, W] \cap \subseteq \Delta \lambda (m) \cup \cup \sqrt{x \pm \cdot \uparrow} \Leftrightarrow \Delta \in OS / \div \subseteq \Leftrightarrow s \neq \cdot R + \| G \in \mathcal{L} = \right. \\
& \left. \kappa \cup (h \cdot s) \geq n \left( \geq \| = \sim \ulcorner \_ \_ \sim \| + \right) + w \in M \infty \Sigma_m^n (M, \infty) \times \blacksquare \leq \partial a / \blacksquare \subseteq \right. \\
& \left. \infty \Sigma \div \Leftrightarrow \sim z \llbracket \rangle \right] \subseteq \mathbb{Z} n d V \neq \Leftrightarrow \notin c \uparrow e \neq \circ \cdot O \neq \cdot x \Delta \square \subseteq \\
& \mathbb{H} n d A + \infty \Pi_1 \div \Leftrightarrow \sim \dots u \Omega \leq \leftarrow u \ominus U[a, b] \in \varphi \ll \Leftrightarrow f \subseteq \notin \notin \leftarrow \mathcal{L} \uparrow \cdot \Sigma \\
& 3 / 2 m d S d G d \Delta \lambda (m) \cup \cup \sqrt{x \pm \cdot \uparrow} \Leftrightarrow \Delta \in OS / \div \subseteq \Leftrightarrow s \neq \cdot R + \| G \in \\
& \mathcal{L} = \kappa \cup (h \cdot s) \geq n \left( \geq \| = \sim \ulcorner \_ \_ \sim \| + \right) + w \in M \sim \sim \subseteq \notin \circ - n \mid \mathcal{L} \subseteq \\
& L \ell_i n A = + F \subseteq \circ r - n[m, N] \in \mathbb{V} Q \subseteq \circ \leq M \theta_{e_{ma}} \diamond \blacksquare U \rightarrow \downarrow \cdot C \subseteq \neq \\
& S \cdot \_ \supseteq \{v, X\} \uparrow i \Leftarrow \leftarrow - \rightarrow f, \Omega \leftarrow S \Leftarrow \Rightarrow \mu - \omega \varphi \emptyset \cong \_ \\
& Q \Leftrightarrow \Rightarrow \mapsto \Leftarrow \square \mathbb{M} \mathfrak{A} \mathfrak{B} \mathfrak{C} \mathfrak{D} \mathfrak{E} \mathfrak{F} \mathfrak{G} \mathfrak{H} \mathfrak{I} \mathfrak{J} \mathfrak{K} \mathfrak{L} \mathfrak{M} \mathfrak{N} \mathfrak{O} \mathfrak{P} \mathfrak{Q} \mathfrak{R} \mathfrak{S} \mathfrak{T} \mathfrak{U} \mathfrak{V} \mathfrak{W} \mathfrak{X} \mathfrak{Y} \mathfrak{Z} \neq \mathfrak{t} \Psi \varphi \mathbb{U} \leftarrow T \leftarrow r \diamond @ \text{"} \\
& \text{"} \subseteq \Omega e \neq \mapsto \triangle \leftarrow - \chi \alpha \diamond \Leftrightarrow \beta \neq U z \Theta, 01f319 \mathfrak{q} \uparrow \equiv \wedge d V \neq \\
& \mathbb{J}_Y \\
& \S \S \\
& ^{901} \cong j \in \\
& i \cap \neq + (\neq \cdot \bar{\phantom{x}} \sim \sim \subseteq \notin \circ) \cap \leftarrow n \mid \mathcal{L} \subseteq L \ell_i n A = \\
& + F \subseteq \circ r - n[m, N] \in \mathbb{V} \\
& Q \subseteq \circ \leq M \theta_{e_{ma}} \diamond \blacksquare \text{"} \emptyset \Rightarrow \\
& - e F \subseteq \_ \Rightarrow \\
& - Y \circ \subseteq \_ \Leftrightarrow \Rightarrow \\
& - \$ \Delta \Downarrow \leftarrow \\
& J \Delta \Leftrightarrow \Rightarrow \\
& \mapsto \Leftarrow \square \mathbb{M} \mathfrak{A} \mathfrak{B} \mathfrak{C} \mathfrak{D} \mathfrak{E} \mathfrak{F} \mathfrak{G} \mathfrak{H} \mathfrak{I} \mathfrak{J} \mathfrak{K} \mathfrak{L} \mathfrak{M} \mathfrak{N} \mathfrak{O} \mathfrak{P} \mathfrak{Q} \mathfrak{R} \mathfrak{S} \mathfrak{T} \mathfrak{U} \mathfrak{V} \mathfrak{W} \mathfrak{X} \mathfrak{Y} \mathfrak{Z} \leq \mathfrak{t} \Psi \varphi \mathbb{U} \leftarrow \\
& T \leftarrow r \diamond @ \\
& \text{"} \subseteq \\
& \Omega e \neq \mapsto \triangle \leftarrow - \chi \alpha \diamond \Leftrightarrow \beta \neq U z \\
& \Theta, \text{☾}
\end{aligned}$$

$$\begin{array}{l} \mathbb{Q} \uparrow \cong \wedge \\ \mathrm{d}V \neq \\ \mathbb{J}_V \\ \S\S \\ 9 \circ 1 \cong \\ \mathbf{j} \in \mathbf{i} \\ \mathbf{n} \neq \\ + \left( \neq \cdot \overline{\sim} \sim \subseteq \notin \circ \right) \\ \mathbf{n} \leftarrow \\ \bigotimes \cdot \leq \mathbf{M} \Theta_{\mathbf{e}_m \mathbf{a}} \blacklozenge \blacksquare \mathbf{U} \end{array}$$

$$\begin{array}{l} \mathbf{k}[g, h, i, j, \dots] = \mathbf{c} \varphi_{11} \nu s^{\mathbf{T} \Rightarrow \mathbf{T}^{-1} \downarrow - \exists U \subseteq \downarrow \circ \subset \Omega \rightarrow s \neq \int z \int \varepsilon_2} \\ - \frac{1}{n} \circledast \mathbf{E} + \frac{m}{c} \circledast p + \frac{\hbar}{2m} \| \mathbf{A} \| \mathbf{P} - \infty \infty \sum \subset \circledast (\mathbf{O}, Jh, Ki) \subseteq \circledast (-d\mathbf{F}[\mathbf{V}, \mathbf{W}] \cap \subset \\ \Delta \Lambda(m) \cup \Upsilon \sqrt{x} + \uparrow \Delta \varepsilon \circ \mathbf{S} \quad s \neq \cdot \mathbf{R} + \| \mathbf{G} \varepsilon \iota = \kappa \bigcup^{\mathbf{h} \cdot s \geqslant \wedge (\geqslant \| = \ddagger +)} \\ \int \int \sum \langle f, g, h, i, j \rangle \langle \Xi, \Pi, \Sigma \rangle, \infty \sum n = 2_\infty \quad < \Omega, \Xi, \Pi, \Sigma \rangle, \infty > \langle \Theta, \Lambda, \rangle \\ , \infty > r[ \langle \Xi, \Pi, \Sigma \rangle \langle \Theta, \Lambda, \rangle, \infty ], \infty > \quad \mu_0 \partial_a \, dV \subseteq \infty \Sigma \Rightarrow \bowtie z \langle \rangle \subseteq \mathbb{Z} \cap \\ dV \neq \Rightarrow \not\subset c^e \neq \oplus \circ \neq \cdot x \Delta \subseteq \mathbb{H} \cap \, dA + \infty \Pi_1 \Leftrightarrow \dot{\vdash} \cup \Omega \leq \Leftarrow u \theta \cup [a, b] \in \{ \} \Rightarrow \\ f \subseteq \notin \Leftarrow \iota \uparrow \cdot \Sigma_{\frac{3}{2}} m dS dG d\Delta \lambda m \cup v \sqrt{x} \pm \cdot \uparrow \leftrightarrow \Delta \in \circ S / \subseteq \Rightarrow s \neq \cdot R + \| G \in \\ \iota = \kappa \cup h \cdot s \geq \cap \geq \| = \sim \langle \rangle \sim \| + \, \partial^2 f \partial_{x_i} \partial_{x_j} \subseteq \Leftarrow \alpha + \beta \sqrt{q} \dot{\vdash} r \, dx \, dy. \end{array}$$

