

Effects of ski piste preparation on alpine vegetation

SONJA WIPF*†, CHRISTIAN RIXEN*†, MARKUS FISCHER‡,
BERNHARD SCHMID† and VERONIKA STOECKLI*

*WSL Swiss Federal Institute for Snow and Avalanche Research SLF, Flüelastr. 11, 7260 Davos Dorf, Switzerland;

†Institute of Environmental Sciences, University of Zurich, Winterthurerstr. 190, 8057 Zurich, Switzerland; and

‡Institute of Biochemistry and Biology, University of Potsdam, Villa Liegnitz, Lennéstr. 7a, 14471 Potsdam, Germany

Summary

1. Ski resorts increasingly affect alpine ecosystems through enlargement of ski pistes, machine-grading of ski piste areas and increasing use of artificial snow.

2. In 12 Swiss alpine ski resorts, we investigated the effects of ski piste management on vegetation structure and composition using a pairwise design of 38 plots on ski pistes and 38 adjacent plots off-piste.

3. Plots on ski pistes had lower species richness and productivity, and lower abundance and cover of woody plants and early flowering species, than reference plots. Plots on machine-graded pistes had higher indicator values for nutrients and light, and lower vegetation cover, productivity, species diversity and abundance of early flowering and woody plants. Time since machine-grading did not mitigate the impacts of machine-grading, even for those plots where revegetation had been attempted by sowing.

4. The longer artificial snow had been used on ski pistes (2–15 years), the higher the moisture and nutrient indicator values. Longer use also affected species composition by increasing the abundance of woody plants, snowbed species and late-flowering species, and decreasing wind-edge species.

5. *Synthesis and applications.* All types of ski piste management cause deviations from the natural structure and composition of alpine vegetation, and lead to lower plant species diversity. Machine-grading causes particularly severe and lasting impacts on alpine vegetation, which are mitigated neither by time nor by revegetation measures. The impacts of artificial snow increase with the period of time since it was first applied to ski piste vegetation. Extensive machine-grading and snow production should be avoided, especially in areas where nutrient and water input are a concern. Ski pistes should not be established in areas where the alpine vegetation has a high conservation value.

Key-words: artificial snow, biodiversity, functional groups, machine-grading, snow ecology, Switzerland

Journal of Applied Ecology (2005) **42**, 306–316

doi: 10.1111/j.1365-2664.2005.01011.x

Introduction

The structure of alpine vegetation is affected by altitude, aspect and inclination. Some alpine habitats are extraordinarily rich in species, while communities in the highest and most extreme regions consist of only a few specialists (Ellenberg 1988). Furthermore, most communities are characterized by specific proportions of functional groups, such as dwarf shrubs, grasses and herbs. The communities can consist of species from special eco-

logical groups, for example snowbed species, indicating depressions with a long-lasting snow cover and a very short vegetation period, or wind-edge species, indicating a long vegetation period and extreme winter temperatures (Körner 1999).

Alpine ecosystems are sensitive and susceptible to changes in land use and climate (Chapin & Körner 1995; Fischer & Wipf 2002; Laiolo *et al.* 2004). Perturbations of the alpine habitats and changes in snow cover characteristics, for example through skiing, are likely to have impacts on species composition, diversity and productivity of alpine vegetation, which in turn may have negative effects on ecosystem functioning and stability (Tilman 1996; Mulder, Uliassi & Doak 2001). As ski tourism is

Correspondence: Sonja Wipf, Swiss Federal Institute for Snow and Avalanche Research SLF, Flüelastr. 11, 7260 Davos Dorf, Switzerland (fax + 41 81 417 01 10; e-mail wipf@slf.ch).

one of the most important economical factors in European alpine regions (Abegg *et al.* 1997; Elsasser & Messerli 2001), the area affected by ski pistes or by constructions related to tourism is still increasing.

Skiing and ski piste preparation by snow-grooming vehicles are likely to cause mechanical damage to the vegetation and the soil (Cernusca *et al.* 1990; Rixen *et al.* 2004). Furthermore, the winter preparation of ski pistes leads to a thin and compressed snow cover with decreased temperature insulation capacity. Soils and vegetation under these ski pistes may experience temperatures lower than $-10\text{ }^{\circ}\text{C}$ (Rixen, Haeberli & Stoeckli 2004), while under undisturbed snow-packs, temperatures rarely fall below $0\text{ }^{\circ}\text{C}$ (Rixen 2002; Rixen, Stoeckli & Ammann 2003). As a result, plant species with insufficient cold hardiness and plants sensitive to mechanical stress may be damaged, resulting in shifts among functional groups and a higher proportion of unvegetated ground. In summer, piste construction measures, such as machine-grading, are carried out to remove obstacles like trees and rocks or to level rough or bumpy soil surfaces. In the process of machine-grading, the upper soil layers and the vegetation are removed or heavily damaged (Pröbstl 1990; Haeberli 1992; Bayfield 1996; Titus & Tsuyuzaki 1999; Ruth-Balaganskaya & Myllynen-Malinen 2000). Seed mixtures are often applied to accelerate revegetation, but the success of revegetation measures declines with height above sea level (a.s.l.) (Urbanska 1997).

In response to ongoing climatic changes, artificial snow production is increasingly used in most ski resorts of the world. Climate-change scenarios predict changes in seasonal snowfall patterns, with the snow season beginning later and ending earlier, and a rise in the snow line (IPCC 1996, 1998, 2001). As a consequence, the minimum altitude termed snow-secure for winter sports in the European Alps would rise from 1200 m today to 1500 m a.s.l. within the next 30 years (Abegg 1996). Winters with little snow have already been a major concern of winter sport resorts since the mid-1980s (Seilbahnen Schweiz 2001). As a strategy to minimize the dependence on natural snowfall, ski areas invest in systems for artificial snowing.

Effects of ski piste manipulation in single resorts and single effects of artificial snowing or machine-grading have been investigated in earlier studies (Urbanska 1997; Kammer 2002). In our comprehensive study, sites covered a wide range of locations, altitudes and aspects in 12 Swiss ski resorts. We investigated plots on ski pistes and adjacent control plots beside pistes. With a total of 76 vegetation records arranged in a factorial design, we addressed the following specific questions. (i) Do productivity, species richness and composition of the vegetation on ski pistes differ from the vegetation off pistes? (ii) What are the impacts of machine-grading on productivity, species richness and composition of ski piste vegetation? (iii) How does the use of artificial snow modify environmental conditions on ski pistes and how do productivity, species richness and composition

of the vegetation respond? (iv) Do the effects of artificial snow differ between graded and ungraded ski pistes?

Materials and methods

STUDY SITES AND DESIGN

We studied the vegetation of plots on and next to ski pistes in 12 ski resorts in the Swiss Alps (see Appendix 1), ranging in altitude between 1750 and 2550 m a.s.l. and representing a wide range of vegetation types (mostly alpine grasslands on acidic or calcareous bedrock and dwarf shrub heath), time since grading and years of application of artificial snow. The study sites were chosen on machine-graded or ungraded ski pistes with either natural or artificial snow, resulting in four different ski piste types: natural snow/ungraded (ns), artificial snow/ungraded (as), natural snow/graded (nsg) and artificial snow/graded (asg). Sites were selected in consultation with the local ski piste managers to ensure that the pistes had received the same management continuously in previous years.

At each study site, we randomly chose a plot on a ski piste and an undisturbed control plot in the nearest off-piste area to allow for pairwise comparisons. The distance between piste and control plot was 15–50 m and the pair of plots did not differ significantly in altitude, inclination, aspect and management. All ski pistes were prepared by snow-grooming vehicles and were used annually by skiers from approximately mid-November/early December to Easter (mid-April). Artificial snow had been produced in all ski resorts for 2–15 years ($n = 21$ pistes, mean 7.1 years). On 17 pistes in nine resorts, machine-grading (i.e. construction of ski pistes during which vegetation and topsoil are removed) had been carried out 4–30 years ago ($n = 17$ pistes, mean 18.7 years). On 10 machine-graded plots, revegetation had been encouraged by sowing seed, but details of seed mixtures and sowing techniques were not available.

VEGETATION DATA

From 1 July to 24 August 2000, we visited each site to collect vegetation records. In 4×4 -m plots, we noted the presence of each vascular plant species and estimated its cover in seven classes according to Braun-Blanquet (Müller-Dombois & Ellenberg 1974). We assigned a cover value to each class for further calculations (0.01% to 'r', 0.1% to '+', midpoint to other classes). To compare site conditions between plots, we calculated weighted indicator values for moisture (F), soil reaction (R), nutrients (N) and light (L) per plot according to Landolt (1977). For each plot, we also estimated the percentage area not covered by vascular plants.

To assess productivity, we randomly selected three patches of 0.2×0.2 m plot⁻¹, harvested the biomass 3 cm above ground and pooled the samples. In cases where plots had been grazed, we randomly sampled ungrazed

areas with similar vegetation within 5 m distance from the plot. Three grazed plots with no undisturbed vegetation nearby were omitted from biomass sampling. We measured annual productivity as the total biomass excluding litter, dead biomass and woody shoots older than 1 year. The biomass was dried at 70 °C and weighed to the nearest 0.01 g.

For each plot we calculated species number and Shannon's index of diversity as $H = -\sum P_i \ln P_i$, where P_i is the abundance class of species i . We calculated the abundance of four functional groups: graminoids (Poaceae, Cyperaceae, Juncaceae), woody plants, legumes and remaining forbs (including Pteridophyta), to obtain taxonomical and plant functional characteristics of the plots. According to their habitat preference in relation to snow cover and length of growing season (Ellenberg 1988; Wilmanns 1998; Delarze, Gonseth & Galland 1999), we classified 26 of 322 species as inhabiting snowbeds and 30 as inhabiting wind-edges (Wipf *et al.* 2002). Depending on their phenology, we also classified 67 species as early flowering and 94 as late-flowering, according to the flowering period indicated in Lauber & Wagner (2001). For the four functional, two ecological and two phenological groups, we calculated the proportions relative to total species number and total vegetation cover. See Appendix 2 for a list of species and their classification.

STATISTICAL ANALYSIS

We analysed our data with hierarchical analysis of variance (ANOVA) models. Differences in the type of bedrock, aspect, slope and hours of sunshine between pairs were taken into account by the pairwise design of piste and off-piste plots. Moreover, we considered altitude as a covariate in all analyses. To mirror the study design with pairs of plots on and next to pistes, we fitted the type of plot pairs, i.e. the type of piste treatment of the piste plot in the pair. Next we fitted the factor piste, which indicated differences between plots on and next to pistes within the pairs, across all types of piste treatment. Then, we fitted the interactions of type of pair (machine-grading or years of artificial snow) and piste, which indicated whether the difference between plots on and next to the piste depended on the type of piste treatment. We considered machine-grading, which is applied only once, as a binary factor, and artificial snowing as a continuous variable (time since start of application of artificial snow) because of its potentially cumulative impacts. Three-way interactions between resort, type of plot pair and piste indicated whether effects of machine-grading and artificial snow differed between resorts. To analyse the effect of succession and revegetation measures on the vegetation of machine-graded ski pistes, we performed the same analysis on a reduced data set (pairs of graded piste and control plots) and included years since machine-grading as a continuous variable and sown as a binary factor. If necessary, dependent variables were transformed prior to analysis to reach

normality. Residuals were checked visually for normal distribution. Analyses were carried out with SPSS 10.0.5 (SPSS 1999).

Results

IMPACTS OF SKI PISTES VS. NO PISTE

The indicator values of the vegetation composition showed an increased supply in moisture and nutrients available for plants (indicator value F, $P = 0.098$; N, $P = 0.003$; Tables 1 and 2 and Fig. 1) and a tendency for higher base content of the soil (R, $P = 0.059$; Tables 1 and 2) on ski pistes. Despite the favourable environmental conditions for growth, plant productivity on ski pistes was lower than beside the pistes (Tables 1 and 3 and Fig. 2). The mean number of species per plot was 11% lower on piste plots than on off-piste plots ($P = 0.047$; Tables 1 and 3 and Fig. 2). The pistes thus had a negative effect on the species richness in alpine grassland and dwarf shrub vegetation.

Woody plants covered 24.3% of unmanipulated control plots, but were significantly less frequent on ski pistes (10.5% cover, $P = 0.003$; Tables 1 and 4 and Fig. 3), mainly because of their low abundance on machine-graded pistes (1.7% cover, see next section). The same pattern was found in the proportion of woody species (Tables 1 and 5 and Fig. 3). Legume species were more abundant and had higher cover on piste plots than on control plots. Both cover and species number of early flowering species were lower on piste than on control plots ($P = 0.005$; Tables 1 and 6 and Fig. 4). Thus, the snow compaction changed the plot composition of functional and ecological groups.

MACHINE-GRADING OF SKI PISTES

The proportion of ground not covered by vegetation was almost five times higher on graded than on ungraded piste plots ($P = 0.008$; Tables 1 and 3 and Fig. 2) but was unaffected by revegetation measures (sowing) or time since machine-grading. Corresponding with the high proportion of ground not covered by vegetation, the light input for the plants (expressed as indicator value L) was higher on graded piste plots than on ungraded ones ($P = 0.002$; Tables 1 and 2 and Fig. 1). Nutrient availability to the vegetation (expressed as indicator value N), which was generally increased on ski pistes, reached even higher values on graded pistes than on ungraded ones ($P = 0.07$; Tables 1 and 2 and Fig. 1). Whereas the productivity on ungraded piste plots was similar to that of off-piste plots, it was reduced 4.6 times on graded piste plots ($P = 0.011$; Tables 1 and 3 and Fig. 2). Machine-graded pistes that had been revegetated by sowing supported fewer species ($P = 0.098$). The Shannon index was lower for all graded piste plots ($P = 0.03$; Tables 1 and 3 and Fig. 2).

Machine-grading also affected the vegetation composition: woody plants, which were generally less common

Table 1. The effects of ski piste treatments on the parameters describing the vegetation in 12 Swiss ski resorts: Predicted means and slopes of the correlation with years of artificial snow. The *n* in parentheses refer to the biomass measurements (*n* lower because of missing values)

	Total <i>n</i> = 76 (73) Mean	Off-piste <i>n</i> = 38 (36) Mean	Piste <i>n</i> = 38 (37) Mean	Piste graded <i>n</i> = 17 (16) Mean	Piste ungraded <i>n</i> = 21 Mean	Correlation with years of artificial snow Slope
Indicator values						
F*	2.85	2.83	2.87	2.88	2.87	0.000
R†	2.55	2.50	2.59	2.61	2.58	0.007
N‡	2.46	2.38	2.54	2.56	2.53	0.003
L§	3.80	3.76	3.85	4.03	3.71	0.01
Diversity and productivity						
Species number	36.4	38.5	34.3	31.7	36.5	0.30
Shannon index	3.6	3.8	3.4	2.8	3.8	-0.02
Annual productivity (g m ⁻²)	158.0	173.1	143.3	46.9	216.7	-0.25
Proportion of cover (%)						
Grasses	41.4	38.8	44.0	38.7	48.4	-0.93
Forbs	24.2	26.5	22.0	20.5	23.1	0.13
Woody plants	17.4	24.3	10.5	1.6	17.7	0.38
Legumes	5.1	2.9	7.4	10.6	4.8	0.15
Snowbed species	8.4	7.3	9.6	14.0	6.0	1.36
Wind-edge species	7.5	7.7	7.4	8.0	6.9	-0.27
Early flowering species	35.6	40.8	30.5	22.9	36.6	-1.26
Late-flowering species	30.5	28.0	32.9	34.1	32.0	0.03
Bare ground	11.8	7.0	16.1	28.6	6.0	0.26
Proportion of species (%)						
Grasses	27.9	27.9	27.8	27.2	28.2	-0.13
Forbs	60.2	59.8	60.6	63.2	58.5	0.09
Woody plants	11.9	13.0	10.8	6.9	14.0	-0.23
Legumes	7.8	6.6	9.1	10.9	7.6	0.21
Early flowering species	33.4	35.7	31.1	25.5	35.7	-0.71
Late-flowering species	31.0	29.9	32.2	32.5	31.9	0.25

*F = 2, medium dryness; 3, medium humidity.

†R = 2, acid soils; 3, neutral or weakly alkaline soils.

‡N = 2, nutrient-poor soils; 3, medium- to nutrient-poor soils.

§L = 3, half-shaded conditions; 4, full-light conditions.

Table 2. ANOVA table for the indicator values of the vegetation for moisture (F), soil acidity (R), nutrient (N) and light availability (L) measured on 38 ski piste plots and 38 corresponding off-piste plots in 12 Swiss ski resorts. The type of piste treatment (machine-grading and years of artificial snowing) was fitted for pairs of piste/off-piste plots. The factor ski piste indicates the difference between piste and off-piste plots. The dependent variables were (ln(*x* + 1))-transformed prior to analysis. Three plots were excluded from biomass sampling. The sample size for productivity is therefore 73, and the degrees of freedom are indicated in parentheses. MS, mean squares; *F*, variance ratio; subscripts refer to source of variation (*)*P* < 0.1, **P* < 0.05, ***P* < 0.01, ****P* < 0.001

Source of variation	Skeleton analysis						
	MS	<i>F</i>	d.f.	Variance ratio			
				F-value	R-value	N-value	L-value
Altitude	MS _E	MS _E /MS _{Rest}	1	3.14	5.87(*)	70.23**	37.19**
Resort	MS _R	MS _R /MS _{Rest}	11	3.34	4.87(*)	4.43(*)	1.83
Type of plot pair	MS _T	MS _T /MS _{RT}	3	0.16	0.50	0.25	0.75
Piste	MS _P	MS _P /MS _{RP}	1	3.27(*)	4.42(*)	14.64**	9.46*
Grading × piste	MS _{LP}	MS _{LP} /MS _{RLP}	1	0.32	2.68	4.51(*)	24.36**
Duration of snowing × piste	MS _{AP}	MS _{AP} /MS _{RAP}	1	3.68(*)	4.80(*)	3.26(*)	0.13
Grading × duration of snowing × piste	MS _{LAP}	MS _{LAP} /MS _{Rest}	1	1.02	0.03	0.03	0.06
Resort × type of plot pair	MS _{RT}	MS _{RT} /MS _{Rest}	23	1.42	1.52	3.97(*)	0.87
Resort × piste	MS _{RP}	MS _{RP} /MS _{Rest}	11	0.51	0.45	0.93	0.24
Resort × grading × piste	MS _{RLP}	MS _{RLP} /MS _{Rest}	7	0.37	0.44	1.31	0.18
Resort × duration of snowing × piste	MS _{RAP}	MS _{RAP} /MS _{Rest}	11	0.34	0.51	0.44	0.15
Rest	MS _{Rest}		4				
Total			76				




Fig. 1. Differences in indicator values between ski piste plots and corresponding control plots. (a) Mean differences in indicator values for the four ski piste treatments (ns, ungraded pistes with natural snow; as, ungraded pistes with artificial snow; nsg, graded pistes with natural snow; asg, graded pistes with artificial snow; F, moisture indicator value; R, reaction (soil acidity) indicator value; N, nutrient indicator value; L, light indicator value). (b-d) The relationship between the differences in (b) F (moisture), (c) R (soil acidity) and (d) N (nutrient) indicator values and the time since conversion to artificial snow.




Fig. 2. Differences in productivity and diversity between ski piste plots and corresponding control plots. (a) Mean differences in productivity, species number, Shannon index and the proportion of open ground for the four ski piste treatments (ns, ungraded pistes with natural snow; as, ungraded pistes with artificial snow; nsg, graded pistes with natural snow; asg, graded pistes with artificial snow). (b) The relationship between the differences in annual productivity and the time since conversion to artificial snow.




Fig. 3. Differences in species composition between ski piste plots and corresponding control plots. Mean differences in the percentage cover of the four functional groups for the four ski piste treatments (ns, ungraded pistes with natural snow; as, ungraded pistes with artificial snow; nsg, graded pistes with natural snow; asg, graded pistes with artificial snow).

on piste plots, were reduced by 91% on graded piste plots in comparison with ungraded ones ($P = 0.03$; Tables 1 and 4 and Fig. 3). Whereas only 6.9% of the species on graded piste plots were woody species, they accounted

Table 3. ANOVA table for the proportion of bare ground, annual productivity, species number and Shannon index measured on 38 ski piste plots and 38 corresponding off-piste plots in 12 Swiss ski resorts. The type of piste treatment (machine-grading and years of artificial snowing) was fitted for pairs of piste/off-piste plots. The factor ski piste indicates the difference between piste and off-piste plots. Three plots were excluded from biomass sampling. The sample size for productivity therefore is 73, and the degrees of freedom are indicated in parentheses. (*) $P < 0.1$, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$

Source of variation	Skeleton analysis			Variance ratio			
	MS	F	d.f.				
				Bare ground (%)	Productivity	Species	Shannon number
Altitude	MS _E	MS _E /MS _{Rest}	1	254.41***	79.96**	5.59(*)	0.58
Resort	MS _R	MS _R /MS _{Rest}	11	9.53*	2.13	5.06(*)	8.00*
Type of plot pair	MS _T	MS _T /MS _{RT}	3	3.18	4.32(*)	1.33	1.62
Piste	MS _P	MS _P /MS _{RP}	1	8.81*	9.78**	5.02*	4.21(*)
Grading × piste	MS _{LP}	MS _{LP} /MS _{RLP}	1	13.64**	10.71*	0.35	7.27*
Duration of snowing × piste	MS _{AP}	MS _{AP} /MS _{RAP}	1	0.02	0.01	0.23	0.26
Grading × duration of snowing × piste	MS _{LAP}	MS _{LAP} /MS _{Rest}	1	2.02	1.83	9.81*	9.99*
Resort × type of plot pair	MS _{RT}	MS _{RT} /MS _{Rest}	23	7.74*	1.01	1.90	2.21
Resort × piste	MS _{RP}	MS _{RP} /MS _{Rest}	11	8.18*	0.90	1.09	2.68
Resort × grading × piste	MS _{RLP}	MS _{RLP} /MS _{Rest}	7	6.05(*)	0.93	1.16	1.85
Resort × duration of snowing × piste	MS _{RAP}	MS _{RAP} /MS _{Rest}	11 (9)	2.74	1.60	0.85	2.27
Rest	MS _{Rest}		4 (3)				
Total			76 (73)				

Table 4. ANOVA table for the proportion of the four functional groups, grasses, woody plants, forbs and legumes, at the plot area measured on 38 ski piste plots and 38 corresponding off-piste plots in 12 Swiss ski resorts. The type of piste treatment (machine-grading and years of artificial snowing) was fitted for pairs of piste/off-piste plots. The factor ski piste indicates the difference between piste and off-piste plots. Three plots were excluded from biomass sampling. The sample size for productivity therefore is 73, and the degrees of freedom are indicated in parentheses. (*) $P < 0.1$, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$

Source of variation	Skeleton analysis			Variance ratio			
	MS	F	d.f.				
				Cover grass (%)	Cover woody plants (%)	Cover forbs (%)	Cover legumes (%)
Altitude	MS _E	MS _E /MS _{Rest}	1	40.49**	0.30	3.28	0.44
Resort	MS _R	MS _R /MS _{Rest}	11	6.37*	3.76	6.70*	9.95*
Type of plot pair	MS _T	MS _T /MS _{RT}	3	1.02	1.58	0.61	2.38
Piste	MS _P	MS _P /MS _{RP}	1	1.76	13.87**	2.47	5.39*
Grading × piste	MS _{LP}	MS _{LP} /MS _{RLP}	1	0.72	7.67*	0.00	1.54
Duration of snowing × piste	MS _{AP}	MS _{AP} /MS _{RAP}	1	29.86***	9.20*	0.69	0.20
Grading × duration of snowing × piste	MS _{LAP}	MS _{LAP} /MS _{Rest}	1	0.00	0.12	0.99	0.81
Resort × type of plot pair	MS _{RT}	MS _{RT} /MS _{Rest}	23	5.21(*)	1.01	2.69	4.66(*)
Resort × piste	MS _{RP}	MS _{RP} /MS _{Rest}	11	3.23	0.70	1.77	5.30(*)
Resort × grading × piste	MS _{RLP}	MS _{RLP} /MS _{Rest}	7	1.53	1.19	1.02	2.83
Resort × duration of snowing × piste	MS _{RAP}	MS _{RAP} /MS _{Rest}	11	0.26	0.40	0.54	4.14(*)
Rest	MS _{Rest}		4				
Total			76				

for 14% of the species on ungraded piste plots ($P = 0.02$; Tables 1 and 5). The reactions of forb and grass species were not uniform: in six resorts the proportion of forb species was higher on graded compared with ungraded pistes (resort × grading × piste, $P = 0.02$). The proportion of grass species depended on the time since machine-grading: compared with ungraded control plots, the proportion was higher in recently graded pistes, and lower the older the graded pistes were ($P = 0.016$). Species that flower early in the year were reduced in abundance by 28% ($P = 0.018$; Tables 1 and 6) and covered 37%

less ground on graded piste plots than on ungraded ones ($P = 0.004$; Tables 1 and 6).

ARTIFICIAL SNOW

The longer artificial snow had been applied on ski pistes, the higher the moisture and nutrient availability, and the lower the soil base content (indicator values F, N and R: $P = 0.081$, 0.099 and 0.051, respectively; Tables 1 and 2 and Fig. 1). However, we could not detect any effects of artificial snow on plant productivity (Fig. 2).

Table 5. ANOVA table for the proportion of the four functional groups, grasses, woody plants, forbs and legumes, of the total species number per plot measured on 38 ski piste plots and 38 corresponding off-piste plots in 12 Swiss ski resorts. The type of piste treatment (machine-grading and years of artificial snowing) was fitted for pairs of piste/off-piste plots. The factor ski piste indicates the difference between piste and off-piste plots. The dependent variables of were $(\ln(x + 1))$ -transformed prior to analysis. Three plots were excluded from biomass sampling. The sample size for productivity therefore is 73, and the degrees of freedom are indicated in parentheses. (*) $P < 0.1$, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$

Source of variation	Skeleton analysis				Variance ratio			
	MS	MS	d.f.	Grass species (%)	Woody species (%)	Forb species (%)	Legume species (%)	
Altitude	MS _E	MS _E /MS _{Rest}	1	2.72	0.08	22.07**	8.59*	
Resort	MS _R	MS _R /MS _{Rest}	11	11.33*	2.01	18.79**	2.44	
Type of plot pair	MS _T	MS _T /MS _{RT}	3	0.29	0.45	0.26	1.99	
Piste	MS _p	MS _p /MS _{RP}	1	0.30	5.72 *	0.24	7.31*	
Grading × piste	MS _{LP}	MS _{LP} /MS _{RLP}	1	1.28	10.18 *	0.31	3.19	
Duration of snowing × piste	MS _{AP}	MS _{AP} /MS _{RAP}	1	0.02	0.01	0.04	0.08	
Grading × duration of snowing × piste	MS _{LAP}	MS _{LAP} /MS _{Rest}	1	0.04	0.09	3.22	0.01	
Resort × type of plot pair	MS _{RT}	MS _{RT} /MS _{Rest}	23	3.03	2.36	11.63*	1.11	
Resort × piste	MS _{RP}	MS _{RP} /MS _{Rest}	11	6.93*	0.90	6.21	0.36	
Resort × grading × piste	MS _{RLP}	MS _{RLP} /MS _{Rest}	7	4.19(*)	0.58	17.73**	1.13	
Resort × duration of snowing × piste	MS _{RAP}	MS _{RAP} /MS _{Rest}	11	3.44	0.79	11.84*	0.53	
Rest	MS _{Rest}	MS _E /MS _{Rest}	4					
Total			76					

Table 6. ANOVA table for the proportion of snowbed and wind-edge, as well as early flowering and late-flowering species, of the total vegetation cover in a plot (%) measured on 38 ski piste plots and 38 corresponding off-piste plots in 12 Swiss ski resorts. The type of piste treatment (machine-grading and years of artificial snowing) was fitted for pairs of piste/off-piste plots. The factor ski piste indicates the difference between piste and off-piste plots. Three plots were excluded from biomass sampling. The sample size for productivity therefore is 73, and the degrees of freedom are indicated in parentheses. (*) $P < 0.1$, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$

Source of variation	Skeleton analysis				Variance ratio			
	MS	MS	d.f.	Snowbed species (%)	Wind-edge species (%)	Early flowering (%)	Late-flowering (%)	
Altitude	MS _E	MS _E /MS _{Rest}	1	85.42***	34.45**	51.38**	5.29(*)	
Resort	MS _R	MS _R /MS _{Rest}	11	3.45	3.79	2.45	1.93	
Type of plot pair	MS _T	MS _T /MS _{RT}	3	2.18	1.04	1.24	0.61	
Piste	MS _p	MS _p /MS _{RP}	1	3.14	0.07	13.90**	3.92(*)	
Grading × piste	MS _{LP}	MS _{LP} /MS _{RLP}	1	0.29	0.01	17.93**	2.12	
Duration of snowing × piste	MS _{AP}	MS _{AP} /MS _{RAP}	1	13.19**	6.68*	5.68*	0.09	
Grading × duration of snowing × piste	MS _{LAP}	MS _{LAP} /MS _{Rest}	1	0.43	0.84	7.35(*)	1.21	
Resort × type of plot pair	MS _{RT}	MS _{RT} /MS _{Rest}	23	4.09(*)	0.83	2.87	2.13	
Resort × piste	MS _{RP}	MS _{RP} /MS _{Rest}	11	0.79	0.22	1.68	0.74	
Resort × grading × piste	MS _{RLP}	MS _{RLP} /MS _{Rest}	7	2.77	2.28	0.86	0.72	
Resort × duration of snowing × piste	MS _{RAP}	MS _{RAP} /MS _{Rest}	11	1.56	0.55	1.20	1.74	
Rest	MS _{Rest}	MS _E /MS _{Rest}	4					
Total			76					

The impact of artificial snow on diversity was ambiguous. While the use of artificial snow had negative effects on species number and Shannon index on graded pistes, the opposite was found for ungraded pistes (three-way interaction between piste, grading and duration of snowing, $P = 0.034$; Tables 1 and 3).

The application of artificial snow affected the relative proportion of plant functional groups. The longer artificial snow had been used, the more it had reduced the negative effect of the ski piste treatment on the

cover of woody plants per plot ($P = 0.011$; Tables 1 and 4). However, the proportion of a plot covered by grass species declined the longer a ski piste had been treated with artificial snow ($P < 0.001$; Tables 1 and 4). The effect of artificial snow on the species proportion of forbs and the cover of legumes was not uniform but differed between ski resorts (three-way interaction resort × artificial snow × ski piste, $P = 0.014$ and $P < 0.083$, respectively; Tables 5 and 6). The longer artificial snow had been applied to a ski piste, the higher the cover of




Fig. 4. Differences in ecological groups between ski piste plots and corresponding control plots. (a) Mean differences in the proportion of plot covered by ecological groups for the four ski piste treatments (ns, ungraded pistes with natural snow; as, ungraded pistes with artificial snow; nsg, graded pistes with natural snow; asg, graded pistes with artificial snow). (b–e) The relationship between the differences in (b) percentage cover of snowbed species, (c) percentage cover of wind-edge species, (d) proportion of early flowering species and (e) proportion of late-flowering species and the time since conversion to artificial snow.

snowbed species ($P = 0.004$; Tables 1 and 6 and Fig. 4). Wind-edge species showed the opposite reaction. They covered less ground the longer a ski piste had been treated with artificial snow ($P = 0.024$; Tables 1 and 6 and Fig. 4). The longer artificial snow had been applied, the lower the cover of early flowering plants on piste plots ($P = 0.036$; Tables 1 and 6 and Fig. 4). On the other hand, the relative number of late-flowering species tended to increase the longer artificial snow had been used ($P < 0.096$; Tables 1 and 6 and Fig. 4). Overall, the potentially snow-sensitive groups were affected most by the production of artificial snow.

Discussion

IMPACTS OF SKI PISTES VS. NO PISTE

The changes in indicator values for soil nutrients, moisture and reaction (base content) under ski pistes may be considered beneficial for plant growth (in terms of enhancing growth). However, other non-beneficial factors, including disturbance, seem to prevail because overall productivity and species richness on ski pistes are decreased. The negative effects may be the direct result of disturbance of the vegetation via snow

compaction under skiing and via mechanical abrasion. Although some types of disturbance can increase biodiversity by suppressing dominant species in productive environments (Connell 1979; Vujnovic, Wein & Dale 2002), abiotic stress reduces productivity and dominance in alpine habitats and therefore reverses any positive effects of disturbance on plant diversity (Kammer & Möhl 2002).

The changes in the cover of woody plants on the ski pistes may also be explained by mechanical disturbance. Woody plants (mainly dwarf shrubs of the *Ericaceae*) are particularly sensitive to mechanical injuries because of their size, longevity and vulnerability of their hibernation buds (Körner 1999).

In contrast to woody plants, legumes were more abundant on pistes than next to pistes, perhaps because of decreased competition from shrubs. Legumes may have been responsible for the increased nutrient indicator values on piste plots as a consequence of their ability to fix nitrogen. The reduced abundance of early flowering species on pistes in our study was most probably the result of a narrowed temporal niche for those species. In field surveys, the flowering time of early seasonal species was postponed after soil frosts on pistes with natural snow (Baiderin 1981; Köck *et al.* 1989). A delay in flowering often decreases fecundity and, consequently, can negatively affect the abundance of early flowering species (Kudo 1993; Stanton & Galen 1997).

IMPACTS OF MACHINE-GRADING

Vegetation responses were highly pronounced on graded pistes, most probably because machine-grading of ski pistes is a drastic measure where soil and plants are removed in summer during active plant growth. There was a strong compositional shift from woody plants to grasses or forbs, the extent of which depended partly on the resort and partly on the successional stage of the ski piste. The increased indicator values for nutrients and light may be the result of the disturbance itself (through mobilized nutrients and opened vegetation cover), or of characteristics of the following pioneer species, which often grow with high light and nutrient levels (Grime & Jarvis 1975; Grime 1977). Graded ski pistes with high proportions of bare ground are particularly prone to increased surface runoff and erosion during heavy rain (Löhmannsröben & Cernusca 1987). Revegetation processes at high altitudes are slow, even if revegetation measures such as sowing and planting are applied (Bayfield, Urquart & Rothery 1984; Urbanska 1995; Titus & Tsuyuzaki 1999; Fattorini 2001). In our study, the proportion of vegetation cover on graded pistes was negatively correlated with altitude, but positively affected by neither sowing nor time since machine-grading. This indicates how difficult it is to achieve revegetation at high altitudes. Machine-grading therefore is a particularly damaging management activity, the consequences of which are more severe and long lasting at higher altitudes.

IMPACTS OF ARTIFICIAL SNOW

Application of artificial snow increased the moisture indicator values of the vegetation on piste plots. Artificially snowed pistes contained twice as much snow mass as normal pistes at our study sites in the winter of 1999–2000 (Stoekli & Rixen 2000). Because water for artificial snow making is taken from lakes, rivers or ground water, it usually contains minerals and other chemical compounds (Kammer & Hegg 1990; Kammer 2002), and thus provides a nutrient input during spring melting for about 4 weeks in late spring. Accordingly, the vegetation on piste plots with artificial snow had an increased nutrient indicator value. Adding artificial snow to the ski pistes appeared to mitigate the mechanical disturbance of skiing because woody species became more abundant the longer artificial snow had been applied. Artificial snow is usually produced at the beginning of the winter and lasts beyond the end of the skiing season in spring.

The increase in snowbed and late-flowering species, and the decrease in wind-edge and early flowering species, on pistes with artificial snow probably reflect the shortened growth period resulting from a delay in snowmelt of 2–3 weeks (Newesely 1997; Stoekli & Rixen 2000). Similar to the effects on normal pistes with their compacted natural snow, the narrowed temporal niche in spring presumably conferred a competitive advantage to the snowbed and late-flowering species over wind-edge and early flowering species (Cernusca *et al.* 1990; Newesely 1997; Rixen *et al.* 2001).

Plant diversity reacted differently to artificial snow on the graded and ungraded pistes. The increase in diversity on the ungraded pistes might be because of the decreased mechanical disturbance. The decrease in diversity on the graded pistes, on the other hand, might be a consequence of a decelerated revegetation process because of the shortened growing period. These effects can be enhanced if ice layers in the artificial snow cover decrease gas permeability (Newesely, Cernusca & Bodner 1994).

Whether the impact of artificial snow on alpine vegetation is considered positive or negative depends on the current state of the vegetation and the environmental objectives of a specific ski resort. If mechanical disturbance through snow-grooming vehicles or ski edges is a major problem, the increased protection afforded by artificial snow can be considered beneficial. However, in the case of endangered habitats poor in nutrients, like oligotrophic fens or nutrient-poor grasslands, the additional nutrients input by the melt water of artificial snow is clearly a negative impact.

We have shown that ski pistes in general, and machine-grading and artificial snow production in particular, cause deviations from natural plant species composition and decrease plant species richness. Machine-grading constitutes the most drastic vegetation disturbance on ski pistes. It should be avoided wherever possible, as it causes lasting damage that cannot be overcome even by

revegetation measures, particularly at higher altitudes. Impacts of ski pistes in general, and of artificial snowing in particular, appear comparatively moderate, but are by no means negligible. With the ongoing intensification of ski resorts, the use of artificial snow will become more prevalent and the vegetation will change over an increasing area. Moreover, impacts of artificial snowing are cumulative and will become even more pronounced in the long term. In summary, mountain regions with a high proportion of areas with extensive outdoor recreation activities, like the European Alps, are facing continuous change of their traditional unique environment and vegetation. Therefore, we recommend that environmental goals in ski resort management should be established and respected. In particular, we recommend carefully recording the vegetation in a specific area before any intensification of use as ski piste, and complete avoidance of areas with vegetation of particularly high conservation value. Moreover, we recommend that large-scale machine-grading should be avoided, and that long-term snow production should be banned in areas where any increase in the supply of nutrients and water is a concern. Overall, the establishment of ski pistes should not be allowed in areas where any changes in vegetation composition or any decrease in plant species richness cannot be tolerated.

Acknowledgements

The study was supported by the Swiss Agency for the Environment, Forests and Landscape (SAEFL, Switzerland) and the cantons of Valais and Grisons. We thank the ski resort managers for their collaboration, Katharina Schudel for help in the field, and Dominik Kulakowski and the anonymous referees for comments that helped improving the manuscript.

Supplementary material

The following material is available from <http://www.blackwellpublishing.com/products/journals/suppmat/JPE/JPE1011/JPE1011sm.htm>.

Appendix 1. List of study sites

Appendix 2. List of species

References

- Abegg, B. (1996) *Klimaänderung und Tourismus: Klimafolgenforschung am Beispiel des Wintertourismus in den Schweizer Alpen*. Vdf Hochschulverlag AG, Zürich, Switzerland.
- Abegg, B., Koenig, U., Buerki, R. & Elsasser, H. (1997) Climate impact assessment in tourism. *Die Erde*, **128**, 105–116.
- Baiderin, V.V. (1981) Winter recreation and subnival plant development (translated from Russian). *Ecologiya*, **5**, 3–8.
- Bayfield, N.G. (1996) Long-term changes in colonization of bulldozed ski pistes at Cairn Gorm, Scotland. *Journal of Applied Ecology*, **33**, 1359–1365.
- Bayfield, N.G., Urquart, U.H. & Rothery, P. (1984) Colonization of bulldozed track verges in the Cairngorm Mountains, Scotland. *Journal of Applied Ecology*, **21**, 343–354.
- Cernusca, A., Angerer, H., Newesely, C. & Tappeiner, U. (1990) Auswirkungen von Kunstschnee: eine Kausalanalyse der Belastungsfaktoren. *Verhandlungen der Gesellschaft für Ökologie*, **19**, 746–757.
- Chapin, F.S. III & Körner, C. (1995) *Arctic and Alpine Biodiversity: Patterns, Causes and Ecosystem Consequences*. Springer, Berlin, Germany.
- Connell, J.H. (1979) Intermediate-disturbance hypothesis. *Science*, **204**, 1345–1345.
- Delarze, R., Gonseth, Y. & Galland, P. (1999) *Lebensräume der Schweiz. Ökologie, Gefährdung, Kennarten*, 1st edn. Ott Verlag, Thun, Switzerland.
- Ellenberg, H. (1988) *Vegetation Ecology of Central Europe*, 4th edn. Cambridge University Press, Cambridge, UK.
- Elsasser, H. & Messerli, P. (2001) The vulnerability of the snow industry in the Swiss Alps. *Mountain Research and Development*, **21**, 335–339.
- Fattorini, M. (2001) Establishment of transplants on machine-graded ski runs above timberline in the Swiss Alps. *Restoration Ecology*, **9**, 119–126.
- Fischer, M. & Wipf, S. (2002) Effect of low-intensity grazing on the species-rich vegetation of traditionally mown subalpine meadows. *Biological Conservation*, **104**, 1–11.
- Grime, J.P. (1977) Evidence for the existence of three primary strategies in plants and its relevance to ecological and evolutionary theory. *American Naturalist*, **111**, 1169–1194.
- Grime, J.P. & Jarvis, B.C. (1975) Shade avoidance and shade tolerance in flowering plants II. Effects of light on the germination of species of contrasted ecology. *Light as an Ecological Factor II* (eds G.C. Evans, R. Bainbridge & O. Rackham), pp. 525–532. Blackwell Scientific Publications, Oxford, UK.
- Haeberli, W. (1992) Construction, environmental problems and natural hazards in periglacial mountain belts. *Permafrost and Periglacial Processes*, **3**, 111–124.
- IPCC (1996) *Climate Change 1995: Impacts, Adaptations, and Mitigation of Climate Change. Scientific-Technical Analysis. Contribution of Working Group II to the Second Assessment Report of the Intergovernmental Panel on Climate Change*. Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge and New York.
- IPCC (1998) *The Regional Impacts of Global Change. An Assessment of Vulnerability. Special Report of IPCC Working Group II*. Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge and New York.
- IPCC (2001) *Climate Change 2001: Impacts, Adaptations and Vulnerability*. Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge and New York.
- Kammer, P.M. (2002) Floristic changes in subalpine grasslands after 22 years of artificial snowing. *Journal for Nature Conservation*, **10**, 109–123.
- Kammer, P. & Hegg, O. (1990) Auswirkungen von Kunstschnee auf subalpine Rasenvegetation. *Verhandlungen der Gesellschaft für Ökologie*, **19**, 758–767.
- Kammer, P.M. & Möhl, A. (2002) Factors controlling species richness in alpine plant communities: an assessment of the importance of stress and disturbance. *Arctic, Antarctic, and Alpine Research*, **34**, 398–407.
- Köck, L., Holaus, K., Schnitzer, R. & Walch, A. (1989) Quantitative und qualitative Veränderungen der Vegetation durch den Einfluss des Skisportes auf Pisten und Langlaufloipen. *Landesanstalt für Pflanzenzucht und Samenprüfung in Rinn 1939–89*, pp. 59–79. Eigenverlag, Innsbruck, Rinn, Austria.
- Körner, C. (1999) *Alpine Plant Life*. Springer Verlag, Berlin, Germany.

- Kudo, G. (1993) Relationship between flowering time and fruit-set of the entomophilous alpine shrub, *Rhododendron aureum* (Ericaceae), inhabiting snow patches. *American Journal of Botany*, **80**, 1300–1304.
- Laiolo, P., Dondero, F., Ciliento, E. & Rolando, A. (2004) Consequences of pastoral abandonment for the structure and diversity of the alpine avifauna. *Journal of Applied Ecology*, **41**, 389–397.
- Landolt, E. (1977) Ökologische Zeigerwerte zur Schweizer Flora. *Veröffentlichungen des Geobotanischen Institutes der Eidgenössischen Technischen Hochschule, Stiftung Rübel, Zürich*, **64**, 1–208.
- Lauber, K. & Wagner, G. (2001) *Flora Helvetica*, 3 edn. Paul Haupt, Bern, Switzerland.
- Löhmansröben, R. & Cernusca, A. (1987) Ursachen für Erosionsprozesse im Skigebiet Meran 2000. *Verhandlungen der Gesellschaft für Ökologie*, **15**, 189–196.
- Mulder, C.P.H., Uliassi, D.D. & Doak, D.F. (2001) Physical stress and diversity–productivity relationships: the role of positive interactions. *Proceedings of the National Academy of Sciences (USA)*, **98**, 6704–6708.
- Müller-Dombois, D. & Ellenberg, H. (1974) *Aims and Methods of Vegetation Ecology*. Wiley, New York, NY.
- Newesely, C. (1997) *Auswirkungen der künstlichen Beschneidung von Schipisten auf Aufbau, Struktur und Gasdurchlässigkeit der Schneedecke, sowie auf den Verlauf der Bodentemperatur und das Auftreten von Bodenfrost*. PhD Thesis. Leopold Franzens Universität, Innsbruck, Switzerland.
- Newesely, C., Cernusca, A. & Bodner, M. (1994) Entstehung und Auswirkung von Sauerstoffmangel im Bereich unterschiedlich präparierter Schipisten. *Verhandlungen der Gesellschaft für Ökologie*, **23**, 277–282.
- Pröbstl, U. (1990) *Skisport und Vegetation*. Stöppel, Weilheim, Germany.
- Rixen, C. (2002) *Artificial snow and snow additives on ski pistes: interactions between snow cover, soil and vegetation*. PhD Thesis. University of Zurich, Zurich, Switzerland.
- Rixen, C., Casteller, A., Schweingruber, F.H. & Stoeckli, V. (2004) Age analysis helps to estimate plant performance on ski pistes. *Botanica Helvetica*, **114**, 127–138.
- Rixen, C., Haerberli, W. & Stoeckli, V. (2004) Ground temperatures under ski pistes with artificial and natural snow. *Arctic, Antarctic, and Alpine Research*, **36**, 403–411.
- Rixen, C., Stoeckli, V. & Ammann, W. (2003) Does artificial snow production affect soil and vegetation of ski pistes? A review. *Perspectives in Plant Ecology, Evolution and Systematics*, **5**, 219–230.
- Rixen, C., Stoeckli, V., Huovinen, C. & Huovinen, K. (2001) The phenology of four subalpine herbs in relation to snow cover characteristics. *Proceedings of the Sixth IAHS Symposium 2001. Soil–Vegetation–Atmosphere Transfer Schemes and Large Scale Hydrological Models* (eds A.J. Dolman, A.J. Hall, M.L. Kavvas, T. Oki & J.W. Pomeroy), pp. 359–362. Publication No. 270. IAHS, Maastricht, the Netherlands.
- Ruth-Balaganskaya, E. & Myllynen-Malinen, K. (2000) Soil nutrient status and revegetation practices of downhill skiing areas in Finnish Lapland: a case study of Mt Ylläs. *Landscape and Urban Planning*, **50**, 259–268.
- Seilbahnen Schweiz (2001) *Wirtschaftsbericht 2001*. Paul Haupt, Bern, Switzerland.
- SPSS (1999) *SPSS for Windows*. SPSS Inc., Chicago, IL.
- Stanton, M.L. & Galen, C. (1997) Life on the edge: adaptation versus environmentally mediated gene flow in the snow buttercup, *Ranunculus adoneus*. *American Naturalist*, **150**, 143–178.
- Stoeckli, V. & Rixen, C. (2000) Characteristics of artificial snow and its effect on vegetation. *Conference Proceedings of the International Snow Science Workshop* (eds K. Birkeland, E. Adams & F. Johnson), pp. 468–471. American Avalanche Association, Bozeman USA, Big Sky, Montana, USA, 1–6 October 2000.
- Tilman, D. (1996) Biodiversity: population versus ecosystem stability. *Ecology*, **77**, 350–363.
- Titus, J.H. & Tsuyuzaki, S. (1999) Ski slope vegetation of Mount Hood, Oregon, USA. *Arctic, Antarctic, and Alpine Research*, **31**, 283–292.
- Urbanska, K.M. (1995) Biodiversity assessment in ecological restoration above the timberline. *Biodiversity and Conservation*, **4**, 679–695.
- Urbanska, K.M. (1997) Restoration ecology research above the timberline: colonization of safety islands on a machine-graded alpine ski run. *Biodiversity and Conservation*, **6**, 1655–1670.
- Vujnovic, K., Wein, R.W. & Dale, M.R.T. (2002) Predicting plant species diversity in response to disturbance magnitude in grassland remnants of central Alberta. *Canadian Journal of Botany–Revue Canadienne de Botanique*, **80**, 504–511.
- Wilmanns, O. (1998) *Ökologische Pflanzensoziologie*, 6th edn. Quelle und Meyer, Wiesbaden, Germany.
- Wipf, S., Rixen, C., Freppaz, M. & Stoeckli, V. (2002) Ski piste vegetation under artificial and natural snow: patterns in multivariate analysis. *Interdisciplinary Mountain Research* (eds R. Bottarin & U. Tappeiner), pp. 170–179. Blackwell Science, Berlin, Germany.

Received 5 June 2004; final copy received 22 December 2004