Spin State Controlled Photodissociation of Iron(III) Azide to Iron(V) Nitride Complex

Erik Andris,^[a] Rafael Navrátil,^[a] Juraj Jašík,^[a] Gerard Sabenya,^[b] Miquel Costas,^[b] Martin Srnc^[c] and Jana Roithová^{*[a]}

Abstract: We address the generation of the iron(V) nitride complexes, targets of biomimetic chemistry. Temperature dependent ion spectroscopy demonstrate that generation of iron(V) nitrides is governed by the spin state population of their iron(III) azide precursors and can be tuned by temperature. Complex $[(\text{MePy}_2\text{TACN})\text{Fe}(\text{N}_3)]^{2+}$ exists as a mixture of sextet and doublet spin states at 300 K, whereas only the doublet state is populated at 3 K. Photofragmentation of the sextet state complex leads to reduction of the iron center. The doublet state complex photodissociates to the desired iron(V) nitride complex. We show analogous behaviour also for complexes with cyclam-based ligands.

High-valent iron enzymes catalyse highly challenging reactions such as methane oxidation.^[1,2] The aim to understand and to mimic such reactivity led to a boom in bioinorganic chemistry of iron complexes, especially of iron-oxo complexes.^[3–5] Much less is known about hypervalent iron nitrides.^[6–9] The scarcity of information about pseudo-octahedral iron nitrides stems from their thermal instability.^[10–16] They can be prepared by photolysis of their iron azide precursors. The photodissociation has three possible pathways: desired elimination of N_2 occurs in competition with eliminations of N_3^- and N_3^+ . Temperature and irradiation wavelength determine which of these three pathways occurs preferentially.^[17] Most of the successful preparations of iron nitrides were done in frozen solutions,^[17–19] which implied that a solid matrix and low temperatures are required for their generation.^[19,20] Later however, ultrafast experiments revealed the formation of iron nitrides at room temperature as well.^[21–23]

Here, we study photodissociation of iron(III) azides at molecular level with ion spectroscopy at variable temperature (3 K – 310 K).^[24–27] The experiments provide absorption spectra of mass-selected ions separated according to the individual dissociation pathways. Photodissociation of mass-selected iron azide complexes **1**^[28] in the visible range leads either to the photoreduction associated with elimination of N_3^-

(photoreduction) or to the formation of desired iron(V) nitride complex **3** concomitant with the elimination of N_2 (photooxidation) (Scheme 1, experimental details are in the Supporting Information). The ratio of the reaction pathways critically depends on the wavelength of the incident light (Figure 1). The N_3^+ elimination channel displays a maximum at 455 nm (blue trace in Figure 1a) and the N_2 elimination attains two maxima at 530 and 600 nm (red trace in Figure 1a).

Scheme 1. Photodissociation of the iron azide $[(\text{MePy}_2\text{TACN})\text{Fe}(\text{N}_3)]^{2+}$ (**1**, MePy_2TACN = *N*-methyl-*N,N*-bis(2-picolyl)-1,4,7-triazacyclononane) leads either to photoreduction to $[(\text{MePy}_2\text{TACN})\text{Fe}]^{2+}$ (**2**) or to photooxidation to iron nitride $[(\text{MePy}_2\text{TACN})\text{FeN}]^{2+}$ (**3**).

The internal energy of the precursor complexes **1** can be tuned by the temperature of the ion trap in which the experiments are performed. Photodissociation experiments in dependence on this temperature show that the N_3^+ elimination is almost completely suppressed below 200 K. Below 50 K, the photodissociation of **1** at longer wavelengths starts to be less efficient and the absorption maximum at 600 nm diminishes. The temperature effect can be a consequence of low temperature (**1**) enabling other relaxation processes than the photodissociation or (**2**) triggering a change in the precursor complex that modifies its absorption spectrum. In order to disentangle this dilemma, we measured the absorption spectrum of **1** by the helium tagging method at 3 K (Figure 1b). In this experiment we monitored elimination of the helium atom from the $[\text{1} \cdot \text{He}]$ complex. Upon absorption of a photon, the complex eliminates helium regardless of any other possible relaxation processes due to its extremely low binding energy (on the order of tenth of kJ mol^{-1}).

The helium tagging spectrum shows the same profile as the high temperature spectra determined for the N_2 elimination channel with two maxima at 530 and 600 nm and lacks the maximum at 455 nm (*cf.* Figures 1a and 1b). This means that species giving rise to the photofragmentation band at 455 nm on one hand and to the bands at 530 and 600 nm on the other are different. The absorption of the former species is not present at low temperatures. Hence, we can conclude that **1** has two isomers with different absorption spectra and different preferred photofragmentation pathways.

[a] E. Andris, R. Navrátil, Dr. J. Jašík, Prof. J. Roithová
Department of Organic Chemistry
Faculty of Science
Charles University
Hlavova 2030/8, 128 43 Prague 2 (Czech Republic)
E-mail: roithova@natur.cuni.cz

[b] G. Sabenya, Prof. M. Costas
Departament de Química and Institute of Computational Chemistry and Catalysis (IQCC)
University of Girona
Campus Montilivi, Girona 17071 (Spain)

[c] Dr. M. Srnc
J. Heyrovsky Institute of Physical Chemistry of the CAS, v. v. i.
Dolejškova 2155/3, 182 23 Prague 8 (Czech Republic)

Figure 1. Visible photodissociation (visPD) spectra of mass-selected complex **1** at different temperatures. (a) VisPD spectra obtained from monitoring of the fragment formation (N_2 elimination in red, N_3 elimination in blue) in dependence on the photon wavelength (see Figure S4 in the Supporting Information for the unprocessed data). (b) Helium-tagging visPD spectrum of **1** at 3 K obtained by monitoring attenuation of the helium complexes of **1**. (c) Theoretical TD-DFT spectra of the doublet (red trace) and the sextet (blue trace) iron(III) azide complex **1** calculated at 0 K (both spectra were redshifted by 25 nm to fit the experiment, see Figure S1 in the Supporting Information for all spin states without the shift).

Interconversion of the isomers reveals itself as temperature dependent changes in the branching of the photofragmentation pathways. Experiments performed at 530 nm and 600 nm (Figures 2a and 2b) show that the total photodissociation yield increases with the decreasing temperature (except of the range below 20 K, where it decreases due to other relaxation processes enabled by a longer lifetime of the excited state - see also the Supporting Information). While at low temperatures, photooxidation (N_2 elimination) is far the prevailing process; above 150–200 K, the N_3 loss starts to contribute to the total photofragmentation cross section at the expense of photooxidation. This N_3 elimination originates from the low-energy tail of the 455 nm absorption (Figure 1a). The mutual

dependence of the N_2 and N_3 elimination yield demonstrates that both fragmentation pathways originate from single or interconverting precursors.

The absorption maximum at 455 nm is present only in the vis spectrum of the isomer that is populated at high temperatures and that photoreduces upon irradiation (Figure 1a). Accordingly, the photofragmentation yield at 455 nm decreases with the temperature from 300 K down to 200 K and is dominated by the N_3 loss (Figure 2c). Below 200 K, the increase of the photofragmentation intensity is due to the high energy tail of the photooxidation band at the 530 nm.

Multireference *ab initio* method CASPT2 (complete active space perturbation theory) calculations revealed that the doublet and sextet states are almost isoenergetic and both should be experimentally observable (the ground state is predicted either doublet or sextet depending on whether the zero point vibrational energy is included or not – in either case the difference is on the order of 1 kcal mol^{−1}, see Table S1 in the Supporting Information). The quartet spin state is more than 25 kcal mol^{−1} higher in energy and therefore is not further considered.

Time-dependent density functional theory (TD-DFT) calculations of the vis spectra of **1** in the doublet and sextet spin states (Figure 1c) reveal that the dominant electronic transitions correspond mainly to the ligand-to-metal charge transfer between the azide ligand and the iron centre. The theoretical spectrum of the doublet spin state of **1** reproduces the photodissociation spectra corresponding to the loss of N_2 (Figure 1a, red trace) and the theoretical spectrum of the sextet spin state of **1** reproduces photodissociation spectra corresponding to the loss of the N_3 radical (Figure 1a, blue trace).

Temperature dependent spectra as well as the theoretical calculation lead us to conclude that the photodissociation reactivity of **1** is governed by its spin states. The doublet spin state of **1** steers the photodissociation towards the N_2 elimination, whereas the sextet spin state opens a pathway for the N_3 loss. The photoproduct **3** has a doublet ground state (Table S1 in the Supporting Information), which might indicate its preferential formation from the doublet state of **1**. At room temperature both doublet and sextet states of **1** are populated. With decreasing temperature, we observe a spin relaxation of the sextet state to the doublet state^[29,30] that is dominant below 200 K.

We have further studied the structure of mass-selected complexes **1** by helium tagging infrared photodissociation spectra (IRPD).^[24,31] The helium-tagging IRPD spectrum ($T = 3$ K) of **1** shows the prominent azide group antisymmetric stretch at 2016 cm^{−1} and the symmetric stretch at 1199 cm^{−1} (Figure 3a, black trace). The origin of these bands was confirmed by ¹⁵N isotopic labelling (a mixture of Fe–NN¹⁵N and Fe–¹⁵NNN in a 1:1 ratio, Figure 3a, grey trace). Both azide bands split into two red-shifted bands: the antisymmetric azide stretches emerge at 2013 cm^{−1} and 1991 cm^{−1} and the symmetric stretches at 1187 cm^{−1} and 1170 cm^{−1}. The helium tagging IRPD spectrum agrees best with the theoretical

Figure 2. Temperature dependent photofragmentation of **1**. Photodissociation yields of N_2 elimination (in red), N_3 elimination (in blue) and the total photodissociation yield (in purple) at the absorption maxima at (a) 530 nm, (b) 600 nm and (c) 455 nm as a function of temperature. See Figure S13 in the Supporting Information for additional experimental data.

prediction for the doublet spin state of **1**.

Infrared multiphoton dissociation (IRMPD) spectrum of **1** recorded at 300 K by monitoring the N_2 elimination channel (Figure 3b) features two bands in the region of the antisymmetric azide stretches. The 2016 cm^{-1} (3 K) band is red-shifted to 2009 cm^{-1} and a new band appears at 2052 cm^{-1} (*cf.* Figures 3a and 3b). The same changes are observed in the region of the symmetric azide stretches. The original (3 K) band at 1199 cm^{-1} blue-shifts to 1201 cm^{-1} and a new band appears at 1316 cm^{-1} . Additionally, the 300 K spectrum features a new band at 1024 cm^{-1} , which is theoretically predicted only for the sextet state. Both spin state complexes are able to eliminate N_2 , because this is the lowest energy fragmentation and fast spin-isomerization is expected. Based on the observed frequency shifts and the DFT-aided assignments, we deduce that both sextet and doublet states are present at 300 K, in accord with the results of the vis photodissociation experiments.

The remaining question is whether the spin relaxation and spin governed photodissociation are general features of octahedral iron(III) azides. To address it, we conducted temperature dependent photofragmentation studies with other complexes with cyclam-based ligands. Two of them have a doublet ground state: [(cyclam)Fe(N_3) $_2$] $^+$ (**4**, cyclam = 1,4,8,11-tetraazacyclotetradecane, Figure 4a) $^{[18]}$ and [(cyclam-ac)Fe(N_3)] $^+$ (**5**, cyclam-ac = 1,4,8,11-tetraazacyclotetradecane-1-acetate, Figure 4b) $^{[19]}$ whereas the third complex has a sextet ground state: [(Me3cyclam-ac)Fe(N_3)] $^+$ (**6**, Me3cyclam-ac = 4,8,11-trimethyl-1,4,8,11-tetraazacyclotetradecane-1-acetate,

Figure 4. Branching ratios between the photoreduction (N_3^- loss) and the photooxidation (N_2 loss) for complexes (a) **4** ($S = 1/2$), (b) **5** ($S = 1/2$) and (c) **6** ($S = 5/2$). The brackets indicate the ground states of the complexes. The branching ratios were determined at absorption maxima measured at 3 K; the wavelength is specified in the figure. Elimination of HN_3 from complexes **4** and **5** and elimination of MeN_3 from **6** were excluded from the sum (complex **4** photooxidizes also in a consecutive process of HN_3 and N_2 elimination - this process is included in the photooxidation channel). All data can be found the Supporting Information (Figures S7–S12).

Figure 3. (a) Helium-tagging infrared photodissociation spectra of **1** at 3 K, (b) infrared multiphoton dissociation spectra of **1** at 300 K (grey traces in both spectra correspond to the ^{15}N -labelling, note that **1** (^{15}N) contains unsymmetrically labelled azide ligand, therefore isobaric mixture of [(MePy $_2$ TACN)Fe-(^{15}N)] $^{2+}$ and [(MePy $_2$ TACN)Fe-(^{15}N)] $^{2+}$ in the 1:1 ratio was measured). (c) DFT-predicted spectra of **1**. Bands corresponding to the symmetric and antisymmetric azide stretches (v_s and v_{as}) of the doublet and the sextet state are highlighted with blue and red colours respectively.

Figure 4c). $^{[32]}$ Photofragmentation of the complexes with the doublet ground state **4** and **5** leads to photoreduction and photooxidation as observed for **1**, but also to the elimination of HN_3 . The elimination HN_3 is a gas-phase analogy of the N_3^- elimination that was observed for these complexes in the condensed phase. The absorption spectra obtained for photooxidation and photoreduction have different profiles and different absorption maxima (Figures S7 and S9). This indicates that different electronic transitions are responsible for these two processes. For complexes **4** and **5**, the relative abundance of photooxidation with respect to photoreduction decreases with increasing temperature (Figures 4a, 4b, S8 and S10). The results are thus analogous to those obtained for complex **1** and confirm the suggested rationale. Complex **6** has the sextet ground state 32 and therefore should reveal the opposite

behaviour. Accordingly, we observe almost no photo-oxidation at low temperatures, and it only occurs at increased temperatures (Figure 4c, S11 and S12). Photooxidation remains, however, a minor reaction channel even at the highest studied temperatures. This is a clear indicator that population of the doublet state is energetically as well as entropically disfavoured.

The experiments clearly show that the photodissociation of octahedral iron(III) azides is governed by their spin states. Detailed analysis of complex **1** reveals that its low spin ($S = 1/2$)

state and the high spin ($S = 5/2$) state are almost isoenergetic and that both are populated at room temperature. Photodissociation of the doublet state (absorption maximum at $\lambda_{\text{max}} = 530$ nm) leads preferentially to photooxidation and formation of the iron(V) nitride complex **3**. The sextet state photodissociates ($\lambda_{\text{max}} = 455$ nm) via the loss of the $\text{N}_3^{\cdot -}$ radical (photoreduction). Population of the spin states of complex **1** can be tuned by temperature: the population of the sextet state is depleted with decreasing temperature. Below 200 K, the doublet state population becomes dominant, and it is exclusive at low temperatures (3 K). Two other iron(III) azide complexes **4** and **5** with the $S = 1/2$ ground state showed analogous photochemical behavior. Photodissociation of complex **6** with the $S = 5/2$ ground state has an opposite dependency on temperature: photoreduction is the dominant pathway at low temperature.

Hence, all these results demonstrate that the outcome of the photodissociation of the iron(III) azide complexes is mainly determined by their spin state. Previously suggested effects of the medium of the reaction or the reaction conditions (frozen matrix or liquid solution) are not the origin of the observed temperature effects. The conditions, however, can affect the position and width of the temperature interval in which the spin populations are changing.^[33]

The photoreactivity of the iron(III) azide is analogous to the reactivity of iron(III) hydroperoxides, models for Rieske dioxxygenases.^[34] Their reactivity is governed by the spin state as well: The low-spin $\text{Fe}^{\text{III}}\text{--O--OH}$ complexes have a weaker O–O bond than their high-spin analogues and therefore provide highly reactive iron(V)-oxo complexes.^[35] The high spin complexes, on the other hand, have a higher redox potential that opens other oxidation reaction pathways without the necessity for O–O bond cleavage.^[36] We bring tools to investigate this type of reactivity – in the ground state as well as the excited state of the potential energy surface – that can ultimately provide a deep insight into the electronic structure and reactivity of this group of compounds. The method is sufficiently versatile to be universally applicable to photodissociation processes at the level of isolated molecules.

Acknowledgements

The project was supported by the European Research Council (ERC CoG No. 682275), the Czech Science Foundation (15-10279Y) and the COST action ECOSTBio. Dr. M. S. is also grateful to the Czech Academy of Sciences for the Purkyně fellowship.

Keywords: hypervalent metal complexes, ion spectroscopy, iron(V) nitride, photooxidation, spin crossover

- [1] M. M. Abu-Omar, A. Loaiza, N. Hontzeas, *Chem. Rev.* **2005**, *105*, 2227–2252.
- [2] E. I. Solomon, K. M. Light, L. V. Liu, M. Srnc, S. D. Wong, *Acc. Chem. Res.* **2013**, *46*, 2725–2739.
- [3] B. Meunier, S. P. de Visser, S. Shaik, *Chem. Rev.* **2004**, *104*, 3947–3980.
- [4] W. N. Oloo, L. Que, *Acc. Chem. Res.* **2015**, *48*, 2612–2621.
- [5] X. Engelmann, I. Monte-Perez, K. Ray, *Angew. Chem. Int. Ed.* **2016**, *55*, 7632–7649.
- [6] J. F. Berry, *Comm. Inorg. Chem.* **2009**, *30*, 28–66.
- [7] M. D. Fryzuk, S. A. Johnson, *Coord. Chem. Rev.* **2000**, *200–202*, 379–409.
- [8] B. A. MacKay, M. D. Fryzuk, *Chem. Rev.* **2004**, *104*, 385–402.
- [9] B. M. Hoffman, D. Lukoyanov, Z.-Y. Yang, D. R. Dean, L. C. Seefeldt, *Chem. Rev.* **2014**, *114*, 4041–4062.
- [10] A. Dey, A. Ghosh, *J. Am. Chem. Soc.* **2002**, *124*, 3206–3207.
- [11] J. F. Berry, E. Bill, E. Bothe, S. DeBeer George, B. Mienert, F. Neese, K. Wieghardt, *Science* **2006**, *312*, 1937–1941.
- [12] J. J. Scepaniak, C. S. Vogel, M. M. Khusniyarov, F. W. Heinemann, K. Meyer, J. M. Smith, *Science* **2011**, *331*, 1049–1052.
- [13] J. M. Smith, D. Subedi, *Dalton Trans.* **2012**, *41*, 1423–1429.
- [14] J. Hohenberger, K. Ray, K. Meyer, *Nature Commun.* **2012**, *3*, 720.
- [15] W.-D. Wagner, K. Nakamoto, *J. Am. Chem. Soc.* **1989**, *111*, 1590–1598.
- [16] T. Petrenko, S. DeBeer George, N. Aliaga-Alcalde, E. Bill, B. Mienert, Y. Xiao, Y. Guo, W. Sturhahn, S. P. Cramer, K. Wieghardt, F. Neese, *J. Am. Chem. Soc.* **2007**, *129*, 11053–11060.
- [17] J. Torres-Alacan, P. Vöhringer, *Int. Rev. Phys. Chem.* **2014**, *33*, 521–553.
- [18] K. Meyer, E. Bill, B. Mienert, T. Weyhermüller, K. Wieghardt, *J. Am. Chem. Soc.* **1999**, *121*, 4859–4876.
- [19] C. A. Grapperhaus, B. Mienert, E. Bill, T. Weyhermüller, K. Wieghardt, *Inorg. Chem.* **2000**, *39*, 5306–5317.
- [20] O. Krahe, E. Bill, F. Neese, *Angew. Chem. Int. Ed.* **2014**, *53*, 8727–8731.
- [21] J. Torres-Alacan, U. Das, A. C. Filippou, P. Vöhringer, *Angew. Chem. Int. Ed.* **2013**, *52*, 12833–12837.
- [22] H. Vennekate, D. Schwarzer, J. Torres-Alacan, P. Vöhringer, *J. Am. Chem. Soc.* **2014**, *136*, 10095–10103.
- [23] J. Torres-Alacan, P. Vöhringer, *Chem. Eur. J.* **2017**, *23*, 6746–6751.
- [24] J. Roithová, A. Gray, E. Andris, J. Jašík, D. Gerlich, *Acc. Chem. Res.* **2016**, *49*, 223–230.
- [25] J. Jašík, R. Navrátil, I. Němec, J. Roithová, *J. Phys. Chem. A* **2015**, *119*, 12648–12655.
- [26] E. Andris, R. Navrátil, J. Jašík, T. Terencio, M. Srnc, M. Costas, J. Roithová, *J. Am. Chem. Soc.* **2017**, *139*, 2757–2765.
- [27] R. Navrátil, J. Jašík, J. Roithová, *J. Mol. Spectrosc.* **2017**, *332*, 52–58.
- [28] G. Sabenya, L. Lazaro, I. Gamba, V. Martin-Diaconescu, E. Andris, T. Weyhermüller, F. Neese, J. Roithová, E. Bill, J. Lloret-Fillol, M. Costas, *J. Am. Chem. Soc.* **2017**, *139*, 9168–9177.
- [29] A. Bousseksou, G. Molnár, L. Salmon, W. Nicolazzi, *Chem. Soc. Rev.* **2011**, *40*, 3313–3335.
- [30] D. J. Harding, P. Harding, W. Phonsri, *Coord. Chem. Rev.* **2016**, *313*, 38–61.
- [31] M. A. Johnson in *Encyclopedia of Mass Spectrometry* (Eds.: Gross, M. L. & Caprioli, R.), Vol. 1 (Vol. Ed. Armentrout, P. B.), Elsevier, Amsterdam, **2003**, pp. 271–276.
- [32] J. F. Berry, E. Bill, E. Bothe, T. Weyhermüller, K. Wieghardt, *J. Am. Chem. Soc.* **2005**, *127*, 11550–11551.
- [33] W. Phonsri, P. Harding, L. Liu, S. G. Telfer, K. S. Murray, B. Moubaraki, T. M. Ross, G. N. L. Jameson, D. J. Harding, *Chem. Sci.* **2017**, *8*, 3949–3959.
- [34] K. Chen, L. Que, Jr., *J. Am. Chem. Soc.* **2001**, *123*, 6327–6337.
- [35] N. Lehnert, R. Y. N. Ho, L. Que, Jr., E. I. Solomon, *J. Am. Chem. Soc.* **2001**, *123*, 12802–12816.
- [36] L. V. Liu, S. Hong, J. Cho, W. Nam, E. I. Solomon, *J. Am. Chem. Soc.* **2013**, *135*, 3286–3299.

Entry for the Table of Contents

COMMUNICATION

Photodissociation of iron(III) azide complexes is governed by their spin state. The complexes with the $S = 1/2$ spin state undergo photooxidation and thus yield iron(V) nitride complexes. The $S = 5/2$ complexes are prone to photoreduction. Variation of the temperature induces spin crossover in the complexes and can be used to steer the photodissociation either to the oxidation or to the reduction.

Erik Andris, Rafael Navrátil, Dr. Juraj Jašík, Gerard Sabenya, Prof. Miquel Costas, Dr. Martin Srnec and Prof. Jana Roithová*

Page No. – Page No.

**Spin State Controlled
Photodissociation of Iron(III) Azide to
Iron(V) Nitride Complex**