

## Synthetic Photochemistry

International Edition: DOI: 10.1002/anie.201709571  
German Edition: DOI: 10.1002/ange.201709571

## Radical-Based C–C Bond-Forming Processes Enabled by the Photoexcitation of 4-Alkyl-1,4-dihydropyridines


Luca Buzzetti, Alexis Prieto, Sudipta Raha Roy, and Paolo Melchiorre\*

**Abstract:** We report herein that 4-alkyl-1,4-dihydropyridines (alkyl-DHPs) can directly reach an electronically excited state upon light absorption and trigger the generation of C(sp<sup>3</sup>)-centered radicals without the need for an external photocatalyst. Selective excitation with a violet-light-emitting diode turns alkyl-DHPs into strong reducing agents that can activate reagents through single-electron transfer manifolds while undergoing homolytic cleavage to generate radicals. We used this photochemical dual-reactivity profile to trigger radical-based carbon–carbon bond-forming processes, including nickel-catalyzed cross-coupling reactions.

The chemical reactivity of electronically excited molecules differs fundamentally from that in the ground state. This is the underlying reactivity concept of photochemistry,<sup>[1]</sup> which has traditionally allowed the development of unique chemical transformations that are not achievable through conventional ground-state pathways.<sup>[2]</sup> For example, an excited-state molecule is both a better electron donor (i.e., a better reductant) and a better electron acceptor (i.e., a better oxidant) than in the ground state.<sup>[3]</sup> This explains why the light excitation of organic molecules can unlock unconventional reactivity manifolds. In this context, our laboratory has been exploring the potential of some chiral organocatalytic intermediates to directly reach an electronically excited state upon visible-light absorption to then switch on novel catalytic functions that are unavailable to ground-state organocatalysis.<sup>[4]</sup> For example, we showed that electron-rich enamines **I**, which act as nucleophiles in the ground state, could become strong reductants upon light excitation and trigger the formation of radicals through single-electron transfer (SET) reduction of

electron-poor organic halides (Figure 1a).<sup>[4a,b]</sup> In the same vein, 1,4-dihydropyridine derivatives **II** (H-DHP, Figure 1b) are primarily understood as hydride (H<sup>−</sup>) sources in their ground state.<sup>[5]</sup> Other researchers recently demonstrated that exciting them with visible light affords strong photoreductants that can productively engage in SET manifolds.<sup>[6]</sup>

These findings led us to wonder whether 4-alkyl-1,4-dihydropyridines (alkyl-DHPs, **1**), which closely resemble the structure of H-DHP **II**, might unveil a similar reactivity behavior upon excitation, becoming photoreductants in the excited state (Figure 1c). If so, the ensuing SET event with a suitable electron acceptor would generate intermediate **1<sup>•+</sup>**. The radical cation **1<sup>•+</sup>**, which is generally accessed by SET oxidation of **1** using an external photoredox catalyst<sup>[7]</sup> or a stoichiometric oxidant,<sup>[8]</sup> is prone to a fast homolytic cleavage, eventually serving as a radical precursor.<sup>[7,8]</sup> Overall, the proposed photochemical mechanism would generate C(sp<sup>3</sup>)-centered radicals, which could then trigger carbon–carbon bond-forming processes under mild conditions and only relying upon the photochemistry of readily available alkyl-DHPs **1**, which are prepared from aldehydes in one



**Figure 1.** Radical generation strategies enabled by the excited-state reactivity of organic molecules. a, b) Previous studies demonstrating that light excitation turns enamines (a) and 1,4-dihydropyridines **II** (H-DHP; b) into strong reductants and the resulting SET-based radical generation mechanisms. c) The proposed strategy exploits the ability of 4-alkyl-1,4-dihydropyridines (alkyl-DHPs, **1**) to become both photoreductants and precursors of alkyl radicals upon excitation. EWG = electron-withdrawing group; SET = single-electron transfer; filled dark blue circle represents a bulky substituent on the chiral amine.

[\*] Prof. Dr. P. Melchiorre  
ICREA—Catalan Institution for Research and Advanced Studies  
Passeig Lluís Companys 23, 08010 Barcelona (Spain)  
L. Buzzetti, Dr. A. Prieto, Dr. S. R. Roy, Prof. Dr. P. Melchiorre  
ICIQ—Institute of Chemical Research of Catalonia the Barcelona  
Institute of Science and Technology  
Avenida Països Catalans 16, 43007 Tarragona (Spain)  
E-mail: pmelchiorre@icq.es  
Homepage: [http://www.icq.org/research/research\\_group/prof-paolo-melchiorre/](http://www.icq.org/research/research_group/prof-paolo-melchiorre/)


Supporting information and the ORCID identification number(s) for the author(s) of this article can be found under:  
<https://doi.org/10.1002/anie.201709571>.

© 2017 The Authors. Published by Wiley-VCH Verlag GmbH & Co. KGaA. This is an open access article under the terms of the Creative Commons Attribution Non-Commercial NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial, and no modifications or adaptations are made.

step.<sup>[8,9]</sup> Herein, we demonstrate how this idea was translated into experimental reality.

To test the feasibility of our photochemical plan, we initially studied the photophysical behavior of diethyl 4-benzyl-1,4-dihydropyridine-3,5-dicarboxylate (Bn-DHP, **1a**), which was selected as the model substrate. The optical absorption spectrum of **1a** dissolved in CH<sub>3</sub>CN confirmed its ability to absorb in the visible frequency region, up to 420 nm (Figure 2a, blue line). We found that **1a** was fluorescent when excited at 405 nm, which allowed us to record the emission spectrum (red dotted line in Figure 2a,  $\lambda_{\text{max}} = 470$  nm) and to measure the fluorescence lifetime in CH<sub>3</sub>CN using a picosecond pulsed laser.<sup>[10]</sup> The reconvolution fitting shows a double exponential decay (Figure 2b), affording the following values:  $\tau_0 = 0.28$  ns (below the instrument's detection limit) and  $\tau_1 = 4.1$  ns. Interestingly, the excited-state lifetime of Bn-DHP is comparable with the lifetime of unsubstituted 1,4-dihydropyridine of type **II** (H-DHP in Figure 1b), which was measured to be 320 ps.<sup>[6b]</sup> Considering the ability of H-DHP to act as a strong reducing agent in the excited state,<sup>[6]</sup> these photophysical studies encouraged us to evaluate the photochemistry of the excited Bn-DHP (**1a**\*).


We conducted our experiments in CH<sub>3</sub>CN under irradiation by a high-power single light-emitting diode (HP single LED,  $\lambda_{\text{max}} = 405$  nm) with an irradiance of 50 mW cm<sup>-2</sup>, as controlled by an external power supply (full details of the illumination set-up are reported in the Supporting Information). Direct illumination of a CH<sub>3</sub>CN solution of **1a** led to the formation of toluene, 1,2-diphenylethane, and the corre-



**Figure 2.** a) Absorption of **1a** (blue line,  $\lambda_{\text{max}} = 343$  nm) and its emission (excitation at 405 nm, red dotted line,  $\lambda_{\text{max}} = 470$  nm) in CH<sub>3</sub>CN. b) Fluorescence decay trace of 600  $\mu\text{M}$  **1a** in CH<sub>3</sub>CN after picosecond photoexcitation at 405 nm. c) EPR spectrum of the benzyl radical generated from **1a** at 77 K after 10 minutes (orange line) and 60 minutes (green line) of light irradiation at 405 nm. d) Photochemical behavior of a solution of **1a** in CH<sub>3</sub>CN under 405 nm irradiation. Yield measured by <sup>1</sup>H NMR analysis using trichloroethylene as the internal standard.

sponding pyridine (Figure 2d). On repeating the reaction in the presence of 2,2,6,6-tetramethyl-1-piperidinyloxy (TEMPO, 1 equiv), a radical scavenger, we detected the formation of the benzyl-TEMPO adduct in 55% yield (details in the Supporting Information). Overall, these experiments indicate that the simple photoexcitation of **1a** can trigger the formation of benzyl radicals. This notion was further corroborated by EPR studies of a solution of **1a** in MeCN, conducted at 77 K and with the same illumination system used in the experiments in Figure 2c. The EPR spectrum showed a strong isotropic X-band absorption and afforded a g-value of 2.0035 (Figure 2c), which is consistent with reported data for the characterization of benzyl radicals.<sup>[11]</sup>

After establishing the ability of **1a** to generate C(sp<sup>3</sup>)-centered radicals upon excitation, we evaluated its propensity to act as a strong reducing agent in the excited state. To test this possibility, we selected the aromatic *ipso*-substitution of cyanoarenes as the model reaction.<sup>[7a,12]</sup> Indeed, this process classically proceeds through a radical coupling mechanism that requires the concomitant formation of a C(sp<sup>3</sup>)-centered radical and a persistent cyanoarene radical anion **III** (Scheme 1a). The ensuing C–C bond formation affords




**Scheme 1.** a) Mechanism of the aromatic *ipso*-substitution of cyanoarenes, governed by the coupling of an alkyl radical and the persistent cyanoarene radical anion **III**. Benzylation of 4-cyanopyridine **2a** (b) and tetracyanobenzene **2b** (c) enabled by the direct photoexcitation of **1a**.

a cyclohexadienyl anion **IV** that is prone to rapid rearomatization through the elimination of cyanide to afford the arene alkylation product. When mixing either 4-cyanopyridine (**2a**, Scheme 1b) or tetracyanobenzene (**2b**, Scheme 1c) with **1a** under light irradiation, the corresponding products **3a** and **3b** were generated in fairly good yields. These results are consonant with the excited state of **1a** acting both as a strong reducing agent, affording the cyanoarene radical anion of type **III** upon SET reduction of substrates **2**, and as a source of a benzyl radical, generated upon homolytic cleavage of the radical cation **1a**<sup>+</sup> that emerges from the SET. The reduction potential of the excited Bn-DHP **1a** [ $E_{\text{red}}^*(\mathbf{1a}^*/\mathbf{1a}^+)$ ] was estimated as  $-2.0$  V vs. Ag/Ag<sup>+</sup> in CH<sub>3</sub>CN on the basis of electrochemical and spectroscopic measurements. This establishes the thermodynamic feasibility of the SET reduction of both **2a** ( $E_{\text{red}} = -1.87$  V vs. SCE in CH<sub>3</sub>CN)<sup>[13a]</sup> and **2b** ( $E_{\text{red}} = -0.64$  V vs. SCE in CH<sub>3</sub>CN).<sup>[13b]</sup>

These studies demonstrate that alkyl-DHPs **1** possess a dual reactivity profile, since excitation with a violet-light-emitting diode turns them into strong reducing agents that are able to activate reagents through SET manifolds while undergoing homolytic cleavage to directly generate alkyl radicals. We reasoned that this unique reactivity behavior could be fully exploited in the framework of a photochemical nickel-catalyzed cross-coupling method. Merging nickel catalysis<sup>[14]</sup> and photoredox catalysis<sup>[15]</sup> has recently emerged as a versatile platform for developing highly enabling cross-coupling methods.<sup>[16]</sup> The success of this approach relies on the ability of Ni<sup>0</sup> complexes to undergo facile oxidative addition with aromatic halides while engaging in radical-capture mechanisms. This sequence results in the formation of Ni<sup>III</sup> complexes, which afford the cross-coupled products through rapid reductive elimination. At the same time, the chemistry relies on the reactivity of visible-light-activated photocatalysts that, upon excitation, can generate alkyl radicals through SET activation of non-traditional cross-coupling nucleophiles. Crucially, the photoredox catalyst is also involved in modulating the oxidation state of nickel complexes by SET reduction, an essential step to restore the original Ni<sup>0</sup> catalyst. Since alkyl-DHPs **1** can act as both photoreductants and precursors of alkyl radicals upon excitation, we surmised that their photochemistry could enable a nickel-catalyzed C(sp<sup>2</sup>)-C(sp<sup>3</sup>) cross-coupling process *without the need for any external photoredox catalyst*.<sup>[17]</sup> The proposed mechanism, depicted in Scheme 2, would start with the photoexcitation of alkyl-DHPs **1**. In the first catalytic cycle, the resulting excited-state intermediate **1\*** ( $E_{\text{red}}^*$  (**1a\***/**1a<sup>+</sup>**) = -2.0 V vs. Ag/Ag<sup>+</sup> in CH<sub>3</sub>CN) would reduce the Ni<sup>II</sup> precatalyst through two discrete SET events to afford the active Ni<sup>0</sup> intermediate ( $E_{\text{red}}$  Ni<sup>II</sup>/Ni<sup>0</sup> = -1.2 V versus SCE in DMF).<sup>[18]</sup> The concomitant formation of the radical cation **1<sup>+</sup>** would secure the formation of the alkyl radical, which is then intercepted by the Ni<sup>0</sup> complex. The emerging Ni<sup>I</sup> organometallic species would then undergo oxidative addition with the aryl bromide **4**. This would lead to the Ni<sup>III</sup> intermediate, which, after reductive elimination, forges the desired C(sp<sup>2</sup>)-C(sp<sup>3</sup>) bond in the cross-coupled product **5**. Finally, SET reduction of the generated Ni<sup>I</sup> complex by the excited alkyl-

DHPs **1\*** would complete the nickel catalytic cycle while regenerating the C(sp<sup>3</sup>) radical.


Pleasingly, the proposed light-triggered cross-coupling process turned out to be feasible at ambient temperature in CH<sub>3</sub>CN using NiCl<sub>2</sub>·DME as the metal catalyst (5 mol%), bipyridine (10 mol%) and 2,6-lutidine as the base (1 equiv). Control experiments established the necessity of light irradiation (using high-powered LED,<sup>[19]</sup>  $\lambda_{\text{max}}$  = 405 nm, with an irradiance of 50 mW cm<sup>-2</sup>), with the process being completely inhibited in the dark. As depicted in Figure 3, a variety of aryl



**Figure 3.** Survey of the aryl bromides **4** and the 4-alkyl-1,4-dihydropyridines **1** that can participate in the nickel-catalyzed cross-coupling process enabled by the photochemical activity of **1**. Reactions performed on a 0.5 mmol scale using 1.5 equiv of **1** and an irradiance of 50 mW cm<sup>-2</sup>. Yields refer to isolated products. [a] The NMR yield is given, since the compound was isolated together with 10% of a byproduct. [b] Reaction time 72 hours. [c] 0.1 mmol scale reaction performed over 16 hours (average of two runs).

bromides **4** were successfully coupled with Bn-DHP **1a** to afford the corresponding benzylated products **5a-g** in good to high yields. The process tolerated a wide range of functional groups, including cyano, aldehyde, ester, and chloride moieties. Moreover, a *ortho* substituent did not hinder the reaction, yielding the coupling product **5f** in good yield. To glean further insight into this process, we measured the quantum yield ( $\Phi$ ) of the photochemical cross-coupling reaction leading to adduct **5a**. A value of 0.0034 ( $\lambda$  = 405 nm in CH<sub>3</sub>CN, average of three runs) was determined, which is consonant with the mechanism proposed in Scheme 2.


We then evaluated the possibility of using alkyl-DHP substrates **1** bearing different alkyl fragments to benzyl at the C4-position. Both linear and cyclic fragments could be successfully coupled with aryl bromides (adducts **5h-l**). An alkene, an acetal moiety, and an oxygen-based heterocycle were all well-tolerated, leading to the corresponding products **5h**, **5j** and **5k**, respectively. In addition, a primary radical successfully engaged in this coupling when adorned with a stabilizing  $\alpha$ -oxygen atom (adduct **5l**). These results are mechanistically relevant because they demonstrate that



**Scheme 2.** Proposed reaction mechanism for the nickel-catalyzed C(sp<sup>2</sup>)-C(sp<sup>3</sup>) cross-coupling driven by the photochemical activity of alkyl-DHPs **1**.

different 4-alkyl-1,4-dihydropyridines **1** can be excited upon visible-light irradiation and their photochemical activity successfully integrated into a nickel catalytic cycle to enable  $C(sp^2)-C(sp^3)$  cross-coupling processes.

We then wondered whether the same photochemical method could be successfully translated to the nickel-mediated cross-coupling of **1** with acyl chlorides **6**.<sup>[20]</sup> As shown in Figure 4, an array of dialkyl and aryl/alkyl ketones **7** could be efficiently prepared under mild conditions and relying only on the visible-light excitation of alkyl-DHP substrates **1**.



**Figure 4.** Survey of the acyl chlorides **6** and the 4-alkyl-1,4-dihydropyridines **1** that can participate in the nickel-catalyzed cross-coupling process to afford ketones **7**. Reactions performed on a 0.1 mmol scale using 1.5 equiv of **1** and an irradiance of  $50 \text{ mWcm}^{-2}$ . Yields refer to isolated products (average of two runs). [a] 0.5 mmol scale reaction performed over 24 hours.

In summary, we have demonstrated that 4-alkyl-1,4-dihydropyridines can unlock rich photochemistry upon light excitation. Selective absorption of violet light turns them into strong reducing agents that activate reagents through single-electron transfer manifolds while undergoing homolytic cleavage to generate  $C(sp^3)$ -centered radicals. This light-triggered dual-reactivity profile was integrated into a nickel catalytic cycle to enable  $C(sp^2)-C(sp^3)$  cross-coupling reactions without the need for an external photoredox catalyst. We are conducting ongoing studies to further exploit the photochemical activity of alkyl-DHPs in other radical-based carbon-carbon bond-forming processes.

## Acknowledgements

Financial support was provided by the Generalitat de Catalunya (CERCA Program), MINECO (CTQ2016-75520-P), and the European Research Council (ERC-2015-CoG 681840—CATA-LUX). L.B. thanks MINECO for a predoctoral fellowship (CTQ2013-45938-P). The authors thank Dr. Fernando Bozoglian for assistance with photophysical studies

and Dr. Maria Dolores Gonzalez Candela for assistance with EPR experiments.

## Conflict of interest

The authors declare no conflict of interest.

**Keywords:** cross-coupling · dihydropyridines · nickel catalysis · photochemistry · synthetic methods

- [1] a) V. Balzani, P. Ceroni, A. Juris, in *Photochemistry and Photophysics*, Wiley-VCH, Weinheim, **2014**; b) P. Klán, J. Jacob, in *Photochemistry of Organic Compounds*, Wiley, Hoboken, **2010**.
- [2] A. Albini, M. Fagnoni, in *Handbook of Synthetic Photochemistry*, Wiley-VCH, Weinheim, **2010**.
- [3] N. J. Turro, V. Ramamurthy, J. C. Scaiano, in *Modern Molecular Photochemistry of Organic Molecules*, University Science Books, Sausalito, CA, **2010**, Chapter 7.
- [4] a) M. Silvi, E. Arceo, I. D. Jurberg, C. Cassani, P. Melchiorre, *J. Am. Chem. Soc.* **2015**, *137*, 6120–6123; b) G. Filippini, M. Silvi, P. Melchiorre, *Angew. Chem. Int. Ed.* **2017**, *56*, 4447–4451; *Angew. Chem.* **2017**, *129*, 4518–4522; c) M. Silvi, C. Verrier, Y. P. Rey, L. Buzzetti, P. Melchiorre, *Nat. Chem.* **2017**, *9*, 868–873; d) G. Filippini, M. Nappi, P. Melchiorre, *Tetrahedron* **2015**, *71*, 4535–4542.
- [5] C. Zheng, S.-L. You, *Chem. Soc. Rev.* **2012**, *41*, 2498–2518.
- [6] a) S. Fukuzumi, K. Hironaka, T. Tanaka, *J. Am. Chem. Soc.* **1983**, *105*, 4722–4727; b) J. Jung, J. Kim, G. Park, Y. You, E. J. Cho, *Adv. Synth. Catal.* **2016**, *358*, 74–80; c) M. A. Emmanuel, N. R. Greenberg, D. G. Oblinsky, T. K. Hyster, *Nature* **2016**, *540*, 414–417; d) L. I. Panferova, A. V. Tsymlar, V. V. Levin, M. I. Struchkova, A. D. Dilman, *Org. Lett.* **2016**, *18*, 996–999; e) W. Chen, H. Tao, W. Huang, G. Wang, S. Li, X. Cheng, G. Li, *Chem. Eur. J.* **2016**, *22*, 9546–9550.
- [7] Examples of 4-alkyl-1,4-dihydropyridines serving as radical precursors under the action of external photoredox catalysts: a) K. Nakajima, S. Nojima, K. Sakata, Y. Nishibayashi, *ChemCatChem* **2016**, *8*, 1028–1032; b) Á. Gutiérrez-Bonet, J. C. Tellis, J. K. Matsui, B. A. Vara, G. A. Molander, *ACS Catal.* **2016**, *6*, 8004–8008; c) W. Chen, Z. Liu, J. Tian, J. Ma, X. Cheng, G. Li, *J. Am. Chem. Soc.* **2016**, *138*, 12312–12315; d) K. Nakajima, S. Nojima, Y. Nishibayashi, *Angew. Chem. Int. Ed.* **2016**, *55*, 14106–14110; *Angew. Chem.* **2016**, *128*, 14312–14316; For a review, see: e) W. Huang, X. Cheng, *Synlett* **2017**, *28*, 148–158.
- [8] Á. Gutiérrez-Bonet, C. Remeur, J. K. Matsui, G. A. Molander, *J. Am. Chem. Soc.* **2017**, *139*, 12251–12258.
- [9] N. Tewari, N. Dwivedi, R. Tripathi, *Tetrahedron Lett.* **2004**, *45*, 9011–9014.
- [10] Our attempts to perform Stern–Volmer quenching studies using different electron acceptors, including the electron-poor cyano substrate **2a**, met with failure. For all the tested 4-alkyl-1,4-dihydropyridines, including **1a**, we observed an increase in fluorescence intensity upon multiple irradiation of the substrate alone. This phenomenon is likely ascribable to the formation of a strongly emitting compound, which is generated upon photodegradation of **1**. This emission behavior, which is further detailed in the Supporting Information (Section B4, Figure S6), precluded a reliable quenching measurement.
- [11] K. H. Kim, X. Bai, R. C. Brown, *J. Anal. Appl. Pyrolysis* **2014**, *110*, 254–263.
- [12] For selected examples, see: a) C. Pac, A. Nakasone, H. Sakurai, *J. Am. Chem. Soc.* **1977**, *99*, 5806–5808; b) R. M. Borg, D. R.

- Arnold, T. S. Cameron, *Can. J. Chem.* **1984**, *62*, 1785–1802; c) M. Mella, M. Fagnoni, A. Albini, *J. Org. Chem.* **1994**, *59*, 5614–5622; d) A. McNally, C. K. Prier, D. W. C. MacMillan, *Science* **2011**, *334*, 1114–1117.
- [13] a) P. McDevitt, B. M. Vitimberga, *J. Heterocycl. Chem.* **1990**, *27*, 1903–1908; b) J. D. Bagnato, W. W. Shum, M. Strohmeier, D. M. Grant, A. M. Arif, J. S. Miller, *Angew. Chem. Int. Ed.* **2006**, *45*, 5322–5326; *Angew. Chem.* **2006**, *118*, 5448–5452.
- [14] S. Z. Tasker, E. A. Standley, T. F. Jamison, *Nature* **2014**, *509*, 299–309.
- [15] M. H. Shaw, J. Twilton, D. W. C. MacMillan, *J. Org. Chem.* **2016**, *81*, 6898–6926.
- [16] For a review: a) J. Twilton, C. Le, P. Zhang, M. H. Shaw, R. W. Evans, D. W. C. MacMillan, *Nat. Rev. Chem.* **2017**, *1*, 0052; Selected examples of synergistic nickel/photoredox catalysis: b) J. C. Tellis, D. N. Primer, G. A. Molander, *Science* **2014**, *345*, 433–436; c) Z. Zuo, D. T. Ahneman, L. Chu, J. A. Terrett, A. G. Doyle, D. W. C. MacMillan, *Science* **2014**, *345*, 437–440; d) V. Corcé, L.-M. Chamoreau, E. Derat, J.-P. Goddard, C. Ollivier, L. Fensterbank, *Angew. Chem. Int. Ed.* **2015**, *54*, 11414–11418; *Angew. Chem.* **2015**, *127*, 11576–11580; e) M. Jouffroy, D. N. Primer, G. A. Molander, *J. Am. Chem. Soc.* **2016**, *138*, 475–478.
- [17] 4-Alkyl-1,4-dihydropyridines were recently used as alkyl radical precursors in nickel-catalyzed cross-coupling reactions; the reported methods, however, require the use of an external photoredox catalyst; see Refs. [7b,d].
- [18] a) P. Johnston, R. T. Smith, S. Allmendinger, D. W. C. MacMillan, *Nature* **2016**, *536*, 322–325; b) M. Durandetti, M. Devaud, J. Perichon, *New J. Chem.* **1996**, *20*, 659–667.
- [19] The use of an LED strip ( $\lambda_{\text{max}} = 405 \text{ nm}$ ) severely reduced the reactivity of the nickel cross-coupling reaction. Details of the importance of the light intensity and the use of HP LEDs are reported in Section D2 in the Supporting Information.
- [20] Previous methods required the use of an external photoredox catalyst and alkyltrifluoroborates as radical precursors: a) J. Amani, E. Sodagar, G. A. Molander, *Org. Lett.* **2016**, *18*, 732–735; b) J. Amani, G. A. Molander, *J. Org. Chem.* **2017**, *82*, 1856–1863.

Manuscript received: September 15, 2017

Accepted manuscript online: October 6, 2017

Version of record online: ■■ ■■, ■■■■

## Communications




## Synthetic Photochemistry

L. Buzzetti, A. Prieto, S. R. Roy,  
P. Melchiorre\*



Radical-Based C–C Bond-Forming  
Processes Enabled by the Photoexcitation  
of 4-Alkyl-1,4-dihydropyridines



**Powered by Light:** 4-alkyl-1,4-dihydropyridines **1** are primarily understood as hydride sources in their ground state. Excitation with a violet-light-emitting diode transforms them into strong reducing agents that can activate

reagents through single-electron transfer while undergoing homolytic cleavage to generate alkyl radicals. This process was used to trigger radical-based C–C bond-forming processes, including nickel-catalyzed cross-coupling reactions.