

Chapter 101

Potential Cellular Functions of *N*-Ethylmaleimide Sensitive Factor in the Photoreceptor

Shun-Ping Huang and Cheryl M. Craft

Keywords Arrestin 1 • *N*-ethylmaleimide sensitive factor • Retinitis pigmentosa protein 2 • Phototransduction • Oguchi disease • NSF attachment proteins receptor • Retinitis pigmentosa • Synapses • ATPase • Protein trafficking

101.1 Introduction

Vesicle traffic and fusion are essential, not only for cellular homeostasis but also for neuronal signal transmission across the synaptic junction of nerves, cell growth, and membrane repair. The basic fusion process is mediated by vesicle (v)-soluble *N*-ethylmaleimide sensitive factor attachment proteins receptors

S.-P. Huang

Mary D. Allen Laboratory for Vision Research, Doheny Eye Institute,
Keck School of Medicine of the University of Southern California,
Los Angeles, CA 90033-9224, USA

Department of Ophthalmology, Keck School of Medicine of the University
of Southern California, Los Angeles, CA 90033-9224, USA

C.M. Craft (✉)

Mary D. Allen Laboratory for Vision Research, Doheny Eye Institute,
Keck School of Medicine of the University of Southern California,
Los Angeles, CA 90033-9224, USA

Department of Ophthalmology, Keck School of Medicine of the University
of Southern California, Los Angeles, CA 90033-9224, USA

Department of Cell and Neurobiology, Keck School of Medicine of the University
of Southern California, Los Angeles, CA 90033-9224, USA
e-mail: ccraft@usc.edu




Fig. 101.1 Structure of NSF domains. Each NSF monomer is comprised of three domains: an N-terminal domain (amino acid 1–205) that is responsible for the interaction with a-SNAP and SNAREs; and two homologous ATP-binding domains, D1 (amino acids 206–477), in which the hydrolytic activity is associated with NSF-driven SNARE complex disassembly, and D2 (amino acids 478–744), which is responsible for hexamer formation. Critical residues involved in post-translational processing include Cys21, 91 and 264, Tyr83, Ser237, and Ser569

(SNAP receptors; SNAREs) on the secretory vesicle with their cognate target (t)-SNAREs on the target membrane (Sollner et al. 1993), which assemble in trans-configuration into four-helix bundle complexes, bringing the membranes into close proximity (Jahn and Scheller 2006; Rizo and Rosenmund 2008; Martens and McMahon 2008), which then directly or indirectly leads to fusion. After membrane fusion, all SNAREs constituting one complex are anchored in a relaxed *cis*-configuration to one membrane. Disassembly of these SNAREs complexes for subsequent vesicle transport and recycling is achieved by the concerted action of α -SNAP and NSF.

N-ethylmaleimide sensitive factor (NSF) is a homo-hexameric member of the ATPase-associated family with various cellular activities protein (AAA) family, required for intracellular membrane fusion. NSF functions as a SNAP receptor (SNARE) chaperone that binds through soluble NSF attachment proteins (SNAPs) to SNARE complexes and utilizes the energy of ATP hydrolysis to dissociate SNARE complexes after membrane fusion, thus facilitating SNAREs recycling.

Each NSF protomer contains an N-terminal domain (NSF-N) and two AAA-domains: a catalytic NSF-D1 and a structural NSF-D2 (Fig. 101.1). The amino acid residues 1–205 in the NSF-N domain are required for SNAP–SNARE binding; residues 206–477 in the NSF-D1 domain are responsible for the majority of the ATP hydrolysis, while the carboxy-terminal residues 478–744 in the NSF-D2 domain are required for hexamerization (Tagaya et al. 1993; Nagiec et al. 1995). Within the N-terminal subdomain of both NSF-D1 and NSF-D2, there is a highly conserved region called Second Region of Homology (SRH), which is highly conserved in AAA proteins (Hanson and Whiteheart 2005). By using detailed mutagenesis analysis, Zhao and collaborators (Zhao et al. 2010) showed that a positively charged surface on NSF-N, bounded by R67 and K105, and the conserved central

pore motifs in NSF-D1 (Y296 and G298) are involved in SNAP–SNARE binding but not basal ATP hydrolysis. Sensor 1 is at the N-terminus of the SRH and is important for basal ATPase activity and nucleotide binding. At its carboxy-terminus are two arginine residues, termed Arginine Fingers, which are critical for ATP hydrolysis by the NSF hexamer. Sensor 2 comes from C-terminal helical subdomain and plays a role in ATP- and SNAP-dependent SNARE complex binding and disassembly.

NSF binds to SNARE complexes via its adaptor protein, α -SNAP, only in the presence of ATP (Nagiec et al. 1995). The intrinsic ATPase activity of NSF is low (Tagaya et al. 1993). NSF binding to immobilized α -SNAP stimulates the ATPase activity (Morgan et al. 1994), and maximal stimulation of ATPase activity is achieved when both α -SNAP and SNARE complexes are included (Matveeva and Whiteheart 1998).

101.2 NSF in Photoreceptor Synaptic Regulation

In the G-protein-coupled receptor (GPCR) phototransduction cascade, visual Arrestin1 (Arr1) binds to and deactivates phosphorylated, light-activated opsins, a process that is critical for effective recovery and normal vision. We discovered a novel synaptic protein–protein interaction between Arr1 and NSF that is enhanced in a dark environment when photoreceptors are depolarized and the rate of exocytosis is elevated compared to a light environment (Huang et al. 2010; Huang 2010) (Fig. 101.2a, b). In Figs. 101.2 and 101.3, representative images of sections from either dark- or light-adapted retinas highlight differences in immunohistochemical retinal dual localization in mouse of Arr1 and NSF, respectively (Figs. 101.2a–d and 101.3a–d). We also provided convincing *in vitro* evidence supporting the interaction of Arr1 with NSF to modulate its ATPase activity and to drive disassembly of the SNARE complex. Furthermore, we observed with *in vivo* studies using FM-143 imaging that synaptic vesicle recycling in photoreceptors is dramatically decreased with depolarization and photopic electroretinogram (ERG) b-wave analysis in *Arr1*^{-/-} compared to control mice is defective in light adaptation (Brown et al. 2010). Not only does Arr1 bind to the junction of NSF N-terminal and the first ATPase domains in an ATP-dependent manner, but Arr1 also enhances both NSF ATPase and NSF disassembly activities. In mouse retinas with no Arr1 expression, the expression levels of NSF and other synapse-enriched genes are markedly reduced and lead to a substantial decrease in the exocytosis rate. These cumulative findings demonstrate that normal photoreceptor synaptic function involves the ability of Arr1 to regulate and to enhance the dark-associated activity of NSF in the photoreceptor synapse.




Fig. 101.2 NSF and Arr1 localization in dark-adapted retinas. Immunohistochemical fluorescent labeling of retinal sections with dual localization of NSF and Arr1 staining in the WT mice dark-adapted (DA) overnight and killed in the dark. Adult WT mouse retina frozen sections were triple labeled fluorescently with the antimouse Arr1 MAb D9F2 (red, 1:20,000), antirabbit NSF PAB (green, 1:2,500, Abcam), and secondary antibodies conjugated to Alexa Fluor 488 or 568, respectively (1:500, invitrogen), and DAPI for the nuclei (blue, (a, b)). NSF immunoreactive staining pattern of NSF is mainly in the OPL and IPL in dark-adapted retinas (c, d). Arr1 immunoreactive pattern is predominantly in the inner segment, perinuclear area and a fraction in the photoreceptor terminal in DA retinas ((c), higher magnification; large arrows). OS outer segment; IS inner segments; ONL outer nuclear layer; OPL outer plexiform layer; INL inner nuclear layer; IPL inner plexiform layer; GLC ganglion cell layer. Scale bar, 20 μm in upper panels; 10 μm in lower panels

101.3 NSF in Photoreceptor Membrane Protein Trafficking

While the major function of NSF is involved in vesicle transport and recycling, it also interacts with other proteins such as the AMPA receptor subunit (Osten et al. 1998), β -arrestin1 (McDonald et al. 1999), GluR2 (Nishimune et al. 1998), and β 2-AR (Cong et al. 2001) and is thought to affect their trafficking pattern. More evidence suggests that NSF may be regulated by transient posttranslational modifications such as phosphorylation and nitrosylation. These modifications are ideal mechanisms for reversible regulation of membrane trafficking.




Fig. 101.3 NSF and Arr1 localization in light-adapted retinas. Immunohistochemical fluorescent labeling of NSF and Arr1 in retinal sections from WT mouse exposed to light for 2 h prior to killing (light-adapted [LA]). Adult WT mouse retina frozen sections were triple labeled fluorescently as in Fig. 101.2 ((a, b) DAPI stains nuclei blue). NSF immunoreactive staining pattern of NSF is mainly in the OPL and IPL in light-adapted retinas (c, d). Arr1 immunoreactive pattern is translocated to the outer segment in LA retinas. Arr1 immunoreactivity is extensively dual localized with NSF immunological staining and limited punctate dual staining in the OPL in LA retina ((d) higher magnification; *smaller arrows*). Abbreviations are listed in Fig. 101.2. Scale bar, 20 μm in *upper* panels; 10 μm in *lower* panels

Recently, Molday and his collaborators observed that NSF specifically interacts with RP2 in the retina (Holopainen et al. 2010), which is a ubiquitously expressed protein encoded by a gene associated with X-linked retinitis pigmentosa (Breuer et al. 2002; Miano et al. 2001; Schwahn et al. 1998; Sharon et al. 2003). RP2 is known to bind to the GTP-bound form of ADP ribosylation factor like 3 (Arl3), a member of the Arl subfamily of Ras-related GTP-binding proteins (Bartolini et al. 2002; Kuhnel et al. 2006; Veltel et al. 2008) and acts as an efficient GTPase-activating protein (GAP) for Arl3 (Veltel et al. 2008). NSF and RP2 was shown to colocalize in the ciliary region, inner segment, outer nuclear layer, and synaptic region in the photoreceptor. Furthermore, RP2 binds to the N-terminal domain of NSF and this binding is abolished for the E138G and Δ I137 mutation of RP2 known

to cause X-linked retinitis pigmentosa. These data may indicate RP2 binding to NSF, and Arl3 may play a crucial role in the vesicle trafficking of proteins to the photoreceptor ciliary region as well as in the photoreceptor synapse.

101.4 Summary

Recent work has established potential new functional roles for NSF in the photoreceptor. First, the interaction of Arr1 and NSF is ATP-dependent, and the N-terminal domain of Arr1 interacts with the N and D1 junctional domains of NSF. The Arr1–NSF interactions are greater in the photoreceptor synaptic terminal in the dark. Furthermore, Arr1 enhances the NSF ATPase activity and increases the NSF disassembly activities, which are critical for NSF functions in sustaining a higher rate of exocytosis in the photoreceptor synapses and the compensatory endocytosis to retrieve vesicle membrane and vesicle proteins for vesicle recycling. These data demonstrate the Arr1 and NSF interaction are necessary for both maintenance and modulation of normal photoreceptor synaptic regulation. Second, NSF colocalizes and specifically binds to RP2, especially in the ciliary and synaptic region of the photoreceptor, and NSF–RP2 interaction may play an important role in membrane protein trafficking in the photoreceptor.

Inherited retinal degeneration affects about 1 in 2,000–3,000 individuals in the world and is the leading cause of visual loss in young people and accounts for a large proportion of blindness in adult life. These studies accelerate our ability to gain insight into the diverse roles of the NSF in the photoreceptor cells and enable us to understand more precisely the molecular mechanisms underlying night blindness associated with clinically diagnosed Oguchi disease or other forms of retinitis pigmentosa.

Acknowledgments We thank members of the Mary D. Allen Laboratory for scientific discussions, Bruce M. Brown for his technical expertise, Lawrence Rife for ERG analysis, Ernesto Barron for preparation of figures, and Jeannie Chen for the *Arr1*^{-/-} mice. CMC is the Mary D. Allen Chair in Vision Research, DEI, and a Research to Prevent Blindness (RPB) Senior Scientific Investigator. This work was supported, in part, by NIH Grant EY015851 (CMC), EY03040 (DEI), RPB (DEI & CMC), Dorie Miller, Tony Gray Foundation, Mary D. Allen Foundation (Dr. Richard Newton Lolley Memorial Scholarship [SPH]), and a RD2010 Travel Award (SPH).

References

- Bartolini F, Bhamidipati A, Thomas S et al (2002) Functional overlap between retinitis pigmentosa 2 protein and the tubulin-specific chaperone cofactor C. *J Biol Chem* 277:14629–14634
- Breuer DK, Yashar BM, Filippova E et al (2002) A comprehensive mutation analysis of RP2 and RPGR in a North American cohort of families with X-linked retinitis pigmentosa. *Am J Hum Genet* 70:1545–1554
- Brown BM, Ramirez T, Rife, L et al (2010) Visual arrestin 1 contributes to cone photoreceptor survival and light adaptation. *Invest Ophthalmol Vis Sci* 51:2372–2380

- Cong M, Perry SJ, Hu LA et al (2001) Binding of the beta2 adrenergic receptor to N-ethylmaleimide-sensitive factor regulates receptor recycling. *J Biol Chem* 276:45145–45152
- Hanson PI, Whiteheart SW (2005) AAA+ proteins: have engine, will work. *Nat Rev Mol Cell Biol* 6:519–529
- Holopainen JM, Cheng CL, Molday LL et al (2010) Interaction and Localization of the Retinitis Pigmentosa Protein RP2 and NSF in Retinal Photoreceptor Cells. *Biochemistry* 49:7439–7447
- Huang SP (2010) Exploring Alternative Roles of Visual Arrestin 1 in Photoreceptor Synaptic Regulation and Deciphering the Molecular Pathway of Retinal Degeneration Using Mouse Knockout Technology. Ph.D. dissertation. University of Southern California
- Huang SP, Brown BM, Craft CM (2010) Visual Arrestin 1 acts as a modulator for N-ethylmaleimide-sensitive factor in the photoreceptor synapse. *J Neurosci* 30:9381–9391
- Jahn R, Scheller RH (2006) SNAREs--engines for membrane fusion. *Nat Rev Mol Cell Biol* 7:631–643
- Kuhnel K, Veltel S, Schlichting I, Wittinghofer A (2006) Crystal structure of the human retinitis pigmentosa 2 protein and its interaction with Arl3. *Structure* 14:367–378
- Martens S, McMahon HT (2008) Mechanisms of membrane fusion: disparate players and common principles. *Nat Rev Mol Cell Biol* 9:543–556
- Matveeva E, Whiteheart SW (1998) The effects of SNAP/SNARE complexes on the ATPase of NSF. *FEBS Lett* 435:211–214
- McDonald PH, Cote NL, Lin FT et al (1999) Identification of NSF as a beta-arrestin1-binding protein. Implications for beta2-adrenergic receptor regulation. *J Biol Chem* 274:10677–10680
- Miano MG, Testa F, Filippini F et al (2001) Identification of novel RP2 mutations in a subset of X-linked retinitis pigmentosa families and prediction of new domains. *Hum Mutat* 18:109–119
- Morgan A, Dimaline R, Burgoyne RD (1994) The ATPase activity of N-ethylmaleimide-sensitive fusion protein (NSF) is regulated by soluble NSF attachment proteins. *J Biol Chem* 269:29347–29350
- Nagiec EE, Bernstein A, Whiteheart SW (1995) Each domain of the N-ethylmaleimide-sensitive fusion protein contributes to its transport activity. *J Biol Chem* 270:29182–29188
- Nishimune A, Isaac JT, Molnar E et al (1998) NSF binding to GluR2 regulates synaptic transmission. *Neuron* 21:87–97
- Osten P, Srivastava S, Inman GJ et al (1998) The AMPA receptor GluR2 C terminus can mediate a reversible, ATP-dependent interaction with NSF and alpha- and beta-SNAPs. *Neuron* 21:99–110
- Rizo J, Rosenmund C (2008) Synaptic vesicle fusion. *Nat Struct Mol Biol* 15:665–674
- Schwahn U, Lenzner S, Dong J et al (1998) Positional cloning of the gene for X-linked retinitis pigmentosa 2. *Nat Genet* 19:327–332
- Sharon D, Sandberg MA, Rabe VW et al (2003) RP2 and RPGR mutations and clinical correlations in patients with X-linked retinitis pigmentosa. *Am J Hum Genet* 73:1131–1146
- Sollner T, Bennett MK, Whiteheart SW et al (1993) A protein assembly-disassembly pathway in vitro that may correspond to sequential steps of synaptic vesicle docking, activation, and fusion. *Cell* 75:409–418
- Tagaya M, Wilson DW, Brunner M et al (1993) Domain structure of an N-ethylmaleimide-sensitive fusion protein involved in vesicular transport. *J Biol Chem* 268:2662–2666
- Veltel S, Gasper R, Eisenacher E et al (2008) The retinitis pigmentosa 2 gene product is a GTPase-activating protein for Arf-like 3. *Nat Struct Mol Biol* 15:373–380
- Zhao C, Matveeva EA, Ren Q et al (2010) Dissecting the N-ethylmaleimide-sensitive factor: required elements of the N and D1 domains. *J Biol Chem* 285:761–772