

**Addressing cumulative effects, maritime conflicts and ecosystem services threats through MSP-oriented geospatial webtools.**

Stefano Menegon<sup>1,\*</sup>, Daniel Depellegrin<sup>1,\*</sup>, Giulio Farella, Alessandro Sarretta, Chiara Venier, Andrea Barbanti

CNR - National Research Council of Italy, ISMAR - Institute of Marine Sciences Arsenal - Tesa 104, Castello 2737/F, I-30122 Venice, Italy

<sup>1</sup> Joined first authors

\* Corresponding authors: [stefano.menegon@ismar.cnr.it](mailto:stefano.menegon@ismar.cnr.it) (Stefano Menegon), [daniel.depellegrin@ve.ismar.cnr.it](mailto:daniel.depellegrin@ve.ismar.cnr.it) (Daniel Depellegrin)

**Keywords:** Webtools, Geoplatfrom, Cumulative effects assessment, maritime use conflict analysis, marine ecosystem services, MSP

**Abstract**

To solve conservation and planning challenges in the marine environment, researchers are increasingly developing geospatial tools to address impacts of anthropogenic activities on marine biodiversity. The paper presents a comprehensive set of built-in geospatial webtools to support Maritime Spatial Planning (MSP) and environmental management objectives implemented into the Tools4MSP interoperable GeoPlatform. The webtools include cumulative effects assessment (CEA), maritime use conflict (MUC) analysis, MSFD pressure-driven CEA and a CEA-based marine ecosystem service threat analysis (MES-Threat) . The tools are tested for the Northern Adriatic (NA) Sea, one of the most industrialized sea areas of Europe using a case study driven modelling strategy. Overall results show that coastal areas within 0-9 nm in the Gulf of Trieste, Grado-Marano and Venice lagoon and Po Delta outlet are subjected to intense cumulative effects and high sea use conflicts mainly from port activities, fishery, coastal and maritime tourism and maritime shipping. Linking MES into CEA provided novel information on locally threatened high MES supporting and provisioning habitats such *Cymodocea* beds and infralittoral fine sands, threats to cultural MES are most pronounced in coastal areas. Results are discussed for their geospatial relevance for regional planning, resource management and their applicability within MSP and environmental assessment.

**1. Introduction**

Current conservation and planning challenges of the marine environments require flexible tools that ensure to different types of user the access, management, sharing, processing and visualization of a multitude of spatial and non-spatial dataset. Ideally, these datasets are stored within platforms capable to organize a multitude of data and convey them into easily and quickly accessible graphical user interfaces (GUI). The use of Maritime Spatial Planning (MSP, Directive 2014/89/EU) as practical process to achieve environmental, social and economic objectives and minimize conflicts (Hansen et al., 2017) in European seas has posed novel demands to amount, quality and sources of data. Despite the ongoing governance process, considerable work has been done by the scientific community for the development of Spatial Data Infrastructure (SDI; Fowler et al., 2010) in support of a knowledge-based implementation of national and regional plans.

In recent years the application of cumulative effects assessment and sea use conflict analysis have emerged as common analytical tool to support decision-makers in the development of spatial plans and in support of the ecosystem-based management of marine resources. This is also reflected in an emerging number of decision support tools enabling user to perform cumulative effects assessment in various contexts. An extended review of decision support systems performed by Krueger and

Schouten-de Groot (2011) showed that out of the 118 tools in support of MSP, about 46 (39%)  
35 implement CEA models whether serving decision support or scenario analysis development and  
management priorities identification. Examples of MSP oriented decision support system include  
sector specific tools such as Windspeed (Spatial Development of Offshore Wind Energy in Europe;  
www.windspeed.eu) for the identification of suitable areas for wind energy in the North Sea, MARA  
(Marine Aggregate Extraction Risk Assessment framework; www.mara-framework.org.uk) for  
40 probabilistic environmental risk assessment or the Isis-fish (Krueger and Schouten-De Groot, 2011), a  
predictive tool of fish population development under different management scenarios. Other tools that  
allow more comprehensive CEA analysis include the HARMONY tool (Development and  
demonstration of Marine Strategy Framework Directive tools for harmonization of the initial  
assessment in the eastern parts of the Greater North Sea sub-region; Andersen et al., 2013) for human  
45 impact assessment in the eastern North Sea sub-region or the SYMPHONY tool (MSP Platform,  
2016).

Sea use conflict analysis has been extensively applied in different geographical contexts (Hadjimitsis  
et al., 2015; White et al., 2012) based on different decision support systems, such as the GRID tool  
(Georeferenced Interactions Database; Gramolini et al., 2010) providing a platform to spatialize use-  
50 use conflict in sea areas, the MaRS geotool (Marine Resource System;  
www.thecrownstate.co.uk/mars) to support identification and resolution of spatial conflicts and  
AquaSpace that enables integrated assessment of risks and opportunities for proposed aquaculture  
sites (Gimpel et al., 2018).

Also the recent growth of ecosystem services research contributed to the development of several  
55 geospatial tools in support of decision making in coastal and marine environments, such as the habitat  
risk assessment (HRA) tool from Marine InVEST toolset (Integrated Valuation of Ecosystem Services  
and Tradeoffs), that enables user to assess risks to marine ecosystems generated by different human  
activities (Wyatt et al., 2017), the SOLVES tool (Social Values for Ecosystem Services) for the  
analysis and mapping of non-market values of cultural ecosystem services (van Riper et al., 2012) or  
60 the MIMES model (Multi-Scale Integrated Models of Ecosystem Services) which supports MSP for  
tradeoff analysis among competing uses (Center for Ocean Solutions, 2011).

The very diverse suites and packages of geospatial tools poses considerable opportunities in the  
development of new generation decision-support systems for strategic planning and environmental  
conservation in the marine domain. However this diversity is source of difficulties in identifying  
65 suitable tools addressing specific decision-making objectives, may produce a fragmented utilization of  
several tools leading to input and outputs procedures that can require a costly data treatment for  
harmonizing the processing workflow.

In this research we present the functionalities of three webtools implemented in the Tools4MSP  
Geoplatfrom (tools4msp.eu), namely Cumulative Effects Assessment (CEA), Maritime Use Conflict  
70 (MUC) analysis and a Marine Ecosystem Services Threat analysis (MES-Threat). The webtools were  
tested in a case study for the Northern Adriatic (NA) Sea, one of the most crowded sea areas of  
Europe. The application of webtools is presented using a stepwise workflow based on a structured  
case study driven modelling strategy. The paper is organized as follows: Section 2 presents the overall  
workflow, the theoretical background of the webtools and the stepwise procedure for the webtools'  
75 setup, in Section 3 the geospatial and geostatistical results of the model setup are presented and  
Section 4 discusses the results for their relevance in MSP, environmental management and the  
applicability of the webtool along EIA and SEA.

## **2. Materials and Methods**

### 80 2.1. The Tools4MSP Geoplatform

The Tools4MSP Geoplatform is a community-based, open-source portal, based on GeoNode (GeoNode Development Team, 2018), a web-based Content Management System (CMS) for developing geospatial information systems (GIS) and for deploying spatial data infrastructures (SDI). The aim of the Geoplatform is to provide an operational set of webtools that can assist decision-  
85 makers and strategists in undertaking MSP-oriented case studies and support the development of environmental management strategies.

The webtools are integrated as GeoNode Plugin into the Geoplatform, that provides a graphical user interface (GUI) facilitating the usability of the Tools4MSP core functionalities for different user communities (Menegon, 2018b). The Plugin reflects the Tools4MSP modelling framework  
90 (Depellegrin et al., 2017, Menegon et al., 2016), a python-based Free and Open Source Software (FOSS) which combines several FOSS projects for geodata processing and scientific modelling: (1) NumPy and SciPy for efficient numerical computation (van der Walt, 2011); (2) Pandas and GeoPandas for data structures manipulation and data analysis (McKinney, 2010); (3) OWSLib which implements the client-side for OGC web services standard interfaces (OWSLib Development Team,  
95 2018); (4) Rectifiedgrid for efficient 2D grid-based analysis (Menegon, 2018a) and (5) the interactive visualization of the Tools4MSP results are created through Bokeh (Bokeh Development Team, 2018). The Tools4MSP software package can be freely downloaded from github (<https://github.com/CNR-ISMAR/tools4msp>).

In order to demonstrate the functionalities of the Tools4MSP Geoplatform, we present four  
100 operational steps for its utilization (Fig. 1): (Step 0) Webtool selection depending on the scope and objectives of the analysis; (Step 1) case study area selection, available for different geospatial scales (from seabasin to regional level); (Step 2) dataset configuration defining human uses, environmental components and MSFD-pressures themes used for modelling and (Step 3) generation of geospatial and statistical outputs to be used for data curation and re-analysis within a dedicated GIS software  
105 such as Quantum GIS (QGIS Development Team, 2018).




Fig. 1. Workflow for webtools application in the Tool4MSP Geoplatform: Step 0 - Webtool selection; Step 1 - Case study area selection; Step 2 - Study area selection & Dataset configuration and Step 3 - Geospatial and statistical outputs.

110 *2.2. Step 0: Webtools selection*

This step allows user to select a comprehensive set of webtools available in the Geoplatform (Fig. 2) namely a Cumulative Effects Assessment (CEA), Maritime Use Conflict (MUC) analysis and Marine Ecosystem Services Threat analysis (MES-Threat). In Fig. 2 (right) the buttons to prompt user to the webtool model run. In the following section a detailed description of theoretical and methodological

115 background of the webtools is provided.




Fig. 2. GUI for the webtools selection within the Tools4MSP Geoplatform. The interface includes also background information of the CEA, MUC and MES-Threat webtool.

120

### 2.2.1. Cumulative Effects Assessment (CEA)

The Tools4MSP Geoplatform implements a Cumulative Effects Assessment (CEA) for the analysis of cumulative effects generated by anthropogenic activities on marine environmental components. Its implementation is based on archetypical CEA implementations proposed in various geographical scales (Halpern et al., 2008; Andersen et al., 2013). In detail, we define CEA as a systematic procedure for identifying and evaluating the significance of effects from multiple pressures and/or activities on single or multiple receptors (Judd et al., 2015). The CEA incorporates two major improvements, such as the modulation of propagation of pressures through a distance model  $M(U_i, P_j, E_k)$  based on 2D Gaussian spatial convolution and the distinction of sensitivity scores ( $s_{j,k}$ ) into sensitivity values combined with use-specific relative pressure weight ( $w_{i,j,k}$ ). The CEA algorithm implemented in the Geoplatform is described in Eq.1. The algorithm takes into account an additive effects combination, meaning that cumulative effects correspond to the sum of individual effects on an environmental component (CEAA-ACEE, 2016), and considers a linear response of the

125

130

135 environmental component to the pressure. The CEA score on a single grid cell is calculated as follows:

$$CEA = \sum_{k=1}^n d(E_k) \sum_{j=1}^m s_{j,k} \text{eff}(P_j, E_k) \quad (1)$$

where *eff* is the effect of pressure *P* over the environmental component *E*,

$$\text{eff}(P_j, E_k) = \left( \sum_{i=1}^l w_{i,j,k} i(U_i, M_{i,j,k}) \right)'$$

140

whereas,

*U* = *i*-th human use

*P* = *j*-th pressures derived from the MSFD (EC, 2008)

*E* = *k*-th environmental components

145 *eff*(*P<sub>j</sub>*, *E<sub>k</sub>*) = effect exerted by the pressure *P<sub>j</sub>* over the *k*-th environmental component, *E<sub>k</sub>*.

*s*(*P<sub>j</sub>*, *E<sub>k</sub>*) = sensitivity of the environmental component *E<sub>k</sub>* to the *j*-th pressure *P<sub>j</sub>*

*w<sub>i,j,k</sub>* = use-specific relative pressure weight

*d*(*E<sub>k</sub>*) = Intensity or presence/absence of the *k*-th environmental component on the cell (*x*, *y*), which is 1 for fixed *E* (seabed habitats), and varies from 0 to 1 for mobile special features (turtles, marine mammals and seabirds).

150 *i*(*U<sub>i</sub>*, *M*(*U<sub>i</sub>*, *P<sub>j</sub>*, *E<sub>k</sub>*)) = distance model propagating *j*-th pressure caused by *i*-th activity over the *k*-th environmental component

*M*(*U<sub>i</sub>*, *P<sub>j</sub>*, *E<sub>k</sub>*) = 2D gaussian kernel function used for convolution considers buffer distances at 1 km, 5 km, 10 km, 20 km and 50 km

' (tick) = effect rescaling operator (from 0 to 1)

Further details on proposed CEA algorithm can be obtained from Menegon et al. (2018a).

155

### 2.2.2. Maritime Use Conflict (MUC) Analysis

Maritime Use Conflict (MUC) analysis is based on a methodology developed within the COEXIST Project – Interaction in European coastal waters: A roadmap to sustainable integration of aquaculture and fisheries (COEXIST, 2013). In particular the methodology presented by Gramolini et al. (2010) enables the identification of current/potential human uses and assesses their interaction in terms of conflicts. The algorithm implemented for the MUC score on a single grid is presented in Eq. 2:

160

$$MUC = \sum_{i=1}^l \sum_{j=i+1}^l c_{i,j} p(U_i) p(U_j) \quad (2)$$

165

where,

*c<sub>ij</sub>* = potential conflict score between *U<sub>i</sub>* and *U<sub>j</sub>*

*p*(*U<sub>i</sub>*) = presence (1) or absence (0) of the *i*-th human use in the cell

*p*(*U<sub>j</sub>*) = presence (1) or absence (0) of the *j*-th human use in the cell

170

The potential conflict score (*c<sub>ij</sub>*) between two uses *U<sub>i</sub>* and *U<sub>j</sub>* can vary from 0 (no conflict) to 6 (very high conflict score) and was calculated following the COEXIST methodology: i) application of an expert judgment approach to characterize each human uses through four attributes (vertical scale, spatial domain, temporal domain and mobility); ii) automatic assessment of the potential score for each use combination based on uses characterization and COEXIST rules application; iii) supervised expert based adjustment of the *c<sub>ij</sub>* coefficients to take into account legal and practical constraints between uses (see Appendix 1 for human uses characterization and COEXIST rules). Appendix 2 presents the potential conflict score matrix representing the *c<sub>ij</sub>* coefficients applied for each use

175

180 combination. For further details on the methodology applied in the study area we refer to Barbanti et al. (2015) and Depellegrin et al. (2017).

### 2.2.3. Threat analysis to Marine ecosystem services (MES-Threat)

185 The MES-Threat assessment builds on existing theoretical and practical approaches for the integrated analysis and mapping of stressor/pressure effects on MES supply units within the Great Lakes Restoration Initiative (Allan et al., 2013), North Sea (Hooper et al., 2017) or the ODEMM linkage framework (Options for Delivering Ecosystem-Based Marine Management; White et al., 2013). We define as MES-Threat the risk of MES reduction, partial or permanent loss of provision or impairment of use due to single or multiple anthropogenic effects targeting the MES providing ecological  
190 components (Worm et al., 2006; Maron et al., 2017). The tool incorporates an expert based MES capacity scoring and mapping procedure implemented for the Adriatic Sea's EUNIS habitats (see Appendix 3) with CEA modelling capabilities of the Geoplatform. The MES-Threat algorithm is presented in Eq. 3:

$$195 \quad MES - Threat = CEA \times MES = CEA \times \sum_{k=1}^n cap_k p(E_k) \quad (3)$$

where,

*CEA* = cumulative effects assessment model as described in Eq. 1

*cap<sub>k</sub>* = marine ecosystem services supply capacity (0-2, see Appendix 3)

200 *p(E<sub>k</sub>)* = Presence/absence of the *k*-th EUNIS habitat on the cell (*x, y*)

### 2.2. Step 1: Case study selection

After selecting the webtool to be applied, the system prompts the user to a pre-selected case study list. At the current stage three different geospatial domains are available, namely Mediterranean Seabasin  
205 level, macro-regional level for the Adriatic Sea and regional level for Emilia-Romagna Region (Fig. 3). Each case study represents a pre-configured set of webtool-specific data, with consistent spatial coverage of human uses and environmental components and incorporating all other necessary parameters for the model run. For the tools application in the case study a pre-configured grid resolution of 500 m x 500 m was applied.

**Step 1: Case Study selection**

In Step 1 you can select a case study available for three domains: Mediterranean Sea (2018), Adriatic Sea (2017) and Emilia-Romagna Region (2017). Each of the case study provides a pre-configured datasets, time reference and grid resolution to run the webtools.

Case Study Domain	Resolution	Uses	Envs
Mediterranean Sea 2018	1000.0 m	13	20
Adriatic Sea 2017	500.0 m	13	28
Emilia-Romagna Region RITMARE 2017 - CEA (Measu...)	500.0 m	13	9

210

Fig. 3. GUI illustrating the case study selection including case study domain, time reference and applicable grid resolution. Currently three study domains can be selected: Mediterranean Sea (2018), Adriatic Sea (2017) and Emilia-Romagna Region (2017).

## 215 2.3. Step 2: Case study setup

### 2.3.1 Study area definition

After selecting the case study, the Geoplatform prompts the user to the case study setup using an interactive web mapping application with a polygon selector tool (Fig. 4a). The Northern Adriatic (NA) Sea was selected as area of analysis based on the biogeographic boundaries defined by Bianchi (2004). The NA biogeographic region covers about 22,500 km<sup>2</sup> and is delimited by the Conero Regional Park to the southern tip of Istrian Peninsula (Bianchi, 2004). The NA is relatively shallow, with depth not exceeding 50 m (Turk and Odorico, 2009). From an administrative point of view, the NA embraces three countries and five coastal regions, including Italy (Emilia-Romagna, Veneto and Friuli-Venezia-Giulia Region), Slovenia (Coastal Karst Region) and Croatia (Istria Region). The NA is an extremely complex environment as it combines intensive anthropogenic activities (e.g. maritime transport, commercial fishery, aquaculture, coastal and maritime tourism), with sensitive coastal and marine ecosystems (e.g. essential fish habitats, nursery and spawning grounds of species of high commercial interest, seabirds and hotspots of Species of Community Interest such as *Caretta caretta* turtles and marine mammals, mainly *Tursiops truncatus*).

230






Fig. 4. Example of GUI illustrating the CEA setup for the Northern Adriatic Sea: a) Study area selection, dataset configuration for (b) human uses, (c) environmental components and (d) MSFD pressure selection. On the top-right (e) the button for initiating CEA model run is provided.

235

### 2.3.2. Dataset configuration

After the selection of the study area, the user can select the configuration of the geospatial dataset to be modelled (Fig. 4b-d). The Geoplatform incorporates a stocktake of over 65 MSP relevant geospatial layers for viewing, querying and download (Table 1). The user can select the human uses, the environmental components and the MSFD pressures to be included in the case study development. The dataset configuration is a key element for a case study development strategy as it allows to customize model outputs. The most updated version of the dataset can be freely download at Menegon et al. 2018b (<https://doi.org/10.5281/zenodo.1173764>).

245

Table 1. Geospatial layers and indicators implemented (P/A = presence/absence; I = normalized intensity indicator; PR = proxy; wP/A weighted presence/absence), adopted from Depellegrin et al., 2017. Note: a detailed version of this table and the most updated version of the dataset can be freely download at Menegon et al. 2018. <https://doi.org/10.5281/zenodo.1173764>.

Dataset	Indicator
---------	-----------

<i>Human uses</i>	
Aquaculture <sup>1,2</sup> , Cables and Pipelines <sup>2</sup> , Coastal Defence Work <sup>2</sup> , Dumping area for dredging <sup>2</sup> , LNGs <sup>3</sup> , Military areas <sup>2,5</sup> , Off-shore sand deposit <sup>1,4,5,7</sup> , Oil and Gas Extraction <sup>2,8,9,10</sup> , Oil and Gas Research <sup>2,8,9,10</sup> , Renewable Energy facilities (Offshore Wind farms) <sup>6</sup>	P/A
Coastal and Maritime Tourism <sup>11</sup>	I/PR – distance from the marinas and number of boats/marinas
Naval Based Activities <sup>15</sup>	I/PR – distance from the cargo ports and port capacity
Maritime Transport <sup>4</sup>	I – Traffic density (vessels/year)
Small Scale Fishery <sup>4</sup>	I – fishing effort expressed in 5 classes of intensity: from very low to high
Trawling <sup>12</sup>	I – hours of activities calculate through Vessel Monitoring System (VMS)
<i>Environmental components</i>	
Seabed habitats <sup>15</sup>	P/A
Marine Mammals <sup>13</sup> , Giant Devil Ray <sup>13</sup> , Turtles <sup>13</sup> , Seabirds <sup>13</sup> , Nursery Habitats <sup>14</sup>	wP/A

<sup>1</sup> Veneto Region ([www.regione.veneto.it](http://www.regione.veneto.it)); <sup>2</sup> SHAPE – Shaping a Holistic Approach to Protect the Adriatic Environment between coast and sea ([www.shape-ipaproject.eu](http://www.shape-ipaproject.eu)); <sup>3</sup> OGS – Istituto Nazionale di Oceanografia e di Geofisica Sperimentale ([www.ogs.trieste.it](http://www.ogs.trieste.it)); <sup>4</sup> CNR-ISMAR – Italian National Research Council – Institute of Marine Sciences ([www.cnr-ismar.it](http://www.cnr-ismar.it)); <sup>5</sup> MIPAAF – Italian Ministry of Agriculture, Food and Forests ([www.politicheagricole.it](http://www.politicheagricole.it)); <sup>6</sup> Emilia Romagna Region ([www.regione.emilia-romagna.it](http://www.regione.emilia-romagna.it)); <sup>7</sup> Arenaria S.r.l. ([www.arenariasabbie.com](http://www.arenariasabbie.com)); <sup>8</sup> MEDTRENDS-The Mediterranean Sea, Trends, Threats and Recommendations ([www.medtrends.org](http://www.medtrends.org)); <sup>9</sup> CHA – Croatian Hydrocarbons Agency ([www.azu.hr](http://www.azu.hr)); <sup>10</sup> MISE – Italian Ministry for Economic Development ([www.sviluppoeconomico.gov.it](http://www.sviluppoeconomico.gov.it)); <sup>11</sup> modeled on the basis of <https://www.pagineazzurre.com> data; <sup>12</sup> Blue Hub, JRC in-house platform to exploit big data in the maritime domain ([www.bluehub.jrc.ec.europa.eu](http://www.bluehub.jrc.ec.europa.eu)); <sup>13</sup> UNEP-MAP-RAC/SPA, Regional Activity Center for Specially Protected Areas; <sup>14</sup> MEDISEH MAREA Project ([www.mareaproject.net/medviewer](http://www.mareaproject.net/medviewer)); <sup>15</sup> EMODnet Seabed Habitats ([www.emodnet-seabedhabitats.eu](http://www.emodnet-seabedhabitats.eu)); <sup>15</sup> modeled on the basis of Eurostat data ([http://ec.europa.eu/eurostat/statistics-explained/index.php/Maritime\\_ports\\_freight\\_and\\_passenger\\_statistics](http://ec.europa.eu/eurostat/statistics-explained/index.php/Maritime_ports_freight_and_passenger_statistics)).

In total 13 layers of human uses were available in the Tools4MSP Geoplatform. The dataset is used by all three webtools (CEA, MUC and MES-Threat). Sources of the dataset are multiple and include: EU wide datasets (e.g. EMODnet Data Portals, European Atlas of the Seas, EEA map services), project portals specific datasets (e.g. the SHAPE Adriatic Atlas, COCONET WebGIS), data made available by research institutions (e.g. HCMR - Hellenic Centre for Marine Research; CNR-ISMAR – Italian National Research Council – Institute of Marine Sciences) and from national (e.g. OTE S.A. – Hellenic Telecommunication Organization; MIPAAF – Italian Ministry of Agriculture, Food and Forests) and regional authorities (Veneto and Emilia-Romagna).

The dataset of environmental components is used for the CEA and MES-Threat analysis and is based on 20 layers. Marine habitat layers include 15 distinct habitats, which were derived from EUNIS classification based EUSeaMap dataset (Populus, 2017). Layers for marine mammals, Loggerhead turtles and Giant Devil Ray densities were obtained from UNEP-MAP-RAC/SPA (Fortuna et al., 2015) and are based on a weighted presence/absence (wP/A) in terms of individuals per 20 km x 20 km. The nursery areas of 33 valuable commercial fishery species, including European pilchard (*Sardina pilchardus*), common sole (*Solea solea*), Norway lobster (*Nephrops norvegicus*), red mullet (*Mullus barbatus*) were obtained from the MEDISEH MAREA (Mediterranean sensitive habitats; [www.mareaproject.net/medviewer](http://www.mareaproject.net/medviewer)) Project. These layers are available as dummy indicator of presence/absence (P/A).

280 The CEA model implements 15 MSFD pressures out of 18 provided by the MSFD (EC, 2008). In the description of Table 2, pressures are grouped into three pressure themes, according to MSFD amended version (EC, 2017, Annex 4, Table 2): biological (2 pressures), physical (5 pressures) and a mixed substances-litter-energy (8 pressures) theme. The three pressures related to significant changes in salinity regime, introduction of radio-nuclides and introduction of microbial pathogens were omitted from the pressure dataset due to lack of reference and expert judgement.

285 The marine ecosystem services component necessary for the MES-Threat analysis is based on a MES capacity matrix (Appendix 3) rescaled for the NA Sea according to an initial assessment by Depellegrin et al. (2017). The MES capacity matrix adopts a qualitative indicator of the potential ES supply of the habitat ranging from 0 (neglectable capacity) to 2 (high capacity) adopted from a methodology proposed by Galparsoro et al. (2014) and Salomidi et al. (2012). In the NA the matrix implements 12 MES (x-axes) and 15 EUNIS habitats (y-axes) grouped into four MES categories (provisioning, regulating, cultural and supporting).

290

### 2.3.3. Case study strategy development

295 In order to provide meaningful assessment for environmental management and planning in the NA Sea, the webtools were applied by operating seven distinct dataset configurations resulting into three webtool base runs (CEA/MUC/MES-Threat), three CEA/MUC sector-specific runs (maritime traffic, commercial fishery and coastal and maritime tourism), three MSFD pressure themes driven CEA runs (biological, physical and substance-litter-energy pressures) and four MES-Threat model runs, one for each MES category (provisioning, regulating, cultural and supporting). In Table 2 a summary of the dataset requirements and setup for each model run is presented.

300 Table 2. Overview of webtools, dataset selection & requirements for model runs and outputs based on case study configuration strategy. Note: U - human uses; E - environmental components; P - MSFD pressures; MES - marine ecosystem services.


Webtool	Dataset selection	Outputs	Description
CEA	All U, P and E	CEA base run	Overall CEA model run includes all 13 human uses, 15 MSFD pressures and 20 environmental components
	Sectorial U, all P and E	Maritime traffic Commercial fishery (small scale fishery, bottom and pelagic trawling) Coastal and maritime tourism	Sectorial model run includes 3 human uses, 15 MSFD-Pressures and 20 environmental components
	MSFD themes, all U and E	CEA - biological	CEA model run for two pressure composing the biological theme: (1) Introduction of non-indigenous species and translocation; (2) Selective extraction of species, including incidental non-target catches
		CEA - physical	CEA model run for five pressures composing the physical theme: (1) Smothering; (2) sealing; (3) changes in siltation; (4) selective extraction; (5) abrasion
		CEA - substances-litter-energies	CEA model run for eight pressures composing the substances-litter-energies theme: (1) underwater noise; (2) marine litter; (3) inputs of fertilisers and other nitrogen and phosphorus-rich substances; (4) introduction of non-synthetic substances and compounds; (5) introduction of other substances; (6) introduction of synthetic compounds; (7) significant changes in thermal regime; (8) organic matter

MUC	All U	MUC base run	Overall MUC model run output includes a conflict analysis on 13 human uses.
	Sectorial U	MUC - Maritime traffic MUC - Commercial fishery (small scale fishery, bottom and pelagic trawling), MUC - Coastal and maritime tourism	Sectorial MUC model run for three distinct human uses.
MES-Threat	All U, P and MES categories (EUNIS habitats)	MES-Threat Base run	Overall MES-Threat model run for 12 MESs provided by 15 EUNIS habitats, 13 human uses and 15 MSFD-Pressures
	Specific MES categories, all U and P	MES-Threat provisioning MES-Threat regulating MES-Threat cultural MES-Threat supporting	MES category driven model output includes four MES-Threat maps, one for each MES categories identified in the MES capacity matrix (Appendix 3)

### *2.5. Step 3: Geospatial and statistical outputs*

305 The Tools4MSP Geoplatform provides to the user a full range of geospatial and statistical results that can be used for further deepening of the analysis. In Fig. 5 a GUI example of the multiple outputs are provided including exploration of geospatial and statistical results (Fig.5 a and d), the view layer functionality to share and download modelling results (geotiff format) with the user community (Fig.5 b) and the complete metadata functionality to compile metadata information on the modelling results (Fig.5 c). In particular the downloaded results can be used for further investigation or re-analysis

310 using dedicated GIS software. In the following Section geospatial visualizations were presented using Quantum-GIS (QGIS Development Team, 2018) and statistical results were presented with Python numeric and scientific libraries including Numpy, Scipy, Pandas, Matplotlib and Seaborn (van der Walt, 2011; McKinney, 2010).



315

Fig. 5. Example of GUI presenting geospatial and statistical results: (a) the geospatial results of CEA run for the NA, (b) view layer button enabling users to download geospatial results, (c) complete metadata button for describing metadata relevant for the understanding of the result and (d) statistical result in terms of CEA score distributions by number of cells per CEA score.

320

### 3. Results

#### 3.1. Cumulative Effects Assessment

In Fig. 6 a-d results from CEA case study development are presented. The CEA base run show that highly impacted sea areas are located mainly in Italian coastal areas such as the Gulf of Trieste and along a coastal segment in front of the Po river outlet (Fig. 6a). On overall the NA Sea reaches a mean CEA score of 3.06. The maximum CEA score of 8.3 is located in proximity of the port of Trieste, in the North-Eastern NA.

The sectorial CEA application for maritime transport is presented in Fig. 6b. CEA scores reach a maximum score of 2.73. Areas of highest CEA score are located mainly offshore, and correspond to high density shipping lanes (up to 400-500 vessels per year) connecting the main Adriatic ports (e.g. Venice, Trieste and Koper) to the Mediterranean Sea potentially affecting valuable hotspots of marine

330

mammals (mainly *T. truncatus*) and loggerhead turtles (*C. caretta*). The sectorial CEA application for coastal and maritime tourism is presented in Fig. 6c. Areas of highest CEA score are located in the Gulf of Trieste (score 1.5) and Venice Lagoon and the Malamocco outlet. The Gulf of Trieste is particularly densely populated with 26 marinas on the Italian coastal areas and 3 on the Slovenian coastal areas. To notice is that on overall the the Italian coastal regions of Friuli-Venezia-Giulia and Veneto have higher CEA scores distributed along the entire coast, compared to Slovenian and Croatian coastal segments. The sectorial CEA score from the commercial fishery is represented in Fig. 6d. The maximum CEA score is 4.9 in proximity of Riccione (Emilia-Romagna Region). Other sea areas of high CEA score (3.8) are located in front of the Po Delta inlet. Both areas are subjected to intense fishery activities along Italian coasts, especially trawling (e.g. bottom otter trawl, pair pelagic trawl), which greatly affects both biological resources and seafloor integrity. Areas of lower intensity CEA scores can be attributed to the 3 nm boundary, where trawling activities are forbidden (EC Regulation 1967/2006).




Fig. 6. CEA model outputs: (a) CEA base run; CEA sectorial runs for (b) maritime transport; (c) coastal and maritime tourism and (d) commercial fishery.

350 Results for the MSFD pressure-specific CEA case study runs are presented in Fig. 7. Geospatial  
 results for the biological theme (Fig. 7a) show that highest CEA score (1.2 - 1.5) have a patchy  
 distribution, mostly corresponding with commercial fishing activities in offshore areas, in front of the  
 Venice Lagoon, Po Delta outlet, Rimini Port (Emilia-Romagna Region), and in offshore areas in front  
 of the coastal settlements of Rovinj (Istria Region) (CEA score = 1.4). The environmental components  
 355 with highest sensitivity to biological pressures refer to commercial fishes nursery habitats, marine  
 mammals and turtles.

The geospatial distribution of the physical pressure theme (Fig. 7b) shows a more homogenous  
 distribution. In particular the offshore areas (about 3-6 nautical miles) in front of the Italian coastal  
 Regions of Veneto, Emilia-Romagna and Marche Region are areas of high CEA scores (2.5 - 2.8) in  
 360 front of the Po Delta outlet and between Port of Rimini and Pesaro, which clearly relate these  
 pressures to intense trawling activities. The environmental components most affected by the physical  
 pressures refer to infralittoral and circalittoral sand a mud habitats, *Cymodocea* beds and infralittoral  
 rock and other hard substrata. The geospatial results for the substances-litter-energy pressure theme  
 (Fig. 7c) has the highest relative score (3.97) among all three pressure themes. High CEA score are  
 365 concentrated in small area in front of the Po Delta outlet. Other high CEA score areas are located in  
 offshore areas in proximity of hotspots of Species of Community Interest (*C. caretta* and *T.*  
*truncatus*).




Fig. 8. CEA model outputs from MSFD pressures themes: (a) biological pressures, (b) physical pressures and (c) substances-litter-energies.

370

### 3.2. Maritime Use Conflict analysis

Results from MUC base run and sectoral conflict analysis are presented in Fig. 8a. Sea areas with  
 MUC score > 20 (298 km<sup>2</sup>) are located in proximity of the Ports of Trieste and Koper, in front of  
 Venice lagoon and Chioggia. Other conflict areas are located in coastal areas of Emilia-Romagna  
 375 region (Porto Garibaldi and Ravenna) and in Marche region (Port of Ancona). Sea areas with MUC  
 score ranging from 20 to 10 (1300 km<sup>2</sup>) are located further outside Port of Trieste, Veneto Region  
 (Jesolo and Caorle) in front of Venice Lagoon, Po Delta outlet, and other hotspots localized along the  
 Emilia-Romagna and Marche Regions. Similarly to the CEA base run, the eastern coast of the NA Sea  
 has lower conflict areas with exception of Koper Port in the Slovenian Coastal Karst Region and




380 coastal segment between the Croatian coastal settlements of Novigrad and Pula (MUC score between 5 and 8).

The analysis of sector-specific MUC runs are presented in Fig. 8b-d: Results for maritime transport (Fig. 8b) show that areas of highest MUC ( $>8$ ; 580 km<sup>2</sup>) are located in front of main port areas in the NA Sea (Trieste, Koper) and northern Po River Delta, Port of Ravenna and Ancona. Highest MUC score ( $> 8$ ; 612 km<sup>2</sup>) for coastal and maritime tourism (Fig. 8c) show a similar pattern to the maritime transport. The MUC run for commercial fishery (Fig. 8d) evidences clear patterns of conflict between the different types of fishing and with maritime transport. Areas of highest MUC scores ( $>15$ ; 179 km<sup>2</sup>) are located in front of Chioggia and Venice lagoon , followed by port of Trieste and a narrow offshore area between 3 and 4 nm in front of Veneto, Emilia-Romagna and Marche region (MUC score 10-15; 983 km<sup>2</sup>) and a widespread offshore area along the main maritime traffic corridors (MUC score 5-10; 6254 km<sup>2</sup>).

385

390




395 Fig. 8. MUC analysis model outputs: (a) MUC base run; sectorial MUC analysis runs for (b) maritime transport; (c) coastal and maritime tourism and (d) commercial fishery.

### 3.3. Comparison of CEA/MUC outputs

400 In Fig. 9a the CEA (by MSFD pressure themes) and MUC contribution as function of distance from coast in nautical miles (nm) is illustrated. In terms of pressure-specific CEA, the cumulative effects coming from input of substances, litter and other forms of energies contributes to 41.1% to the total CEA score, followed by the cumulative effects from physical pressures with 40.9% and the cumulative effects from biological pressures with 18%. Within the 0-3 nm the the CEA Substances-litter-energies contributes to 60.9% to the total CEA, followed by physical pressures with 29.3% and biological pressures with 9.8%. Beyond the 12 nm, the CEA from physical pressures contributes to 41.7%, followed by CEA from Substances-litter-energies (38.3%) and from biological (20%) pressures.

410 The MUC analysis evidences that 50% of the conflict relies within the 9 nm. Peak of conflict is located within the 3-9 nm with contribution of 21% of the total MUC score. On the contrary, about the 70% of the CEA score is almost uniformly distributed between the 3 and 24 nm. Highest CEA score is located between 15-18 nm.



415 Fig. 9. (a) Percent contribution of total CEA (divided into MSFD pressure themes) and MUC scores from base run, according to distance intervals of 3 nautical miles (nm) from coastline. (b) Percent contribution of sectorial CEA/MUC score (commercial fishery, maritime transport, coastal and maritime tourism and other uses) to total CEA/MUC score according to distance intervals of 3 nautical miles (nm) from coastline.

420 Sector-specific CEA/MUC as a function of distance (nm) are presented in Fig. 9b. The overall contribution of maritime traffic, commercial fishery and coastal maritime tourism represents the 95% of the total CEA score and the 74% of the MUC score. Commercial fishery has the highest

425 contribution to CEA (55.4%) and MUC (44%) overall score, followed by maritime transport (CEA = 31.3% and MUC = 20.2%) and coastal and maritime tourism (CEA = 5% and MUC = 10.2%). To notice is that 94 % of CEA score derived from coastal and maritime tourism is concentrated within 0-9 nm, similarly 99% of MUC score contribution comes from this segment.

### 3.4. MES threats analysis

430 In Fig. 10a results from MES-Threat base run are presented. The areas of highest threat (MES-Threat score >25; 165 km<sup>2</sup>) are located in front of Grado-Marano Lagoon coastal area referring and smaller patchy areas in proximity of the Venice Lagoon. Threatened habitats refer to *Cymodocea* beds (A5.531), habitats providing a multitude of MES (e.g. providing nursery, biodiversity, food provisioning, nutrient cycling, water quality; Appendix 3) associated with areas of high cumulative pressures. CEA applied on provisioning MES (Fig. 10b) shows potential threats (MES-Threat score > 10) located in the Gulf Trieste for coastal habitats responsible for food provisioning capacity, such as circalittoral sandy (A5.35), circalittoral fine mud (A5.36) and circalittoral muddy sand (A5.26). CEA applied on regulating MES (Fig. 10c) has a more patchy distribution of high threat areas (MES-Threat score > 3) localized in the Gulf of Trieste, coastal areas of Grado-Marano and the segment from Venice Lagoon to Po Delta. CEA applied on cultural MES (Fig. 10d) shows high threats in nearshore areas along Friuli-Venezia-Giulia Region (MES-Threat score >10), Veneto Region (MES-Threat score > 4) and southern Emilia-Romagna and Marche Region (MES-Threat score > 2). To notice is that in the Eastern NA more extended threat areas occur, but with lower threat score (MES-Threat score > 2). Threats are particular relevant for infralittoral habitats and *Cymodocea* beds.

440

445 Threat analysis for supporting MES (Fig. 10e) responsible for sustaining biodiversity and nursery provision show a threat distribution compared to the MES-Threat base run, with highest threat scores (>12) in proximity of Grado-Marano, Venice lagoon and Conero Promontory.





Fig. 10. MES-Threat analysis outputs for: a) Base run, b) provisioning, c) regulating, d) cultural and e) supporting MES.

450

In Fig. 11 the MES-Threat score contribution in percentage as function of distance (in nm) from coastline were presented. On overall results show that highest threat scores from MES-Threat are located within 3-6 nm (18.7 % of total contribution), where about 50% of the contribution is due threats to provisioning MES, 14% to regulating MES, 4% cultural MES and 33 % to supporting MES.

455

To notice is that threat areas for MES cultural capacity are entirely located within the 0-9 nm from coastline with a 65% contribution within 0-3 nm, 28 % within 3-6 nm and 7% within the 6-9 nm.



460 Fig. 11. Percent contribution of MES-Threat base run (divided into four MES categories) to total MES-Threat score according to distance intervals of 3 nautical miles (nm) from coastline.

#### 4. Discussion

##### 4.1. Overall results

465 The presented webtools embedded into the Tools4MSP Geoplatform in combination with a clearly defined case study modelling strategy exemplified how meaningful geospatial and statistical results can be obtained in support of planning and environmental management considerations. The geospatial webtools in support of planning and environmental management provide set of advantages as largely discussed in literature (Atkinson and Canter, 2011; González Del Campo, 2017; Palomino et al., 2017) such as transparency, objectivity and replicability of model outputs, all considered fundamental within a pragmatic planning process. Particularly relevant are the dynamic functionalities coupled to the Spatial Data Infrastructure (SDI) enabling data pre-processing (normalization and aggregation, rescaling, filtering), access to existing and novel datasets as they become available and republish of spatial outputs for utilization within user communities and possibilities for model re-run.

475 The case study developed for the Northern Adriatic demonstrated the potentialities of the Tools4MSP Geoplatform to be further developed towards an operational Decision Support System for a multitude of marine and coastal environmental management and MSP-oriented planning tasks. In addition to the presented modelling capabilities, the Geoplatform aggregates a multitude of geospatial dataset and formats into already normalized datasets and therefore enhances its usability by reducing time and manpower for extensive data preparation. The tool can be flexibly applied to different spatial scales (from seabasin to regional level) by defining the study area and depending on the availability and quality of dataset, also grid resolution can be customized. The case study for the NA Sea was performed on a resolution 500 m x 500 m grid. A higher resolution of analysis can be considered in combination with local datasets. In the NA, this is particularly required for inland waters such as Venice Lagoon and Grado-Marano Lagoon and the Gulf of Trieste, where anthropogenic activities are particularly intense (Gallmetzer et al., 2017; Malačič et al., 2008; Munaretto and Huitema, 2012).

The geospatial results outline the complexity of anthropogenic impacts and interactions in the study area. CEA and MUC outputs highlight the need for thorough planning measures in order to deal with the intense conflicts occurring in the area, especially in the western segment of study area along Italian nearshore areas.

490 The CEA analysis presented in Fig. 6a-d provides an overview of the spatial distribution of the cumulative and the sectorial effect scores in the study area. The CEA score exerted by commercial fishery has the most evident effects across the study area (Fig. 6d and 9b). The Northern Adriatic Sea is one of the most intensively fished area in Europe, where most of the harvested stocks are overexploited, especially in the western sector, due to intense non-selective fishing activities from  
495 Italian fleets exerted on fish habitats (Colloca et al., 2013; Russo et al., 2015, Bastardie et al., 2017). In nearshore areas (about 3 nm) lower CEA scores refer to areas where towed gears are banned or unsuitable, usually in favour of small scale fisheries (mainly set gears and longlines). Maritime traffic effects are located mainly offshore (Fig 6b), in the central-eastern portion of the study area, where the north-south Adriatic traffic route connects to the Mediterranean. Those areas intersect with  
500 biodiversity hotspots of valuable marine species, such as loggerhead turtles and marine mammals (Fortuna et al., 2015). The cumulative effects from coastal and maritime tourism sector generates main impact phenomena in proximity of coastal areas (Fig. 6c and 9b), where the necessary infrastructure and facility occur (Papageorgiou, 2016) and where the majority of vulnerable ecosystem are present. In the NA, summer recreational resorts from Friuli-Venezia-Giulia, Veneto Region and  
505 Emilia-Romagna, belong to the top 20 most popular tourist destinations on EU level (Eurostat, 2017). The MSFD-driven CEA representation (Fig. 7) shows that the anthropogenic pressures within 0 to 3 nm derive from the MSFD pressure theme substances-litter-energies exerted mainly by land-based activities (Fig. 7c, 9a). The higher scores are linked to riverine discharge from Po and Adige rivers, that are the biggest contributors of freshwater, nutrients and pollutants of the Adriatic Sea (Chiogna et al., 2016; Simonini et al., 2004), commercial traffic in proximity of ports (Venice, Chioggia, Ravenna), coastal tourism and leisure boating. In comparison, the MSFD pressure themes concerning biological and physical pressures close to the coast have lower effects intensity (Fig. 7a,b and 9a) and mainly derive from artisanal fisheries and maritime transport. High CEA scores are evidenced for valuable habitats such as *Cymodocea* beds and nursery areas. A major reason for lower CEA score by  
515 physical pressures are related to limitations of towed gears ban within the 3nm off the coast, with the significant exception of those close to port and marinas or subjected to intense artisanal fisheries with bottom impacting tools. Beyond 3 nm limitations, physical pressures became the most intensive (Fig. 7b and 9a), due to the strong contribution of bottom trawling fisheries on the CEA score, determining high physical (e.g. abrasion) pressures on essential fish and seabed habitats (Pranovi et al., 2000).  
520 Moreover, trawling fisheries are also highly responsible for biological pressures (e.g. extraction of species; Eigaard et al., 2016) and, together with maritime traffic, releases of marine litter and substances, potentially affecting seabed habitats and the populations of marine turtles and mammals. The application of the MUC and its setup for sectoral analysis of conflicts allowed to identify main areas of conflict to compare results among the most relevant sea uses in the Northern Adriatic Sea  
525 (Fig. 8). According to Fig. 9b over 60% of the total MUC score is concentrated within the 12 nm boundary, mainly caused by intense interactions between coastal tourism, maritime transport, fisheries and other activities (e.g. aquaculture), especially close to ports (e.g. Trieste, Koper, Venice and Ravenna) and marinas, while in offshore areas spatial conflicts occur between traffic routes and trawling fishery grounds (Fig. 8). The distribution of conflicts evidences the high demand for sea space in proximity of coastal areas in a relatively small sea space. Soft uses (e.g. fishery and coastal  
530 tourism) and hard uses (e.g. aquaculture, Oil & Gas exploitation or maritime transport) need to trade-

off especially in coastal areas that function as hub aggregating infrastructure and facilities necessary to support maritime economic activities.

535 The MES-Threat analysis demonstrated how CEA based modelling capabilities can be linked with  
MES mapping based on the MES supply capacity of EUNIS habitats. Spatial results provided novel  
insights on the distribution of threats to MES and therefore to the risk of reduction, loss or impairment  
of the MES provisioning capacity of a particular habitat or combination of habitats. From a planning  
perspective, the produced results can be considered as highly integrative to the CEA and MUC  
540 outputs, as they incorporate societal values into the analysis (Maron et al., 2017), can more efficiently  
delineate protection priorities (Werner et al., 2014) and support the design of restoration plans (Allan  
et al., 2013) for coastal areas or habitats (e.g. *Cymodocea* beds) subjected to highest threat from  
anthropogenic stressors (Fig. 10a). Similar to CEA/MUC results, the analysis showed that threats to  
MES are highest within the 0-6 nm, a critical area for MES provision (Fig. 11), but also for intensity,  
545 variability of pressures and conflict areas (Fig. 9). Particularly affected categories are provisioning  
and supporting MES, which are responsible for the provision of fundamental goods and services  
sustaining various components of coastal economies such as commercial fishery, aquaculture and  
tourism.

The use of a case study development strategy demonstrated a high degree of customization of the  
webtools by the user and a flexible adaptation to different MSP stages: First, the webtool can facilitate  
550 data gathering through a community-based approach and interoperable access to EU-level and  
international SDIs, such as EmodNet, EEA, SeaDATANet or International Hydrographic  
Organization (IHO). Current datasets can be flexibly visualized and recombined through interoperable  
view services (i.e WMS, TMS, ArcGIS REST service) in order to create shareable online maps.  
Second, the webtools can be used for the identification of specific planning constraints and current  
555 conditions of the sea space in terms of multiple and single pressure on environmental components and  
existing conflict among uses. Third, the model outputs can be used to evaluate different management  
actions or define alternative scenarios through the comparison of two case studies and understand  
variations of cumulative effects, consequences for sea use conflict and threats to ecosystem service  
supply capacity.

560

#### *4.2. Webtools support to EIA and SEA*

MSP, as an area-based management framework can represent plans, programs or policies that require  
environmental impact assessment (EIA) and strategic environmental assessment (SEA). Common aim  
for MSP, EIA and SEA is to promote sustainable development through the integration of  
565 environmental considerations into the planning process and reduce spatial negatives (IUCN, 2013).

In the context of the EIA (2011/92/EC) and SEA (2001/41/EC) the presented webtools can deliver a  
promising support to address several requirements of both Directives: (1) The presented Geoplatform  
facilitates the access and usage of geospatial datasets that can be relevant to EIA and SEA  
(Vanderhaegen and Muro, 2005). (2) Although on different spatial and implementation scales the  
570 CEA/MUC can be flexibly deployed on sectoral level (Fig. 6 and 8b-d) and MSFD pressure specific  
(Fig. 7) local project as required within EIA [Article 5(1), ANNEX IV], while as requested within  
SEA [Article 3(5) ANNEX II], the tool can flexibly address the cumulative nature of effects on a  
broader scale from regional to national and also transnational level. (3) The implemented CEA  
isotropic distance model (Eq.1) is capable to modulate propagation of pressures and can further  
575 complement the analysis of spatial influence of the proposed project on local, regional, national or  
transboundary level. Although methods for ES-inclusive SEA are still lacking (Slootweg and van  
Beukering 2008; Söderman et al., 2012), the presented marine ES-inclusive threat analysis approach



580 based on EUNIS habitats MES supply scores can be a valid complement to SEA. In particular the ES-driven concept can be flexibly incorporated along a SEA activities and evidence sea areas of socio-ecological importance, possible impacts of present and future plans on key ES and identify solutions and restoration measures to reduce anthropogenic effects (Geneletti, 2011).

#### *4.3. Webtools support to MSP*

585 The relevance of geospatial tools to support MSP implementation has been evidenced by many academic, planning and decision-making communities around Europe (MSP Platform, 2018; JPI Ocean, 2017; WESTMED, 2017). In particular the tools for cumulative effects and associated processes (e.g. strategic environmental assessments) have found vivid development in the last decade, with application in various planning and spatial contexts (Andersen et al., 2013; Depellegrin et al., 2017; Stelzenmüller et al., 2013). At this stage the presented tool need to be seen as a test toolset, with  
590 particular limitations related to data availability and model robustness. Depending on the modelling approach for uses, pressures and the environmental components, its over-simplifications do not allow to take decisions with high socio-economic relevance. Moreover, the connections with provisions from other policy instruments (e.g. MSFD, WFD, CFP, H&BD) are relevant and are only partially explored, while maritime plans have to carefully consider the coordination and compliance with all  
595 relevant policies. Within a typical MSP methodology (Ehler and Douvère, 2009; Barbanti et al., 2015), these tools can be used both in the analysis phase, defining and analysing existing conditions, and in the planning phase, supporting the development of measures and scenarios and the evaluation of their effectiveness. Finally, the presented webtools are part of a wider ecosystem of analytical techniques supporting MSP. In fact CEA, MUC and MES-Threat can be combined with each other  
600 ensuring a multi-functional approach or can play a complementary role in support of tool functionalities, such as Displace for use specific investigations (Bastardie et al., 2017), Marxan with Zones for scenario development (Ban et al., 2013) or Seasketch (2018) for incorporating participatory stakeholder engagement into spatial modelling.

#### *4.3. Datasets*

605 Although the presented webtools benefit from a multitude of geospatial datasets (in total 65) composed by human uses, environmental components and pressures further extension can be considered. In particular, integration of novel datasets can be used for scenario analysis of emerging sectors of the marine economy in the Adriatic Sea, such as potential offshore wind energy farms off  
610 the coastal settlement of Rimini (Emilia-Romagna Region; Schweizer et al., 2016) or in front of the coastal settlement of Pula (Istria Region; Hadžić et al., 2014), extensions of the ports of Ravenna (RER, 2015), Trieste (TMT, 2016) and Koper (Port of Koper, 2015) or increasing aquaculture development to meet fish food demand (Piante and Ody, 2015) should be incorporated and analysed for its environmental impacts and the creation of potential sea use conflicts. Moreover, our case study  
615 shows high variability in CEA/MUC scores between western and eastern coastal areas, this is related to higher number and intensities of human activities along the Italian coasts compared to Slovenian and Croatian ones, but also due to a high heterogeneity in human activities (especially Oil and Gas extraction, aquaculture and shipping) and the number of datasets available from different countries. Concerning the environmental components, higher resolved geospatial datasets on habitats, benthic  
620 communities and species (Certain et al., 2015; Marcotte et al., 2015) should be integrated considering their potential sensitivity towards specific anthropogenic pressures (Eno et al., 2013) and with proper classification schemes.

625 While the current dataset incorporates a multitude of endogenic pressures, generated within the  
system and that can be managed (Elliott, 2011), there is the need to incorporate as well exogenic  
pressures such as climate change in order to align the methodology to other CEA assessments around  
the globe (Halpern et al., 2015; Clarke Murray et al., 2015). This would support the analysis of  
climate change scenarios and its influence on coastal areas, marine ecosystems and interactions with  
human activities (Pinarbasi et al., 2017). In fact, ecological and geomorphological conditions of the  
Northern Adriatic Sea make it particularly sensitive to changing hydrological and oceanographic  
630 conditions (Bosnjakovic and Haber, 2015), inducing harmful algal blooms (Barale et al., 2008), red  
tides (Socal et al., 2011) or different hydrodynamic impacts (e.g. inundations, storm surges or coastal  
erosion).

#### *4.4. Limitations*

635 The tested tools are not free of limitations. Difficulties in the parametrization of the model induced the  
implementation of an additive and linear model, while ongoing research in cumulative assessment  
demonstrate the need of integration of mitigative and antagonistic effects in the pressure -  
environmental components interaction. There is spatio-temporal inhomogeneity among dynamic  
environmental components (seabirds, mammals and turtle datasets) with coarser resolution datasets  
640 compared to EUNIS marine habitats (100 m x 100 m) and differing nominal scales (e.g.  
individuals/km<sup>2</sup> versus presence-absence indicators).

Although an extensive sensitivity and uncertainty analysis has been performed for the Adriatic-Ionian  
Sea (Gissi et al., 2017), an implementation of uncertainty in the webtools is currently absent.  
Uncertainty analysis is an important component within CEA, as it supports realistic knowledge  
645 aggregation optimised methodological procedures which are baseline for MSP (Judd et al., 2015). In  
particular an uncertainty analysis as part of model-based decision support, should be an integral part  
of the webtool results to better identify data gaps (Meyer, 2012), support effective risk assessment  
(Stelzenmüller et al., 2015), take into account the precautionary principle into planning considerations  
and communicate the uncertainty within a participative dialogue (Bijlsma et al., 2011).

650 A limitation of the presented MUC is the absence of representation of synergies in the sea space, as  
the MUC model only considers spatial conflicts focused only on use-use conflicts, without  
considering multiple interaction (within three or more sea uses) or dynamic interaction from highly  
mobile sea uses (e.g. maritime traffic, commercial fishery, coastal and maritime tourism).

We consider the MES-Threat analysis a first methodological approach for integration of the socio-  
655 ecological dimension into cumulative effects assessment. Major challenges remain identifying the  
suitable spatial extent and resolution for quantifying MES and models for taking into account their  
space-time variability. Further research is needed to aggregate other environmental components into  
MES datasets (e.g. marine mammals and turtles), develop pressure specific MES sensitivity charts  
(Hooper et al., 2017), deepen the link of biodiversity attributes with ecosystem services supply  
660 (Harrison et al., 2014) and provide theoretical and methodological integrations of MES into  
environmental and socio-economic impact assessment also for specific pollution risks (Depellegrin  
and Blažauskas, 2013; Song et al., 2017) and emerging new uses, such as renewable energy  
(Papathanasopoulou et al., 2015).

#### **665 5. Conclusions**

The Tools4MSP Geoplatform provides a novel system in support of planning and environmental  
management, incorporating within a single geoplatform three operational webtools: the cumulative  
effects assessment, maritime use conflict analysis and marine ecosystem services threat analysis. This

670 is an added value for decision-makers and planners that seek for a rapid exploratory mapping of  
 human-environment interactions. Modelling outputs were guided by a structured case study  
 development strategy allowing overall analysis and context specific investigations, such as by marine  
 sector, by MSFD pressure themes or by marine ecosystem services categories. The community-based  
 Geoplatform demonstrated to be highly versatile instrument for the spatialization and geostatistical  
 675 evaluation of MSP relevant knowledge applicable in several stages of an MSP process and potentially  
 also supporting EIA and SEA. In particular the Geoplatform can ensure notable support for  
 transparent analysis that can engage a multitude of user communities into decision-making, ensure  
 replicability of the modelling process and iterative data assimilation, as it becomes available. The tool  
 can be used by a broad spectrum of stakeholders, including decision-makers, planners, academics,  
 680 research institutions and the general public.

### **Acknowledgements**

Funding: This work was supported by: the Flagship Project RITMARE - Italian Research for the Sea, coordinated by the Italian National Research Council and funded by the Italian Ministry of Education, University and Research within the National Research Program 2012–2016.

**Declarations of interest:** none

### **Author contributions**

Conceived and designed the experiments: SM, DD.

Data curation: SM, DD, CV.

Performed the experiments: SM.

Analyzed the data: SM, DD, GF, AS.

Contributed reagents/materials/analysis tools: SM, DD, GF.

Founding acquisition: AB.

Wrote the paper: SM, DD, GF, AB.

### **Appendix 1. Human use classification and rules for spatial conflicts according to COEXIST applied in the maritime use conflict (MUC) webtool methodology (Gramolini, 2010).**

*Human uses* can be classified according to five traits: vertical, spatial (horizontal), temporal scale, mobility, and location.

*Rules for spatial conflicts:* rule system to define conflict score for each pair of human uses.


- Rule 1: if vertical domain of activity 1 is different from vertical domain of activity 2 and no one of them interests the whole water column then conflict score is equal to 0;
- Rule 2: If both activities are “mobile” then conflict score is equal to the minimum of temporal domain plus the minimum of spatial domain.
- Rule 3: if Rule1 and Rule2 cannot be applied then the conflict score is equal to the maximum value of temporal domain plus the maximum value of spatial domain.

1	<b>Vertical scale</b> <ul style="list-style-type: none"> <li>● Pelagic</li> <li>● Benthic</li> <li>● Whole water column</li> </ul>	Value = 1 Value = 2 Value = 3
---	--	-------------------------------------

2	<b>Spatial scale</b> <ul style="list-style-type: none"><li>• Small</li><li>• Medium</li><li>• Large</li></ul>	Value = 1 Value = 2 Value = 3
3	<b>Temporal scale</b> <ul style="list-style-type: none"><li>• Short</li><li>• Medium</li><li>• Long/permanent</li></ul>	Value = 1 Value = 2 Value = 3
4	<b>Mobility</b> <ul style="list-style-type: none"><li>• Mobile</li><li>• Fixed</li></ul>	Value = 1 Value = 2
5	<b>Location</b> <ul style="list-style-type: none"><li>• Land</li><li>• Sea</li></ul>	Value = 1 Value = 2

**Appendix 2. MUC conflict matrix for the Northern Adriatic Sea**

It's based on MUC base run, ranging from 5 (very high conflict) to 0 (non conflict).



### Appendix 3. Marine ecosystem services capacity matrix

The matrix is based on EUNIS Habitats extracted for the Northern Adriatic Sea (adopted from Depellegrin et al., 2017). The score ranges from 2 (high capacity) to 0 (no or neglectable capacity). In total 15 EUNIS habitats were extracted for the Northern Adriatic Sea.

Habitat code	Habitat name	km <sup>2</sup>	perc	MES <sub>prov</sub>		MES <sub>reg</sub>			MES <sub>cul</sub>			MES <sub>sup</sub>				Σ
				Food provisioning	Raw material	Air quality	Disturbance protection	Water quality	Cognitive benefits	Leisure	Feel good/warm glove	Photosynthesis	Nutrient cycling	Nursery	Biodiversity	
A3	Infralittoral rock and other hard substrata	24.2	0.1	2	2	2	2	2	2	2	2	2	1	2	2	23
A4	Circalittoral rock and other hard substrata	208.6	0.5	2	2	1	2	2	2	2	2	0	2	2	2	21
A5.13	Infralittoral coarse sediment	29.8	0.1	2	2	0	0	0	0	1	1	0	1	2	1	10
A5.14	Circalittoral coarse sediment	0.1	0.0	2	2	0	0	0	0	0	0	0	1	1	1	7
A5.23	Infralittoral fine sands	2148.5	5.0	2	1	0	0	0	0	1	1	0	1	2	1	9
A5.25	Circalittoral fine sand	5717.8	13.2	2	1	0	0	0	0	0	0	0	1	2	1	7
A5.26	Circalittoral muddy sand	9693.2	22.5	2	1	0	0	1	0	0	0	0	1	1	1	7
A5.33	Infralittoral sandy mud	12.2	0.0	2	0	0	0	1	0	0	0	0	1	1	1	6
A5.34	Infralittoral fine mud	62.9	0.1	1	0	0	0	1	0	0	0	0	1	0	1	4
A5.35	Circalittoral sandy mud	10379.6	24.0	2	0	0	0	1	0	0	0	0	1	1	1	6
A5.36	Circalittoral fine mud	9203.1	21.3	2	0	0	0	1	0	0	0	0	1	1	1	6
A5.38	Med. biocoenosis of muddy detritic bottoms	82.1	0.2	1	0	0	0	1	0	0	0	0	1	0	1	4
A5.39	Med. biocoenosis of coastal terrigenous muds	50.3	0.1	2	0	0	0	1	0	0	0	0	1	1	1	6
A5.46	Med. biocoenosis of coastal detritic bottoms	5442.0	12.6	2	0	0	0	1	0	0	0	0	1	1	2	7
A5.531	<i>Cymodocea</i> beds	106.3	0.2	2	1	2	2	2	2	2	2	2	2	2	2	23

### References

- 52° North, SOS 52° North Development Team, 2018. SOS 52° North: Standardized, web-based upload and download of sensor data and sensor metadata.
- Allan, J.D., McIntyre, P.B., Smith, S.D.P., Halpern, B.S., Boyer, G.L., Buchsbaum, A., Burton, G.A., Campbell, L.M., Chadderton, W.L., Ciborowski, J.J.H., Doran, P.J., Eder, T., Infante, D.M., Johnson, L.B., Joseph, C.A., Marino, A.L., Prusevich, A., Read, J.G., Rose, J.B., Rutherford, E.S., Sowa, S.P., Steinman, A.D., 2013. Joint analysis of stressors and ecosystem services to enhance restoration effectiveness. *PNAS* 110, 372–377. <https://doi.org/10.1073/pnas.1213841110>
- Andersen, J.H., Stock, A., Heinänen, S., Mannerla, M., Vinther, M., 2013. Human uses, pressures and impacts in the eastern North Sea (Technical Report from DCE – Danish Centre for Environment and Energy No. 18).
- Atkinson, S.F., Canter, L.W., 2011. Assessing the cumulative effects of projects using geographic information systems. *Environmental Impact Assessment Review*, SPECIAL

- ISSUE on Cumulative Effects Assessment 31, 457–464.  
<https://doi.org/10.1016/j.eiar.2011.01.008>
- Ban, N.C., Bodtker, K.M., Nicolson, D., Robb, C.K., Royle, K., Short, C., 2013. Setting the stage for marine spatial planning: Ecological and social data collation and analyses in Canada's Pacific waters. *Marine Policy* 39, 11–20. <https://doi.org/10.1016/j.marpol.2012.10.017>
- Barale, V., Jaquet, J.-M., Ndiaye, M., 2008. Algal blooming patterns and anomalies in the Mediterranean Sea as derived from the SeaWiFS data set (1998–2003). *Remote Sensing of Environment, Earth Observations for Marine and Coastal Biodiversity and Ecosystems Special Issue* 112, 3300–3313. <https://doi.org/10.1016/j.rse.2007.10.014>
- Barbanti, A., Campostrini, P., Musco, F., Sarretta, A., Gissi, E., 2015. Developing a Maritime Spatial Plan for the Adriatic-Ionian Region. CNR-ISMAR, Venice, IT.
- Bastardie, F., Angelini, S., Bolognini, L., Fuga, F., Manfredi, C., Martinelli, M., Nielsen, J.R., Santojanni, A., Scarcella, G., Grati, F., 2017. Spatial planning for fisheries in the Northern Adriatic: working toward viable and sustainable fishing. *Ecosphere* 8, n/a-n/a. <https://doi.org/10.1002/ecs2.1696>
- Bianchi, C.N., 2004. Proposta di suddivisione dei Mari Italiani in Settori Biogeografici. *Notiziario della Società di Biologia Marina* 46, 57–59.
- Bijlsma, R., Bots, P., Wolters, H., Hoekstra, A., 2011. An Empirical Analysis of Stakeholders' Influence on Policy Development: the Role of Uncertainty Handling. *Ecology and Society* 16. <https://doi.org/10.5751/ES-03865-160151>
- Bokeh Development Team, 2018. Bokeh: Python library for interactive visualization.
- Bosnjakovic, B., Haber, I.M., 2015. Climate changes and adaptation policies in the baltic and the adriatic regions. *UTMS Journal of Economics* 6, 21.
- CEAA-ACEE (Canadian Environmental Assessment Agency), 2016. Technical Guidance for Assessing Cumulative Environmental Effects Technical Guidance Highlights. Appendix 2: Types of cumulative effects.
- Center for Ocean Solutions, 2011. Decision Guide - Selecting decision support tools for marine spatial planning. Stanford University, California.
- Certain, G., Jørgensen, L.L., Christel, I., Planque, B., Bretagnolle, V., 2015. Mapping the vulnerability of animal community to pressure in marine systems: disentangling pressure types and integrating their impact from the individual to the community level. *ICES J Mar Sci* 72, 1470–1482. <https://doi.org/10.1093/icesjms/fsv003>
- Chiogna, G., Majone, B., Cano Paoli, K., Diamantini, E., Stella, E., Mallucci, S., Lencioni, V., Zandonai, F., Bellin, A., 2016. A review of hydrological and chemical stressors in the Adige catchment and its ecological status. *Science of The Total Environment* 540, 429–443. <https://doi.org/10.1016/j.scitotenv.2015.06.149>
- Clarke Murray, C., Agbayani, S., Ban, N.C., 2015. Cumulative effects of planned industrial development and climate change on marine ecosystems. *Global Ecology and Conservation* 4, 110–116. <https://doi.org/10.1016/j.gecco.2015.06.003>
- COEXIST, 2013. Deliverable D3.9. COEXIST Project. Interaction in Coastal Waters: A Roadmap to Sustainable Integration of Aquaculture and Fisheries.
- Colloca, F., Cardinale, M., Maynou, F., Giannoulaki, M., Scarcella, G., Jenko, K., Bellido, J.M., Fiorentino, F., 2013. Rebuilding Mediterranean fisheries: a new paradigm for ecological

- sustainability. *Fish and Fisheries* 14, 89–109. <https://doi.org/10.1111/j.1467-2979.2011.00453.x>
- Depellegrin, D., Blažauskas, N., 2012. Integrating Ecosystem Service Values into Oil Spill Impact Assessment. *Journal of Coastal Research* 836–846. <https://doi.org/10.2112/JCOASTRES-D-11-00191.1>
- Depellegrin, D., Menegon, S., Farella, G., Ghezzi, M., Gissi, E., Sarretta, A., Venier, C., Barbanti, A., 2017. Multi-objective spatial tools to inform maritime spatial planning in the Adriatic Sea. *Science of The Total Environment* 609, 1627–1639. <https://doi.org/10.1016/j.scitotenv.2017.07.264>
- EC (European Council), 2017. Amending Directive 2008/56/EC of the European Parliament and of the Council as regards the indicative lists of elements to be taken into account for the preparation of marine strategies [WWW Document]. URL <http://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32017L0845&from=EN> (accessed 3.1.18).
- EC (European Council), 2014. Directive 2014/89/EU of the European Parliament and of the Council of the 23 July 2014 establishing a framework for Maritime Spatial Planning. *Off. J. Eur. Union* L257, 135.
- EC (European Council), 2011. Environmental Impact Assessment Directive. *EIA Directive* 2011/92.
- EC (European Council), 2008. Directive 2008/56/EC of the European Parliament of the Council of 17 June 2008 establishing a framework for community action in the field of marine environmental policy (Marine Strategy Framework Directive).
- EC (European Council), 2006. Regulation No. 1967/2006 of 21 December 2006, concerning management measures for the sustainable exploitation of fishery resources in the Mediterranean Sea, amending Regulation (EEC) No. 2847/93 and repealing Regulation (EC) No. 1626/94.
- EC (European Council), 2001. SEA (Strategic Environmental Assessment) Directive, 2001. Directive 2001/42/EC on the assessment of the effects of certain plans and programmes on the environment (SEA Directive) [WWW Document]. URL <http://ec.europa.eu/environment/eia/sea-legalcontext.htm> (accessed 2.19.18).
- Ehler, C., Douvère, F., 2009. Marine spatial planning, a step-by-step approach towards ecosystem-based management. (Report). UNESCO.
- Eigaard, O.R., Bastardie, F., Hintzen, N.T., Buhl-Mortensen, L., Buhl-Mortensen, P., Catarino, R., Dinesen, G.E., Egekvist, J., Fock, H.O., Geitner, K., Gerritsen, H.D., González, M.M., Jonsson, P., Kavadas, S., Laffargue, P., Lundy, M., Gonzalez-Mirelis, G., Nielsen, J.R., Papadopoulou, N., Posen, P.E., Pulcinella, J., Russo, T., Sala, A., Silva, C., Smith, C.J., Vanelslander, B., Rijnsdorp, A.D., 2016. The footprint of bottom trawling in European waters: distribution, intensity, and seabed integrity. *ICES Journal of Marine Science: Journal du Conseil* fsw194. <https://doi.org/10.1093/icesjms/fsw194>
- Elliott, M., 2011. Marine science and management means tackling exogenic unmanaged pressures and endogenic managed pressures – A numbered guide. *Marine Pollution Bulletin* 62, 651–655. <https://doi.org/10.1016/j.marpolbul.2010.11.033>
- Eno, N.C., Frid, C.L.J., Hall, K., Ramsay, K., Sharp, R. a. M., Brazier, D.P., Hearn, S., Dernie, K.M., Robinson, K.A., Paramor, O. a. L., Robinson, L.A., 2013. Assessing the sensitivity of



- habitats to fishing: from seabed maps to sensitivity maps. *J Fish Biol* 83, 826–846. <https://doi.org/10.1111/jfb.1213>
- Eurostat, 2017. Tourism statistics at regional level accessed [WWW Document]. URL [http://ec.europa.eu/eurostat/statistics-explained/index.php/Tourism\\_statistics\\_at\\_regional\\_level](http://ec.europa.eu/eurostat/statistics-explained/index.php/Tourism_statistics_at_regional_level) (accessed 2.2.18).
- Fortuna, C.M., Holcer, D., Mackelworth, P., 2015. Conservation of cetaceans and sea turtles in the Adriatic Sea: status of species and potential conservation measures (Report produced under WP7 of the NETCET project, IPA Adriatic Cross-border Cooperation Programme.).
- Fowler, C., Smith, B., Stein, D., 2010. Building a marine spatial data infrastructure to support marine spatial planning in US waters. *Geographic Technologies Applied to Marine Spatial Planning and Integrated Coastal Zone Management*. 120–128.
- Gallmetzer, I., Haselmair, A., Tomašových, A., Stachowitsch, M., Zuschin, M., 2017. Responses of molluscan communities to centuries of human impact in the northern Adriatic Sea. *PLoS one* 12, e0180820. <https://doi.org/10.1371/journal.pone.0180820>
- Galparsoro, I., Borja, A., Uyarra, M.C., 2014. Mapping ecosystem services provided by benthic habitats in the European North Atlantic Ocean. *Front. Mar. Sci.* 1. <https://doi.org/10.3389/fmars.2014.00023>
- Geneletti, D., 2011. Reasons and options for integrating ecosystem services in strategic environmental assessment of spatial planning. *International Journal of Biodiversity Science, Ecosystem Services & Management* 7, 143–149. <https://doi.org/10.1080/21513732.2011.617711>
- GeoServer Development Team, 2018. GeoServer: open source server for sharing geospatial data.
- Gimpel, A., Stelzenmüller, V., Töpsch, S., Galparsoro, I., Gubbins, M., Miller, D., Murillas, A., Murray, A.G., Pınarbaşı, K., Roca, G., Watret, R., 2018. A GIS-based tool for an integrated assessment of spatial planning trade-offs with aquaculture. *Science of The Total Environment*. <https://doi.org/10.1016/j.scitotenv.2018.01.133>
- Gissi, E., Menegon, S., Sarretta, A., Appiotti, F., Maragno, D., Vianello, A., Depellegrin, D., Venier, C., Barbanti, A., 2017. Addressing uncertainty in modelling cumulative impacts within maritime spatial planning in the Adriatic and Ionian region. *PLOS ONE* 12, e0180501. <https://doi.org/10.1371/journal.pone.0180501>
- González Del Campo, A., 2017. Mapping environmental sensitivity: A systematic online approach to support environmental assessment and planning. *Environmental Impact Assessment Review* 66, 86–98. <https://doi.org/10.1016/j.eiar.2017.06.010>
- Gramolini, R., Grati, F., Fabi, G., Schule, T., 2010. GRID GeoReference Interactions Database. Deliverable D3.9. COEXIST Project. Interaction in coastal waters: A roadmap to sustainable integration of aquaculture and fisheries.
- Hadjimitsis, D.G., Agapiou, A., Themistocleous, K., Mettas, C., Evagorou, E., Papoutsas, C., Nisantzi, A., Mammouri, R., Tzouvaras, M., Lyssandrou, V., 2015. Resolving sea and land conflicts in Cyprus using Marine Spatial Planning, in: *Proceedings of the 14th International Conference on Environmental Science and Technology CEST 2015*. Athens, Greece.

- Hadžić, N., Kozmar, H., Tomić, M., 2014. Offshore renewable energy in the Adriatic Sea with respect to the Croatian 2020 energy strategy. *Renewable and Sustainable Energy Reviews* 40, 597–607. <https://doi.org/10.1016/j.rser.2014.07.196>
- Halpern, B.S., Frazier, M., Potapenko, J., Casey, K.S., Koenig, K., Longo, C., Lowndes, J.S., Rockwood, R.C., Selig, E.R., Selkoe, K.A., Walbridge, S., 2015. Spatial and temporal changes in cumulative human impacts on the world's ocean. *Nature Communications* 6, 7615. <https://doi.org/10.1038/ncomms8615>
- Halpern, B.S., Walbridge, S., Selkoe, K.A., Kappel, C.V., Micheli, F., D'Agrosa, C., Bruno, J.F., Casey, K.S., Ebert, C., Fox, H.E., Fujita, R., Heinemann, D., Lenihan, H.S., Madin, E.M.P., Perry, M.T., Selig, E.R., Spalding, M., Steneck, R., Watson, R., 2008. A Global Map of Human Impact on Marine Ecosystems. *Science* 319, 948–952. <https://doi.org/10.1126/science.1149345>
- Hansen, H.S., Reiter, I.M., Schrøder, L., 2017. A System Architecture for a Transnational Data Infrastructure Supporting Maritime Spatial Planning, in: *Electronic Government and the Information Systems Perspective, Lecture Notes in Computer Science*. Presented at the International Conference on Electronic Government and the Information Systems Perspective, Springer, Cham, pp. 158–172. [https://doi.org/10.1007/978-3-319-64248-2\\_12](https://doi.org/10.1007/978-3-319-64248-2_12)
- Harrison, P.A., Berry, P.M., Simpson, G., Haslett, J.R., Blicharska, M., Bucur, M., Dunford, R., Egoh, B., Garcia-Llorente, M., Geamănă, N., Geertsema, W., Lommelen, E., Meiresonne, L., Turkelboom, F., 2014. Linkages between biodiversity attributes and ecosystem services: A systematic review. *Ecosystem Services* 9, 191–203. <https://doi.org/10.1016/j.ecoser.2014.05.006>
- Hooper, T., Beaumont, N., Griffiths, C., Langmead, O., Somerfield, P.J., 2017. Assessing the sensitivity of ecosystem services to changing pressures. *Ecosystem Services* 24, 160–169. <https://doi.org/10.1016/j.ecoser.2017.02.016>
- Human uses, pressures and impacts in the eastern North Sea - Anderson et al 2013.pdf, n.d.
- IUCN, 2013. An International Instrument on Conservation and Sustainable Use of Biodiversity in Marine Areas beyond National Jurisdiction Exploring Different Elements to Consider. PAPER VII: Relation between Environmental Impact Assessments, Strategic Environmental Assessments and Marine Spatial Planning.
- JPI OCEANS, 2017. JPI Oceans workshop: Cumulative effects of human activities [WWW Document]. URL <http://www.jpi-oceans.eu/news-events/events/jpi-oceans-workshop-cumulative-effects-human-activities> (accessed 2.19.18).
- Judd, A.D., Backhaus, T., Goodsir, F., 2015. An effective set of principles for practical implementation of marine cumulative effects assessment. *Environmental Science & Policy* 54, 254–262. <https://doi.org/10.1016/j.envsci.2015.07.008>
- Krueger, I., Schouten-de Groot, P., 2011. KPP Noordzee: Tools (Deltares Report).
- Malačić, V., Faganeli, J., Malej, A., 2008. Environmental Impact Of Lng Terminals In The Gulf Of Trieste (Northern Adriatic), in: *Integration of Information for Environmental Security, NATO Science for Peace and Security Series C: Environmental Security*. Springer, Dordrecht, pp. 361–381. [https://doi.org/10.1007/978-1-4020-6575-0\\_25](https://doi.org/10.1007/978-1-4020-6575-0_25)

- Marcotte, D., Hung, S.K., Caquard, S., 2015. Mapping cumulative impacts on Hong Kong's pink dolphin population. *Ocean & Coastal Management* 109, 51–63. <https://doi.org/10.1016/j.ocecoaman.2015.02.002>
- Maron, M., Mitchell, M.G.E., Runting, R.K., Rhodes, J.R., Mace, G.M., Keith, D.A., Watson, J.E.M., 2017. Towards a Threat Assessment Framework for Ecosystem Services. *Trends in Ecology & Evolution* 32, 240–248. <https://doi.org/10.1016/j.tree.2016.12.011>
- McKinney, W., 2010. Data structures for statistical computing in python, in: *Proceedings of the 9th Python in Science Conference*. Austin, TX, pp. 51–56.
- Menegon, S., 2018a. Tools4MSP: Geospatial tools to support Maritime Spatial Planning. <https://doi.org/10.5281/zenodo.1186159>
- Menegon, S., 2018b. Rectifiedgrid: Geospatial python library for grid-based analyses. <https://doi.org/10.5281/zenodo.1160651>
- Menegon, S., Depellegrin, D., Farella, G., Gissi, E., Sarretta, A., Venier, C., Barbanti, A., 2018a. Developing a modelling framework for MSP-oriented cumulative effects assessment: A case study for the Italian Adriatic Sea. *Ecological Indicators* (accepted).
- Menegon, S., Depellegrin, D., Venier, C., 2018b. Geospatial dataset for Cumulative Effects Assessment in the Italian Adriatic Sea [WWW Document]. URL <https://doi.org/10.5281/zenodo.1173764> (accessed 2.5.18).
- Menegon, S., Sarretta, A., Barbanti, A., Gissi, E., Venier, C., 2016. Open source tools to support Integrated Coastal Management and Maritime Spatial Planning, in: Marchesini, I., Pierleoni, A. (Eds.), *Proceedings of the 4th Open Source Geospatial Research and Education Symposium (OGRS2016)*. Perugia. [https://doi.org/10.30437/ogrs2016\\_paper\\_22](https://doi.org/10.30437/ogrs2016_paper_22)
- Meyer, R., 2012. Uncertainty as a science policy problem. *Climatic Change* 110, 1–2. <https://doi.org/10.1007/s10584-011-0050-8>
- MSP Platform, 2018. Summary: Cumulative Impact Tools for MSP Expert Roundtable [WWW Document]. MSP Platform. URL [http://msp-platform.eu/sites/default/files/20180214\\_finalcumulativeimpactsroundtable\\_msplatform\\_documentation.pdf](http://msp-platform.eu/sites/default/files/20180214_finalcumulativeimpactsroundtable_msplatform_documentation.pdf) (accessed 2.19.18).
- MSP Platform, 2016. Symphony: Tool to estimate cumulative impacts [WWW Document]. European MSP Platform. URL <http://msp-platform.eu/practices/symphony-tool-estimate-cumulative-impacts> (accessed 2.19.18).
- Munaretto, S., Huitema, D., 2012. Adaptive Comanagement in the Venice Lagoon? An Analysis of Current Water and Environmental Management Practices and Prospects for Change. *Ecology and Society* 17. <https://doi.org/10.5751/ES-04772-170219>
- Palomino, J., Muellerklein, O.C., Kelly, M., 2017. A review of the emergent ecosystem of collaborative geospatial tools for addressing environmental challenges. *Computers, Environment and Urban Systems* 65, 79–92. <https://doi.org/10.1016/j.compenvurbsys.2017.05.003>
- Papageorgiou, M., 2016. Coastal and marine tourism: A challenging factor in Marine Spatial Planning. *Ocean & Coastal Management* 129, 44–48. <https://doi.org/10.1016/j.ocecoaman.2016.05.006>

- Papathanasopoulou, E., Beaumont, N., Hooper, T., Nunes, J., Queirós, A.M., 2015. Energy systems and their impacts on marine ecosystem services. *Renewable and Sustainable Energy Reviews* 52, 917–926. <https://doi.org/10.1016/j.rser.2015.07.150>
- Piante, C., Ody, D., 2015. Blue growth in the Mediterranean Sea: the challenge of good environmental status. *MedTrends Project*. WWF-France 192.
- Pınarbaşı, K., Galparsoro, I., Borja, Á., Stelzenmüller, V., Ehler, C.N., Gimpel, A., 2017. Decision support tools in marine spatial planning: Present applications, gaps and future perspectives. *Marine Policy* 83, 83–91. <https://doi.org/10.1016/j.marpol.2017.05.031>
- Populus, J., Vasquez, M., Albrecht, J., Manca, E., Agnesi, S., Al Hamdani, Z., Andersen, J., Annunziatellis, A., Bekkby, T., Bruschi, A., 2017. EUSeaMap. A European broad-scale seabed habitat map (Final Report).
- Port of Koper, 2015. New Cranes for the Container Terminal [WWW Document]. Luka Koper. URL <https://luka-kp.si/eng/news/single/new-cranes-for-the-container-terminal-4075> (accessed 2.19.18).
- Pranovi, F., Raicevich, S., Franceschini, G., Farrace, M.G., Giovanardi, O., 2000. Rapido trawling in the northern Adriatic Sea: effects on benthic communities in an experimental area. *ICES J Mar Sci* 57, 517–524. <https://doi.org/10.1006/jmsc.2000.0708>
- QGIS Development Team, 2018. QGIS A Free and Open Source Geographic Information System [WWW Document]. URL <https://www.qgis.org/en/site/> (accessed 3.1.18).
- RER (Regione Emilia Romagna), 2015. Piano Regionale Integrato dei Trasporti 2025 (Technical Report).
- Russo, T., Pulcinella, J., Parisi, A., Martinelli, M., Belardinelli, A., Santojanni, A., Cataudella, S., Colella, S., Anderlini, L., 2015. Modelling the strategy of mid-water trawlers targeting small pelagic fish in the Adriatic Sea and its drivers. *Ecological Modelling* 300, 102–113. <https://doi.org/10.1016/j.ecolmodel.2014.12.001>
- Salomidi, M., Katsanevakis, S., Borja, A., Braeckman, U., Damalas, D., Galparsoro, I., Mifsud, R., Mirto, S., Pascual, M., Pipitone, C., Rabaut, M., Todorova, V., Vassilopoulou, V., Vega Fernández, T., 2012. Assessment of goods and services, vulnerability, and conservation status of European seabed biotopes: a stepping stone towards ecosystem-based marine spatial management. *Mediterranean Marine Science*. <https://doi.org/10.12681/mms.23>
- Schweizer, J., Antonini, A., Govoni, L., Gottardi, G., Archetti, R., Supino, E., Berretta, C., Casadei, C., Ozzi, C., 2016. Investigating the potential and feasibility of an offshore wind farm in the Northern Adriatic Sea. *Applied Energy* 177, 449–463. <https://doi.org/10.1016/j.apenergy.2016.05.114>
- Seasketch, 2018. SeaSketch Supports Collaborative Planning for our Oceans. [WWW Document]. URL <https://www.seasketch.org/home.html> (accessed 2.19.18).
- Simonini, R., Ansaloni, I., Bonvicini Pagliai, A.M., Prevedelli, D., 2004. Organic enrichment and structure of the macrozoobenthic community in the northern Adriatic Sea in an area facing Adige and Po mouths. *ICES J Mar Sci* 61, 871–881. <https://doi.org/10.1016/j.icesjms.2004.06.018>
- Slootweg, R., Van Beukering, P., 2008. Valuation of ecosystem services and strategic environmental assessment-Lessons from influential cases.

- Socal, G., Acri, F., Bastianini, M., Bernardi Aubry, F., Bianchi, F., Camatti, E., Conversi, A., De Lazzari, A., Pugnetti, A., 2011. The study and the impact of plankton communities over the northern Adriatic pelagic ecosystem, in: Volume Mare. pp. 141–151.
- Söderman, T., Kopperoinen, L., Shemeikka, P., Yli-Pelkonen, V., 2012. Ecosystem services criteria for sustainable development in urban regions. *J. Env. Assmt. Pol. Mgmt.* 14, 1250008. <https://doi.org/10.1142/S1464333212500081>
- Song, N.-Q., Wang, N., Lin, W.-N., Wu, N., 2017. Use of pollution risk and ecosystem service values in marine management strategies in Bohai Sea, China. *Human and Ecological Risk Assessment: An International Journal* 23, 1041–1055. <https://doi.org/10.1080/10807039.2017.1297675>
- Stelzenmüller, V., Lee, J., South, A., Foden, J., Rogers, S.I., 2013. Practical tools to support marine spatial planning: A review and some prototype tools. *Marine Policy* 38, 214–227. <https://doi.org/10.1016/j.marpol.2012.05.038>
- Stelzenmüller, V., Vega Fernández, T., Cronin, K., Röckmann, C., Pantazi, M., Vanaverbeke, J., Stamford, T., Hostens, K., Pecceu, E., Degraer, S., Buhl-Mortensen, L., Carlström, J., Galparsoro, I., Johnson, K., Piwowarczyk, J., Vassilopoulou, V., Jak, R., Louise Pace, M., van Hoof, L., 2015. Assessing uncertainty associated with the monitoring and evaluation of spatially managed areas. *Marine Policy* 51, 151–162. <https://doi.org/10.1016/j.marpol.2014.08.001>
- TMT, 2016. Trieste Marine Terminal [WWW Document]. Trieste Marine Terminal. URL <http://www.trieste-marine-terminal.com/it> (accessed 2.21.18).
- Turk, R., Odorico, R., 2009. Marine protected areas in the Northern Adriatic. *Varstvo narave* 22, 33–45.
- van der Walt, S., Colbert, S.C., Varoquaux, G., 2011. The NumPy Array: A Structure for Efficient Numerical Computation. *Computing in Science Engineering* 13, 22–30. <https://doi.org/10.1109/MCSE.2011.37>
- van Riper, C.J., Kyle, G.T., Sutton, S.G., Barnes, M., Sherrouse, B.C., 2012. Mapping outdoor recreationists' perceived social values for ecosystem services at Hinchinbrook Island National Park, Australia. *Applied Geography* 35, 164–173. <https://doi.org/10.1016/j.apgeog.2012.06.008>
- Vanderhaegen, M., Muro, E., 2005. Contribution of a European spatial data infrastructure to the effectiveness of EIA and SEA studies. *Environmental Impact Assessment Review* 25, 123–142. <https://doi.org/10.1016/j.eiar.2004.06.011>
- Werner, S.R., Spurgeon, J.P.G., Isaksen, G.H., Smith, J.P., Springer, N.K., Gettleson, D.A., N'Guessan, L., Dupont, J.M., 2014. Rapid prioritization of marine ecosystem services and ecosystem indicators. *Marine Policy* 50, 178–189. <https://doi.org/10.1016/j.marpol.2014.03.020>
- WESTMED, 2017. Findings and Recommendations from stakeholder involvement in Phase 1 [WWW Document]. URL [http://www.westmed-initiative.eu/wp-content/uploads/2016/07/WestMed-Maritime-Initiative-Report2\(public\).pdf](http://www.westmed-initiative.eu/wp-content/uploads/2016/07/WestMed-Maritime-Initiative-Report2(public).pdf) (accessed 2.19.18).

- White, C., Halpern, B.S., Kappel, C.V., 2012. Ecosystem service tradeoff analysis reveals the value of marine spatial planning for multiple ocean uses. *PNAS* 109, 4696–4701. <https://doi.org/10.1073/pnas.1114215109>
- White, L., Koss, R., Knights, A., Eriksson, A., Robinson, L., 2013. ODEMM linkage framework userguide (Version 2). ODEMM Guidance Document Series.
- Worm, B., Barbier, E.B., Beaumont, N., Duffy, J.E., Folke, C., Halpern, B.S., Jackson, J.B.C., Lotze, H.K., Micheli, F., Palumbi, S.R., Sala, E., Selkoe, K.A., Stachowicz, J.J., Watson, R., 2006. Impacts of Biodiversity Loss on Ocean Ecosystem Services. *Science* 314, 787–790. <https://doi.org/10.1126/science.1132294>
- Wyatt, K.H., Griffin, R., Guerry, A.D., Ruckelshaus, M., Fogarty, M., Arkema, K.K., 2017. Habitat risk assessment for regional ocean planning in the U.S. Northeast and Mid-Atlantic. *PLOS ONE* 12, e0188776. <https://doi.org/10.1371/journal.pone.0188776>