



Research Article

IN-VITRO ANTIHELMINTHIC ACTIVITY, PHYTOCHEMICAL SCREENING AND TLC STUDIES OF ETHANOL-WATER EXTRACTION ON *IPOMEA CARNEA* FLOWER USING IN - STATE FESTIVAL OF TELANGANA (BATHUKAMMA)P. Sravan Kumar ^{1*}, A. Santhosh ², K. Manasa, M. Swapna ³, M. Sreevani ³, M. Bhavusingh ³, M. Mamatha ³¹ Associate Professor, Department of Pharmacology, Avanthi Institute of Pharmaceutical Sciences, Gunthapally-501512, Hyderabad, Telangana State, INDIA.² Assistant Professor, Department of Pharmacognosy, Avanthi Institute of Pharmaceutical Sciences, Gunthapally-501512, Hyderabad, Telangana State, INDIA.³ B. Students, Department of Pharmacy, Avanthi Institute of Pharmaceutical Sciences, Gunthapally-501512, Hyderabad, Telangana State, INDIA.

Received on: 29-03-2018; Revised and Accepted on: 21-04-2018

ABSTRACT

India Being a Rich and Varied Flora of Medicinal Plants. It's used in Traditional Festivals of India. The Present Study deals with the In-Vitro Antihelmintic Activity, Phytochemical Screening and TLC Studies of Ethanol:Water (80:20) Extraction on *Ipomea Carnea* Flower using in - State Festival of Telangana (Bathukamma). Ethanol:Water (80:20) Extraction on *Ipomea Carnea* Flower was used evaluation of Phytochemical Screening Determination by some chemical tests and Thin Layer Chromatographic Study was carried out by using Various Solvent System of varying Polarity of Hexane, Ethyl Acetate, Acetone. Phytochemical Screening Reflects Presence of like Carbohydrates, Alkalides, Phenols, Tannins, Phytosterols, Glycosides, Flavonoids, Tannins Shows Methanol Solvents Extracts. Thin Layer Chromatographic Studies of the *Ipomea Carnea* Flower Parts Constituted Different Colored Phytochemical Compounds with different R_f Values. Ethanol:Water (80:20) Extractions of *Ipomea Carnea* Flower Various Concentrations (25, 50, 100mg/ml) of all Extracts were Tested and Results were Expressed in Terms of time for Paralysis and time for Death of Worms. Piperazine Citrate (10 mg/ml) was used as a Reference Standard and Distilled Water as a Control Group. Treatment with Concluded that the Ethanol:Water (80:20) Extract of *Ipomea Carnea* Flowers showed Potent Anthelmintic Activity and was Equipotent to Standard Anthelmintic drug. The Potent Anthelmintic Activity could be due to Presence of Glycosides, Flavonoids and Sterols. So, from the above Findings, it was Concluded that Ethanol:Water (80:20) Extract of *Ipomea Carnea* Flowers Posses's Significant Wormicide Activity Property.

KEYWORDS: *Ipomea Carnea*, Anthelmintic, Phytochemicals, TLC Profile.

INTRODUCTION

Herbal medicines have recently attracted much attention as alternative medicines useful for treating or prevent in life style related disorders and relatively very little knowledge is available about their mode of action. There has been a growing interest in the analysis of plant products which has stimulated intense research on their potential health benefits ^[1].

The earth is created with many incredible things which are supporting the life of human beings; we found our life in the nature which gives everything we want like water, food, and shelter etc. and everyone pray for these things to remain forever with us. This nature looks more beautiful with different flowers, each flower has a unique fragrance which attracts everyone and these flowers were used for celebrations and other occasions, we also see various plants in the nature and each plant has its own importance ^[2].

Grate India is known for its traditions and celebrations of festivals. Every festival has a scientific reason to support its celebration.

in this regard, present study is planned to evaluate pharmacological activity of flowers used in bathukamma festival which has been recently declared as state festival of newly formed telangana state ^[3]. It is a floral festival in which every day various colored flowers are arranged row after row in a brass plate, called as bathukamma, placed in front of diety and daily worshipped for a week. In the evening it is carried to nearby pond or any water body and released in it.

Helminthes parasite infections are global problems. The diseases affect the health status of a large fraction of the human population as well as animals. Some type of dangerous helminthes infections like filariasis has only a few therapeutic modalities at present ^[4]. In addition, after treatment with albendazole or mebendazole, several side effects ^[5]. Helminths are the most common infectious agents of humans and produce a global burden of disease and contribute to the prevalence of malnutrition, anaemia, eosinophilia, and pneumonia. The disease is highly prevalent particularly in poor countries. Plant derived drug serve as a prototype to develop more effective and less toxic medicines ^[5-7].

In India many times used in traditional medicine system. It is a common weed throughout india and used in traditional medicine for *ipomea carnea* flower is most important flower used in this festival. It is commonly known pink morning glory, is a species of morning glory, besharam in hindi, thutu kada in telugu, morning glory flowers are also known as ipomea, it is belongs to family convolvulaceae ^[8].

The recent studies displayed that *ipomea carnea* possessed a wide range of therapeutic activities which were proved that this plant have a potential regenerator capacity of various cells, glycosidase inhibitory activities, anti-inflammatory activity, antioxidant activity, wound healing activity, antidiabetic activity, immunomodulatory

Corresponding author:*P. Sravan Kumar**Associate Professor,
Department of Pharmacology,
Avanthi Institute of Pharmaceutical Sciences,
Gunthapally-501512, Hyderabad, Telangana State, INDIA.

* E-Mail:

DOI:

activity, antimicrobial activity, antibacterial activity, cardiovascular activity, embryotoxic effect, antifungal activity, hepatoprotective activity, anti hiv activity, anxiolytic activity. The wide range of therapeutic potentialities of *ipomea carnea* are mainly due to the presence of various bioactive molecules in flowers, roots, stems, leaves and herbs plant *ipomea carnea* of family convolvulaceae are all ornamental and colourful^[8,9].

Resent study was shows flower tops have been used many pharmacological activity. To support the significance of immersing these flowers in water reservoirs, their anthelmintics potential was studied in the present investigation.

MATERIALS AND METHODS

1. Plant Material:

Fresh flowers of *ipomea carnea* were collected from herbal garden of gunthapally village, near ramoji film city in month of october and november-2017. This plant material was identified at dept. of botany, Dr. S. baburaj, botanist, M.S college of arts and science, kodad. Around 700 gms of fresh flowers were collected and washed with fresh water. The flowers were then dried under shade and homogenized to get a coarse powder. This powder was stored in an air tight container and used for further solvent extraction^[10-12].

2. Preparation of Extract By Sox Halation:

Dried *Ipomoea Carnea* flowers powdered (250 gm) materials were of soxhlation by ml of 500 ml of ethanol:water (80:20) extracted with 6 hrs. The solution was then filtered using sterilized cotton and buchner funnel^[10-12]. The filtrate was concentrated to evaporate solvent using rotary evaporator at 40 °c and 50 rpm. Finally, 16.37g (yield 8.84 %) of dried extract was obtained and this crude extract was used for phytochemical screening, thin layer chromatography & anthelmintica activity studies.

3. Qualitative Phytochemical Analysis:

The powdered material and extract of the plants were subjected to different kinds of chemical tests to investigate the presence of secondary metabolites such as saponins, tannins, flavonoids, phenol, anthraquinones cyanogenic glycosides, cardiac glycosides and alkaloids etc using standard procedures^[13-15]. Results Are Shown In (**Table-1**)

4. Test For Proteins:

4.1. Millon's Test:

Crude extract when mixed with 2ml of millon's reagent, white precipitate appeared which turned red upon gentle heating that confirmed the presence of protein.

5. Test For Carbohydrates:

5.1. Fehling's Test:

Equal volume of fehling a and fehling b reagents were mixed together and 2ml of it was added to crude extract and gently boiled. A brick red precipitate appeared at the bottom of the test tube indicated the presence of reducing sugars.

6. Iodine Test:

Crude extract was mixed with 2ml of iodine solution. A dark blue or purple coloration indicated the presence of the carbohydrate.

7. Test for Phenols and Tannins:

Crude extract was mixed with 2ml of 2% solution of fecl₃. A blue-green or black coloration indicated the presence of phenols and tannins.

8. Test for Flavonoids:

To 1 ml of extract, 1ml of 10% lead acetate solution was added. The formation of a yellow precipitate was taken as a positive test for presence of flavonoids.

9. Alkaline Reagent Test:

Crude extract was mixed with 2ml of 2% solution of naoh. An intense yellow colour was formed which turned colourless on addition of few drops of diluted acid which indicated the presence of flavonoids.

9.1. Test for Phytosterol:

The extract was refluxed with solution of alcoholic potassium hydroxide till complete saponification takes place. The mixture was diluted and extracted with ether. The ether layer was evaporated and the residue was tested for the presence of phytosterol. The residue was dissolved in few drops of diluted acetic acid; 3 ml of acetic anhydride was added followed by few drops of concentrated H₂SO₄. Appearance of bluish green colour showed the presence of phytosterol.

9.2. Test for Triterpenoids:

10mg of the extract was dissolved in 1 ml of chloroform, 1 ml of acetic anhydride was added followed by addition of 2 ml of concentrated H₂SO₄. Formation of reddish violet colour indicates the presence of triterpenoids.

9.3. Test for Tannins:

About 2 ml of aqueous extract was added to 2 ml of 1% HCl and the mixture was boiled. Deposition of a red precipitate was taken as an evidence for the presence of tannins.

About 2 ml of the extract was stirred with 2ml of distilled water and few drops of ferric chloride (FeCl₃) solution were added. Formation of green precipitate was indication of presence of tannins.

9.4. Test for Saponins:

Crude extract was mixed with 5ml of distilled water in a test tube and it was shaken vigorously. The formation of stable foam was taken as an indication for the presence of saponins.

9.5. Test for Glycosides:

Liebermann's Test: Crude extract was mixed with each of 2ml of chloroform and 2ml of acetic acid. the mixture was cooled in ice. Carefully concentrated h₂so₄ was added. A colour change from violet to blue to green indicated the presence of steroidal nucleus, that is, glycone portion of glycoside.

9.6. Keller-Kilani Test:

Crude extract was mixed with 2ml of glacial acetic acid containing 1-2 drops of 2% solution of fecl₃. The mixture was then poured into another test tube containing 2ml of concentrated H₂SO₄. A brown ring at the interphase indicated the presence of cardiac glycosides.

9.7. Test for Steroid:

Crude extract was mixed with 2ml of chloroform and concentrated h₂so₄ was added sidewise. a red colour produced in the lower chloroform layer indicated the presence of steroids. Another test was performed by mixing crude extract with 2ml of chloroform. 2ml of each of concentrated H₂SO₄ and acetic acid were poured into the mixture. The development of a greenish coloration indicated the presence of steroids.

Test for Steroids A. Salkowski, S Test: A red color produced in the lower chloroform layer when 2 ml of organic extract was dissolved in 2 ml of chloroform and 2 ml concentrated H₂SO₄ was added in it, indicates the presence of steroids.

Liebermann Burchard Test: Development of a greenish color when 2 ml of the organic extract was dissolved in 2 ml of chloroform and treated with concentrated H₂SO₄ and acetic acid indicates the presence of steroids.

9.8. Test for Terpenoids:

Crude extract was dissolved in 2ml of chloroform and evaporated to dryness. To this, 2ml of concentrated H₂SO₄ was added and heated for about 2 minutes. A grayish colour indicated the presence of terpenoids.

9.9. Test for Alkaloids:

Crude extract was mixed with 2ml of 1% hcl and heated gently. Mayer's and Wagner's reagents were then added to the mixture. Turbidity of the resulting precipitate was taken as evidence for the presence of alkaloids.

Thin Layer Chromatography:

Thin Layer Chromatography of extracts was done by using standard procedures and is mainly used for the detection of the nature of phytoconstituents present. Thin layer chromatography is a very effective technique for the separation of chemical constituents of an extract and for their identification. The history of TLC has been reviewed by various authors. TLC profiles developed for an extract from a defined solvent system and other parameters could be used as a fingerprint in comparative qualitative evaluation of herbal drugs.

The trend of evaluation by this method is becoming popular in view of its simplicity and reproducibility. TLC is an important analytical tool in the separation, identification and estimation of different classes of natural products. In this technique, the different components are separated by the differential migration of solute between two phases – a stationary phase and a mobile phase. Here, the principle of separation is adsorption and the stationary phase acts as an adsorbent. depending on the particular type of stationary phase, its preparation and use with different solvents, separation can be achieved on the basis of partition or a combination of partition and adsorption in this methods used artificial aluminum oxide tlc plate was using, the plates were activated by heating at 100°C for 10 minutes.

Solvent systems used in TLC different solvent system [hexane: acetic acid (9:1)] solvent system ii [hexane: ethyl acetate: acetic acid (5:4:1)], in solvent system iii [hexane: ethyl acetate: acetic acid (4:4:2)], in solvent system IV [hexane: ethyl acetate: acetic acid (3:6:1)] used. After pre-saturation with mobile phase for 20 min for development were used. After the run plates are dried and TLC plate place the developing jar containing the iodine were used to detect staining process the bands on the TLC plates. The movement of the active compound was expressed by its retention factor (R_f), values were calculated for different sample [15].

$$R_f = \frac{\text{Distance Traveled By Solutes}}{\text{Distance Traveled By Solvent Front By TLC}}$$

Anthelmintic Activity:

The anthelmintic activity was performed according to the method of on adult indian earthworm *pheritima posthuma* as it has anatomical and physiological resemblances with the intestinal roundworm parasites of human beings. Five groups of approximately equal sized indian earthworms consisting of six earthworms in each group were released into 50ml of desired formulation.

Group first serve as control, receive only normal saline, group second serve as standard, receive standard drug piperazine citrate (10mg/ml) group third serve as test dose-1, receive concentrations of (25 mg/ml) eweic, group fourth serve as test dose -2, receive concentrations of (50 mg/ml) eweic, group fifth serve as test dose -3, receive different concentrations of (100 mg/ml) eweic and observations were made for the time taken to paralyse or death of individual worms. Six worms (same type) in each were placed in it. Time for paralysis was noted when no movement observed except when the worms were shaken vigorously. time for death of worms were recorded after ascertaining that the worms neither moved when shaken vigorously nor when dipped in warm water (50°C). Paralysis was said to occur when the worms do not receive even in normal saline. Death was concluded when the worms lose their motility followed with fading away of their body colour [16,17]. Results are shown in (Table-1).

Group I (Normal Control)	: Normal Saline
Group II (Standard Treated)	: Piperazine citrate (10mg/MI)
Group III (EWEIC Dose-1)	: (25 mg/ml) concentrations of EWEIC
Group IV (EWEIC Dose-2)	: (50 mg/ml) concentrations of EWEIC
Group V (EWEIC Dose-3)	: (100 mg/ml) concentrations of EWEIC

RESULTS AND DISCUSSION

Table No. 1: Preliminary Phytochemical Screening of the Ethanol:Water (80:20) Extract of *Ipomoea Carnea* Flowers

Plant Constituents Test	Preliminary Phytochemical Screening
1. Test For Alkaloids	
A) Dragendroff's Test	+
B) Mayer's Test	+
2. Test For Carbohydrate	
A) Fehling's Test	+
B) Iodine Test	+
3. Test For Flavonoids	
A) Lead Acetate Test	+
B) Alkaline Reagent Test	+
4. Test For Fixed Oils	
A) Spot's Test	-
B) Saponification's Test	
5. Test For Phytosterols	
A) Salkowski Test	-
B) Liebermann-Burchard Test	-
6. Test For Glycosides	
A) Legal's Test	+
B) Borntrager's Test	+
7. Test For Proteins	
A) Millon's Test	+
8. Test For Phenol & Tannins	
A) 5% FeCl ₃ Solution	+
9. Test For Triterpenoids	
A) Salkowaski Test	+
B) Libermann-Burchard Test	+
10. Test For Amino Acids	
A) Ninhydrin Test	+
11). Test For Saponines	-

"+" Indicates Presence "-" Indicates Absence

Table No. 2: Effect of Ethanol: Water Extract of *Ipomoea Carnea* Flower on Indian Earth Worms (*Pheritima Posthuma*)

Group	Treatment (Dose)	Time for Paralysis of Worms (Mins)	Time for Death of Worms (Mins)
I(Normal Control)	Saline	-	-
II(Standard Treated)	10 mg/ml of piperazine citrate	22.10±2.10	45.07±1.33
III(EWEIC Dose-1)	25 mg/ml of EWEIC	46.20±1.15	63.15±2.10
IV (EWEIC Dose-2)	50 mg/ml of EWEIC	35±0.50	55.12±2.50
V (EWEIC Dose-3)	100 mg/ml of EWEIC	28.12±1.44	51.04±0.61

All Values Represent Mean +SD; N= 3 In Each Groups

Table No. 3: R_f Values of TLC with Respect to Ethanol: Water Extract of *Ipomoea Carnea* Flower selected using Different Mobile Phase Solvent Systems

S. No.	Solvent System	No of Spots (EWEIC)	R _f Values (EWEIC)
1.	Hexane: Acetic Acid (9:1)	3	0.33,0.45&0.86
2.	Hexane: Ethyl Acetate: Acetic Acid (5:4:1)	4	0.84,0.89,0.92&0.98
3.	Hexane: Ethyl Acetate: Acetic Acid (4:4:2)	1	0.92
4	Hexane: Ethyl Acetate: Acetic Acid (3:6:1)	2	0.55&0.91



Fig. 1: Preliminary Phytochemical Screening of the Ethanol: Water Extract of *Ipomoea Carnea* Flower




Fig. 2: Effect of Ethanol: Water Extract of *Ipomoea Carnea* Flower on Indian Earth Worms (*Pheritima Posthuma*)




Fig. 3: The predominant effect of piperazine citrate on worm Paralysis and death by various concentrations




Fig. 4: TLC With Respect to Ethanol: Water Extract of *Ipomoea Carnea* (spot-B) flowers Selected Using Different Mobile Phase Solvent Systems

Phytochemical Screening:

The results of preliminary phytochemical analysis of methanol extract *celosia cristata* flowers showed the presence of various phytochemical constituents like carbohydrates, alkaloids, phenols, tannins, phytosterols, glycosides, flavonoids, tannins (Table-1 & Fig. 1).

Anthelmintic activity of ethanol: water extract of *ipomoea carnea* flowers is confirmed by examining the time taken for paralysis (p) and death (d) for *pheretima posthuma* worms were reported in table 1. Papaya contains many biologically active compounds. The assay was performed on adult indian earthworm, *pheretima posthuma* due to its anatomical and physiological resemblance with the intestinal roundworm parasite of human beings, because of easy availability. Piperazine citrate by increasing chloride ion conductance of worm muscle membrane produces hyperpolarisation and reduced excitability that leads to muscle relaxation and flaccid paralysis, as shown in (Table-2 & Fig. 2) concentrated ethanol:water extract of *ipomoea carnea* (eweic) exhibited anthelmintic activity in dose dependent manner taking shortest time for paralysis (p) and death (d) with 100 mg/ml concentration. Hence ethanol:water extract of *ipomoea carnea* (eweic) in its different concentration shows anthelmintic activity.

It show shortest time of paralysis (p=28.12min) and death (d=51.04min) in 100mg/ml concentration, while the time of paralysis and death will increase in 50mg/ml concentration (p=35min & d=55.12min) and in 25mg/ml concentration (p=46.20min & d=63.15min) respectively as compare to piperazine citrate (10mg/ml) used as standard reference (p= 22.10min & d= 45.07min) and distilled water as control. The predominant effect of piperazine citrate on worm is to cause a flaccid paralysis those results in expulsion of the worm by peristalsis. Thus ethanol:water extract of *ipomoea carnea* (eweic) flowers showed significant anthelmintic activity as compare to standard reference and control (Fig. 3).

Thin Layer Chromatographic studies a large number of solvent systems were tried to achieve a good resolution. Finally, the solvents hexane: ethyl acetate: acetic acid was used. thin layer chromatographic studies of the hexane extract of selected mixed herbal powder solvent system i (hexane: acetic acid (9:1), 0 spots (a) were visible. In solvent system ii (hexane: ethyl acetate: acetic acid (5:4:1)), 1 spot detected r_f value 0.67. in solvent system iii (hexane: ethyl acetate: acetic acid (4:4:2)), 1 spot detected r_f value 0.85. in solvent system iv (hexane: ethyl acetate: acetic acid (3:6:1)), 2 spots were visible r_f values 0.40 and 0.82 (Table -3 & Fig. 4).

CONCLUSION

The present study revealed that a result of ethanol:water extract of *ipomoea carnea* (eweic) contains phytochemical constituents like flavonoids, glycosides, carbohydrates, phenols, tannins, alkaloids compounds by phytochemical investigation with respect to chemical tests and tlc chromatographic techniques. Treatment with concluded that the ethanol:water extract of *ipomoea carnea* (eweic) showed potent anthelmintic activity and was equipotent to standard anthelmintic drug. The potent anthelmintic activity could be due to presence of glycosides, flavonoids and sterols. So, from the above findings, it was concluded that ethanol:water extract of *ipomoea carnea* (eweic) posse's significant wormicidal activity property. Further carried out to isolate, purify, and

characterize the active constituents responsible for the activity of these plants and also to evaluate the exact mechanism of action.

ACKNOWLEDGEMENT

The authors are grateful to avanthi institute of pharmaceutical sciences, hyderabad and also thankful to Dr. T. Rama Rao garu, Head, Animal Ethical Committe for providing all essential laboratory facilities.

REFERENCES:

1. Rajesh SV, Rajkapoor B, Senthil Kumar R, Raju K. Effect of *Clausena Dentata* (Willd) M. Roem, Against Paracetmol Induced Hepatoprotectity in Rat. Pak J Pharm Sci **2009**;22(1):90-93.
2. Madhusudhan L. Floral Festival: A Culture of Telangana, Arts and Social Sci J **2016**;7:5.
3. Flower Festival, www. The Hindu.Com.
4. Abdul Jabbar A, Muhammad AZ, Zafar IA, Muhammad YB, Asim S. Anthelmintic Activity of *Chenopodium Album* (L.) and *Caesalpinia Crista* (L.) Against *Trichostrongylid* Nematodes of Sheep. J Ethnopharmacol **2007**;114(1):86-91.
5. Korakot A, Weerah W, Puttinan M, Prasat K, Prasert S, Ann BE, Phil JW. In Vitro Screening for Anthelmintic and Antitumour Activity of Ethnomedicinal Plants from Thailand. J Ethnopharmacol **2009**;123(3):475-482.
6. Sureka Shruti, Goyal Swapnil A, Shrivastva Kratika B, Panwar Mangal Singh B, Dave Kapilesh C and Singh Narendra. Evaluation of In-Vitro Anthelmintic Activity of Hydro Alcoholic Extract of Leaves of *Nyctanthes Arbotristis* Linn. J Global Pharm Tech **2009**;1(1):101-104.
7. Neha Shekhawat, Rekha Vijayvergia. To Study of Anthelmintic Activity of Extracts of Some Medicinal Plants. Int J Computat Sci and Math **2011**;3:183-187.
8. Sravan Kumar P, Vanam Priyanka. A Review on Phytomorphological, Phytochemical and Phyto pharmacological Profile of *Ipomoea Carnea* and Future Perspective. IAJPS **2018**;05(01):371-378.
9. Sharma A, RK. Bachheti. A Review on *Ipomoea Carnea*. Int J Pharm and Bio Sci **2013**;4(4):363-377.
10. Khatiwora E, Adsul VB, Kulkarni MM, Deshpande NR, Kashalkar RV. Spectroscopic Determination of Total Phenol and Flavonoid Contents of *Ipomoea Carnea*. Int J Chem Tech Res **2010**;2(3).
11. Sahayaraj K, Ravi C. Preliminary Phytochemistry of *Ipomea Carnea* Jacq and *Vitex Negundo* Linn. Leaves. Int J Chem Sci **2008**;6(1):1-6.
12. Brain KR, Turner TD. Wright - Scientecnica. 1st Ed. Bristol: Practical Evaluation of Phytopharmaceuticals. Eur J Medi Plants **2013**;69-380.
13. Harborne JB. A Guide to Modern Technique of Plant Analysis. London: Chapman and Hill. Phytochemical Methods **1973**;279.
14. Anthony E. Ojeh, Ese C. Adegor and Ewhre O. Lawrence. Preliminary Phytochemical Screening, Analgesic and Anti-Inflammatory Properties of *Celosia Isertii*. Eur J Medi Plants **2013**;3(3):369-380.

15. V. Satyanarayana. Preliminary Phytochemical Screening and TLC Profile of Selected Four Plants of Tirupati Hills in Chittoor District, Andhra Pradesh. J Pharmacog and Phytochem **2016**;5(2):259-264.
16. Satyavathi K, Lakshmi Kanta Kantha, Prasenjit Monda, Somnath De, Soma Jana, S. Aneela. Evaluation of Anthelmintic Activity of *Carica Papaya Latex* using *Pheritima Posthuma* Anthelmintic Activity, IJLSPS **2012**;2.
17. Neha Shekhawat. Anthelmintic Activity of Extracts of Some Medicinal Plants. Int J Computat Sci and Math **2011**;3:183-187.

How to cite this article:

P. Sravan Kumar et al. IN-VITRO ANTIHELMINTHIC ACTIVITY, PHYTOCHEMICAL SCREENING AND TLC STUDIES OF ETHANOL-WATER EXTRACTION ON *IPOMEA CARNEA* FLOWER USING IN - STATE FESTIVAL OF TELANGANA (BATHUKAMMA). J Pharm Res 2018;7(4):50-55.

Conflict of interest: The authors have declared that no conflict of interest exists.

Source of support: Nil