

ments into the marsh. Because hurricanes are so important to the inorganic sediment budget, other factors must be considered to understand how to reduce wetland losses and further their restoration. Changes in the in situ accumulation of organics, rather than the reduction of inorganic sediments arriving via overbank flooding, are implicated as a causal agent of wetland losses on this coast. This is illustrated by the fact that the soil volume occupied by organic sediments plus water in healthy saltmarsh wetlands is >90% (10) and is certainly the same or higher in wetlands of lower salinity. This organic portion plays a major role in wetland soil stability and hence in wetland ecosystem health (19).

References and Notes

- R. H. Baumann, R. E. Turner, *Environ. Geol. Water Res.* **15**, 189 (1990).
- T. A. Morton, J. C. Bernier, J. A. Barras, N. F. Ferina, *USGS Open-File Rep.* 2005-1215 (2005).
- G. W. Stone, X. Zhang, A. Sheremet, *J. Coastal Res.* **44**, 40 (2005).
- J. M. Rybczyk, D. R. Cahoon, *Estuaries* **25**, 985 (2002).
- M. L. Parsons, *J. Coastal Res.* **14**, 939 (1998).
- D. R. Cahoon *et al.*, *J. Coastal Res.* **21** (special issue), 280 (1995).
- J. P. Morgan, L. G. Nichols, M. Wright, *Morphological Effect of Hurricane Audrey on the Louisiana Coast* (contribution no. 58-3, Coastal Studies Institute, Louisiana State Univ., Baton Rouge, LA, 1958).
- J. A. Nyman, R. D. DeLaune, H. H. Roberts, W. H. Patrick Jr., *Mar. Ecol. Prog. Ser.* **96**, 269 (1992).
- Materials and methods are available as supporting material on *Science* Online.
- R. E. Turner, E. M. Swenson, C. S. Milan, in *Concepts and Controversies in Tidal Marsh Ecology*, M. Weinstein, D. A. Kreeger, Eds. (Kluwer, Dordrecht, Netherlands, 2001), pp. 583–595.
- The peak water measured by U.S. Geological Survey water-level gage 30174809020900, at Pass Manchac Turtle Cove near Ponchatoula, LA, in the western Lake Pontchartrain watershed, was about 6 cm higher during Hurricane Rita than during Hurricane Katrina, even though the hurricane path was over 300 km further away.
- Storm surge model results for Hurricanes Katrina and Rita are available at the Louisiana State University Hurricane Center Web sites (http://hurricane.lsu.edu/floodprediction/katrina/deadly_funnel1.jpg and http://hurricane.lsu.edu/floodprediction/rita24/images/adv24_SurgeTX_LA.jpg).
- G. J. Chakrapani, *Curr. Sci.* **88**, 569 (2005).
- R. H. Baumann, thesis, Louisiana State Univ., Baton Rouge, LA (1980).
- D. J. Reed, N. De Luca, A. L. Foote, *Estuaries* **20**, 301 (1997).
- G. A. Snedden, J. E. Cable, C. Swarzenski, E. Swenson, *Estuar. Coastal Shelf Sci.*, in press.
- R. E. Turner, *Estuaries* **14**, 139 (1991).
- K. W. Wheelock, thesis, Louisiana State Univ., Baton Rouge, LA (2003).
- R. E. Turner, E. M. Swenson, C. S. Milan, J. M. Lee, T. A. Oswald, *Ecol. Res.* **19**, 29 (2004).
- R. H. Kesel, *Environ. Geol. Water Res.* **11**, 271 (1988).
- C. M. Belt Jr., *Science* **189**, 681 (1975).
- Kesel (20) estimated the sediment load in the spring floods using water records from 1950 to 1983 and determined the amount of sediment that would be available from unconfined overbank flooding if the levees were not there. The estimate is actually an overestimate of the amount available, because the artificial constriction of the river channel throughout the basin raised the flood stage for the same-sized discharge event (21).
- Supported by the NSF Division of Geomorphology and Land-Use Dynamics (award EAR-061250), the National Oceanic and Atmospheric Administration Coastal Ocean Program MULTISTRESS (award no. NA16OP2670), and a Louisiana Board of Regents Fellowship (J.S.S.). We thank J. Gore for assistance in sampling from an airboat; E. Babin for assistance in finding photographs; H. Hampp, our deft, safe, and considerate helicopter pilot; and B. Sen Gupta, N. N. Rabalais, and three anonymous reviewers for constructive reviews of the manuscript.

Supporting Online Material

www.sciencemag.org/cgi/content/full/1129116/DC1
Materials and Methods
SOM Text
Fig. S1
Table S1
References and Notes

24 April 2006; accepted 1 September 2006
Published online 21 September 2006;
10.1126/science.1129116
Include this information when citing this paper.

A Combined Mitigation/Geoengineering Approach to Climate Stabilization

T. M. L. Wigley

Projected anthropogenic warming and increases in CO₂ concentration present a twofold threat, both from climate changes and from CO₂ directly through increasing the acidity of the oceans. Future climate change may be reduced through mitigation (reductions in greenhouse gas emissions) or through geoengineering. Most geoengineering approaches, however, do not address the problem of increasing ocean acidity. A combined mitigation/geoengineering strategy could remove this deficiency. Here we consider the deliberate injection of sulfate aerosol precursors into the stratosphere. This action could substantially offset future warming and provide additional time to reduce human dependence on fossil fuels and stabilize CO₂ concentrations cost-effectively at an acceptable level.

In the absence of policies to reduce the magnitude of future climate change, the globe is expected to warm by ~1° to 6°C over the 21st century (1, 2). Estimated CO₂ concentrations in 2100 lie in the range from 540 to 970 parts per million, which is sufficient to cause substantial increases in ocean acidity (3–6). Mitigation directed toward stabilizing CO₂ concentrations (7) addresses both problems but presents considerable economic and technological challenges (8, 9). Geoengineering (10–17) could help reduce the future extent of climate change due to warming but does not address the problem of ocean acidity. Mitigation is therefore necessary, but geoengineering could provide additional time to address the economic and

technological challenges faced by a mitigation-only approach.

The geoengineering strategy examined here is the injection of aerosol or aerosol precursors [such as sulfur dioxide (SO₂)] into the stratosphere to provide a negative forcing of the climate system and consequently offset part of the positive forcing due to increasing greenhouse gas concentrations (18). Volcanic eruptions provide ideal experiments that can be used to assess the effects of large anthropogenic emissions of SO₂ on stratospheric aerosols and climate. We know, for example, that the Mount Pinatubo eruption [June 1991 (19, 20)] caused detectable short-term cooling (19–21) but did not seriously disrupt the climate system. Deliberately adding aerosols or aerosol precursors to the stratosphere, so that the loading is similar to the maximum loading from the Mount Pinatubo eruption, should therefore present minimal climate risks.

Increased sulfate aerosol loading of the stratosphere may present other risks, such as through its influence on stratospheric ozone. This particular risk, however, is likely to be small. The effect of sulfate aerosols depends on the chlorine loading (22–24). With current elevated chlorine loadings, ozone loss would be enhanced. This result would delay the recovery of stratospheric ozone slightly but only until anthropogenic chlorine loadings returned to levels of the 1980s (which are expected to be reached by the late 2040s).

Figure 1 shows the projected effect of multiple sequential eruptions of Mount Pinatubo every year, every 2 years, and every 4 years. The Pinatubo eruption-associated forcing that was used had a peak annual mean value of -2.97 W/m² (20, 21). The climate simulations were carried out using an upwelling-diffusion energy balance model [Model for the Assessment of Greenhouse gas-Induced Climate Change (MAGICC) (2, 25, 26)] with a chosen climate sensitivity of 3°C equilibrium warming for a CO₂ doubling (2 × CO₂). Figure 1 suggests that a sustained stratospheric forcing of ~-3 W/m² (the average asymptotic forcing for the biennial eruption case) would be sufficient to offset much of the anthropogenic warming expected over the next century. Figure 1 also shows how rapidly the aerosol-induced cooling disappears once the injection of material into the stratosphere stops, as might become necessary should unexpected environmental damages arise.

Three cases are considered to illustrate possible options for the timing and duration of aerosol injections. In each case, the loading of the stratosphere begins in 2010 and increases linearly to

National Center for Atmospheric Research, Post Office Box 3000, Boulder, CO 80307-3000, USA. E-mail: wigley@ucar.edu

-3 W/m² over 30 years. The options depart from each other after this date (Fig. 2). These geoengineering options are complemented by three future CO₂ emissions scenarios: a central “no climate policy” scenario from the Special Report on Emissions Scenarios (SRES) (27) data set, namely the A1B scenario; an ambitious scenario known as WRE450 (7) in which CO₂ concentration stabilizes at 450 ppm (the present level is ~380 ppm); and an overshoot scenario in which CO₂ concentration rises to 530 ppm in 2080 before declining to 450 ppm. [Because an atmospheric CO₂ concentration of 450 ppm “produces both calcite and aragonite undersaturation in most of the deep ocean” (4), a level even less than this may ultimately be desirable.]

CO₂ concentrations and corresponding fossil fuel emissions for these three CO₂ scenarios are shown in Fig. 3. Emissions for the stabilization cases were calculated with the use of an inverse version of MAGICC, which accounted for climate



feedbacks on the carbon cycle. The WRE450 scenario is an archetypal mitigation-only case, stabilizing at a level that many researchers believe would avoid “dangerous anthropogenic interference” with the climate system (28). The overshoot scenario is introduced here to be considered in conjunction with the three geoengineering options. It allows much larger CO₂ emissions and a much slower departure from the A1B no-policy scenario baseline. Although the rate of decline of emissions in the mid- to late 21st century is more rapid in the overshoot scenario than in WRE450, these reductions begin 15 to 20 years later in the former scenario, allowing additional time both to phase out existing CO₂-emitting fossil fuel energy technologies and to develop and deploy energy sources that have net-zero CO₂ emissions (7–9).

Figure 4 shows global mean temperature and sea-level projections for the no-policy scenario (A1B), the mitigation-only scenario (WRE450),

and the overshoot CO₂ scenario combined with the three alternative geoengineering options (HIGH GEO, MID GEO, and LOW GEO). For the decades immediately after 2010, changes in aerosol forcing in all three GEO options occur more rapidly than forcing changes for the CO₂ scenarios, so the net effect is cooling. After 2040, the HIGH GEO-associated cooling tends to balance the warming from the overshoot CO₂ stabilization scenario, eventually leading to a slight cooling that would bring global mean temperatures back to near their preindustrial level. The MID and LOW GEO options lead to temperatures stabilizing at approximately 1° and 2°C relative to temperatures in 2000 (29). After 2100, the LOW GEO option (where injection into the stratosphere is decreased to zero by 2090) closely matches the WRE450 mitigation-only scenario but requires less-stringent emissions reductions.



The sea-level results (Fig. 4B), derived from models used in the Third Assessment Report

Fig. 1. (left) Global mean temperature response to multiple volcanic eruptions. The standard eruption used was that of Mount Pinatubo [forcing data from Ammann *et al.* (20, 21)], and eruptions were assumed to occur every 4 years (top curve), every 2 years (middle curve), or every year (bottom curve). The results shown are annual mean values plotted year by year. In the 2- and (especially) 4-year cases, the forcing varies considerably from year to year, leading to noticeable interannual temperature variations. These appear as bands of values because the abscissa scale in the graph is insufficient to resolve these rapid variations. A climate sensitivity of 3°C



equilibrium warming for 2 × CO₂ is assumed. **Fig. 2. (right)** Radiative forcing scenarios for the three geoengineering options considered. The HIGH GEO option corresponds approximately to the steady-state forcing that would result from eruptions of Mount Pinatubo every 2 years.

Fig. 3. (left) (A) CO₂ concentration projections used in the analysis together with (B) corresponding fossil fuel emissions. The overshoot scenario was used in conjunction with the three geoengineering options shown in Fig. 2. A climate sensitivity of 3°C equilibrium warming for 2 × CO₂ is assumed. CO₂ emissions results depend on the climate sensitivity because of climate feedbacks on the carbon cycle. GtC, gigatons of carbon. **Fig. 4. (right)** Global mean temperature (A) and sea-level (B) changes for the A1B scenario, the WRE450 scenario, and three scenarios combining both mitigation and geoengineering. The latter cases use the overshoot scenario (Fig. 3) and the three increasingly strong geoengineering options (Fig. 2). A climate sensitivity of 3°C equilibrium warming for 2 × CO₂ is assumed.



of the Intergovernmental Panel on Climate Change (30, 31), show the much larger inertia of this part of the climate system. The LOW GEO option and WRE450 scenario again are similar, with neither tending toward stabilization. Even the HIGH GEO option shows a continuing (but slow) rise in sea level toward the end of the study period, but the rate of rise is small, even relative to changes observed over the 20th century (30, 32).

A combined mitigation/geoengineering approach to climate stabilization has a number of advantages over either alternative used separately. A relatively modest geoengineering investment (33, 34) corresponding to the present LOW GEO option could reduce the economic and technological burden on mitigation substantially, by deferring the need for immediate or near-future cuts in CO₂ emissions. More ambitious geoengineering, when combined with mitigation, could even lead to the stabilization of global mean temperature at near present levels and reduce future sea-level rise to a rate much less than that observed over the 20th century: aspects of future change that are virtually impossible to achieve through mitigation alone.

As a guide to the amount of SO₂ required, the eruption of Mount Pinatubo injected about 10 teragrams of sulfur (TgS) into the stratosphere (35, 36), and the analysis here suggests that an annual flux of half that amount would have a substantial influence. Smaller diameter aerosols would have longer lifetimes and require still smaller injection rates (15). Five TgS/year is only ~7% of current SO₂ emissions from fossil fuel combustion (37, 38). Further analysis is required to assess (i) the technological feasibility of the suggested injections of SO₂ [or of more radiatively efficient material (34)] into the stratosphere, (ii) the economic costs of this option relative to the reduced costs of mitigation that an overshoot CO₂-stabilization pathway would allow, and (iii) the detailed effects of the proposed SO₂ injections and CO₂ concentration changes on climate [compare with (39)] and stratospheric chemistry.

References and Notes

- U. Cubasch, G. A. Meehl, in *Climate Change 2001: The Scientific Basis*, J. T. Houghton et al., Eds. (Cambridge Univ. Press, Cambridge, 2001), pp. 525–582.
- T. M. L. Wigley, S. C. B. Raper, *Science* **293**, 451 (2001).
- K. Caldeira, M. E. Wickett, *Nature* **425**, 365 (2003).
- K. Caldeira, M. E. Wickett, *J. Geophys. Res.* **110**, 10.1029/2004JC002671 (2005).
- J. C. Orr et al., *Nature* **437**, 681 (2005).
- A. J. Andersson, F. T. Mackenzie, A. Lerman, *Global Biogeochem. Cycles* **20**, 10.1029/2005GB002506 (2006).
- T. M. L. Wigley, R. Richels, J. A. Edmonds, *Nature* **379**, 240 (1996).
- M. I. Hoffert et al., *Nature* **395**, 881 (1998).
- M. I. Hoffert et al., *Science* **298**, 981 (2002).
- Geoengineering refers to engineering aimed at counteracting the undesired side effects of other human activities (11). Two classes may be distinguished: climate engineering (12–15) and carbon cycle engineering [for example, through modifying ocean alkalinity (16, 17)]. Here, the word geoengineering refers specifically to climate engineering (i.e., direct and intentional management of the climate system).
- D. W. Keith, *Annu. Rev. Energy Environ.* **25**, 245 (2000).
- S. H. Schneider, *Nature* **409**, 417 (2001).
- National Academy of Sciences, *Policy Implications of Greenhouse Warming: Mitigation, Adaptation and the Science Base* (National Academy Press, Washington, DC, 1992), pp. 433–464.
- B. P. Flannery et al., in *Engineering Response to Global Climate Change: Planning a Research and Development Agenda*, R. G. Watts, Ed. (Lewis Publishers, Boca Raton, FL, 1997), pp. 379–427.
- P. J. Crutzen, *Clim. Change* **76**, 10.1007/s10584-006-9101-y (2006).
- H. S. Khesghi, *Energy* **20**, 915 (1995).
- K. S. Lackner, *Annu. Rev. Energy Environ.* **27**, 193 (2002).
- M. I. Budyko, *Climate Changes* (Engl. transl.) (American Geophysical Union, Washington, DC, 1977).
- A. Robock, *Rev. Geophys.* **38**, 191 (2000).
- C. M. Ammann, G. A. Meehl, W. M. Washington, C. S. Zender, *Geophys. Res. Lett.* **30**, 10.1029/2003GL016875 (2003).
- T. M. L. Wigley, C. M. Ammann, B. D. Santer, S. C. B. Raper, *J. Geophys. Res.* **110**, 10.1020/2004JD005557 (2005).
- X. Tie, G. Brasseur, *Geophys. Res. Lett.* **22**, 3035 (1995).
- S. Solomon et al., *J. Geophys. Res.* **101**, 6713 (1996).
- R. W. Portmann et al., *J. Geophys. Res.* **101**, 22991 (1996).
- MAGICC is a suitable tool because it reproduces the results from more complex coupled ocean/atmosphere general circulation models for both volcanic time-scale (21) and long-term (26) forcing.
- S. C. B. Raper, J. M. Gregory, T. J. Osborn, *Clim. Dyn.* **17**, 601 (2001).
- N. Nakicenović, R. Swart, Eds., *Special Report on Emissions Scenarios* (Cambridge Univ. Press, Cambridge, 2000).
- Avoiding dangerous anthropogenic interference with the climate system is one of the primary guidelines for climate policy espoused in Article 2 of the United Nations Framework Convention on Climate Change.
- In all options, there is a residual warming tendency arising from the emissions of non-CO₂ gases (such as CH₄, N₂O, halocarbons, and tropospheric aerosols). Emissions from these sources are assumed to follow the A1B scenario to 2100 and then remain constant, leading to a slow but long-term increase in radiative forcing.
- J. A. Church, J. M. Gregory, in *Climate Change 2001: The Scientific Basis*, J. T. Houghton et al., Eds. (Cambridge Univ. Press, Cambridge, 2001), pp. 639–694.
- T. M. L. Wigley, S. C. B. Raper, *Geophys. Res. Lett.* **32**, 10.1029/2004GL021238 (2005).
- A. Cazenave, R. S. Nerem, *Rev. Geophys.* **42**, 10.1029/2003RG000139 (2004).
- Teller et al. (34) point out that sulfate aerosols are “grossly nonoptimized” as scatterers of short-wave radiation and that metallic or resonant scatterers offer large mass savings. Although emplacement costs for such scatterers would be higher, they estimate net costs (for metals) to be “as much as five times less” than for sulfate aerosols.
- E. Teller, R. Hyde, L. Wood, *Active Climate Stabilization: Practical Physics-Based Approaches to Prevention of Climate Change* (Preprint UCRL-JC-148012, Lawrence Livermore National Laboratory, Livermore, CA, 2002); www.llnl.gov/global-warm/148012.pdf.
- G. J. S. Bluth, S. D. Doiron, C. C. Schnetzler, A. J. Krueger, L. S. Walter, *Geophys. Res. Lett.* **19**, 151 (1992).
- S. Guo, G. J. S. Bluth, W. I. Rose, I. M. Watson, A. J. Prata, *Geochem. Geophys. Geosys.* **5**, 10.1029/2003GC000654 (2004).
- S. J. Smith, H. Pitcher, T. M. L. Wigley, *Global Planet. Change* **29**, 99 (2001).
- At steady-state conditions, an injection of 5 TgS/year into the stratosphere would increase the flux into the troposphere by the same amount, with larger fluxes in high latitudes. Given current emissions, the consequences of this extra flux are likely to be minor, but this aspect warrants further attention.
- B. Govindasamy, K. Caldeira, *Geophys. Res. Lett.* **17**, 2151 (2000).
- The National Center for Atmospheric Research is supported by NSF.

26 June 2006; accepted 6 September 2006
Published online 14 September 2006;
10.1126/science.1131728
Include this information when citing this paper.

Dendritic Cell Stimulation by Mycobacterial Hsp70 Is Mediated Through CCR5

R. Andres Floto,^{1*} Paul A. MacAry,^{1,2*} Jessica M. Boname,¹ Tan Suet Mien,³ Beate Kampmann,⁴ James R. Hair,¹ Oh Seen Huey,² Edith N. G. Houben,⁵ Jean Pieters,⁵ Cheryl Day,⁶ Wulf Oehlmann,⁷ Mahavir Singh,⁷ Kenneth G. C. Smith,¹ Paul J. Lehner^{1†}

An effective host immune response to mycobacterial infection must control pathogen dissemination without inducing immunopathology. Constitutive overexpression of mycobacterial heat shock protein (myHsp70) is associated with impaired bacterial persistence, but the immune-mediated mechanisms are unknown. We found that myHsp70, in addition to enhancing antigen delivery to human dendritic cells, signaled through the CCR5 chemokine receptor, promoting dendritic cell aggregation, immune synapse formation between dendritic cells and T cells, and the generation of effector immune responses. Thus, CCR5 acts as a pattern-recognition receptor for myHsp70, which may have implications for both the pathophysiology of tuberculosis and the use of myHsps in tumor-directed immunotherapy.

Mycobacterium tuberculosis infects one-third of the world's population and is responsible for two million deaths annually. A tightly controlled T cell response to *M. tuberculosis* results in granuloma formation, which limits mycobacterial replication and controls the immunopathological conse-

quences of infection (1). The immune response and granuloma formation are regulated by host and mycobacterial factors (2); one of these, mycobacterial heat shock protein 70 (myHsp70), stimulates dendritic cells (DCs) to release pro-inflammatory mediators (3). Hsps are conserved between microorganisms and mammalian cells



A Combined Mitigation/Geoengineering Approach to Climate Stabilization
T. M. L. Wigley
Science **314**, 452 (2006);
DOI: 10.1126/science.1131728

This copy is for your personal, non-commercial use only.

If you wish to distribute this article to others, you can order high-quality copies for your colleagues, clients, or customers by [clicking here](#).

Permission to republish or repurpose articles or portions of articles can be obtained by following the guidelines [here](#).

The following resources related to this article are available online at www.sciencemag.org (this information is current as of April 6, 2015):

Updated information and services, including high-resolution figures, can be found in the online version of this article at:

<http://www.sciencemag.org/content/314/5798/452.full.html>

A list of selected additional articles on the Science Web sites **related to this article** can be found at:

<http://www.sciencemag.org/content/314/5798/452.full.html#related>

This article **cites 23 articles**, 2 of which can be accessed free:

<http://www.sciencemag.org/content/314/5798/452.full.html#ref-list-1>

This article has been **cited by** 56 article(s) on the ISI Web of Science

This article has been **cited by** 22 articles hosted by HighWire Press; see:

<http://www.sciencemag.org/content/314/5798/452.full.html#related-urls>

This article appears in the following **subject collections**:

Atmospheric Science

<http://www.sciencemag.org/cgi/collection/atmos>