

Similarity measures of bipolar neutrosophic sets and their application to multiple criteria decision making

Vakkas Uluçay¹  · Irfan Deli² · Mehmet Şahin¹

Received: 5 March 2016 / Accepted: 6 July 2016
© The Natural Computing Applications Forum 2016

Abstract In this paper, we introduced some similarity measures for bipolar neutrosophic sets such as; Dice similarity measure, weighted Dice similarity measure, Hybrid vector similarity measure and weighted Hybrid vector similarity measure. Also we examine the propositions of the similarity measures. Furthermore, a multi-criteria decision-making method for bipolar neutrosophic set is developed based on these given similarity measures. Then, a practical example is shown to verify the feasibility of the new method. Finally, we compare the proposed method with the existing methods in order to demonstrate the practicality and effectiveness of the developed method in this paper.

Keywords Neutrosophic sets · Bipolar neutrosophic set · Dice similarity measure · Hybrid vector similarity · Decision making

1 Introduction

As a generalization of classical sets, fuzzy set [51] and intuitionistic fuzzy set [3], neutrosophic set was presented by Smarandache [32, 33] to capture the incomplete, indeterminate and inconsistent information. The neutrosophic set has three completely independent parts, which are truth-membership degree, indeterminacy-membership degree and falsity-membership degree; therefore, it is applied to

many different areas, such as decision-making problems [1, 2, 4, 7–11, 19, 23, 31, 37–39, 42, 52]. In additionally, since the neutrosophic sets are hard to be applied in some real problems because of the truth-membership degree, indeterminacy-membership degree and falsity-membership degree lie in $]^{-}0, 1^{+}[$, single-valued neutrosophic sets introduced by Wang et al. [40].

Recently, Lee [21, 22] proposed notation of bipolar fuzzy set and their operations based on fuzzy sets. A bipolar fuzzy set has two completely independent parts, which are positive membership degree $T^{+} \rightarrow [0, 1]$ and negative membership degree $T^{-} \rightarrow [-1, 0]$. Also the bipolar fuzzy models have been studied by many authors including theory and applications in [12, 17, 25, 34, 35, 50]. After the definition of Smarandache's neutrosophic set, neutrosophic sets and neutrosophic logic have been applied in many real applications to handle uncertainty. The neutrosophic set uses one single value in $]^{-}0, 1^{+}[$ to represent the truth-membership degree, indeterminacy-membership degree and falsity-membership degree of an element in the universe X . Then, Deli et al. [18] introduced the concept of bipolar neutrosophic sets, as an extension of neutrosophic sets. In the bipolar neutrosophic sets, the positive membership degree $T^{+}(x)$, $I^{+}(x)$, $F^{+}(x)$ denotes the truth membership, indeterminate membership and false membership of an element $x \in X$ corresponding to a bipolar neutrosophic set A and the negative membership degree $T^{-}(x)$, $I^{-}(x)$, $F^{-}(x)$ denotes the truth membership, indeterminate membership and false membership of an element $x \in X$ to some implicit counter-property corresponding to a bipolar neutrosophic set A .

Similarity measure is an important tool in constructing multi-criteria decision-making methods in many areas such

✉ Vakkas Uluçay
vulucay27@gmail.com

¹ Gaziantep University, 27310 Gaziantep, Turkey

² Department of Mathematics, Kilis 7 Aralık University, 79000 Kilis, Turkey

as medical diagnosis, pattern recognition, clustering analysis, decision making and so on. Similarity measures under all sorts of fuzzy environments including single-valued neutrosophic environments have been studied by many researchers in [5, 6, 13, 14, 15, 20–24, 26–30, 43–49]. Also, Deli and Şubas’s [16] and Şahin et al. [36] presented some similarity measures on bipolar neutrosophic sets based on correlation coefficient similarity measure and Jaccard vector similarity measure of neutrosophic set and applied to a decision-making problem, respectively. This paper is constructed as follows. In Sect. 2, some basic definitions of neutrosophic sets and bipolar neutrosophic sets are introduced. In Sect. 3, we propose some similarity measures for bipolar neutrosophic sets such as; Dice similarity measure, weighted Dice similarity measure, Hybrid vector similarity measure and weighted Hybrid vector similarity measure and investigate their several properties. In Sect. 4, a multi-criteria decision-making method for bipolar neutrosophic set is developed based on these given similarity measures and a practical example is given. In Sect. 5, we compare the proposed method with the existing methods in order to demonstrate the practicality and effectiveness of the developed method in this paper. In Sect. 6, the conclusions are summarized.

2 Preliminary

In the section, we give some concepts related to neutrosophic sets and bipolar neutrosophic sets.

Definition 1 [32] Let E be a universe. A neutrosophic set A over E is defined by

$$A = \{ \langle x, (T_A(x), I_A(x), F_A(x)) \rangle : x \in E \}.$$

where $T_A(x)$, $I_A(x)$ and $F_A(x)$ are called truth-membership function, indeterminacy-membership function and falsity-membership function, respectively. They are, respectively, defined by

$$T_A : E \rightarrow]^{-}0, 1^{+}[, \quad I_A : E \rightarrow]^{-}0, 1^{+}[, \quad F_A : E \rightarrow]^{-}0, 1^{+}[$$

such that $0^{-} \leq T_A(x) + I_A(x) + F_A(x) \leq 3^{+}$.

Definition 2 [40] Let E be a universe. A single-valued neutrosophic set (SVN-set) A over E is defined by

$$A = \{ \langle x, (T_A(x), I_A(x), F_A(x)) \rangle : x \in E \}.$$

where $T_A(x)$, $I_A(x)$ and $F_A(x)$ are called truth-membership function, indeterminacy-membership function and falsity-

membership function, respectively. They are respectively defined by

$$T_A : E \rightarrow [0, 1], \quad I_A : E \rightarrow [0, 1], \quad F_A : E \rightarrow [0, 1]$$

such that $0 \leq T_A(x) + I_A(x) + F_A(x) \leq 3$.

Definition 3 [18] A bipolar neutrosophic set (BNS) A in X is defined as an object of the form

$$A = \{ \langle x, T^{+}(x), I^{+}(x), F^{+}(x), T^{-}(x), I^{-}(x), F^{-}(x) \rangle : x \in X \}.$$

where

$$T^{+}, I^{+}, F^{+} : E \rightarrow [0, 1], \quad T^{-}, I^{-}, F^{-} : X \rightarrow [-1, 0].$$

The positive membership degree $T^{+}(x)$, $I^{+}(x)$, $F^{+}(x)$ denotes the truth membership, indeterminate membership and false membership of an element $x \in X$ corresponding to a bipolar neutrosophic set A and the negative membership degree $T^{-}(x)$, $I^{-}(x)$, $F^{-}(x)$ denotes the truth membership, indeterminate membership and false membership of an element $x \in X$ to some implicit counter-property corresponding to a bipolar neutrosophic set A .

Definition 4 [18] Let $A_1 = \langle x, T_1^{+}(x), I_1^{+}(x), F_1^{+}(x), T_1^{-}(x), I_1^{-}(x), F_1^{-}(x) \rangle$ and $A_2 = \langle x, T_2^{+}(x), I_2^{+}(x), F_2^{+}(x), T_2^{-}(x), I_2^{-}(x), F_2^{-}(x) \rangle$ be two BNSs in a universe of discourse X . Then the following operations are defined as follows:

1. $A_1 = A_2$ if and only if $T_1^{+}(x) = T_2^{+}(x), I_1^{+}(x) = I_2^{+}(x), F_1^{+}(x) = F_2^{+}(x)$ and $T_1^{-}(x) = T_2^{-}(x), I_1^{-}(x) = I_2^{-}(x), F_1^{-}(x) = F_2^{-}(x)$.
2. $A_1 \cup A_2 = \{ \langle x, \max(T_1^{+}(x), T_2^{+}(x)), \frac{I_1^{+}(x)+I_2^{+}(x)}{2}, \min(F_1^{+}(x), F_2^{+}(x)), \min(T_1^{-}(x), T_2^{-}(x)), \frac{I_1^{-}(x)+I_2^{-}(x)}{2}, \max(F_1^{-}(x), F_2^{-}(x)) \rangle \forall x \in X$.
3. $A_1 \cap A_2 = \{ \langle x, \min(T_1^{+}(x), T_2^{+}(x)), \frac{I_1^{+}(x)+I_2^{+}(x)}{2}, \max(F_1^{+}(x), F_2^{+}(x)), \max(T_1^{-}(x), T_2^{-}(x)), \frac{I_1^{-}(x)+I_2^{-}(x)}{2}, \min(F_1^{-}(x), F_2^{-}(x)) \rangle \forall x \in X$.
4. $A^c = \{ \langle x, 1 - T_A^{+}(x), 1 - I_A^{+}(x), 1 - F_A^{+}(x), 1 - T_A^{-}(x), 1 - I_A^{-}(x), 1 - F_A^{-}(x) \rangle \}$
5. $A_1 \subseteq A_2$ if and only if $T_1^{+}(x) \leq T_2^{+}(x), I_1^{+}(x) \leq I_2^{+}(x), F_1^{+}(x) \geq F_2^{+}(x)$ and $T_1^{-}(x) \geq T_2^{-}(x), I_1^{-}(x) \geq I_2^{-}(x), F_1^{-}(x) \leq F_2^{-}(x)$.

Definition 5 [45] Let $A = \langle T_A(x_i), I_A(x_i), F_A(x_i) \rangle$ and $B = \langle T_B(x_i), I_B(x_i), F_B(x_i) \rangle$ be two SVNNSs in a universe of discourse $X = (x_1, x_2, \dots, x_n)$. Then Dice similarity measure between SVNNSs A and B in the vector space is defined as follows:

$$D(A, B) = \frac{1}{n} \sum_{i=1}^n \times \left(\frac{2(T_A(x_i)T_B(x_i) + I_A(x_i)I_B(x_i) + F_A(x_i)F_B(x_i))}{[((T_A)^2(x_i) + (I_A)^2(x_i) + (F_A)^2(x_i)) + ((T_B)^2(x_i) + (I_B)^2(x_i) + (F_B)^2(x_i))]} \right)$$

It satisfies the following properties:

1. $0 \leq D(A, B) \leq 1$;
2. $D(A, B) = D(B, A)$;
3. $D(A, B) = 1$ for $A = B$ i.e. $T_A(x_i) = T_B(x_i), I_A(x_i) = I_B(x_i), F_A(x_i) = F_B(x_i) \quad (i = 1, 2, \dots, n) \quad \forall x_i (i = 1, 2, \dots, n) \in X$.

Definition 6 [29] Let $A = \langle T_A(x_i), I_A(x_i), F_A(x_i) \rangle$ and $B = \langle T_B(x_i), I_B(x_i), F_B(x_i) \rangle$ be two SVN-sets in a universe of discourse $X = \{x_1, x_2, \dots, x_n\}$. Then

$$E(X, Y) = \lambda \left(\frac{2(T_A(x_i)T_B(x_i) + I_A(x_i)I_B(x_i) + F_A(x_i)F_B(x_i))}{[((T_A)^2(x_i) + (I_A)^2(x_i) + (F_A)^2(x_i)) + ((T_B)^2(x_i) + (I_B)^2(x_i) + (F_B)^2(x_i))]} \right) + (1 - \lambda) \left(\frac{T_A(x_i)T_B(x_i) + I_A(x_i)I_B(x_i) + F_A(x_i)F_B(x_i)}{[\sqrt{((T_A)^2(x_i) + (I_A)^2(x_i) + (F_A)^2(x_i))} \cdot \sqrt{((T_B)^2(x_i) + (I_B)^2(x_i) + (F_B)^2(x_i))}]} \right)$$

It satisfies the following properties:

1. $0 \leq D(A, B) \leq 1$;
2. $D(A, B) = D(B, A)$;
3. $D(A, B) = 1$ for $A = B$ i.e.

3 Similarity measures of bipolar neutrosophic sets

In this section, we introduce some similarity measures for bipolar neutrosophic sets including Dice similarity measure, weighted Dice similarity measure, Hybrid vector similarity measure and weighted Hybrid vector similarity measure by extending the studies in [29, 45].

Definition 7 Let $A = \langle T_A^+(x_i), I_A^+(x_i), F_A^+(x_i), T_A^-(x_i), I_A^-(x_i), F_A^-(x_i) \rangle$ and $B = \langle T_B^+(x_i), I_B^+(x_i), F_B^+(x_i), T_B^-(x_i), I_B^-(x_i), F_B^-(x_i) \rangle$

$I_B^-(x_i), F_B^-(x_i) \rangle$ be two BNSs in the set $X = \{x_1, x_2, \dots, x_n\}$. Then, Dice similarity measure between BNS A and B, denoted $D(A, B)$, is defined as;

$$D(A, B) = \frac{1}{n} \sum_{i=1}^n \times \left(\frac{[(T_A^+(x_i)T_B^+(x_i) + I_A^+(x_i)I_B^+(x_i) + F_A^+(x_i)F_B^+(x_i)) - (T_A^-(x_i)T_B^-(x_i) + I_A^-(x_i)I_B^-(x_i) + F_A^-(x_i)F_B^-(x_i))]}{[((T_A^+)^2(x_i) + (I_A^+)^2(x_i) + (F_A^+)^2(x_i)) + ((T_B^+)^2(x_i) + (I_B^+)^2(x_i) + (F_B^+)^2(x_i)) - ((T_A^-)^2(x_i) + (I_A^-)^2(x_i) + (F_A^-)^2(x_i)) - ((T_B^-)^2(x_i) + (I_B^-)^2(x_i) + (F_B^-)^2(x_i))]} \right)$$

Example 1 Suppose that $A = \langle 0.5, 0.2, 0.6, -0.4, -0.3, -0.7 \rangle$, $B = \langle 0.7, 0.1, 0.3, -0.2, -0.2, -0.1 \rangle$ be two BNSs in the set. Then,

1. $0 \leq D_w(A, B) \leq 1$;
2. $D_w(A, B) = D_w(B, A)$;

$$D(A, B) = \frac{1}{6} \times \left(\frac{[(0.5)(0.7) + (0.2)(0.1) + (0.6)(0.3)] - [(-0.4)(-0.2) + (-0.3)(-0.2) + (-0.7)(-0.1)]}{[(0.5)^2 + (0.2)^2 + (0.6)^2] + [(0.7)^2 + (0.1)^2 + (0.3)^2] - [(-0.4)^2 + (-0.3)^2 + (-0.7)^2] - [(-0.2)^2 + (-0.2)^2 + (-0.1)^2]} \right) = 0.1382$$

Definition 8 Let $A = \langle T_A^+(x_i), I_A^+(x_i), F_A^+(x_i), T_A^-(x_i), I_A^-(x_i), F_A^-(x_i) \rangle$ and $B = \langle T_B^+(x_i), I_B^+(x_i), F_B^+(x_i), T_B^-(x_i), I_B^-(x_i), F_B^-(x_i) \rangle$ be two BNSs in the set $X = \{x_1, x_2, \dots, x_n\}$ and $w_i \in [0, 1]$ be the weight of each element x_i for $i = 1, 2, \dots, n$ such that $\sum_{i=1}^n w_i = 1$. Then, weighted Dice similarity measure between BNS A and B, denoted $D_w(A, B)$, is defined as;

3. $D_w(A, B) = 1$ for $A = B$ i.e., $T_A^+(x_i) = T_B^+(x_i), I_A^+(x_i) = I_B^+(x_i), F_A^+(x_i) = F_B^+(x_i), T_A^-(x_i) = T_B^-(x_i), I_A^-(x_i) = I_B^-(x_i), F_A^-(x_i) = F_B^-(x_i)$ ($i = 1, 2, \dots, n$) $\forall x_i$ ($i = 1, 2, \dots, n$) $\in X$.

Proof

1. It is clear from Definition 8.

$$D_w(A, B) = \sum_{i=1}^n w_i \times \left(\frac{[(T_A^+(x_i)T_B^+(x_i) + I_A^+(x_i)I_B^+(x_i) + F_A^+(x_i)F_B^+(x_i)) - (T_A^-(x_i)T_B^-(x_i) + I_A^-(x_i)I_B^-(x_i) + F_A^-(x_i)F_B^-(x_i))]}{[(T_A^+)^2(x_i) + (I_A^+)^2(x_i) + (F_A^+)^2(x_i)] + [(T_B^+)^2(x_i) + (I_B^+)^2(x_i) + (F_B^+)^2(x_i)] - [(T_A^-)^2(x_i) + (I_A^-)^2(x_i) + (F_A^-)^2(x_i)] - [(T_B^-)^2(x_i) + (I_B^-)^2(x_i) + (F_B^-)^2(x_i)]} \right).$$

Example 2 Suppose that $A = \langle 0.8, 0.2, 0.5, -0.2, -0.3, -0.4 \rangle$, $B = \langle 0.6, 0.4, 0.3, -0.1, -0.2, -0.3 \rangle$ be two BNSs in the set and $w = 0.3$. Then,

$$D_w(A, B) = 0.3 \times \left(\frac{[(0.8)(0.6) + (0.2)(0.4) + (0.5)(0.3)] - [(-0.2)(-0.1) + (-0.3)(-0.2) + (-0.4)(-0.3)]}{[(0.8)^2 + (0.2)^2 + (0.5)^2] + [(0.6)^2 + (0.4)^2 + (0.3)^2] - [(-0.2)^2 + (-0.3)^2 + (-0.4)^2] - [(-0.1)^2 + (-0.2)^2 + (-0.3)^2]} \right) = 0.1378$$

Proposition 1 Let $D_w(A, B)$ be a weighted Dice similarity measure between BNSs A and B. Then, we have

2.

$$\begin{aligned}
 D_w(A, B) &= \sum_{i=1}^n w_i \\
 &\times \left(\frac{[(T_A^+(x_i)T_B^+(x_i) + I_A^+(x_i)I_B^+(x_i) + F_A^+(x_i)F_B^+(x_i)) - (T_A^-(x_i)T_B^-(x_i) + I_A^-(x_i)I_B^-(x_i) + F_A^-(x_i)F_B^-(x_i))]}{2[(T_A^+)^2(x_i) + (I_A^+)^2(x_i) + (F_A^+)^2(x_i)] + [(T_B^+)^2(x_i) + (I_B^+)^2(x_i) + (F_B^+)^2(x_i)]} \right. \\
 &\quad \left. - [(T_A^-)^2(x_i) + (I_A^-)^2(x_i) + (F_A^-)^2(x_i)] - [(T_B^-)^2(x_i) + (I_B^-)^2(x_i) + (F_B^-)^2(x_i)] \right) \\
 &= \sum_{i=1}^n w_i \\
 &\times \left(\frac{[(T_B^+(x_i)T_A^+(x_i) + I_B^+(x_i)I_A^+(x_i) + F_B^+(x_i)F_A^+(x_i)) - (T_B^-(x_i)T_A^-(x_i) + I_B^-(x_i)I_A^-(x_i) + F_B^-(x_i)F_A^-(x_i))]}{2[(T_B^+)^2(x_i) + (I_B^+)^2(x_i) + (F_B^+)^2(x_i)] + [(T_A^+)^2(x_i) + (I_A^+)^2(x_i) + (F_A^+)^2(x_i)]} \right. \\
 &\quad \left. - [(T_B^-)^2(x_i) + (I_B^-)^2(x_i) + (F_B^-)^2(x_i)] - [(T_A^-)^2(x_i) + (I_A^-)^2(x_i) + (F_A^-)^2(x_i)] \right) \\
 &= D_w(B, A)
 \end{aligned}$$

3. Since $T_A^+(x_i) = T_B^+(x_i), I_A^+(x_i) = I_B^+(x_i), F_A^+(x_i) = F_B^+(x_i), T_A^-(x_i) = T_B^-(x_i), I_A^-(x_i) = I_B^-(x_i), F_A^-(x_i) = F_B^-(x_i)$ ($i = 1, 2, \dots, n$) $\forall x_i$ ($i = 1, 2, \dots, n$) $\in X$, we have $D_w(A, B) = 1$.

The proof is completed. \square

Definition 9 Let $A = \langle T_A^+(x_i), I_A^+(x_i), F_A^+(x_i), T_A^-(x_i), I_A^-(x_i), F_A^-(x_i) \rangle$ and $B = \langle T_B^+(x_i), I_B^+(x_i), F_B^+(x_i), T_B^-(x_i), I_B^-(x_i), F_B^-(x_i) \rangle$ be two BNSs in the set $X = \{x_1, x_2, \dots, x_n\}$. Then, hybrid vector similarity measure between BNS A and B, denoted $HybV(A, B)$, is defined as;

$$\begin{aligned}
 HybV(A, B) &= \lambda \frac{1}{n} \sum_{i=1}^n \\
 &\times \left(\frac{[(T_A^+(x_i)T_B^+(x_i) + I_A^+(x_i)I_B^+(x_i) + F_A^+(x_i)F_B^+(x_i)) - (T_A^-(x_i)T_B^-(x_i) + I_A^-(x_i)I_B^-(x_i) + F_A^-(x_i)F_B^-(x_i))]}{2[(T_A^+)^2(x_i) + (I_A^+)^2(x_i) + (F_A^+)^2(x_i)] + [(T_B^+)^2(x_i) + (I_B^+)^2(x_i) + (F_B^+)^2(x_i)]} \right. \\
 &\quad \left. - [(T_A^-)^2(x_i) + (I_A^-)^2(x_i) + (F_A^-)^2(x_i)] - [(T_B^-)^2(x_i) + (I_B^-)^2(x_i) + (F_B^-)^2(x_i)] \right) \\
 &+ (1 - \lambda) \frac{1}{n} \sum_{i=1}^n \\
 &\times \left(\frac{[(T_A^+(x_i)T_B^+(x_i) + I_A^+(x_i)I_B^+(x_i) + F_A^+(x_i)F_B^+(x_i)) - (T_A^-(x_i)T_B^-(x_i) + I_A^-(x_i)I_B^-(x_i) + F_A^-(x_i)F_B^-(x_i))]}{2[\sqrt{(T_A^+)^2(x_i) + (I_A^+)^2(x_i) + (F_A^+)^2(x_i)} \times \sqrt{(T_B^+)^2(x_i) + (I_B^+)^2(x_i) + (F_B^+)^2(x_i)}]} \right. \\
 &\quad \left. - \sqrt{(T_A^-)^2(x_i) + (I_A^-)^2(x_i) + (F_A^-)^2(x_i)} \times \sqrt{(T_B^-)^2(x_i) + (I_B^-)^2(x_i) + (F_B^-)^2(x_i)} \right)
 \end{aligned}$$

Example 3 Suppose that $A = \langle 0.6, 0.3, 0.4, -0.1, -0.3, -0.4 \rangle$, $B = \langle 0.5, 0.2, 0.3, -0.4, -0.2, -0.5 \rangle$ be two BNSs in the set. Then,

3. $HybV_w(A, B) = 1$ for $A = B$ i.e., $T_A^+(x_i) = T_B^+(x_i)$, $I_A^+(x_i) = I_B^+(x_i)$, $F_A^+(x_i) = F_B^+(x_i)$, $T_A^-(x_i) = T_B^-(x_i)$, $I_A^-(x_i) = I_B^-(x_i)$, $F_A^-(x_i) = F_B^-(x_i)$ ($i = 1, 2, \dots, n$) $\forall x_i$ ($i = 1, 2, \dots, n$) $\in X$.

$$HybV(A, B) = (0.4) \times \frac{1}{6} \times \left(\frac{[(0.6)(0.5) + (0.3)(0.2) + (0.4)(0.3)] - [(-0.1)(-0.4) + (-0.3)(-0.2) + (-0.4)(-0.5)]}{2[(0.6)^2 + (0.3)^2 + (0.4)^2 + ((0.5)^2 + (0.2)^2 + (0.3)^2) - ((-0.1)^2 + (-0.3)^2 + (-0.4)^2) - ((-0.4)^2 + (-0.2)^2 + (-0.5)^2]} \right) + (1 - (0.4)) \times \frac{1}{6} \times \left(\frac{[(0.6)(0.5) + (0.3)(0.2) + (0.4)(0.3)] - [(-0.1)(-0.4) + (-0.3)(-0.2) + (-0.4)(-0.5)]}{2[\sqrt{((0.6)^2 + (0.3)^2 + (0.4)^2)} \times \sqrt{((0.5)^2 + (0.2)^2 + (0.3)^2)} - \sqrt{((-0.1)^2 + (-0.3)^2 + (-0.4)^2)} \times \sqrt{((-0.4)^2 + (-0.2)^2 + (-0.5)^2)}]} \right) = 0.1421$$

Definition 10 Let $A = \langle T_A^+(x_i), I_A^+(x_i), F_A^+(x_i), T_A^-(x_i), I_A^-(x_i), F_A^-(x_i) \rangle$ and $B = \langle T_B^+(x_i), I_B^+(x_i), F_B^+(x_i), T_B^-(x_i), I_B^-(x_i), F_B^-(x_i) \rangle$ be two BNSs in the set $X = \{x_1, x_2, \dots, x_n\}$ and $w_i \in [0, 1]$ be the weight of each element x_i for $i = 1, 2, \dots, n$ such that $\sum_{i=1}^n w_i = 1$. Then, weighted hybrid vector similarity measure between BNS A and B, denoted $HybV_w(A, B)$, is defined as;

4 BN-multi-criteria decision-making methods

In this section, we introduce applications of weighted similarity measures in multi-criteria decision making problems under bipolar neutrosophic environment.

Definition 11 [36] Let $U = (u_1, u_2, \dots, u_n)$ be a set of alternatives, $A = (a_1, a_2, \dots, a_m)$ be the set of attributes,

$$HybV_w(A, B) = \lambda \sum_{i=1}^n w_i \times \left(\frac{[(T_A^+(x_i)T_B^+(x_i) + I_A^+(x_i)I_B^+(x_i) + F_A^+(x_i)F_B^+(x_i)) - (T_A^-(x_i)T_B^-(x_i) + I_A^-(x_i)I_B^-(x_i) + F_A^-(x_i)F_B^-(x_i))]}{2[((T_A^+)^2(x_i) + (I_A^+)^2(x_i) + (F_A^+)^2(x_i)) + ((T_B^+)^2(x_i) + (I_B^+)^2(x_i) + (F_B^+)^2(x_i)) - ((T_A^-)^2(x_i) + (I_A^-)^2(x_i) + (F_A^-)^2(x_i)) - ((T_B^-)^2(x_i) + (I_B^-)^2(x_i) + (F_B^-)^2(x_i))]} \right) + (1 - \lambda) \sum_{i=1}^n w_i \times \left(\frac{[(T_A^+(x_i)T_B^+(x_i) + I_A^+(x_i)I_B^+(x_i) + F_A^+(x_i)F_B^+(x_i)) - (T_A^-(x_i)T_B^-(x_i) + I_A^-(x_i)I_B^-(x_i) + F_A^-(x_i)F_B^-(x_i))]}{2[\sqrt{(T_A^+)^2(x_i) + (I_A^+)^2(x_i) + (F_A^+)^2(x_i)} \times \sqrt{(T_B^+)^2(x_i) + (I_B^+)^2(x_i) + (F_B^+)^2(x_i)} - \sqrt{(T_A^-)^2(x_i) + (I_A^-)^2(x_i) + (F_A^-)^2(x_i)} \times \sqrt{(T_B^-)^2(x_i) + (I_B^-)^2(x_i) + (F_B^-)^2(x_i)}]} \right)$$

Proposition 2 Let $HybV_w(A, B)$ be a weighted hybrid vector similarity measure between bipolar neutrosophic sets A and B. Then, we have

1. $0 \leq HybV_w(A, B) \leq 1$;
2. $HybV_w(A, B) = HybV_w(B, A)$;

$w = (w_1, w_2, \dots, w_n)^T$ be the weight vector of the attributes C_j ($j = 1, 2, \dots, n$) such that $w_j \geq 0$ and $\sum_{j=1}^n w_j = 1$ and $b_{ij} = \langle T_{ij}^+, I_{ij}^+, F_{ij}^+, T_{ij}^-, I_{ij}^-, F_{ij}^- \rangle$ be the decision matrix whose entries are the rating values of the alternatives. Then,

$$[b_{ij}]_{m \times n} = \begin{matrix} & a_1 & a_2 & \cdots & a_n \\ u_1 & \begin{pmatrix} b_{11} & b_{12} & \cdots & b_{1n} \\ b_{21} & b_{22} & \cdots & b_{2n} \\ \vdots & \vdots & \vdots & \vdots \\ b_{m1} & b_{m2} & \cdots & b_{mn} \end{pmatrix} \\ u_2 & \\ \vdots & \\ u_m & \end{matrix}$$

is called an NB-multi-attribute decision-making matrix of the decision maker.

Algorithm

- Step 1 Give the decision-making matrix $[b_{ij}]_{m \times n}$; for decision;
- Step 2 Compute the positive ideal (or negative ideal) bipolar neutrosophic solution for $[b_{ij}]_{m \times n}$;
- Step 3 Calculate the weighted hybrid vector (or Dice) similarity measure between positive ideal (or negative ideal) bipolar neutrosophic solution b_j^* and $b_i = [b_{ij}]_{1 \times n}$ for all $i = 1, 2, \dots, m$ and $j = 1, 2, \dots, n$ as;

$$HybV_w(b_j^*, b_i) = \lambda \sum_{j=1}^n w_j \times \left(\frac{[(T_j^*)^+(T_{ij}^*)^+ + (I_j^*)^+(I_{ij}^*)^+ + (F_j^*)^+(F_{ij}^*)^+ - ((T_j^*)^-(T_{ij}^*)^- + (I_j^*)^-(I_{ij}^*)^- + (F_j^*)^-(F_{ij}^*)^-)]}{2[(((T_j^*)^+)^2 + ((I_j^*)^+)^2 + ((F_j^*)^+)^2) + ((T_{ij}^*)^+)^2 + (I_{ij}^+)^2 + (F_{ij}^+)^2] - (((T_j^*)^-)^2 + ((I_j^*)^-)^2 + ((F_j^*)^-)^2) - ((T_{ij}^-)^2 + (I_{ij}^-)^2 + (F_{ij}^-)^2)} \right) + (1 - \lambda) \sum_{j=1}^n w_j \times \left(\frac{[(T_j^*)^+(T_{ij}^*)^+ + (I_j^*)^+(I_{ij}^*)^+ + (F_j^*)^+(F_{ij}^*)^+ - (T_j^*)^-(T_{ij}^*)^- + (I_j^*)^-(I_{ij}^*)^- + (F_j^*)^-(F_{ij}^*)^-]}{2[\sqrt{((T_j^*)^+)^2 + ((I_j^*)^+)^2 + ((F_j^*)^+)^2} \times \sqrt{(T_{ij}^+)^2 + (I_{ij}^+)^2 + (F_{ij}^+)^2} - \sqrt{((T_j^*)^-)^2 + ((I_j^*)^-)^2 + ((F_j^*)^-)^2} \times \sqrt{(T_{ij}^-)^2 + (I_{ij}^-)^2 + (F_{ij}^-)^2}]} \right) \tag{3}$$

Also; positive ideal bipolar neutrosophic solution $u^* = (b_1^*, b_2^*, \dots, b_n^*)$ is the solution of decision matrix $[b_{ij}]_{m \times n}$ where every component of has the following form:

$$b_j^* = \langle \max_i\{T_{ij}^+\}, \min_i\{I_{ij}^+\}, \min_i\{F_{ij}^+\}, \min_i\{T_{ij}^-\}, \max_i\{I_{ij}^-\}, \max_i\{F_{ij}^-\} \rangle (j = 1, 2, \dots, n) \tag{1}$$

and negative ideal bipolar neutrosophic solution $\bar{u}^* = (\bar{b}_1^*, \bar{b}_2^*, \dots, \bar{b}_n^*)$ is the solution of decision matrix $[b_{ij}]_{m \times n}$ where every component of has the following form:

$$\bar{b}_j^* = \langle \min_i\{T_{ij}^+\}, \max_i\{I_{ij}^+\}, \max_i\{F_{ij}^+\}, \max_i\{T_{ij}^-\}, \min_i\{I_{ij}^-\}, \min_i\{F_{ij}^-\} \rangle \tag{2}$$

Now we give an algorithm as;

- Step 4. Determine the nonincreasing order of $s_i = HybV_w(b_j^*, b_i)$ and select the best alternative.

Example 4 Let us consider the decision-making problem given in [41] for bipolar neutrosophic set. “Global environmental concern is a reality, and an increasing attention is focusing on the green production in various industries. A car company is desirable to select the most appropriate green supplier for one of the key elements in its manufacturing process.” [41]. After pre-evaluation, four suppliers(alternatives) are taken into consideration, which are denoted by u_1, u_2, u_3 and u_4 . Three criteria are considered including a_1 is product quality; a_2 is technology capability; a_3 is pollution control. Assume the weight vector of the

three criteria is $W = \{w_1, w_2, w_3\}^T = \{0.2, 0.5, 0.3\}^T$. In order to determine the decision information, an expert has gathered the criteria values for the four possible alternatives under bipolar neutrosophic environment. Then

Algorithm

Step 1 The decision-making matrix $[b_{ij}]_{m \times n}$ is given by a expert as;

$$\begin{matrix}
 & a_1 & a_2 & a_3 \\
 u_1 & \langle 0.4, 0.5, 0.3, -0.6, -0.4, -0.5 \rangle & \langle 0.6, 0.1, 0.2, -0.4, -0.3, -0.2 \rangle & \langle 0.8, 0.6, 0.5, -0.3, -0.2, -0.1 \rangle \\
 u_2 & \langle 0.6, 0.4, 0.2, -0.4, -0.5, -0.7 \rangle & \langle 0.6, 0.2, 0.3, -0.5, -0.2, -0.3 \rangle & \langle 0.7, 0.4, 0.5, -0.1, -0.3, -0.4 \rangle \\
 u_3 & \langle 0.7, 0.2, 0.4, -0.2, -0.6, -0.4 \rangle & \langle 0.9, 0.3, 0.6, -0.2, -0.2, -0.5 \rangle & \langle 0.6, 0.1, 0.5, -0.2, -0.4, -0.6 \rangle \\
 u_4 & \langle 0.8, 0.6, 0.5, -0.5, -0.3, -0.6 \rangle & \langle 0.6, 0.4, 0.3, -0.1, -0.3, -0.4 \rangle & \langle 0.9, 0.6, 0.4, -0.5, -0.3, -0.6 \rangle
 \end{matrix}$$

Step 2 The positive ideal bipolar neutrosophic solutions for are computed as; $u^* = [\langle 0.8, 0.2, 0.2, -0.6, -0.3, -0.4 \rangle, \langle 0.9, 0.1, 0.2, -0.5, -0.2, -0.2 \rangle, \langle 0.9, 0.1, 0.4, -0.5, -0.2, -0.1 \rangle]$.

Step 3 The weighted hybrid vector similarity measure between positive ideal (or negative ideal) bipolar neutrosophic solution for alternative $u_j \in U$ are computed and selected the best alternative.

Step 4 Ranking the alternatives (Table 1).

5 Comparative analysis and discussion

In this subsection, a comparative study is presented to show the flexibility and feasibility of the introduced NB-multi-attribute decision-making method. Therefore, different methods are used to solve the same NB-multi-attribute

decision-making problem with the bipolar neutrosophic information given by Şahin et al. [36], Deli et al. [18], Deli and Şubaş [16]. Şahin et al. [36] present a method by using Jaccard vector similarity and weighted Jaccard vector similarity measure and Deli and Şubaş [16] present a method by using correlation measure based on multi-criteria decision making for bipolar neutrosophic sets. Also,

Deli et al. [18] contains two major phrases. The method firstly use score, certainty and accuracy functions to compare the bipolar neutrosophic sets. Secondly, he use bipolar neutrosophic weighted average operator and bipolar neutrosophic weighted geometric operator to aggregate the bipolar neutrosophic information. The ranking results obtained by different methods are summarized in Table 2.

In Table 2, there are some differences between the ranking results obtained by the methods. The optimal alternative is u_3 obtained by the proposed methods except the result obtained by the method of Deli et al.'s method [18] and the proposed method $HybV_w$ with $\lambda = 0.9$. The reason may be score, certainty and accuracy functions and weighted average operator and bipolar neutrosophic weighted geometric operator in Deli et al.'s method [18] and parameter λ in the proposed method $HybV_w$. Generally, the proposed methods can effectively overcome the

Table 1 Results for different values of λ

Similarity measure	Values	Measure value	Ranking order
$HybV_w(u^*, u_i)$	$\lambda = 0.25$	$HybV_w(u^*, u_1) = 0.24683$	$u_3 \succ u_1 \succ u_4 \succ u_2$
		$HybV_w(u^*, u_2) = 0.117780$	
		$HybV_w(u^*, u_3) = 0.27833$	
		$HybV_w(u^*, u_4) = 0.21136$	
$HybV_w(u^*, u_i)$	$\lambda = 0.3$	$HybV_w(u^*, u_1) = 0.27063$	$u_3 \succ u_1 \succ u_4 \succ u_2$
		$HybV_w(u^*, u_2) = 0.19497$	
		$HybV_w(u^*, u_3) = 0.30222$	
		$HybV_w(u^*, u_4) = 0.22904$	
$HybV_w(u^*, u_i)$	$\lambda = 0.6$	$HybV_w(u^*, u_1) = 0.41342$	$u_3 \succ u_1 \succ u_4 \succ u_2$
		$HybV_w(u^*, u_2) = 0.29803$	
		$HybV_w(u^*, u_3) = 0.44555$	
		$HybV_w(u^*, u_4) = 0.33510$	
$HybV_w(u^*, u_i)$	$\lambda = 0.9$	$HybV_w(u^*, u_1) = 0.55620$	$u_1 \succ u_3 \succ u_4 \succ u_2$
		$HybV_w(u^*, u_2) = 0.40109$	
		$HybV_w(u^*, u_3) = 0.54313$	
		$HybV_w(u^*, u_4) = 0.44116$	

Table 2 The ranking results of different methods

Methods		Ranking results
The proposed method $HybV_w$ with $\lambda = 0.25$		$u_3 \succ u_1 \succ u_4 \succ u_2$
The proposed method $HybV_w$ with $\lambda = 0.3$		$u_3 \succ u_1 \succ u_4 \succ u_2$
The proposed method $HybV_w$ with $\lambda = 0.6$		$u_3 \succ u_1 \succ u_4 \succ u_2$
The proposed method $HybV_w$ with $\lambda = 0.9$		$u_1 \succ u_3 \succ u_4 \succ u_2$
The proposed method $D_w(u^*, u_i)$		$u_3 \succ u_1 \succ u_4 \succ u_2$
Deli and Şubas's method [16]		$u_3 \succ u_1 \succ u_4 \succ u_2$
Deli et al.'s method [18]		$u_2 \succ u_3 \succ u_4 \succ u_1$
Şahin et al.'s method [36]		$u_3 \succ u_4 \succ u_2 \succ u_1$

decision-making problems which contain bipolar neutrosophic information. So, we think the proposed methods developed in this paper is more suitable to handle this application example.

6 Conclusion

This paper developed a multi-criteria decision-making method for bipolar neutrosophic set is developed based on these given similarity measures. To get the comprehensive values, some similarity measures for bipolar neutrosophic sets such as; Dice similarity measure, weighted Dice similarity measure, Hybrid vector similarity measure and weighted Hybrid vector similarity measure are introduced. In the future, it shall be significant to research some special kinds of bipolar neutrosophic measures.

References

1. Ali M, Smarandache F (2016) Complex neutrosophic set. *Neural Comput Appl* 25:1–18
2. Ali M, Deli I, Smarandache F (2016) The theory of neutrosophic cubic sets and their applications in pattern recognition. *J Intell Fuzzy Syst* 30(4):1957–1963
3. Atanassov K (1986) Intuitionistic fuzzy sets. *Fuzzy Sets Syst* 20:87–96
4. Athar KA (2014) Neutrosophic multi-criteria decision making method. *New Math Nat Comput* 10(02):143–162
5. Aydoğdu A (2015) On similarity and entropy of single valued neutrosophic sets. *Gen Math Notes* 29(1):67–74
6. Broumi S, Smarandache F (2013) Several similarity measures of neutrosophic sets. *Neuro Sophic Sets Syst* 1(1):54–62
7. Broumi S (2013) Generalized neutrosophic soft set. *Int J Comput Scie Eng Inf Technol(IJCSEIT)* 3/2:17–30
8. Broumi S, Talea M, Bakali A, Smarandache F (2016) Single valued neutrosophic graphs. *J New Theory* 10:86–101
9. Broumi S, Talea M, Bakali A, Smarandache F (2016) On bipolar single valued neutrosophic graphs. *J New Theory* 11:84–102
10. Broumi S, Smarandache F, Talea M, Bakali A (2016) An introduction to bipolar single valued neutrosophic graph theory. *Appl Mech Mater* 841:184–191
11. Broumi S, Bakali A, Talea M, Smarandache F (2016) Isolated single valued neutrosophic graphs. *neutrosophic Sets Syst* 11:74–78
12. Balasubramanian A, Prasad KM, Arjunan K (2015) Bipolar interval valued fuzzy subgroups of a group. *Bull Math stat Res* 3(3):234–239
13. Broumi S, Smarandache F (2013) Several similarity measures of neutrosophic sets. *Neutrosophic Sets Syst* 1(1):54–62
14. Broumi S, Deli I, Smarandache F (2014) Distance and similarity measures of interval neutrosophic soft sets. In: *Critical review, center for mathematics of uncertainty, Creighton University, USA, vol 8, pp 14–31*
15. Broumi S, Deli I (2016) Correlation measure for neutrosophic refined sets and its application in medical diagnosis. *Palest J Math* 5(1):135–143
16. Deli I, Subas YA (2016) Multiple criteria decision making method on single valued bipolar neutrosophic set based on correlation coefficient similarity measure. In: *International conference on mathematics and mathematics education (ICMME-2016), Frat University, May 12–14, Elazg, Turkey*
17. Chen J, Li S, Ma S, Wang X (2014) m-Polar fuzzy sets: an extension of bipolar fuzzy sets. *Sci World J*. doi:10.1155/2014/416530
18. Deli I, Ali M, Smarandache F (2015) Bipolar neutrosophic sets and their application based on multi-criteria decision making problems. In: *2015 International conference on advanced mechatronic systems (ICAMechS). IEEE.(2015, August), pp 249–254*
19. Deli I, Broumi S (2015) Neutrosophic soft matrices and NSM-decision making. *J Intell Fuzzy Syst* 28:2233–2241
20. Karaaslan F (2016) Correlation coefficient between possibility neutrosophic soft sets. *Math Sci Lett* 5(1):71–74
21. Lee KM (2000) Bipolar-valued fuzzy sets and their operations. In: *Proceedings of international conference on intelligent technologies, Bangkok, Thailand, pp 307–312*
22. Lee KJ (2009) Bipolar fuzzy subalgebras and bipolar fuzzy ideals of BCK/BCI-algebras. *Bull Malays Math Sci Soc* 32(3):361–373
23. Ma YX, Wang JQ, Wang J, Wu XH (2016) An interval neutrosophic linguistic multi-criteria group decision-making method and its application in selecting medical treatment options. *Neural Comput Appl*. doi:10.1007/s00521-016-2203-1
24. Maji PK (2013) Neutrosophic soft set. *Ann Fuzzy Math Inf* 5(1):157–168
25. Majumdar SK (2012) Bipolar valued fuzzy sets in -semigroups. *Math Aeterna* 2(3):203–213
26. Majumdar P, Smanata SK (2014) On similarity and entropy of neutrosophic sets. *J Intell Fuzzy Syst* 26(3):1245–1252
27. Mondal K, Pramanik S (2015) Neutrosophic tangent similarity measure and its application to multiple attribute decision making. *Neutrosophic Sets Syst* 9:85–92
28. Mukherjee A, Sarkar S (2014) Several similarity measures of neutrosophic soft sets and its application in real life problems. *Ann Pure Appl Math* 7(1):1–6
29. Pramanik S, Biswas P, Giri BC (2015) Hybrid vector similarity measures and their applications to multi-attribute decision making under neutrosophic environment. *Neural Comput Appl* 1–14. doi:10.1007/s00521-015-2125-3
30. Pramanik S, Mondal K (2015) Cosine similarity measure of rough neutrosophic sets and its application in medical diagnosis. *Global J Adv Res* 2(1):212–220
31. Peng JJ, Wang JQ, Zhang HY, Chen XH (2014) An outranking approach for multi-criteria decision-making problems with simplified neutrosophic sets. *Appl Soft Comput* 25:336–346
32. Smarandache F (1998) *A unifying field in logics neutrosophy: neutrosophic probability, set and logic*. American Research Press, Rehoboth

33. Smarandache F (2005) Neutrosophic set, a generalisation of the intuitionistic fuzzy sets. *Int J Pure Appl Math* 24:287–297
34. Santhi V, Shyamala G (2015) Notes On Bipolar-Valued Multi Fuzzy Subgroups Of A Group. *Int J Math Arch (IJMA)*. 6(6). ISSN 2229–5046
35. Saeid AB (2009) Bipolar-valued fuzzy BCK/BCI-algebras. *World Appl Sci J* 7(11):1404–1411
36. Şahin M, Deli I, and Ulucay V (2016) Jaccard vector similarity measure of bipolar neutrosophic set based on multi-criteria decision making. In: *International conference on natural science and engineering (ICNASE'16)*, March 19–20, Kilis
37. Tian ZP, Wang J, Wang JQ, Zhang HY (2016) Simplified neutrosophic linguistic multi-criteria group decision-making approach to green product development. *Group Decis Negot* 1–31. doi:10.1007/s10726-016-9479-5
38. Tian ZP, Wang J, Wang JQ, Zhang HY (2016) An improved MULTIMOORA approach for multi-criteria decision-making based on interdependent inputs of simplified neutrosophic linguistic information. *Neural Comput Appl*. doi:10.1007/s00521-016-2378-5
39. Tian ZP, Zhang HY, Wang J, Wang JQ, Chen XH (2015) Multi-criteria decision-making method based on a cross-entropy with interval neutrosophic sets. *Int J Syst Sci*. doi:10.1080/00207721.1102359
40. Wang H, Smarandache FYQ, Zhang Q, Sunderraman R (2010) Single valued neutrosophic sets. *Multispace Multistructure* 4:410–413
41. Wu J, Cao QW (2013) Same families of geometric aggregation operators with intuitionistic trapezoidal fuzzy numbers. *Appl Math Model* 37:318–327
42. Wu XH, Wang JQ, Peng JJ, Chen XH (2016) Cross-entropy and prioritized aggregation operator with simplified neutrosophic sets and their application in multi-criteria decision-making problems. *Int J Fuzzy Syst*. doi:10.1007/s40815-016-0180-2
43. Xu X, Zhang L, Wan Q (2012) A variation coefficient similarity measure and its application in emergency group decision making. *Syst Eng Proc* 5:119–124
44. Ye J (2014) Vector similarity measures of simplified neutrosophic sets and their application in multicriteria decision making. *Int J Fuzzy Syst* 16(2):204–215
45. Ye J (2014) Improved correlation coefficients of single valued neutrosophic sets and interval neutrosophic sets for multiple attribute decision making. *J Intell Fuzzy Syst* 27:24532462
46. Ye J, Zhang QS (2014) Single valued neutrosophic similarity measures for multiple attribute decision making. *Neutrosophic Sets Syst* 2:48–54
47. Ye S, Ye J (2014) Dice similarity measure between single valued neutrosophic multisets and its application in medical diagnosis. *Neutrosophic Sets Syst* 6:48–52
48. Ye J, Fu J (2016) Multi-period medical diagnosis method using a single valued neutrosophic similarity measure based on tangent function. *Comput Methods Program Biomed* 123:142–149
49. Ye J (2015) Single-valued neutrosophic similarity measures based on cotangent function and their application in the fault diagnosis of steam turbine. *Soft Comput* 1–9. doi:10.1007/s00500-015-1818-y
50. Jun YB, Kang MS, Kim HS (2009) Bipolar fuzzy structures of some types of ideals in hyper BCK-algebras. *Sci Math Jpn* 70:109–121
51. Zadeh LA (1965) Fuzzy sets. *Inf Control* 8:338–353
52. Zhang HY, Ji P, Wang JQ, Chen XH (2016) A neutrosophic normal cloud and its application in decision-making. *Cogn Comput*. doi:10.1007/s12559-016-9394-8