The Journal of the American Medical Association

Published Under the Auspices of the Board of Trustees

VOL. 70, No. 4

CHICAGO, ILLINOIS

JANUARY 26, 1918

A RARE TYPE OF BLADDER ULCER

FURTHER NOTES, WITH A REPORT OF
EIGHTEEN CASES *

GUY L. HUNNER, M.D.

Fellow of the American College of Surgeons
BALTIMORE

In 1914, I described a rare type of bladder ulcer in women.¹ Further work with this type of bladder inflammation has not revealed new material of special value, but I hope that a new publication with illustrations may receive a wider hearing and familiarize physicians with this exceedingly important group of bladder cases.

The fact that ten new cases can be added in less than three years demonstrates that we are dealing with a condition which is not so rare as may have been inferred from the former report, and which will be discovered and relieved much more promptly when the profession awakens to its existence.²

There is no department of gynecology or of urology that has baffled our attention more than the complaints commonly classified as "irritable bladder," "neuralgia or neurosis of the bladder," or "neurasthenia with the mind centered on the urinary apparatus." We have used such terms as a diagnostic entity, or as a hazy cloak for our ignorance, in seeking to classify that large group of patients who persist for years in complaining of the severe symptoms usually associated with cystitis, but in whom we can find but little evidence of cystitis by the ordinary urinary and cystoscopic examinations, and who fail to answer to the usual methods of bladder and urethral treatments.

By discovering that some of these cases are due to a chronic urethritis, which in turn is secondary to a distant focal infection, I have demonstrated a definite pathologic basis for a large group of these cases.³ Such patients may suffer for months or years with the most distressing bladder symptoms, even to the extreme degree of a complete incontinence, and may indefinitely resist all forms of bladder and urethral treatments, and then promptly recover without further local treatment after the removal of infected tonsils, or abscessed teeth, or the drainage of infected sinuses.

The demonstration of the bladder lesion under discussion has removed another considerable group of cases from the vague classification of "neurosis of the bladder," and placed them on a definite pathologic

basis, enabling us to treat them in a logical and successful manner.

DESCRIPTION OF THE LESION AND DISCUSSION OF NOMENCLATURE

In my initial report, this lesion was spoken of as "simple" ulcer of the bladder for the reason that it is unassociated with any known infection. I contrasted this ulcer with the so-called simple ulcer of Fenwick.⁴ Fenwick's terminology was based on the resemblance of his ulcer to the "simple ulcer" of the stomach, and he was apparently unaware that both Rokitsansky and Tait⁵ had forestalled him in this comparison.

This terminology was in use before the day of accurate bacteriologic work on the bladder infections, and it is a question now just what ulcers, if any, should be classified as "simple." In more recent years Buerger⁶ has used the term "simple callous ulcer" to describe what is apparently the simple ulcer of Fenwick; but he reports one of his cases as beginning in pregnancy and yielding a *Staphylococcus albus* culture. Le Fur⁷ states that simple chronic ulcers of the vesical mucosa do not always present themselves as round ulcers with thickened callous edges simulating ulcer of the stomach, but may offer very irregular forms or special aspects. He holds that they may be so small and superficial as to be called microscopic or histologic ulcers, as contrasted with the usual macroscopic ulcer.

Unfortunately, he illustrates the latter point by referring⁸ to some of his results in animal experimentation, wherein histologic ulcers were caused after inoculation of the blood with organisms and then injuring of the bladder wall. Such conditions might conceivably result in what eventually might be called a simple ulcer; but these small fresh lesions in a recently inoculated animal would not correspond to the usual conception of a simple ulcer as one developing without demonstrable cause.

It is apparent to any one reviewing the literature of bladder ulcer that some organization, such as the American Urological Association or the American Association of Genito-Urinary Surgeons, could perform a valuable service by appointing a small commission to review the subject from every point of view, and adopt a nomenclature that would serve as a working basis for our present knowledge.

4. Fenwick: The Clinical Significance of the Simple Ulcer of the Urinary Bladder, Brit. Med. Jour., 1896, **1**, 1133; Ulcer of the Bladder, London, J. and A. Churchill, 1900, p. 8.

5. Tait: On the Cure of the Chronic Perforating Ulcer of the Bladder by the Formation of an Artificial Vesicovaginal Fistula, Lancet, London, 1870, **54**, 738.

6. Buerger, Leo: Pathology and Treatment of Callous Ulcer of the Bladder, Med. Rec., New York, April 12, 1913; Ulcer of the Bladder, THE JOURNAL A. M. A., Feb. 8, 1913, p. 419.

7. Le Fur: Des ulcérations vésicales et en particulier de l'ulcère simple de la vessie, Paris, G. Steinheil, 1901.

8. Le Fur (Footnote 7), p. 205.

* From the Gynecological Department of the Johns Hopkins University.

1. Hunner, G. L.: A Rare Type of Bladder Ulcer in Women, with Report of Eight Cases, Tr. South. Surg. and Gynec. Assn., 1914, **27**; Boston Med. and Surg. Jour., 1915, **172**, 660.

2. Four new cases have been seen between June and December, 1917.

3. Hunner, G. L.: Chronic Urethritis and Chronic Ureteritis Caused by Tonsillitis, THE JOURNAL A. M. A., April 1, 1911, p. 937.

Geraghty⁹ has proposed the designation of "paracystitis" for the lesion under discussion; but this conveys the idea of an inflammation of the tissues surrounding the bladder, and is therefore inadequate.

My associate, Dr. Cullen, after a review of the clinical and pathologic features of this lesion, has suggested

Fig. 1.—Location of multiple minute ulcers on the right vertex, and the surrounding edema area found at operation in Case 3.

the term "elusive ulcer," and this comprehends satisfactorily the difficulties in locating the ulcer part of the lesion, and its changing characters while under observation and treatment; but in common with any simple term, it fails to describe adequately the widespread character of the chronic inflammatory involvement of the bladder walls.

The lesion under discussion is a chronic inflammation of all coats of the bladder wall; and by the time these cases have reached me for treatment, the lesion has become widespread.

In the two least extensive cases, the lesion in Case 3 covered an area 3 by 5 cm., and in Case 11 an area 2 by 3 cm. In the more extensive cases the lesion has necessitated the removal of more than the upper three fourths of the bladder. In spite of this widespread lesion, there is remarkably little to be found by cystoscopy, and this is one reason that many of these patients have suffered fifteen years or more, and have traveled from clinic to clinic in an effort to find relief.

The crucial test in cystoscopy is the finding of a small abrasion on the mucosa surface which, if not bleeding on discovery, will easily bleed on being touched with an instrument or with a cotton pledget. It is this minute area that gives rise to the urinary findings of a few leukocytes or a few erythrocytes or both when the urine is carefully settled or centrifuged. Occasionally the distention of the bladder by air as the patient assumes the knee-breast posture causes this area to split, and a tiny stream of blood flows to the vertex. In such a case one first notices the blood-tinged urine in the vertex in striking contrast to the

perfectly clear urine that has just been drawn by catheter; and by sweeping the speculum around the equator of the bladder, one finds the blood streak and can trace it to the source. The ulcer is usually found in the vertex or free portion of the bladder, as contrasted with the location of the Fenwick ulcer on the base or fixed portion (Figs. 1 and 2). Its frequent location on the anterior superior wall is another reason for its having escaped diagnosis, this being a difficult area to examine with either the direct or the indirect methods of cystoscopy.

With the proper knee-breast posture and good relaxation of the patient, we can examine every portion of the bladder mucosa, including the anterior wall back of the symphysis. When there is any difficulty in a free examination of the anterior wall, this can easily be brought into vision by sliding the free hand beneath the mons veneris and gently lifting the anterior wall of the bladder into line with the examining cystoscope.

The actual abrasions of the mucosa may be single or multiple. If multiple, they may be in a small group (Fig. 1) or they may be widely separated, one ulcer being found in the vertex, perhaps close behind the symphysis on the anterior wall, and another near the base on the posterior wall (Fig. 2).

These ulcers are always small, varying in diameter from 2 to 5 mm., and they appear to be very superficial. I have never seen them covered with fibrin or with a deposit of urinary salts, and have never seen them present a picture suggesting malignancy.

They are exceedingly sensitive on being touched, and it is most gratifying on discovering one of these ulcers and touching it to have the patient who has been through many fruitless cystoscopies exclaim, "There: you have found the right spot at last!" The preliminary bimanual examination of the bladder will often suggest the location of the ulcer because of this tenderness. It is probably this preliminary palpation that

Fig. 2.—Site of the inflammatory area near the posterior pole in Case 8. In addition to this area there was a small ulcer back of the symphysis on the left anterior wall, and at operation an area 10 by 4 cm. was removed to include all the congested area.

starts the bleeding in some cases after one has just catheterized clear urine.

On cystoscopy one's attention is not infrequently first arrested by a glazed, dead white appearance of a portion of the bladder mucosa. In such an infiltrated area the vessels may have a broken or "choppy"

9. Geraghty: Infections of the Bladder, with Special Reference to Localized Resistant Areas of Cystitis, Surg., Gynec. and Obst., 1917, 24, 655.

appearance, only short segments of vessels appearing on the extra pale surface. One may see a dead white scar area with a small congested area in the immediate neighborhood; and while one is examining this area, or after one has touched it, the congestion becomes marked and may even begin to ooze blood (Fig. 3).

Fig. 3.—Cystoscopic appearances of an area near the posterior pole region in Case 8. The vascular changes from *A* to *B* occurred spontaneously while the artist was viewing the field through the speculum.

Sometimes during the cystoscopy an area of edema appears, the mucosa taking on a deep red congested appearance and being raised above the general mucosa level. This area of edema may occur immediately about the ulcer area or it may appear on a portion of the wall at some distance from the ulcer. It may be absent at several consecutive examinations and then appear at a subsequent examination. It is particularly likely to be seen if the patient is in the midst of an unusually bad period of bladder symptoms with strangury. The significance of this edema area will be fully discussed under the section on treatment.

ETIOLOGY

The cause of this type of bladder inflammation remains a mystery. There is no evidence from the histories of these patients that the condition began as an infection in the urine. The patients, except in Cases 7 and 16, gave no history of having had pus or blood in the urine at any stage of their cystitis history. In Case 7, the patient said she had seen blood in the urine at times during the sixteen years of her symptoms. In Case 16, the patient was convinced that her bladder trouble of eighteen years' duration began after an operation, and was due to faulty catheterization. She was known to have had at that time a typical generalized cystitis with pus in the urine.

With this one exception, none of the cases, so far as could be determined by the histories, followed operation, childbearing or gonorrhea.

The urine is always sterile on our ordinary culture mediums. Attempts to grow anaerobic cultures from the removed tissues have resulted in failure, as have many attempts to find organisms in the tissues by numerous methods of staining.

In my early cases the long duration and the resistance to treatment made me suspect primary tuberculosis of the bladder; but the excised specimens examined microscopically did not bear out this view.

Because my work has demonstrated the probable relationship between infections of the tonsils, adenoids, sinuses and teeth, and conditions of chronic infiltration in the ureters and urethra, I have suspected that this chronic infiltration in the bladder wall may have a similar focal origin. These patients all have an infiltration, granulation, and extreme hypersensitivity of the urethra. The tenderness and scar tissue contraction was so marked in most of the cases as to require thorough cocainizing and careful manipulation in order that satisfactory cystoscopy might be done. Fenwick emphasizes this urethral hyperesthesia in most cases of nonmalignant ulceration of the bladder, and advises chloroform anesthesia for cystoscopic examination. It is possible that the bladder ulceration and the inflammatory condition of the urethra may have a common cause, or it may be that the frequency of voiding and straining bring about a hypertrophy and hypersensitivity of the sphincter region and an inflammation of the urethral tissues. I am inclined to take the latter view from the fact that these patients have no symptoms referable to the urethra after recovering from the excision of the bladder lesion. The location of these lesions in the vertex or free portion of the bladder, instead of in the base where there is the richer blood and lymph supply, probably argues against the focal infection theory.

The fact that many of the patients begin with their bladder symptoms in early life (seven of them dating the onset at 20 years and under) would tend to bear out the focal infection theory. On the other hand, the histories and examinations of these patients have

Fig. 4.—From Case 7 (Pathologic No. 29137): *a*, gradual fading of transitional epithelium; *b*, ulcer area, made up largely of blood vessels; *c*, muscle in mucosa layer; *d*, blood vessels packed with leukocytes even in the deeper layer of mucosa.

not borne out the theory of a distant focus of infection in the striking manner we find to be the rule in most ureteral stricture cases.

We have failed to make the systematic notes that should have been made on the condition of the teeth.

In Cases 4 and 7, the patients thought their bladder trouble began soon after the infectious fevers of child-

hood. In Case 3, the patient, who had suffered with bladder symptoms "all her life," had tonsillitis as a child. In Case 8, the patient had a severe attack of tonsillitis soon after the bladder trouble began at 15 years of age. In Case 16, the patient, whose bladder trouble began after an operation at 21 years of age, had had her tonsils guillotined when she was 12 years old. In Case 18, the patient, aged 28 years, had suffered with her tonsils, and had had them removed five years after the bladder symptoms began. In

Fig. 5.—From Case 3 (Pathologic No. 20722): *a*, abrupt ending of stratified epithelium at edge of healing ulcer; *b*, single layer of cuboidal cells; attempt at repair over the healing area; *c*, so-called cysts or glands, with mucous or goblet cell lining; *d*, smooth muscle in mucosa layer; *e*, blood vessels filled with leukocytes; *f*, collection of small round cells and leukocytes. General rarefaction due to edema.

Case 15, the patient, aged 30 years, said that her bladder trouble had begun three years previously while she was suffering with a severe "cold." In Cases 8 and 10, the sinuses were diseased. In Cases 2, 5 and 9, the patients had rheumatism.

Sex.—My eighteen cases have all occurred in women, my work being confined to this sex. Dr. Geraghty has recently been finding these cases in men, and he thinks that their clinic has overlooked many of this type in the past and sent them away unimproved, after long courses of various forms of bladder and urethral treatment.

Age.—In my first eight cases, reported before, the average age of the patient on consultation was 37 years; the average duration of symptoms was seventeen years, making the average onset age 20 years. In my last ten cases the average age at consultation was 44.2 years, and the average duration of symptoms ten and four-fifths years, making the average age of onset 33.4 years.

Childbearing does not seem to have any relation to the disease, nine of the eighteen patients having been unmarried.

SYMPTOMS

The chief symptom associated with this type of bladder ulcer is pain. Associated with the pain, the other symptoms of cystitis occur in varying degree, namely, frequency day and night, strangury, burning and smarting. The pain is often of the most extreme grade, the patient complaining of a jabbing or stabbing knifelike pain or of a sensation of a jagged, sharp stick in the bladder. In Case 8 the patient often had such extreme urgency that she had to leave a street-car in order to enter the nearest house and ask for permission to void. In Case 6 the patient expressed the urgency of her pain by saying that if she were crossing the street, and an automobile were bearing down on her at the time of getting one of her knife-

like stabs in the bladder, she would have to drop in her tracks and allow the automobile to run over her. In Case 3, the patient, when at her worst, could not allow people to walk across the room in which she was sitting because of the increased pain from the vibration of the floor. One is at a loss to describe the misery of these patients without selecting such descriptions from the patients themselves.

The pain is often referred to some other portion of the pelvis, particularly to the rectum. The pain is often increased after eating, probably owing to increased peristalsis. It is increased when there is "gas on the bowels" or when the patient is constipated, and just before and during stools, probably because of the fecal matter's passing by the tender peritoneum located over the diseased area of the bladder wall.

Pain referred to the hip corresponding to the side of the open ulcer is not uncommon. Pain in the perineum is common. The patients often say they must sit down in a particular manner, resting mostly on one hip, or sitting on the edge of the chair. Turning in bed at night may bring on severe pain and awaken the victim. In other words, these patients are in a constant state of muscular tension in an effort to "splint" the bladder and give it no undue movement or jarring.

The rectal symptoms, together with the failure of a diagnosis, sometimes convince the patient that a cancer of the bowel is developing. It is largely because of these referred pains, and the failure to make a diagnosis, that these patients are so often operated on for other supposed lesions.

Urine Examination.—The diagnostic features of the Fenwick type of ulcer are pain and hemorrhage. Only one of our eighteen patients reported having seen blood in the urine.

The urine is macroscopically crystal clear except for a mucin cloud in suspension; a hurried microscopic examination may readily miss the few erythrocytes practically always present. In a suspected case one

Fig. 6.—From Case 3 (Zeiss DD, 4.3 mm., reduced about one half): *a*, the horizontal arrangement and "hornified" appearance of the epithelial cells; *b*, single cuboidal layer over healing area; *c*, muscle cells of the mucosa layer; *d*, capillaries filled with leukocytes.

should always settle, pipet and centrifuge the urine. We still hear the expression "an occasional leukocyte and an occasional red blood corpuscle, not enough to indicate any trouble." As long as clinicians take this attitude, these cases will be undiagnosed.

While treating these patients we can occasionally get the ulcer to heal over temporarily; at such times

we may be unable by the most careful microscopic search to find a pathologic element in the urine. At the patient's next visit a week later we are very likely to find the usual few leukocytes or erythrocytes or both.

PATHOLOGY ¹⁰

The histologic study of specimens removed at operation shows a fairly uniform picture in all cases. It is a picture of chronic inflammation involving all coats

Fig. 7.—From Case 8 (Zeiss AA, 17 mm., reduced about one half): *a*, absence of epithelium except in protected depressions; *b*, bridge of organized tissue covering depression; *c*, blood vessel in bridge; *d*, unusual thickening of basement membrane; *e*, prominent muscularis mucosae.

of the bladder and extending over the wide area which at operation shows the edematous thickening. At the site of the minute ulcer there is loss of the epithelial coat, and the underlying mucosa shows the granulation tissue characteristic of an active ulcer (Fig. 4). Other sections, taken from what appears to be an ulcer in the gross specimen, show an abrupt ending of the epithelial layer at the edge of an area that is evidently undergoing healing, and here the mucosa layer is less richly supplied with capillaries, and is much like the mucosa beneath the epithelial covering, showing a preponderance of connective tissue and infiltration of small round cells and leukocytes. Such a healing area may be covered by a single layer of cuboidal cells (Figs. 5 and 6).

Figure 7 shows another type of attempt at repair of the epithelial layer. In several of our sections we find depressed areas in the mucosa which have apparently become filled with fibrin or serum and in the bottom of which new epithelium is growing. This specimen shows, extending across the depression, a protecting bridge in which can be traced organized tissue, including blood vessels.

In general, the transitional type of epithelium has changed to a flattened horizontal type. Compare the normal transitional type of epithelium in Figures 8 and 9 with the changed types in Figures 5, 6 and 11.

It is interesting to find on the histologic examination that broad areas of the mucosa may be denuded of the epithelial coat even though the cystoscopic picture showed only a small ulcer area. It is possible that this wide loss of epithelium is an artefact due to the

trauma during operation or during the preparation of the specimen. Great care has been used, however, during the operation not to injure the epithelium, and I think that a glance at Figures 5 and 6, showing the abrupt ending of the epithelial layer on the edge of an area apparently not in active ulceration, will convince one that this was the condition in vivo. The gradual shading off of the epithelial coat in Figure 4 *a* at the edge of an ulcer area is similar to the gradual ending of the epithelium in Figure 12 *a*, where the denuded area is covered by an intact thickened basement membrane and shows no special evidence of ulcer.

A thickening of the basement membrane is a prominent feature in some cases (Figs. 12 and 7). Apparently this basement membrane thickening is not dependent on the absence of the epithelial layer, but appears as well beneath the areas covered with epithelium, as seen in Figure 12.

In Case 3 there occur many of the so-called "glands of the bladder" lined with goblet cells (Fig. 5 *c*). The significance of these will be discussed in our future full pathologic report.

The mucosa layer in the neighborhood of the ulcer shows an increase in the number and size of the capillaries, varying in the different specimens. The capillaries are often stuffed with leukocytes, as are many of the lymph spaces. There are some enlarged lymph spaces crowded with lymphoid cells and leukocytes, and where these are seen lying beneath the normal epithelium near the ulcer, they at first suggest the picture of tubercle (Fig. 12 *d*). There is an appearance of hyalin degeneration in some areas, and in some of the sections the edema causes rarefaction.

It is important to note that the presence of leukocytes stuffing the vessels in these specimens probably has no pathologic significance. Figures 8, 9 and 10, representing a section of bladder wall which I removed from a patient who had never had a history of cystitis,

Fig. 8.—From normal case (Zeiss AA, 17 mm., reduced one half): *a*, transitional type of epithelium; *b*, indistinct basement membrane; *c*, muscles of mucosa layer; *d*, medium sized vessels of mucosa layer; *e*, large vessels of submucosa region; *f*, muscle layers.

show the presence of a great many leukocytes in spite of the fact that the specimen was removed with as great rapidity and with as little trauma as possible, and was fixed in formaldehyd solution at once.

The muscularis mucosae seems to be hypertrophied in most of the specimens (Figs. 5 *d*, 8 *c*, normal, and 7 *e*).

10. The author hoped to have ready with his associate, Dr. George L. Stickney, a full report on the pathology of this condition. Dr. Stickney was one of the early volunteers to leave for medical service in Europe, and we shall therefore defer the full pathologic report for a future publication.

The muscle coat shows enlarged lymph spaces filled with small round cells; these spaces, as well as some of the blood vessels of the muscle coat, contain many leukocytes. Immediately beneath the ulcer, the muscle in places shows invasion and breaking up by the inflammatory process.

The peritoneum, subtending a diseased area, shows decided thickening.

Fig. 9.—From normal case. (Zeiss DD, 4.3 mm., reduced one half; enlarged from Fig. 8): a, typical transitional epithelium; b, cell of surface layer with double nucleus.

DIAGNOSIS

The diagnosis of this form of bladder inflammation rests on the history, the urine examination, the cystoscopic findings, and the reaction of the disease to treatment.

So far as I know, there is no form of bladder inflammation or ulceration that presents an absolutely characteristic clinical history or cystoscopic picture. The process under discussion is no exception to the rule.

The patient's recital of the insidious onset without apparent antecedent cause, and the long duration in spite of various forms of treatment, make one think at once of tuberculous disease. This is excluded as soon as one finds macroscopically clear urine associated with such a long history, and absence of any sign of disease of the kidney or ureter.

One sometimes gets a history of serious symptoms simulating cystitis from patients suffering with chronic urethritis. The presence of microscopic pus or blood or of both in urine catheterized from women makes one certain that one is dealing with something more than a urethritis. Rarely one gets a few blood cells from trauma to the sphincter region during insertion of a catheter.

I have learned by experience to suspect the presence of this lesion when I am consulted about any patient who has suffered for years with symptoms chiefly referred to the bladder, but varied enough in character to have led to a number of operations without relief of the symptoms. If such an outline of trouble ends with the statement that "there has never been anything of importance in the urinary findings, only at times a few leukocytes or red cells," my suspicion is changed to a practical certainty.

These points may best be emphasized by a brief summary of the efforts at diagnosis and relief in two of my recent cases:

Mrs. W., aged 50, referred in January, 1917, by Dr. Catharine Macfarlane of Philadelphia, had suffered for eighteen years with symptoms chiefly referable to the bladder. She was convinced that these symptoms dated from a cystitis due to faulty catheterization after an Alexander operation.

The patient wrote me in April, three months after operation, to say that she was "perfectly well," and she sent me an interesting file of correspondence concerning her case, showing that she had been under the care of some of the best medical and surgical talent in Philadelphia, Baltimore, New York and Boston; and in her fits of despondency over the failure of these leaders in our profession to benefit her, she had interspersed among these treatments Christian Science, osteopathy, mechanoneurotherapy, and other *fin de siècle* methods in an attempt to get well. The patient had all of the diseases of childhood in a severe form, and at 12 years of age the tonsils were guillotined. She was married in 1896 at 19 years of age, and her first child was born a year later. In May, 1898, she had the Alexander operation, followed by the severe attack of cystitis. About a year later a Boston gynecologist made a cystoscopic examination under anesthesia, and diagnosed a colon bacillus ulcer of the bladder. This was later treated by a Philadelphia gynecologist, and it was apparently completely cured, according to cystoscopic findings. The symptoms, however, continued. In 1900, the sphincter ani was dilated by an osteopath. In 1903, the patient tried Christian Science. In 1905, a prominent Philadelphia gynecologist made an abdominal exploratory operation, and found a few adhesions spreading from the top of the uterus over the posterior wall of the bladder. These were cut, and it was hoped that the cause of the symptoms had been found; but there was general disappointment when the patient again became active and found that her old symptoms were still with her. In 1913, vaccines were used in the hope of clearing up abdominal adhesions. Bladder irrigations were tried, and an examination of the right kidney was made. The patient was sent away for a rest cure, a Philadelphia neurologist having concluded that she needed "a course of nervous and mental training with particular attention not only to control of the bladder function but along other lines." She was prescribed "a period of six or eight months at a place like Lake Placid, or in Canada, with a nurse of sufficient brain power and force of character and training to be able to carry out in detail the directions of the physician, and at the same

Fig. 10.—Section of muscle wall next to peritoneum, from normal case (Zeiss DD, 40.3 mm., reduced one half): a, vessel crowded with leukocytes; b, leukocytes traversing vessel walls. It took perhaps three minutes to strip back the peritoneum and to cut out the specimen, pin it to a thin board, and drop it in formaldehyd solution.

time to direct the mental training when the proper time has arrived." In 1915, after the patient had considerably reduced her weight by diet, the right kidney was found floating, and a fixation was done. In 1916, treatment by the roentgen ray, electricity, and colored lights were used. Pyorrhea was discovered, and two teeth were drawn. Later there was treatment for arthritis. On first consulting me, the patient was

convinced that the fixed right kidney was back of her bladder symptoms, and that it should be removed.

The foregoing is only a brief summary, and does not begin to cover the field of the various diagnoses and treatments during these eighteen years of suffering.

My latest patient, Case 18, a woman, aged 28, unmarried, had been suffering with bladder symptoms for six years, dur-

Fig. 11.—From Case 4 (Zeiss DD, 4.3 mm., reduced one half): *a*, flattening and "hornification" of upper layers of epithelium; *b*, clear spaces at base of epithelial layer, one space containing two leukocytes and one space an epithelial cell with double nucleus.

ing which time she had been in the hands of at least seven different physicians, all of them of excellent standing in their respective specialties. In the first year of her trouble she saw two physicians in Atlanta, one of whom gave her urethral and rectal irrigations, and the other dilated the urethra and "removed a small growth from the external orifice," and reported a negative bladder examination. He also catheterized the right ureter with negative results. Three and a half years before I saw the patient, a prominent Baltimore physician examined carefully with the cystoscope with negative results, dilated the urethra, and treated with radium for vaginismus. Eighteen months before, a Greenville, S. C., physician found an ulcer in the bladder, and after a few bladder treatments the bladder seemed normal, and the urine was reported normal; but the patient's symptoms remained the same. One of her chief symptoms was pain in the left labium. This physician severed the left pudic nerve without benefit. One year before, he did an exploratory celiotomy and "removed a normal looking appendix and a normal looking left ovary, and suspended the uterus." He also removed the left labium minus. Six months before, a Columbus, S. C., physician examined the bladder and kidneys with negative results. Eight months before, tonsillectomy was done. Four months before, a prominent Atlanta specialist treated her for sacro-iliac joint trouble without benefit.

The patient was referred to me in March, 1917, by Dr. Barker. A typical lesion was found, consisting of a small ulcer in the right vertex and a congested area of mixed scar and vascular tissue in the left vertex. At operation the edema area carried us far down the posterior wall and on the base, and at the close of the operation there was just enough flap of bladder wall on the left side to be brought across to the resected edge of the right base and trigonal region. The remaining bladder probably held less than 1 ounce. One month after the patient's return home she reported: "Although my bladder is not quite well yet, the unpleasant symptoms are so slight now that they cause me very little inconvenience. The pain and gas in the stomach have entirely disappeared, my bowels move well without laxatives, and the old rectal discomfort is a thing of the past. My bladder will hold easily 10 ounces; and best of all, I am quite free from those horrible perineal spasms which had puzzled so many doctors

and almost wrecked my nervous system. My appetite is good, I sleep well, and have gained 12 pounds since I left the Hopkins. My strength has returned with amazing rapidity; I ride and walk without pain, and I am sure that you would be pleased to know how much I am able to do in the way of housework."

Nine of the eighteen patients had had a total of sixteen operations directed toward the relief of their symptoms. These operations were largely futile because of a failure to make a proper diagnosis. I operated in Case 2, doing a perineal repair and supra-vaginal hysteromyomectomy, and in Case 4, removing an atrophic appendix, before discovering that the patients' chief symptoms were due to this type of bladder inflammation.

TREATMENT

After an experience with eighteen cases, covering a period of seventeen years since treating the first patient, and a period of nine years since seeing the second patient and beginning to make a special study of this type of bladder inflammation, I believe it safe to say that no form of treatment will suffice in these cases except complete excision of the inflammatory area.

In Case 2, the patient is still under observation after nine years of intermittent treatment, and her cystoscopic picture has remained practically constant, the symptoms being alleviated materially when she persists in having an application of 10 per cent. solution of silver nitrate once in six weeks. From the first there has been a minute area of ulceration in the upper left vertex of the bladder close back of the symphysis. When the symptoms have been aggravated, there has been some edema about this ulcer area, and in addition an area of edema 3 or 4 cm. in diameter in the middle vertex. When the symptoms have been alleviated after a systematic course of treatments, the ulcer area on two or three occasions has seemed to be healed, and could then be identified only by a slight congestion.

Fig. 12.—From Case 6 (Pathologic No. 20073): *a*, gradual fading of transitional epithelium; *b*, marked thickening of basement membrane; *c*, marked connective tissue thickening and small round cell infiltration immediately beneath the epithelium; *d*, focus of small round cells.

On such occasions the urine after careful centrifugation has been entirely free from pathologic elements. This "healed" condition has never been seen on two successive examinations, and the patient has never been entirely free from symptoms. She has had persistent courses with instillations twice a week with mercuric solution, 1:10,000 and with silver nitrate solution, 1:500, but the most effective treatment has

been with applications of 10 per cent. silver nitrate solution about once in four or six weeks. She was 54 years of age when first seen, and then had a repair of the outlet and supravaginal hysteromyomectomy

Fig. 13.—Excised specimen in Case 3. The area of edema and congestion measures about 3 by 5 cm., surrounding the small central ulcers. Naturally the trauma of operation makes the specimen more congested than it appears at cystoscopy.

before the true cause of her symptoms was discovered, and she has since preferred the comparative relief obtained by local treatments to the prospects of another operation.

In two other cases (10 and 13) the patients, aged 68 and 57, respectively, have not been operated on because of their age and general condition. In Case 10 the patient was treated in two consecutive summers by local applications of silver nitrate and has not been seen for a year past, but she still has bladder symptoms. In Case 13 the patient has had systematic treatment for more than a year with instillations given at home by her daughter, 1 ounce of 1:500 silver nitrate solution being used twice a week. In addition, she has had occasional applications of 10 per cent. silver nitrate solution. She has a small ulcer on each lateral wall rather near the base, and an edematous, angry, red area across the anterior wall of the bladder. The ulcers have been temporarily improved and the symptoms apparently benefited by the use on two occasions of the actual cautery wire.

The other fifteen patients have all had excision of the diseased area. In Case 9 the patient died. I had observed in the after-treatment of three successive patients operated on before Case 9 that there was complete incontinence for three or four days after operation, all the urine escaping through the urethra in spite of the suprapubic drainage tube. This led me in Case 9 to omit the drainage tube.

The resected bladder when closed held about 1 ounce. The nurse failed to report that the patient was not having incontinence, and she was in great distress for seven hours until catheterized of 50 c.c. of urine. After the first catheterization the patient had incontinence.

At the operation, the peritoneum was so adherent over the posterior wall of the bladder that it was found necessary to do the transperitoneal operation and remove with the posterior bladder wall an area of peritoneum 2 by 4 cm. in extent. Great care was taken in closing the bladder, a combination interrupted and whipped suture No. 2 chromic twenty-day catgut being used instead of my usual closure with a running lock stitch suture. The first sweep with the interrupted portion of each suture took in all coats of the bladder, and the second sweep or whipped portion of the suture buttressed in the outer coats of the bladder. After closure of the bladder with a series of these sutures, the peritoneum was closed with a running suture, and then the line of bladder sutures was reinforced with a generous layer of fat and fascia. Two gauze and rubber wick drains were placed down to the bladder, and the suprapubic wound was closed about these.

In spite of the overdistention of the bladder on the first day, the patient seemed to do well until on the night of the second day, there being incontinence after the first catheterization and evidence of bladder leakage in a free suprapubic drainage of urine. Thirty-six hours after operation there was bleeding from the suprapubic drainage tract. This seemed to cease after twenty-four hours, the patient losing considerable blood in this time, and the pulse rising from 100 to 130. On the fourth day the bleeding having apparently ceased, we removed the two suprapubic drains and there was a gush of what appeared to be a pint of urine and blood which had been accumulated under pressure in the prevesical space. We considered that this gush of fluid under pressure argued for an intact peritoneal closure. The patient did not improve after removal of the drains, and eight hours later, on the evening of the fourth day, there was considerable distention

Fig. 14.—Section of bladder wall removed in a recent case. Arrow points to edge of mucosa 1 cm. from the sphincter urethrae. The area of edema seen over most of the specimen extended to within 2 cm. of the sphincter; *u, u*, two ulcers in the right and left vertex. An ulcer on the posterior wall at *p* was excised at the operating table and given to the pathologist for maceration and an attempt to grow anaerobic organisms. Bladder wall at *p* and over most of the area of edema about 1 cm. thick as contrasted with the 2 mm. normal thickness at the edges of the specimen.

and eructation, and the patient occasionally brought up brownish fluid. The temperature, 101.6 in the morning when the drains were removed, had dropped to 100 at night, but the pulse seemed weaker. The leukocytes were 30,000.

The patient was rather stout, and we could not determine whether she had a low grade peritonitis or an intestinal ileus. The wound was opened, and through a small forceps opening through the fatty omentum there was found no evidence of peritonitis. On the passing of a stomach tube there was a great gush of air, and almost a quart of brownish fluid with suggestive fecal odor was removed. The stomach was washed again at midnight and on the morning of the fifth day, and after this washing an ounce of castor oil was left in the stomach. Two copious stools at noon showed considerable oil. The patient seemed better until the morning of the sixth day, when she became irrational and unconscious, and she died at 4 p. m. that day.

In the other fourteen excision cases, with the exception of Case 5, the patients have had wonderful recoveries from their previously miserable existence, and they form a group of most appreciative and grateful patients. In Case 5, the edema area disclosed at operation spread low over the base of the bladder, and to have excised beyond the edema line would have necessitated working close to the ureters. This was the first case in which I performed operation, Dr. Kelly having operated in Case 1 seven years previously, after I had made the diagnosis of ulcer and had treated the patient by various methods for more than five years.

Case 5 being my first operative case for this type of ulcer, I did not appreciate the significance of the widespread edema area, and undoubtedly left a portion of this. Our operation experience with these cases and subsequent microscopic studies have taught that the edema is coexistent with the chronic infiltration of the bladder wall. It is probable that the patient will again develop ulcer and symptoms if the excision fails to include the entire edema area (Figs. 13 and 14).

In Case 5, the patient made a splendid recovery, and seven weeks after operation reported: "I have been home three weeks now and am still improving. I get up about three times every night; occasionally it has been only once. When I came home I was voiding only 3 ounces, and now I have increased to 7 ounces." A week later the patient wrote me, in distress, that her old pains had returned and that the bladder capacity had dropped to 4 ounces. She is still suffering, four years after operation, and refuses to come back for another excision, although her physician reports that there is only a "tiny red spot" in the bladder. As incredible as it may seem, this patient was operated on one year after my bladder operation for "a fibroid tumor pressing on the left side of the bladder." Among the four operations she had had before I saw her was one abdominal section for "fibroid tumor pressing on the bladder."

As stated before, the cystoscopic picture in one of these bladder ulcers is not sufficiently characteristic to justify one in operating on a patient on the evidence gained by cystoscopy alone, and I trust that this paper will not lead to the removal by operation of bladder ulcers that should be treated by other methods.

I am becoming rather more radical in advising the removal of other types of ulcer. At times we are able to cure a general cystitis, with the exception of one ulcer area which seems to persist indefinitely in spite of various local applications. If such an ulcer is located in the upper portion of the bladder where it is easily accessible to operation, and if it refuses to heal even after the use of the actual cautery, I consider its excision a more conservative and much more certain method of treatment than the making of a

vesicovaginal fistula. Another type of persistent chronic ulcer, fortunately of rare occurrence, is one which persists for years after the removal of a tuberculous kidney. It is not so uncommon to have cystitis symptoms persist for one, two or three years after the removal of a tuberculous kidney, but this is usually due to infiltration about the ureterovesical region, which eventually heals. On rare occasions a solitary ulcer will persist in the vertex of the bladder and perpetuate the patient's preoperative symptoms of cystitis indefinitely. Such an ulcer should be excised.

Given a patient who has had years of bladder misery in spite of many courses of systematic bladder treatment, who has microscopically clear urine, free from infection, but containing microscopic blood and leukocytes, and whose cystoscopic examination reveals one or more minute ulcers, one is justified in making a diagnosis of this type of ulceration.

Usually patients with a minute bladder ulcer have not had systematic treatments because the ulcer has been overlooked in spite of their ulcer history. For this reason it will be safer for those who have not had experience with these cases first to subject the patient to more conservative forms of treatment before making a final diagnosis. I have thus treated the patients on whom I have not operated: Case 2, nine years; Case 10, during two successive summers; Case 13, over one year. In none of them has there been more than an amelioration of the symptoms, and in none has there been an appreciable change in the bladder lesion. Of the patients on whom I operated, I treated with like results Case 1, intermittently for five years, Cases 7 and 8 systematically for more than a year, Case 11 once a week for two months, and Case 9 intermittently for a year.

Except in Case 1, the diagnosis was made promptly, and the long courses of treatment before operation were due in Case 9 to the patient's reluctance to undergo an operation, and in the other cases to a deliberate postponing in an effort to test the effects of various methods of treatment.

The diagnostic features in this form of bladder inflammation are so clear in most cases that one is not justified in allowing the patient to continue suffering under ordinary methods of treatment unless the patient objects to operation or has some other factor, such as advanced age, or heart or lung disease, to contraindicate operation.

No better evidence can be adduced as to the extent of the suffering of these patients than to see them willingly accept another operation as the possible way out, even when they have been through from one to five operations with similar hopes of relief.

My conservative treatments consisted of instillations, local applications and the use of the high frequency current and the actual cautery wire. The instillations consisted of as strong solution of mercuric chlorid (usually 1:10,000) or of silver nitrate (usually 1:500) as the patient could bear, 1 ounce of the solution being left in the bladder twice a week. This treatment can be carried out by a nurse at the patient's home. Local applications were made of some strong cauterizing agent about every four to six weeks. For these we use pure phenol (carbolic acid), the silver stick or, as has usually proved most satisfactory, the 10 per cent. silver solution. The high frequency current has been tried twice on one patient and once on another, and both patients refused to have further

treatments by this method because of the after-pain. In contrast to this result it is rather curious that the actual cautery wire, used on two of these patients, has given satisfaction as regards the amelioration of symptoms, but we have not seen an ulcer of this type show increased tendency to heal after the use of the cautery; while for the chronic ulcer following an infected cystitis, this has usually proved a valuable method for getting the deep slough necessary for final healing.

When we study the excised specimens microscopically, and see the widespread chronic infiltration and inflammation of all coats of the bladder wall, we are not surprised that excision is the only method of cure.

OPERATIVE TECHNIC

The excision is done through a suprapubic incision. To facilitate the finding and handling of the bladder, it is left full of air if cystoscopy has just been done in the knee-breast posture, or it is distended with sterile fluid just before operation.

If possible, the operation is kept extraperitoneal. This can be accomplished in those cases in which the disease is confined to the vertex and anterior wall. If the disease involves a portion of the bladder covered by peritoneum, it is possible in some cases to separate the involved peritoneum; but more often the attempt at separation results in an opening into the peritoneum or into the bladder.

Experience has taught that the best method of separating the peritoneum is first to brush back the vertex portion as far as it will go with ease; then, instead of persisting in trying to carry this separation farther down the posterior wall from the vertex, one should next free the lateral walls and attempt to get down near the ureter regions on both sides. From near the base in the lateral or ureter regions one can work toward the midline about the base of the bladder; and after separating the peritoneum in the uterovesical region, one can finally come forward on the posterior wall until the entire peritoneum, except the urachus portion, is freed. The freeing of the urachus portion can be done with less danger of entering the peritoneum if done as the final stage in the separation than if undertaken as the first step.

If it is found at any point that the peritoneum cannot be separated from the bladder wall, it is best to continue the separation at other points until all the peritoneum that will separate easily is free. Then one enters the peritoneum with a free opening and packs off the peritoneal contents with an abundance of hot salt solution gauze pads before entering the bladder. One always tries to determine by cystoscopy where the first opening into the bladder should be made in order to avoid the inflammatory area.

If the redness and the edema of the bladder mucosa have extended over the anterior wall of the bladder to within 1 or 2 cm. of the sphincter urethrae, it is well to make the first incision as near the urethra as can safely be done without taking the sphincter.

If the preliminary cystoscopic studies have shown the inflammatory area to be located to the right of the midline, one makes the first incision in the midline or slightly to the left. In spite of the most careful cystoscopic studies, one usually finds at operation that the inflammatory area, as indicated by the edema, is more widespread than anticipated, and one may open through the inflammatory area instead of opening through normal mucosa near one edge of the diseased area as intended.

Every portion of the bladder wall should be carefully inspected before the end of the operation in order to make certain that all the edematous wall is removed.

On opening the bladder, one may first fail to see the one or two small ulcers before seen by cystoscopy, because the edema arising during the preliminary freeing of the bladder may elevate the mucosa about the ulcers in a manner almost to hide them. Usually the preliminary trauma has caused the ulcer or ulcers to bleed so they are easily found on opening the bladder. A suction apparatus should be ready to keep the bladder clean during operation, and sponging should be avoided. The mucosa should be handled as little as possible in order to preserve its microscopic features. In order to preserve its natural outline for microscopic study, the specimen should be pinned to a thin board as soon as removed, being stretched to its former size as nearly as possible, and then dropped at once into the hardening solution.

During the bladder operation, all blood vessels of the wall should be carefully clamped and either tied at once or at the close of the excision. The bladder vessels are liable to retract into the wall and cease bleeding temporarily, only to give serious trouble later. The presence of urine is liable to interfere with coagulation and determine a serious bleeding from a small vessel that would promptly close in another location.

In Case 3, oozing of blood from the suprapubic wound began on the night of the third day, and continued intermittently for four days.

In Case 9, I attribute one of the chief causes of death to hemorrhage which began on the second day and continued for twenty-four hours. It was not the hemorrhage per se that played an important rôle in the patient's death; but the hemorrhage prevented me from starting the drains at a time when we feared that the patient had properitoneal accumulation of bladder fluid. This fluid under pressure probably got through the peritoneal incision, and caused a low grade peritonitis.

After excision of the diseased area, the bladder is closed by bringing the edges together with a running lock stitch suture of twenty-day formaldehyd catgut No. 2, leaving a slight opening in the vertex through which the mushroom retention catheter is carried and sutured to the bladder wall with a No. 2, ten-day catgut. In recent cases, I have used the combination interrupted and whipped suture as explained in the description of the closure in the report of Case 9, in which the patient died. This probably controls bleeding better than the lock stitch suture. The abdominal wall is closed except for a small opening to carry the rubber catheter and two cigaret drains, which are introduced down to the bladder wall.

These cigaret drains are removed at the end of forty-eight hours, and daily irrigations of the bladder with 1:10,000 solution of silver nitrate are carried on through the retention catheter, the strength of the solution being increased as the patient recovers. An ounce or two of the solution is introduced and immediately withdrawn through the catheter until several washings are made. This is best accomplished with the patient on her side.

The retention catheter is withdrawn on the tenth to the fourteenth day, after which the irrigations are given through the urethra daily until the bladder is free from infection.

2305 St. Paul Street.