ascertained that a current of steam passing through the tin pipe simultaneously with a current of drops of water produces no electricity of condensation, provided the air is not admitted at the same time. This result is, as far as the electricity of condensation is concerned, similar to those of (89.), in which the necessity of a good supply of air is fully seen.

I do not here give the details of the above experiments, being willing that they should form part of a paper which I hope shortly to communicate.

I am, Gentlemen,

Your very obedient Servant,

7 Prospect Place, Ball's Pond Road,
March 5, 1850.

REUBEN PHILLIPS.

XXXVIII. *On Quaternions; or on a New System of Imaginaries in Algebra.* By Sir WILLIAM ROWAN HAMILTON, LL.D., M.R.I.A., F.R.A.S., Corresponding Member of the Institute of France, &c., Andrews' Professor of Astronomy in the University of Dublin, and Royal Astronomer of Ireland.

[Continued from vol. xxxv. p. 204.]

88. **T**HE writer desires to put on record, in this place, the following enunciations of one or two other theorems, out of many to which the quaternion analysis has conducted him, respecting the inscription of *gauche* polygons in surfaces of the second order; without yet entering on any fuller account of that analysis itself, than what is given or suggested in some of the preceding articles. See the Numbers of the Philosophical Magazine for August and September 1849. And in the first place he will here transcribe the memorandum of a communication, hitherto unprinted, which was sent by him, in the month last mentioned, to the Mathematical and Physical Section of the Meeting of the British Association at Birmingham.

89. Conceive that any rectilinear (but generally *gauche*) polygon of n sides, $BB_1B_2 \dots B_{n-1}$, has been inscribed in any surface of the second order; and that n fixed points, $A_1, A_2, \dots A_n$, not on that surface, have been assumed on its n successive sides, namely A_1 on BB_1 , A_2 on B_1B_2 , &c. Take then at pleasure any point P upon the same surface, and draw the chords $PA_1P_1, P_1A_2P_2, \dots P_{n-1}A_nP_n$, passing respectively through the n fixed points (A). Again, begin with P_n , and draw, through the same n points (A), n other successive chords, $P_nA_1P_{n+1}, P_{n+1}A_2P_{n+2}, \dots P_{2n-1}A_nP_{2n}$. Again begin with P_{2n} , and draw in like manner the n chords, $P_{2n}A_1P_{2n+1}, P_{2n+1}A_2P_{2n+2}, \dots P_{3n-1}A_nP_{3n}$. Then one or other of the two following

Theorems will hold good, according as the number n is *odd* or *even*.

Theorem I. If n be *odd*, and if we draw *two tangent planes* to the surface at the points P_n, P_{2n} , meeting the two new chords, $PP_{2n}, P_n P_{3n}$, respectively, in two new points, R, R' ; then the *three points* RRR' shall be situated on one straight line.

Theorem II. If n be *even*, and if we describe *two pairs of plane conics on the surface*, each conic being determined by the condition of passing through *three points* thereon, as follows: the first pair of conics passing through BPP_{2n} , and $P_n P_{2n} P_{3n}$; and the second pair through $BP_n P_{3n}$ and $PP_n P_{2n}$; it will then be possible to trace, on the same surface, *two other plane conics*, of which the *first shall touch the two conics of the first pair*, at the two points B and P_n ; while the *second new conic shall touch the two conics of the second pair*, at the two points B and P_{2n} .

90. With respect to the *first* of the two theorems thus communicated, it may be noticed now, that it gives an easy mode of resolving the following *Problem*, analogous to a celebrated problem in plane conics :—To find the *two* (real or imaginary) polygons, $BB_1 B_2 \dots B_{n-1}$ and $B'B'_1 B'_2 \dots B'_{n-1}$, with any given *odd* number n of sides, which can be inscribed in a given *surface* of the second order, so that their n successive sides, namely $BB_1, B_1 B_2, \dots$ for one polygon, and $B'B'_1, B'_1 B'_2, \dots$ for the other polygon thus inscribed, shall pass respectively through n given points $A_1, A_2, \dots A_n$, which are not themselves situated upon the surface. For we have only to assume at pleasure *any* point P upon that surface, and to deduce thence the *two* non-superficial points lately called R and R' , by the construction assigned in the theorem; since by then joining the two points thus found, the *joining line* RR' will cut the given *surface of the second order* in the *two* (real or imaginary) points, B, B' , which are adapted to be, respectively, the *first corners of the two polygons required*.—That there are (in general) *two* such (real or imaginary) polygons, when the number of sides is *odd*, had been previously inferred by the writer, from the quaternion analysis which he employed. Indeed, it may have been perceived to be, through geometrical deformation, a consequence of what was stated in § XIV. of article 87 of this series of papers on Quaternions, for the particular case of the ellipsoid, in the Philosophical Magazine for September 1849. See also the account, in the Proceedings of the Royal Irish Academy, of the author's communication to that body, at the meeting of June 25th, 1849; in which account, indeed, will be found (among many others) both the theorems of the preceding article 89; the second of those theorems being however there enunciated under a *metric*, rather than under a *graphic* form.

[To be continued.]