EROSIVE VULVITIS

THOMAS LATANE DRISCOLL

P. A. Surgeon (Reserve), U. S. P. H. S.

RICHMOND, VA.

The disease of men known as erosive and gangrenous balanitis has been recognized for several years. The same disease occurring in women as erosive and gangrenous vulvitis is not generally recognized.

Three cases of this disease have come under my charge in the Department of Venereal Diseases of the United States Public Health Service. I have successfully isolated the etiologic organisms in each case.

ETIOLOGY

The organisms isolated in each instance were the typical spirochete and vibrio growing in symbiosis as described by Tunncliffe.

The spirochete averages from 5 to 30 microns in length, has very rapid motion, is gram-negative, and takes the ordinary dyes well. It has an especial tendency to grow in the more superficial of the diseased parts.

The vibrio, or fusiform bacillus, is about 2 microns in length and 0.8 micron in width, pointed at each end as one of its names infers. It grows singly or in chains, is gram-positive when carefully decolorized, and takes the ordinary anilin dyes. In my cases this organism seemed to penetrate the deeper tissues, and it was necessary to probe deeply with a capillary pipet in order to obtain a specimen for microscopic study. This would seem to be in accord with the anaerobic nature of the vibrio; i. e., reproduction occurs under anaerobic conditions.

The predisposing causes seem especially to be filth and prostitution, with attending frequent copulation and exposure, and unnatural sexual relations. All three women were inmates of the jail in Richmond, and all had been convicted for prostitution. The diseased genitals were extremely dirty, with a large amount of discharge from the focus as well as from the vagina, and presented ideal conditions for the growth of the specific organisms. In each instance there was extensive ulceration of the part with a slight amount of local edema. Two of the cases presented an inflammatory involvement of Bartholin's gland, while the third gave a history of such an involvement two years before.

SYMPTOMS

The first case (Fig. 1), the least severe of the three, presented an area of ulceration extending from the clitoris posteriorly to the margin of the anus. On the right side it was bounded by the labium minus, but this structure on the left side was partly sloughed off, so that only the anterior portion remained, the ulceration on this side extending laterally on the inner side of the left great labium. This lesion was erosive, with the edges somewhat turned outward, standing above the

Fig. 1 (Case 1).—Area of ulceration extending from the clitoris posteriorly to the margin of the anus.

ulcer itself and the surrounding tissues. A grayish yellow pus exuded from the surface of the ulcer as well as from the vagina, for the erosion had involved this canal for from 1 to 2 cm. inward; the pus was characterized by the same foul odor as that of erosive and gangrenous balanitis. The lesion was dark red, and when it was covered with the characteristic yellow pus it appeared quite similar to the color noted in an ordinary varicose ulcer of the leg. The left labium majorum was somewhat edematous and showed a very evident abscess

in Bartholin's gland, for the skin covering it was tense and shining, and the labium greatly swollen and fluctuating. A second similar erosion, but only about 1 cm. in diameter, appeared on the inner surface of the labium at its anterior end just before it merges into the mons veneris; it would seem from this and from the second case that the organisms were auto-inoculable.

The second case (Fig. 2), while relatively of medium severity, was ulcerated extensively, but the destruction had not advanced to such a stage as that noted in the third case. The main area extended from the interior margin of the symphysis pubis to the anterior margin

Fig. 2 (Case 2).—Area of ulceration extending from the inferior margin of the symphysis pubis to the anterior margin of the anus.

of the anus. Laterally, on each side the labia minora had been destroyed, while the posterior portion of both labia majora had sloughed off and the remaining anterior portion had been extensively undermined, so that in the standing posture these structures hung down slightly, giving somewhat the appearance of a scrotal sac. Besides undermining the labia majora, the ulcer covered the inner side of both labia and posteriorly passed laterally over the ischiorectal fossa. The clitoris was involved and had partially sloughed away, while the vagina was involved inward for a distance of from 1 to 3 cm.,

as in the preceding case. The appearance of the ulcer and the character of the pus were the same in this case as in the first, but here there was a bilateral involvement of Bartholin's glands, which presented a tendency toward suppuration also. Three smaller ulcerations were presented: one on the outer side of the left great labium, and the other two, one on each side just posterior and lateral to the anus, on the medial margins of the buttocks.

The most severe case (Fig. 3) presented an almost complete destruction of the vulva. The labia and clitoris had completely

Fig. 3 (Case 3).—Area of ulceration extending from base of clitoris anteriorly to tip of coccyx posteriorly and laterally on each side as far as outer limits of labia majora.

sloughed away, leaving a slit-like ulcer between the legs, with the vagina and anus quite similar in appearance at their outlet, presenting a depressed area in the slit. The ulcer itself extended from the base of the clitoris anteriorly, to the tip of the coccyx posteriorly, and laterally on each side as far as the outer limits of the labia majora, which had completely sloughed off except for a short distance just posterior to the mons, where they projected backward to about the anterior margin of the vestibule. Bartholin's glands had been totally destroyed on both sides. This ulcer was of the same general char-

acter as noted in the first instance, except for its obviously more destructive nature; the pus had the same characteristics.

In all three cases the inguinal lymphatic glands were involved. On palpation they were hard, nodular and moderately enlarged, but there was no suppuration. No systemic changes of pathologic significance could be noted.

Locally, there were symptoms that pointed to a chronic process. Case 1 had existed for eight months; Case 2 was of two years' duration; while Case 3, the most severe, began three years before. Local pain and discomfort with violent itching seem to be constant factors. Dysuria was more or less marked. All the cases had had a foul smelling leukorrhea since the date of onset, and the discharge preceded the vulvar condition in at least one instance.

Generally, there appeared to be a slight degree of toxemia, and the patients showed some degree of melancholia.

In each of the three cases the Wassermann reaction was negative. There was no response to antisyphilitic treatment, including arsphenamin treatment. In each the characteristic spirochete and fusiform bacilli were isolated from the serum of the ulcers.

DIAGNOSIS

Any persistent ulceration with a negative Wassermann reaction and the general characteristics noted above should be regarded as suspicious, and some of the serum should be obtained for bacteriologic study. If the serum shows spirochetes and fusiform bacilli the diagnosis is conclusive. The infection might, of course, be superimposed on other lesions of the genitals.*

BIBLIOGRAPHY

- Carter: *Lancet*, London **11**:1128, 1910.
 Corbus: *J. A. M. A.* **60**:1769, 1913.
 Daniels: *Lancet* **11**: 1648, 1910.
 Darre and Delaunay: *Gaz. d. hôp.*, Paris **77**:657, 1904.
 Guenther, E. E.: *Am. J. Obst.* **49**:373, 1904.
 Heller: *Arch. f. Dermat. u. Syph.*, Orig. **112**:401, 1912.
 Schiller, H.: *Am. J. Obst.* **77**:398, 1918.
 Szasz, H.: *Monatschr. f. Geburtsh. u. Gynäk.* **17**:994, 1903.

* Thanks to Drs. E. G. Hopkins and Raymond Vonderlehr.