Archives of Dermatology and Syphilology

VOLUME 3

JANUARY, 1921

NUMBER 1

A CONTRIBUTION TO THE STUDY OF EPIDERMOPHYTON INGUINALE

NOTES ON AN INQUIRY INTO THE MORPHOLOGY OF THE CAUSATIVE FUNGI; A CLINICAL AND LABORATORY STUDY OF THIRTY-SEVEN CASES OF DERMAL SCALING

GROVER W. WENDE, M.D., AND KATHARINE R. COLLINS, M.D.
BUFFALO

The higher fungi were credited with being the causative factors in the production of human disease processes long before the recognition of the important rôle played therein by the lower fungi by Langenbeck, 1839; Schoenlein, 1839; Pasteur, 1862, and Koch, 1882. During the era in which the causal relationship of bacteria to human disease was being developed, interest lagged in regard to the etiologic importance of the higher fungi. Since 1900, the importance of the higher fungi in the production of pathologic states has received more attention and their etiologic rôle has been more clearly recognized. That the integumentary system is most often the site for their implantation and growth is obvious and is shown in the large number and variety of skin lesions due to them.

The findings of bacteriologists and dermatologists lead one to consider that a wide distribution of dermatologic lesions due to the higher fungi exists; that many different species are causative factors, and that certain clinical types of such lesions may be produced by species morphologically dissimilar. On the other hand, one is constrained to believe that skin lesions clinically dissimilar may be due to the same species of fungi which at times may be virulent, at other times inert; at times pathogenic, at other times saprophytic. Indeed, Sabouraud claims that a change of country frequently determines a change in virulence, so that a change in local environment may well cause a change in morphologic development. All this may account for the seeming discrepancies in the discussions of various authorities, for the lack of sharp-cut distinctions, and for an amount of indefiniteness and obscurity that has prevented proper appreciation and clear understanding of interesting and easily recognized skin lesions—lesions that fre-

quently are permanently eradicated with great difficulty, that often are chronically recurrent, and that actually are unsightly and annoying but not dangerous nor deforming.

Probably no group of skin lesions has recently excited greater interest and investigation than those produced by tinea fungi. In 1844, Gruby described two varieties of fungi in ring form—large spore and small spore. This observation remained many years forgotten until it was revived and verified by Sabouraud. While *Tinea trichophyton*—large spore—lesions are common in tropical and subtropical regions, they are apparently not so common in temperate regions. Castellani, in 1905, while in Ceylon, described a fungus from human skin lesions that he called *Trichophyton cruris*; the fungus was independently reported in France by Sabouraud in 1907, and was named *Epidermophyton inguinale*. Castellani¹ claims that, according to the laws of nomenclature, this fungus should be called *Epidermophyton cruris*. *Epidermophyton inguinale* differs from the other tinea in that it does not attack the hair or hair follicles and in that the color of the colonies on Sabouraud's medium is lemon or orange yellow.

Since the publication in this country of the papers by Ormsby and Mitchell,² and by White,³ we have been particularly interested in the skin lesions said to be produced by *Epidermophyton inguinale*. We have endeavored to determine: whether the lesions on the flat skin surfaces and those in the skin folds, that clinically are so different, are due to different varieties of the same fungus or to a single variety morphologically modified by the slightly different environment presented by the various sites; and, if due to a fungus, whether the fungus is a modified trichophyton or a modified epidermophyton.

SITE OF EPIDERMOPHYTON INGUINALE LESIONS AND DURATION

From the designation *Epidermophyton inguinale*, the inguinal region would be considered the most frequent site of the lesions, but the few observations made by us indicate that the skin lesions due to the epidermophyton may be found on any part of the body. Our observations corroborate the statement made by White in his report on a many times larger number that in the United States the inguinal region is not so frequently the site as are other parts. In all probability the name was given to this fungus because it was first recovered from lesions occurring in the inguinal region. In our small series we found epi-

1. Lancet 1:945 (May 1) 1920.

2. Ormsby, O. S., and Mitchell, J. H.: Ringworm of Hands and Feet. J. A. M. A. 67:711 (Sept. 2) 1916.

3. White, C. J.: Epidermophyton Infection, J. Cutan. Dis. 37:501 (Aug.) 1919.

Fig. 1.—Vesicles appearing in groups and blebs; slight desquamation; resembles pompholyx.

dermophyton lesions at the following sites: face, one case; axilla, one case; hands, one case; abdomen, one case; nates, one case; anus, two cases; groins, five cases; thighs, two cases; legs, four cases, and feet, eleven cases.

Our findings also seem to corroborate those of White in regard to sex, age and duration. The larger number of lesions were found in males; they occurred during the third and fourth decades; the duration was in periods of months rather than days or years. Our inquiry did not go into the possible relation of occupation or into the possible source of the infection.

APPEARANCE AND TRANSMISSIBILITY

In general, the appearance of the lesions in our series may be summed up as areas of scaliness varying in size from that of a split pea to 4 and 5 inches square or even larger. The lesions were more or less sharp in outline, with thickening of the epidermis without inflammation of adjoining tissue, although underneath the scales there was some redness. In those areas in which the skin was naturally moist, as between the toes, the epidermis was moist and soggy. The color of the lesions in the moist area was usually a dirty white while in the dry areas it was more or less gray, red or brown. Distributed through portions of the lesions deep seated vesicles were frequently seen; often at the inception of the trouble blebs strikingly resembling pompholyx were present. Itchiness and hyperhidrosis were present, while in the folds and moist areas an unpleasant odor, almost a bromidrosis, occurred. Occasionally the process seemed to be influenced by the season, being worse in summer, better in winter.

In none of our cases could it be determined that autoinoculation took place, although the larger number of patients had multiple lesions in widely distributed areas. In regard to the transmissibility of *Epidermophyton inguinale*—we have under observation four members of one family each presenting proved epidermophyton lesions. The father has had patches in the groin and on both plantar surfaces for eighteen years; the mother has had the same condition in the groins and on both plantar surfaces for seven years; one son has had the condition on both plantar surfaces as well as interdigitally for one year, and another son has had it for nine months in the same locations as his brother.

CASES STUDIED BY AUTHORS

In the plan of study cases were selected from the ordinary day's work and short clinical records made. Scales and scrapings were obtained from the various lesions and subjected to microscopic examination for the presence of spores and mycelia; if possible, some of the hairs present in the lesions were included. In cases showing spores

Fig. 2.—Distinct dirty, white, soggy, intertriginous desquamation and thickening; desquamation over plantar surface with deep-seated vesicles.

or mycelia an effort at identification of the fungi was made by means of cultures from the specimens obtained both in vitro and in vivo. In order to check the work and as controls and for differential study specimens were taken from similar lesions due to other known fungi and from lesions of similar character not caused by fungi.

Diagnoses.—All the cases selected for observation presented skin lesions showing scaling with or without vesication, with or without slight redness and with little or no itching. Our series consists of thirty-seven cases, of which these clinical diagnoses were made: acne, two cases; blepharitis, one case; callus, two cases; dermatitis herpetiformis, two cases; eczema, five cases; folliculitis agminate (Hartzell) or hyphomycetes granulosum (Schamberg), one case; epithelioma, one case; neurodermatitis, one case; pityriasis rosea, two cases; tinea capitis, eight cases; tinea cruris, one case; tinea trichophytina, three cases; epidermophyton inguinale, eight cases; not diagnosed, one case.

Of the twenty-one cases which clinically were diagnosed as due to fungi, nineteen were found to present spores or mycelia of which thirteen may be classed as epidermophyton inguinale. Of the eight cases clinically diagnosed as epidermophyton inguinale, the cultures showed six to be due to that fungus. One case diagnosed as eczema proved to be due to the epidermophyton. Of the three cases diagnosed as trichophyton, all were proved to be epidermophyton. The case not diagnosed was shown to be due to the epidermophyton. The case diagnosed as agminate folliculitis proved to be due to epidermophyton. In the case diagnosed as tinea cruris the laboratory reported negative findings. From this analysis the clinical recognition of epidermophyton inguinale lesions is fairly easy, although unquestionably frequently overlooked because the lesions so imitate other skin conditions that careful and more extended examination of the patient is not made with the possibility of epidermophyton as the etiologic factor in mind.

Cultural Characteristics of Epidermophyton Inguinale.—Sabouraud, using the culture medium that bears his name, asserts that the peculiar cultural characteristics of the cryptogam, *Epidermophyton inguinale*, which is but a representative of the trichophyton group, while not making the recognition easy, make it less difficult; he also states that the fungus is isolated with comparative ease. He says that its characters and differences may be classified under three heads: (a) original appearance, (b) rapid pleomorphic transformations, and (c) organs. In his descriptions he states, under the heading appearance, that the culture growths show a powdery surface radiating in form, the center being umbilicated and covered by an embossment of the culture growth, in color a grayish-yellow with an areola of citrin hue, and that these forms are reproduced on proof mediums. Under the heading pleo-

Fig. 3.—Large desquamating yellowish-brown areas on flat surfaces and in moist fields; history of vesication in axillae and under breasts. The condition has existed for twelve years; at times it has been almost absent, at others even more pronounced than shown. This case is not included in the thirty-seven systematically studied, but laboratory findings proved the presence of *Epidermophyton inguinale*.

morphism he says that the growth on sugar or glycerin mediums is especially characteristic of *Epidermophyton inguinale* in that after three or four weeks a white duvet of absorbent cotton-like tufts gradually covers the entire surface of the mediums; and that this degenerating form growth invariably is reproduced on reseeded in sugar mediums, while the original may be rejuvenated by planting in proof mediums like peptone water. Under the heading organs, he says that microscopically the growth presents enclosed clubs or asci, pediculated laterally or terminally on the mycelia filaments and usually arranged in banana-like bunches. Sabouraud further asserts that *Epidermophyton inguinale* is the only trichophyton cryptogam known to present these characteristics.

Technic Used by Authors.—Sabouraud's technic was followed throughout the laboratory study of the thirty-seven specimens examined. The investigation consisted of four steps: (a) the usual microscopic examination of the scales and hairs for spores, mycelial threads and fungal organs; (b) the direct making of two separate plants in Sabouraud's medium of the scales found to contain spores—these plants were designated "primary and secondary"—and the reseeded of each through ten to twelve generations which were designated "subsequent" cultures; (c) the microscopic examination of the mycelia and organs produced from each culture, and (d) the inoculation of guinea-pigs with the culture growth diagnosed as *Epidermophyton inguinale*. When an inoculation was made the hair of the guinea-pig was cut short, and the closely trimmed surface cleansed with alcohol and scarified, after which the macerated culture growth was rubbed in freely. As a control the hair was clipped at another location remote from the inoculation and the cut surface cleansed and scarified but not inoculated.

The stain used for the microscopic examinations from the first twenty-seven cases was methylene blue; for the remaining ten Ehrlich's triple-stain was used. In our opinion Ehrlich's stain gave the better picture.

The microscopic examination of the original scrapings from the thirty-seven cases showed the presence of spores in twenty-six, of mycelia threads in three and of fungal organs (asci) in five; spores were absent in thirteen cases.

Results.—The primary and secondary plantings of the twenty-six cases showing the presence of spores resulted in the production of fungal growths in twenty-four, while in two no growth was secured. The twenty-four molds were identified thus: *Epidermophyton inguinale*, 13; *Microsporon audouini*, 7; *Microsporon minutissimus*, 1; one unidentified mold; *Oidium*, 1 (later observation indicates that this was an epidermophyton); *Aspergillus*, 1 (the aspergillus was a common

contamination in most of our cases; this was corroborated in the animal inoculations; this fungus bears no relation to the etiology of the lesion. Most investigators report some type of the penicillium as a contaminating fungus; with us it was of rare occurrence).

Of the thirteen cases identified as *Epidermophyton inguinale*, the microscopic examination of the original specimens showed the presence of mycelium in three and of asci in five, spores in chains or rows in four and crescentic asci in five, one of which was also septated.

Fig. 4.—The various forms of mycelia and asci seen microscopically in the cultures of the epidermophytos, rubrum and citrin; 1, pyramidal asci; 2, fusiform and crescentic asci.

The colonies in the upwards of 312 “primary and secondary” and “subsequent” cultures all showed a central acuminate growth surrounded by a distinct flattened areola which was again surrounded by the duvet as a lighter fringe; the only exceptions were those derived from one case in the colonies from which the areola was absent. In one case the areola was flat, an atypical growth; this case was off-type all the way through.

The characteristic pleomorphism was demonstrated in the matrass, which in all of the cultures was at first yellow or orange in color. This yellow color persisted in seven of the primary plantings and in four of the secondary plantings. In one both the primary and secondary plantings rapidly became rose red; this case seemed to answer to the description of the "*Epidermophyton rubrum*" of Castellani.¹ One case after several weeks in both primary and secondary plantings became green. In four cases of the primary plantings and in seven of the secondary plantings the matrass became black. In the subsequent cultures the same colors were respectively obtained as in the primary and secondary.

In all cultures the duvet began as an absorbent cotton-like growth, with the exception of one case in which the hyphae presented a

Fig. 5.—Forms of mycelia and asci seen microscopically in cultures of the epidermophytons, noire and noire-gris. The asci are tinted yellow to brown.

bristle-like growth and in one case in which the hyphae were short. The color of the duvet in all cultures at first was white. This persisted throughout in the red type and in the green type. In the yellow type after ten or twelve generations the duvet took on a salmon-pink color. In the black type the duvet passed through a grayish-green to a mouse-gray color, with the exception of one case in which it became soot black. Here also is evidence of pleomorphism.

In all cultures the mycelial threads were septated with occasional nonseptated, highly granular branches. The asci were given off terminally and laterally, often in bunches like bananas; they bore from two to eight or more spores. The mycelium in subsequent cultures was often a more or less homogeneous mass and not well differentiated as

to structure; the same condition was noted in two of the original specimens submitted; in the case in which the duvet became soot black the mycelial threads were clearly differentiated.

Except in two cases, pyramidal forms of asci were present in the primary and secondary plantings. In the seven black type cultures the pyramidal form persisted throughout all subsequent cultures. In the eight yellow type cultures, after a few generations, the pyramidal asci were lost. In twelve cases, fusiform, crescentic and oval asci occurred and were the pleomorphic forms. In the green type case only the oval asci were found. In one yellow type case the pyramidal forms of asci only appeared after the fourth reseedling to disappear in the following one. In seven cases of the red and yellow types the asci at times took on a diminutive form without change in character. One case, on the fourth reseedling, gave small spores, but in the following

Fig. 6.—Forms of mycelia, asci and spores seen microscopically in cultures of an atypical epidermophyton (laboratory case No. 8). The small slender mycelium and spores predominate. The asci are few in number.

ones reverted to large type. In the black cultures the asci took on yellow to brown tints.

The laboratory designation of "*Epidermophyton inguinale*" was based chiefly on the microscopic morphology. The pleomorphism in our series was marked and interesting, and a more careful and extended investigation should be made to account for it. Our systematic examinations were made on Sabouraud's medium. We used proof agar and beer-wort agar, but they did not give different results. All changes herein described took place in the same batch of mediums. Peptone

water was used for keeping the cultures. The formation of oxalate crystals in the medium was never noted in cases of the epidermophyton group, but was frequently observed in cultures of the penicillium and the aspergillus groups.

One interesting development in our work, which we believe has not been noted heretofore by other observers, is the presence of two types of the fungus in the same lesion. The two types gave identical skin reactions in the guinea-pig. Four of our cases showed this double infection.⁴

The inoculations into guinea-pigs with macerated pure cultures from each of the thirteen cases designated "*Epidermophyton inguinale*" gave rise to scaling in different degrees, and the hair did not grow in until after several weeks. Preparations of the scales from the inoculated

Fig. 7.—Microscopic appearance of scales treated with 30 per cent. potassium hydroxid and stained; 1, disintegrated scales—spores in rows and asci; 2, scales, slightly disintegrated, filled with spores; 3, scales, disintegrated, spores and asci and nondifferentiated mycelium; and 4, scales filled with spores and differentiated mycelium.

guinea-pigs showed mycelium and asci. The results from inoculations from both the black-gray type and the yellow type were identical. The findings were the same whether the two types had been isolated from the primary and secondary plantings from the same lesion or from the primary and secondary plantings which gave only one type. The pic-

4. In a recent personal communication from M. Sabouraud, he states that only one type is found in the same case and in the same lesion.—"Dans les cas de Trichophytie on ne trouve qu'une espèce sur un malade et jamais plusieurs Trichophytes sur un même malade et dans la même lésion."

ture presented in guinea-pig controls in which clipping and scarifying alone were done was very different; there was no scaling and the hair grew in quickly. In the guinea-pigs inoculated after the same manner with cultures of aspergillus the hair grew back in two days; there was no scaling. The guinea-pigs inoculated with aspergillus invariably died on the third or fourth day, whereas those inoculated with the "*Epidermophyton inguinale*" lived.

CONCLUSIONS

1. Recognition of the skin lesions of *Epidermophyton inguinale* presents little difficulty; these lesions occur relatively often among scaly skin diseases. Parts of the body other than the inguinal region present original lesions with greater frequency than the name would imply. This condition is well shown in Figure 3 in which there is a general distribution over the entire body; the lesions, however, are confined to the anterior surfaces and are found on the face extending down over the neck, in the axilla, over the shoulder down on the arm to below the elbow, on the trunk below the breasts, about the umbilicus, and on the toes; the groin lesions occurred as three slightly scaly areas, each about the size of a quarter. The designation is, therefore, somewhat misleading when considering lesions at other locations than in the groin.

2. Epidermophyton lesions vary so much in appearance that errors in diagnosis are easily possible unless reasonably full and careful clinical examination and study are made. The character of pleomorphism is as applicable to its lesions as it is to its laboratory growth; environment seems to affect the activity and character of both lesion and growth. In our cases the lesions of flat surfaces and of the folds were due to the same type of fungus.

3. The problems of autoinoculation, sources of infection and conditions favoring human implantation and growth should receive early study and solution, as the frequency of the disease indicates increasing virulence and possibly the development of greater pathologic importance.

4. Laboratory recognition is neither intricate nor difficult. It requires no apparatus not found in every modest clinical laboratory.

5. The characteristic appearance, pleomorphism and biologic reaction are readily secured, and are distinct and easily recognized. The conditions determining the pleomorphism should be further studied.

6. The multiplicity of laboratory growth types without apparent clinical pathologic variability has not before been reported. This presents an extremely interesting problem that should receive further investigation to determine whether it is of any etiologic value as well as mycologic importance. -