HEMOCHROMATOSIS

A REPORT OF THREE CASES AND A DISCUSSION OF THE PATHOGENESIS

T. P. SPRUNT, M.D.

BALTIMORE

Hemochromatosis is the name given by von Recklinghausen¹ to a rare disease of metabolism, the etiology of which remains wrapped in obscurity. In the English literature there are few articles on this subject, which is one of great interest pathologically. Therefore, the three cases of this series are reported in detail with the impressions which I have derived from the study of them, in the hope that they may be of aid in the ultimate solution of the problem.

SUMMARY AND COMPARISON OF CASES

The three cases here reported afford an excellent opportunity for the study of the disease. The first case is one of simple hemochromatosis with the complication of tuberculous peritonitis forming a prominent part of the clinical picture. The other two cases are quite typical of the more frequently described form usually designated as bronzed diabetes. In all three cases there were the associated signs of asthenia, enlargement of the liver, and skin pigmentation which served to make the diagnosis clear in the two cases in which diabetes was present. In Case 1, without diabetes, the diagnosis was made more difficult by the presence of the tuberculous peritonitis, but in this case, too, the association of the signs just mentioned was considered very suggestive of hemochromatosis.

Alcohol cannot be cited as an important factor in the etiology of these cases, although none of the patients was a total abstainer. In Cases 1 and 3 there is a history of attacks of dysentery which occurred, however, at least ten years before there were any symptoms of hemochromatosis.

In none of the cases had the bronzing of the skin or the liver enlargement been noticed by the patient before entering the hospital. The skin pigmentation increased noticeably in all of them while under observation. Pigmentation of the mucous membranes was not present.

The onset in the first case was rather insidious. It was more abrupt in the other two, the first symptoms being those of diabetes. The duration of the disease in each case after the appearance of the first symptoms was twelve, ten and five months, respectively.

It is noteworthy that in Cases 2 and 3 there occurred a bilateral acute parotitis which persisted for only a few days and cleared up entirely

1. Von Recklinghausen: Ueber Hämochromatose, Tagebl. d. Versamml. deutsch. Naturf. u. Aerzte, 1889, p. 324.

before death. In Case 3, there was a terminal *Staphylococcus aureus* septicemia.

The blood-counts show normal numbers of red cells or a slight diminution.

The pathology of the three cases is essentially the same, with minor, though interesting variations. The omission of certain organs in the microscopic descriptions of each case is due to the fact that different tissues were available for study in the different cases. The liver, pancreas, spleen, thyroid, kidneys, adrenals and lymph-nodes are described in all three. In each case, studies were made of other organs which were not available in both the other cases. The iron-containing pigment, hemosiderin, is found in great abundance in the secreting cells of all the glandular organs examined, including the liver, pancreas, salivary glands, stomach, Brünner's glands, serous glands in the bronchi, trachea and larynx, thyroid, parathyroid, glomerular zone of adrenal, prostate, and in certain cells in the kidney. In addition, it appeared in the cardiac fibers, in the connective tissue of those organs after the disintegration of the parenchymatous cells, in the lymph-nodes, in the spleen in small amounts, in the bone-marrow, in certain cells of the cutis, in cartilage cells, and in the endothelium of blood-vessels, especially of the sexual organs.

The abnormal iron-free pigment, hemofuscin, is present in considerable quantities only in Case 3, and is here especially noticeable in the walls of blood-vessels and large ducts and in the capsules and trabeculae of lymph-nodes.

In all the cases there is an increase in those pigments present in less degree physiologically. There is a striking accumulation of fine yellow pigment granules in the tubules of the testes and epididymes and in the vas deferens and seminal vesicles in the third case.

In each case there is an increase in the connective tissue of the liver, pancreas, lymph-nodes and spleen. In addition, this occurs to slight extent in the heart muscle in Case 2 and in the prostate and testicles in Case 3.

Case 2 suggests a slow deposition of pigment for a comparatively long time, while the picture in the third case gives one the impression of a rapid continuous process.

CASE 1.—*Medical No. 24,427. Simple Hemochromatosis without Diabetes. Tuberculous Peritonitis.*

C. M., a white man, a watchman, aged 55, was admitted to the Johns Hopkins Hospital, July 30, 1909, and died Sept. 5, 1909. Complaint: Pains in abdomen and frequency of micturition. Family history negative.

Past History.—The patient has with rare exceptions been well and strong. The periods of ill health, which he recalls most clearly, occurred during his residence in Cuba about ten years ago. He had the usual infectious diseases of childhood and, in addition, small-pox at 3 years of age, and malaria while in Cuba,

ten years ago. Following this malarial attack he was in a military hospital for three months.

Head.—Eyesight has been failing for the past eight years and more rapidly during the past twelve months.

Cardio-Respiratory.—Had some cough, otherwise negative.

Gastro-Intestinal.—Appetite has always been good; there is no nausea nor vomiting, no eructation of gas. Bowels rather irregular. While in Cuba, ten years ago, he suffered from an attack of dysentery during which his stools contained mucus and blood. This attack lasted about six or eight weeks altogether, with intervals of comparative freedom.

Genito-Urinary.—While in Cuba, before the malarial attack, the patient experienced considerable inconvenience from frequency of micturition. He passed large quantities of urine; was obliged to rise four or five times at night for this purpose. His face, legs and ankles became puffy and swollen. The polyuria has never entirely disappeared, but has never again been so marked as at that time.

He denies gonorrhea and syphilis.

Habits.—He drinks whiskey and beer in moderation, but does not get drunk. He has never done any heavy work.

Present Illness.—The onset was rather insidious. Patient says he has had more or less trouble for a year with increased frequency of micturition, passing small quantities of urine each time. There is no pain during the passage and the urine contains no blood. He has considerable abdominal pain, especially on coughing. This pain is not related in any way to the taking of food.

Physical weakness has been especially noticeable during the last three or four weeks. He cannot walk far without sitting down frequently to rest. He has some cough, but no shortness of breath. Bowels are irregular with alternating periods of constipation and diarrhea. The abdominal pain is especially severe on the right side after standing for some time.

Physical Examination.—The patient is a large man. The skin of the face is coppery-red in color, the vessels of the cheeks dilated, the left pupil slightly larger than the right. The breath is foul and the teeth in poor condition.

The skin over the neck and thorax is covered with small, reddish, elevated, discrete papules. The capillaries of the thorax are enlarged, especially in the lower part, where they form a striking band running across the lower front just above the costal margin.

The lungs and heart are normal.

The abdomen is distended and soft; the lateral veins markedly enlarged. There is tympany in the lower central portion, dull tympany and shifting dullness in the dependent flanks. No masses nor areas of tenderness can be demonstrated.

The liver is greatly enlarged; the dullness, beginning above at the fifth rib, extends 9 cm. below the costal margin where the edge is distinctly felt. In the midline it reaches a point half way between the umbilicus and the xyphoid cartilage.

The spleen is readily palpable 4 cm. below the costal margin.

Deep reflexes are demonstrable. The knee-jerk is more marked on the right.

Blood Examination.—August 6, red cells 5,100,000 per c.mm.; leukocytes 9,000; hemoglobin 105 per cent. (Sahli). The Wassermann reaction (blood serum) is negative.

Examination of the stomach contents reveals no special deviation from the normal findings.

On August 11, the pigmentation is noted as extreme over face, hands, forearms, areolæ of nipples, feet and legs, and of a deep coppery-brown color. There was no jaundice.

The patient's temperature, since admission, has ranged from 99 to 100 in the morning, and from 100 to 102 in the evening.

Examination of Stools.—On several occasions the stools were found to be small, semi-solid in consistence and of brown color. There were no macroscopic masses, no parasites, no mucus. The microscope revealed a normal picture. There was no occult blood.

On August 25, the abdominal distention had become so marked that paracentesis was performed and there were obtained 7,500 c.c. of a cloudy, straw-colored fluid, of which the reaction was neutral, specific gravity 1.015, albumin content 15 gm. per liter. The test for occult blood was positive. Microscopic examination of a centrifugalized specimen showed numerous red blood-cells, a few leukocytes and several large multinucleated cells.

At the second tapping, five days later, 550 c.c. of a similar fluid were obtained. A culture from this fluid remained sterile.

On September 2, the patient was drowsy and so weak that he could not feed himself. He complained of pain in the back, tightness of abdomen and difficulty in taking deep breath. The skin pigmentation had definitely increased and was of a bluish, cyanotic tint.

Blood-count showed 3,960,000 red cells, 12,000 leukocytes. A culture from the blood remained sterile. The Wassermann reaction was adjudged negative, though there was partial complement fixation.

The condition of the patient became gradually worse until his death in coma on September 5. During the last week the patient's temperature was intermittent, 97 to 101.

The blood-pressure fell during his stay in the hospital from 105 to 85 mm. of mercury.

Examination of Urine.—Soon after admission, the urine was pale orange in color; acid in reaction; specific gravity 1.023. It contained neither albumin nor sugar, but a few hyaline casts. In subsequent examinations there was occasionally a trace of albumin with hyaline and granular casts and several times there was an atypical and slight reduction of Fehling's and Nylander's solutions. There was never any fermentation with yeast, nor any acetone or diacetic acid present.

Clinical diagnosis was not definitely established. The picture was considered suggestive of hemochromatosis but the fever aroused suspicion of a tuberculous infection with adrenal involvement.

Autopsy No. 3,271 (performed 1½ hours after death).

Anatomical Diagnosis.—Hemochromatosis; pigmentation of liver, pancreas, lymph-nodes, gastric and intestinal mucosa, thyroid, parathyroids, adrenals, and skin over the face, hands and feet; cirrhosis of liver with slight collateral circulation; ascites; fibrosis of pancreas and lymph-nodes; acute tuberculous peritonitis with serohemorrhagic exudate; acute tuberculous pleurisy (right); tubercles in kidneys; atelectasis of lung; chronic purulent bronchitis; chronic passive congestion of kidneys; cloudy swelling of kidneys; cysts of the tunica vaginalis of the testicles; slight chronic fibrous myocarditis; adenoma of pancreas.

Body is that of a slightly emaciated white man, 177 cm. in length. There is an extreme dusky cyanosis of the face, which is only partially obliterated by pressure. After the blood has been expelled from the tissues by pressure, the skin over the face and neck still remains markedly pigmented. This pigmentation has a dusky grayish-brown appearance, resembling to a slight extent that found in argyria. It is very marked over the face and neck; is also found over the hands and forearms, fading gradually toward the elbows, and over the feet and legs becoming less marked towards the thighs. The skin over the entire body is slightly pigmented.

The teeth are carious and there is marked pyorrhea alveolaris.

The abdomen is much distended with distinct bulging of the flanks. The genitalia are slightly pigmented, but here the pigment is not very marked. The fat of the abdominal wall is scanty and the muscles atrophic. On opening the abdominal cavity there escapes about 6 liters of a blood-stained, amber-colored fluid which contains numerous flakes of fibrin. Over the serous surfaces of the

intestines one finds, everywhere, small flakes of fibrin which dull the surface to a slight extent. On separating the intestinal loops from each other the folds of the mesentery are seen to be covered with small, transparent, dew-drop-like nodules only a few millimeters in diameter, alternating with larger more opaque nodules. Between these nodules the serous surface is extremely hemorrhagic so that it has a mottled, reddish and grayish, roughened appearance. This lesion is wide-spread and is especially marked over the parietal peritoneum in the flanks and on the diaphragmatic surface. The surfaces of the pelvic viscera and of the liver and spleen present a similar picture in a less degree. The omentum is everted so that it lies beneath the costal margin extending from one parietal wall to the other without being attached to either. It is short, much thickened, cuts with difficulty and shows throughout numerous hemorrhages.

The liver extends 5.5 cm. below the top of the ensiform in the median line and just reaches the costal margin in the mammillary line. The stomach is slightly dilated.

On removing the sternum, the fat of the anterior mediastinum is found to have a chocolate-brown appearance. There are numerous lymph-glands found in this region which are enlarged and more intensely pigmented than the fat. Most of the glands are slightly firmer than normal and cut with resistance.

The pleural and pericardial cavities are normal.

The heart weighs 320 gm. The cavities and the valves are practically normal. The musculature of the right ventricle shows a slight mottling suggestive of the faded leaf appearance. The heart muscle of the left ventricle is slightly mottled and shows a slight diffuse increase in fibrous tissue. No marked pigmentation is appreciable in gross.

Both lungs show a definite purulent exudate in the smaller bronchioles and a moderate emphysema except in the lower portions, which are atelectatic. On the pleural surface of the right lung there are numerous small, grayish nodules, apparently fresh and surrounded by a slight hemorrhagic zone.

The spleen weighs 570 gm., and measures 21 by 11 by 7 cm. Its capsule is firmly attached to the diaphragm by a mass of adhesions which are torn with difficulty. The surface elsewhere is covered with the exudate already described. The spleen is quite firm. On section one finds the architecture very distinct. There is a marked increase in fibrous tissue. Multiple sections show here and there nodules, which measure twice the size of the average Malpighian corpuscle and which are distinctly yellow and soft.

The stomach and duodenum are somewhat distended, and the mucosa of each viscus is definitely pigmented or stained by a black viscid fluid which is present in large amounts. This material has a fecal odor and suggests regurgitation of intestinal contents. The gastric mucosa is considerably congested. A few small hemorrhages are seen along the rugæ. There are a few dilated venules beneath the mucosa of the esophagus near the cardiac orifice.

The pancreatic and common bile-ducts empty separately into the duodenum. Their orifices are separated only by a thin wall of mucous membrane. Both ducts are patent throughout.

The pancreas presents a very remarkable appearance. Its surface is of a pale chocolate color. The head is relatively large, although apparently not quite so large as normal. The remaining portion of the organ is small. The lobulation is distinct. On section the color of the parenchyma is found to be of a deeper brown color than that of the surface. There is some infiltration of fat between the lobules, but there does not seem to be a great excess of fibrous tissue. There are small grayish spots about the size of a pin head, which are relatively slightly pigmented.

The liver is also very striking. It measures 28 by 18 by 9.5 cm. and weighs 1,650 gm. Its surface is quite irregular, small knobs of parenchyma sticking up everywhere between coarse bands of fibrous tissue which are markedly depressed. The color is a deep brown, but on the liver the pigment is not as striking on the surface as on the pancreas and lymph-nodes surrounding it. The liver is

extremely firm and is cut with marked resistance. The cut surface has a rather striking brownish pigmentation. The normal lobulation is absolutely destroyed. Bands of fibrous tissue varying much in size run through the entire liver tissue dividing it into larger and smaller lobules quite different from the normal structure. The fibrous tissue in many places is stained with bile.

The *gall-bladder* is dilated and contains a large amount of tenacious, tarry bile.

The *right kidney* measures 11 by 5.5 by 4.5 cm. The two kidneys weigh 260 gm., are somewhat congested, and show a few small tubercles.

The *urinary bladder* is small and atrophic. Its mucosa is slightly congested, but otherwise normal.

The *prostate* shows no abnormality.

The *testicles* are atrophic. There are a few small cysts in the tunica vaginalis, which contain a clear fluid.

The *seminal vesicles* seem normal.

The *rectum* is extremely congested and shows numerous hemorrhoids.

The *duodenum* is thickened. Its mucosa is edematous and slightly bile-stained. The entire *intestinal mucosa* shows a deep grayish discoloration.

The retroperitoneal lymph-nodes seem to be very numerous. They are all enlarged and show the same chocolate brown pigmentation described for those in the mediastinum.

Of the neck organs, the *esophagus* is normal except for a slight congestion and the varices found in its lower portion. The thyroid is slightly enlarged, and although the pigment here is less conspicuous than that found in other organs, there is a definite brownish discoloration.

TECHNIC

For histological detail, tissues fixed in Zenker's fluid were cut in paraffin and sections stained with hematoxylin and eosin.

For the microchemical demonstration of iron, various methods were used. Among these may be mentioned Perl's reaction, the conversion of the iron into Prussian blue by treating sections with a 2 per cent. solution of potassium ferrocyanid and 1 per cent. hydrochloric acid; treatment with ammonium sulphid; and a combination method described by Nishimura² working in Kaiserling's laboratory. This last method gave the best results. According to Nishimura's experiments, ammonium sulphid reacts with firmly bound iron compounds which are not affected by Perl's method. The latter, however, gives a much better picture than that produced by the black iron sulphid, especially in the presence of other than iron pigments in the same section. He, therefore, first converts the iron into the sulphid by treating with strong ammonium sulphid for one hour and then subjects the sections to a mixture of 2 per cent. potassium ferrocyanid and 1 per cent. hydrochloric acid in equal parts for one hour, after which the sections are washed in 0.5 per cent. hydrochloric acid followed by distilled water. The sections are then counterstained with carmin.

This procedure gives an excellent and uniform method for the demonstration of iron and is perhaps improved by shortening the time of exposure to the mixture of ferrocyanid and hydrochloric acid to twenty minutes, since, according to A. B. Macallum, this mixture decomposes at the end of that time and is apt to give the tissues a diffuse light blue stain.

Alcohol or formaldehyd solution fixation is generally considered preferable in the study of these pigments and all the reagents used should be fresh.

MICROSCOPICAL EXAMINATION

Liver.—The striking features are excessive deposits of golden-brown pigment granules in the parenchymatous cells and in the connective tissue; a well-marked

2. Nishimura: Vergleichende Untersuchungen über die microchemische Eisenreaktion in menschlichen Lebern, *Centralbl. f. allg. Path. u. path. Anat.*, 1910, xxi, 10.

cirrhosis; and evidence of regeneration in the occurrence of clumps of large, rather slightly pigmented, apparently new liver cells. The normal lobulation is obscured and a new architecture formed by the bands of connective tissue running in different directions throughout the organ and dividing it into lobules of various shapes and sizes.

The liver cells vary widely in size and staining properties in different areas, though these areas may be situated side by side. Almost all contain in striking abundance granules of a golden-brown pigment. Some cells appear to be fairly bursting with this load but even in these extreme cases the nuclei can be seen though staining faintly. The most characteristic cell has a well-staining proto-

Fig. 1.—Liver (Case 1). Low power. Iron pigment had been converted into Prussian blue by Nishimura's method and appears black in the figure. Note the heavy accumulation in the connective tissue (*x*) and the relatively heavy deposits in the periphery of the three liver lobules shown rather than in the center (*a*).

plasm and nucleus with a fair amount of pigment gathered in granules near the center at the furthest point from the blood capillaries as pointed out by Opie³ and others. Only in the few nodes of larger, more deeply staining, fresh appearing liver tissue are there any pigment-free cells. The cells in the periphery of these areas may be deeply pigmented, but in the larger central area, only the

3. Opie: A Case of Hemochromatosis. Tr. Assn. Am. Phys., 1899, xiv, 253.

stellate cells of von Kupfer show the pigment granules. In general, the cells in the centers of the lobules are the least affected, the pigment increasing in amount as the periphery of the lobule and the connective tissue septa are approached (Fig. 1). Where the amount of pigment is increased and the granules crowded together, there is a tendency toward the larger size of the individual pigment body with the formation of globules measuring several microns in diameter.

The fibrous tissue varies much in density in different places; hence, presumably, in age. The greater part and that in closest relation to the disintegrating liver cells is quite loose and cellular and contains many pseudo-bile-ducts. Here the pigment is scattered more uniformly and in smaller granules or globules than in the older, denser, narrower, fibrous strands further removed from the liver cells. The epithelium of most of the numerous pseudo-bile-ducts is heavily loaded with very finely granular yellow pigment, only some of the cells near the end having escaped. The walls of all vessels contain masses of pigment and the endothelium of the veins, both hepatic and portal, show fine yellow granules in abundance. The cells of the peritoneal covering of the organ, where they are preserved in the sections, are swollen and pigmented with finely granular yellow material. Immediately beneath the serous membrane the connective tissue is increased, in places amounting to thick scars in which large bile-ducts are strikingly numerous. None of these ducts are heavily pigmented and many escape altogether. This corresponds to areas described by Hess and Zurhelle⁴ as the adenomatous stage in the growth of the bile-ducts. In their case, too, the pigmentation in these areas was slight.

Practically all the pigment in the liver reacts positively for iron by Perl's Prussian blue test. That in the liver cells, connective tissue, endothelium of vessels, etc., stains quite uniformly, varying in shade with massing of the granules. Only in the pseudo-bile-ducts does the pigment seem to vary in its reaction to the test. Here most of the granules become blue, others bluish-green, a dirty green, or are entirely unaffected. Hess and Zurhelle record these differences as occurring quite widely in their sections and consider them no indication of important distinctions in the pigment. They suggest that the youngest pigment stains bright blue, and, increasing in age, passes through the various stages mentioned, the oldest remaining unstained. This observation and the interpretation as given tend strongly to support the view that these so-called pseudo-bile-ducts are evidences of degeneration rather than regeneration of the liver cells.

Pancreas.—There is distinct though slight cirrhosis of the pancreas, both interlobular and interacinar, which at first glance seems of greater extent because of the deposits of fat, especially in the interlobular tissues. Neither process has become excessive. The connective tissue, everywhere, contains a considerable amount of pigment scattered in its interstices or tightly packed in connective tissue cells. In some areas the new fibrous tissue is quite cellular, containing many well-preserved nuclei, the remnants of parenchyma cells, and many small globules of pigment.

The same golden-brown pigment seen in the liver is deposited to a less extent in all the structures of the pancreas and relatively heavily in the islands of Langerhans, which are large, well defined and quite numerous in all parts of the organ examined. Indeed, in sections stained with hematoxylin and eosin, the heavily pigmented, more lightly staining large islands are the most prominent parts of the field (Fig. 2). The prominence of the pigment granules is, in part, due to the greater contrast between the brown of the pigment and the lighter staining protoplasm of these cells; but that it is not entirely so explained is shown by the Perl stain. Many of the islands contain more blue than the surrounding tissue, and others show a pale background of non-pigmented cells on which stands out a single strand or coil of cells near the center, loaded with blue granules. The cells of the islands are large and well formed and for the

4. Hess and Zurhelle: Klinische und pathologisch-anatomische Beiträge zum Bronzediabetes, Ztschr. f. klin. Med., 1905, lvii, 344.

most part have well-preserved nuclei, though some of them are loaded to the fullest capacity with pigment. Occasionally may be seen an irregular, dark staining mass in the place of the nucleus, perhaps one or two in each island. This nuclear change is more prominent in the main parenchymatous cells, occurring diffusely throughout the sections, sometimes involving several cells in one acinus and again only a few in a whole lobule. These cells vary in size, somewhat in shape and very greatly in their pigment content. Not all the cells are pigmented, but in each small lobule some acini are more or less affected. The pigment may be scattered in fine, almost imperceptible grains throughout the cell protoplasm or may distend the cell membrane and displace the nucleus with large granules crowded closely together. The cell protoplasm, where not obscured by pigment, is seen by high power to contain fine vacuoles. Besides the karyorrhexis already noted, a few of the cells show very faintly staining nuclei. Only the smaller ducts contain pigment in their lining cells. Most of the pigment in the sections reacted strongly for iron by the Perl test. There is a very small

Fig. 2.—Pancreas (Case 1). Hematoxylin and eosin. Note the pigment granules in the island of Langerhans, the thickened capsule and the pigment in the surrounding pancreatic cells.

amount of finely divided yellowish pigment in the walls of some of the vessels which does not give the iron reaction. This corresponds to the hemifuscin of von Recklinghausen in contradistinction to the iron-containing hemosiderin.

Many polymorphonuclear leukocytes, as well as mononuclears, are seen in the blood-vessels, and wandering out in the connective tissue septa. Though carefully searched for, no focal necroses were seen. There are no cells in the lumina of the ducts.

One section passes through a small adenoma about 0.5 mm. in diameter, which resembles an enormous island of Langerhans. It has a distinct fibrous tissue capsule from which septa with vessels enter the tumor. The cells are large and similar in all other respects to those in the islands of Langerhans. Only a few scattered cells contain pigment granules.

Spleen.—There is some increase in connective tissue. Pigmentation is not conspicuous in sections stained by ordinary methods due to the similarity in the color of the granules and the large numbers of red blood cells present in the splenic pulp.

With the Prussian blue method a slight amount of pigment is demonstrated lying along the sinuses, and a few grains here and there in a Malpighian body. The granules are small and are present in rather large phagocytic cells. All the pigment reacts well to iron tests.

The venules are distended with blood. The individual cells differ slightly in size and shape.

The Malpighian corpuscles average about 0.3 mm. in diameter. In several of these and once or twice in the pulp there are seen areas of necrosis with a diameter of about 0.2 mm. They are rather sharply defined, contain a grayish, granular, necrotic-looking material with scattered nuclear remnants. There is very little reaction in the surrounding tissue.

Thyroid Gland.—This shows marked pigmentation. Apparently every cell of every alveolus contains innumerable golden-brown granules. The distribution is remarkably uniform, more so than in the other tissues described. The majority of the cells are affected to the same degree, but here and there are found a row of cells distended and closely packed with pigment and a few of the alveoli contain only a few grains in each cell. The nuclei are well preserved and show no sign of degeneration even in the most heavily loaded cells. The alveoli are natural in size and shape; the colloid is slightly decreased in amount. The connective tissue is practically free from pigment. Only by careful search can a few large grains or, rather, small globules be found in a branching connective tissue cell or free in a lymph space. The blood-vessels, large and small, are distended with blood, the tissues are forced apart by edema, and many free red blood cells are seen in the meshes of the connective tissue. Leukocytes are not prominent, but may be seen occasionally, and here and there one has worked its way in among the pigmented alveolar cells. With the Perl test the pigment reacts positively for iron, but requires longer exposure to the reagents. There is more of the dirty bluish-green color than is usually observed elsewhere. Very little, if any, remains unchanged.

Parathyroid.—Cells with eosin-staining granular protoplasm greatly predominate over the clear non-staining variety. Small colloid vesicles are occasionally seen. There are many pigmented cells scattered throughout the gland. The pigment occurs in fine yellow grains and reacts readily for iron. Most of the pigmented cells are those having a clear non-staining protoplasm.

At one end of the gland there is a cyst measuring about 3 mm. in diameter, almost as large as the remainder of the parathyroid. It contains a heavy, granular precipitate and is lined by a single row of cubical cells having a clear protoplasm. The nucleus, as a rule, is vesicular; in some cells, it is pyknotic with very irregular edges. Many of the cells lining the cyst contain the iron pigment.

Kidney.—There is a very slight grade of chronic nephritis with small increase in fibrous tissue and cloudy swelling of the secreting epithelium. The blood-vessels throughout the sections, glomeruli, and interlobular capillaries of cortex and medulla are congested. The amount of pigment is small, compared with that in the organs just described, and its distribution quite patchy. It occurs as rather large granules in considerable amount if present at all. It does not occur in isolated grains or even a few scattered grains here and there. The sites of predilection are the collecting tubules first, and occasionally, the loops of Henle. It gives the iron reaction readily. No pigment is found in the walls of the blood-vessels or elsewhere in the connective tissue.

Adrenal.—The pigmentation usually seen in the reticular zone of the cortex is much increased, being found in almost all the cells of this layer as scattered granules of light brownish color. Here and there in the outermost cells of the

glomerular zone, the nuclei are surrounded with a very similar pigment in fine granules of quite uniform size and of a slightly darker color than that in the inner layer; for the latter appears brighter even in much larger amounts. With the Perl test most of the glomerular pigment reacts positively, but not completely. Usually the cell protoplasm stains a diffuse blue in which the granules appear as dark green. The reticular pigment remains quite unchanged. Scattered through the medulla may be seen an occasional pigmented cell and a few of these react positively for iron. There was no tissue fixed for study of the chromaffin system.

The lipoids of the cortex were less than is usually found. Their solution leaves the cells of good shape and staining qualities.

Lymph-Nodes.—All sections show heavy pigmentation and increase in connective tissue. The pigment in different sections varies somewhat in situation. As a rule, the sinuses and peripheral parts of the follicles contain almost all of

Fig. 3.—Pancreas (Case 2). Hematoxylin and eosin. Low power. Note the preponderance of fat in the field; the heavily pigmented and sclerotic condition of the pancreatic tissue shown in the right center; and the mass of clear non-staining cells, showing fewer pigment granules.

the pigment, but occasionally, a whole follicle is loaded with it while the surrounding sinuses are relatively free. It occurs in rather large globules, 4 or 5 microns in diameter, which fuse to form larger masses, and lie either free in the interstitial spaces or in phagocytic cells of various kinds. Very often the character of the containing cell is obscured by the amount of pigment present. In other cases they resemble the polyblasts of chronic inflammation. Phagocytic giant cells with two or three nuclei are not uncommon, and occasionally, there are seen large syncytial masses, 80 to 100 microns in diameter, having fifteen or twenty nuclei and enclosing several large clumps of pigment.

The connective tissue is increased, especially between the sinuses, where it forms fairly thick trabeculae. Here and there it is found invading a heavily pigmented follicle or growing out from its central artery. The pigment takes a dark blue color in the iron test whether the ferrocyanid or ferricyanid of potassium is used.

In one of the nodes examined there are three or four young tubercles composed of the typical epithelioid cells and surrounded by lymphocytes. There is no pigment in the tuberculous tissue.

Lung.—The sections show a moderate emphysema, thickening of the smaller vessels, a considerable quantity of coal pigment and a very few conglomerate tubercles of the proliferative type. There is no other pigment than that of anthracosis.

Omentum.—The omentum shows extensive extravasation of blood which occurs in masses throughout the sections together with an exudate of fibrin, mononuclear cells and few polymorphonuclear leukocytes. Scattered uniformly through the exudate are seen typical discrete and conglomerate miliary tubercles with epithelioid, small mononuclear and giant cells.

The parietal peritoneum shows rather acute tuberculous peritonitis with a considerable amount of extravasated blood. There is no pigment in the striated muscle shown in this tissue. The exudate on the visceral layer over the kidney is very similar but contains less blood. The perirenal fat shows many large dilated blood-vessels.

CASE 2.—*Medical No. 20,197. Hemochromatosis with Diabetes; Onset with Diabetic Symptoms Ten Months Before Death.*

D. C. K., white man, a grocer, aged 37 years, entered the Johns Hopkins Hospital September 26, and died Oct. 22, 1906.

Complaint.—Lack of strength and stomach trouble.

Family History.—Negative.

Past History.—The patient has always been robust with the exception of measles, mumps and whooping-cough during childhood and an attack of chills and fever about twenty years ago.

Habits.—He is accustomed to drink three cups of coffee each day, uses tobacco moderately and drinks not more than one glass of beer a week.

Genito-Urinary.—There was no urinary disturbance before the present illness. He denies syphilis and gonorrhea. He has been married ten years and has three children.

Patient has had hemorrhoids for five or six years.

Present Illness.—This began about nine months ago with a severe cold which left the patient with a tired feeling. This cold lasted about one month during which the patient noticed that he was drinking large amounts of water and getting up several times at night to pass urine of crystal clearness. His appetite was then moderate. He had been "excited" but not "worried" over business affairs two or three weeks previous to the polyuria. He began to experience about this time a marked feeling of weakness; he could work no more than fifteen minutes, consecutively.

In July he went to a Cincinnati hospital where his hemorrhoids were cured after three operations. Here he was also placed on a diet, but continued to lose weight.

Since then his appetite has greatly increased; his skin is dry; stools clay-colored; and he has become constipated.

About ten days ago sharp pains developed all over the body, moving from one part to another, and after several days subsided into a dull, aching pain aggravated on motion.

Physical Examination.—The patient is a well-built, poorly nourished man. His skin is harsh, brawny and dry without a trace of perspiration. The face

has a fairly good color. There is no jaundice. The pharynx and soft palate are red, dry and leathery; strings of mucus project down from the posterior nares. The skin of the neck and thorax is negative.

The heart and lungs are normal.

Abdomen.—The liver is enlarged, the relative hepatic dulness begins at the upper border of the sixth rib and extends 4 cm. below the costal margin where the edge is definitely palpable.

The spleen is not felt.

There is no general glandular enlargement.

The knee-jerks are present. The left knee is painful and appears to be slightly swollen, but measures the same as the right knee.

Blood Examination.—September 27, red cells 4,458,000; leukocytes 5,400; hemoglobin (Tallqvist) 90 per cent.

For examination of the urine see Table I. On admission the urine contained 6.5 per cent. sugar.

Fig. 4.—Parathyroid (Case 2). High power. Hematoxylin and eosin. Note the fine granules of pigment in all the cells and the many irregular, dark nuclei.

The patient became gradually more drowsy and apathetic until on October 11 the breath had a distinct odor of acetone and the appetite entirely failed.

On October 16, the patient was seen by Dr. Futeher, who noted that the most striking feature was the very marked grayish-brown bronzing of the hands and lower thirds of the forearms. This was also apparent, but to a less extent, on the face and neck and over the shins, especially the right one. There was no increase of pigmentation in the axillæ or about the nipples or genitalia. The sclerotics and the buccal mucosa were also free from pigment. There was one definite petechial spot on the upper third of the right shin.

The right parotid gland was quite markedly swollen, tense and tender on pressure.

While there was no definite pigmentation of the skin of the body, the color seemed to be a little deeper than normal.

The parotitis cleared up within a few days.

The temperature was subnormal continuously.

The patient died in coma October 22.

Clinical Diagnosis.—Hemochromatosis; diabetes mellitus; acute parotitis; diabetic coma.

TABLE 1.—URINARY FINDINGS, CASE 2

Date.	Total Amount of Urine in c.c.	Specific Gravity.	— Sugar — Per Cent.	Gm.	Acetone.	Diacetic Acid.	Diet.
9/29	3,000	1.032	6.5	195.	..	..	Ward diet.
9/30	1,200+		7.2	86.4+	..	..	Ward diet.
10/ 4	2,600	1.040	4.4	114.4	..	..	Carbohydrate-free, plus one slice of bread.
10/ 5	1,980		4.6	91.	..	..	Carbohydrate-free.
10/ 6	2,630		3.7	97.3	..	..	Carbohydrate-free.
10/ 7	1,560		4.4	68.6	..	..	Carbohydrate-free.
10/ 8	3,020		3.9	117.7	..	..	Carbohydrate-free.
10/12	2,290		3.3	75.5	+	+	Carbohydrates renewed, plus soda bicarbonate.
10/13	3,480		7.2	250.5	+	+	Carbohydrates renewed, plus soda bicarbonate.
10/15	4,740		4.	189.6	+	+	Carbohydrates renewed, plus soda bicarbonate.
10/16	2,240		..		+	+	Carbohydrates renewed, plus soda bicarbonate
10/17	2,140		..		+	+	Carbohydrates renewed, plus soda bicarbonate
10/19	3,000		5.	150	+	+	Carbohydrates renewed, plus soda bicarbonate
10/20	2,820	1.030	2.6	73.3	+	+	Carbohydrates renewed, plus soda bicarbonate.
10/21	1,050+	1.029	2.5	26.2+	+	+	Milk $\frac{3}{4}$ xlvi
10/22	2,120+	1.025	2.6	55.1+	..	..	Nothing retained.

Autopsy No. 2,782 (performed one-half hour after death).

Anatomical Diagnosis.—Hemochromatosis; pigmentation of liver, pancreas, heart, stomach, thyroid, parathyroids, adrenals, lymph-nodes, trachea and skin; chronic interstitial pancreatitis; cirrhosis of liver; chronic splenic tumor; cardiac hypertrophy and dilatation; chronic passive congestion of abdominal viscera; edema of intestines; ascites; degeneration of renal epithelium; hydrothorax and atelectasis of lungs; atrophic testes.

Body is that of a stoutly built man, 180 cm. in length. The subcutaneous fat is small in amount. On opening the peritoneum a large quantity of clear fluid is found, perhaps 2 liters; this examined by Dr. Emerson shows large quantities of sugar. The abdominal organs are in general normally disposed. The intestines are distended, and their walls thick and rather pasty. The peritoneal surfaces are smooth and glistening. The omentum is thickened and adherent here and there to the diaphragm and parietal peritoneum; and in the adhesions there are numerous, large, tortuous veins. These are also abundant in the mesentery but are not especially conspicuous about the round ligament of the liver. The liver does not extend below the costal margin. It is greatly distorted and shows deep fissures about its edge, especially near the gall-bladder. The distended gall-bladder is bent on itself and adherent to the edge of the liver. The spleen is greatly enlarged and not bound by adhesions. The diaphragm reaches the third rib on the right side and the fourth space on the left side.

The pleural cavities contain a small amount of fluid—about 250 c.c. on each side. The surfaces of the lungs are quite smooth and glistening. There are no

adhesions. The pericardial cavity contains very little fluid. Its surfaces are smooth.

The *heart* is somewhat enlarged, weighing 350 gm. The valves and orifices are normal. The root of the aorta shows some small plaques of elevation, and the coronary arteries are similarly flecked with small patches. The heart muscle is, in general, firm. The left ventricle is distinctly dilated. The muscle shows through the endocardium as a rather chestnut brown substance which is very much mottled with lighter and darker areas in the papillary muscles. This is practically the appearance seen in fatty degeneration, but it is not marked on section of the heart. On section the heart muscle shows the same chestnut-brown color throughout, but it is distinctly paler and more translucent toward

Fig. 5.—Bone-marrow (Case 2). Low power. Hematoxylin and eosin. Note the preponderance of fat tissue over the hematopoietic elements. This particular field contains more islands of blood-forming cells than the average.

the inner layer. The muscle of the right ventricle shows the same bright brown color.

The left *lung* shows some atelectasis at the base. The bronchial glands are blackened by coal pigment but not enlarged. The right lung shows a small flat scar at the apex. The bronchial glands are here distinctly brown. Another gland, also on the bronchus, which contains a quantity of black pigment, shows the brown only indistinctly.

The *spleen* weighs 300 gm. and measures 15 by 10 by 5.5 cm. In consistence it is somewhat flabby. On section the splenic substance is dark brownish-red, but

the brownish cast is very slight. The trabeculae and vessels are distinctly visible. The Malpighian bodies are not enlarged, but are easily seen.

There is a minute accessory spleen at the hilum of the spleen and several pigmented lymph-nodes.

The *duodenum* is normal except perhaps for a slight reddening and swelling of the mucosa. The bile can be squeezed slowly from the bile-ducts.

The *stomach* contains only a little mucus. Its wall is corrugated and swollen, dark red, and the mucosa is seen thrown up into thick folds, which are very rough. The veins about the esophagus are not especially dilated.

Along the lesser curvature of the stomach there are several quite large lymph-nodes, which are brilliant, ochre-brown in color. On section the lymph-cords are visible as clear grayish nodules, while the remainder is opaque and yellowish-brown. Some of the nodes are red and show minute masses of pigmentation,

Fig. 6.—Liver (Case 3). Low power. The iron-containing pigment had been changed to Prussian blue by Nishimura's method and appears black in the figure. The section was counterstained in carmin. Note the node of recently regenerated liver cells containing no pigment, except in the Kupfer cells. The older, surrounding liver tissue is heavily pigmented.

while others are uniformly dark brown. It seems possible that the non-pigmented nodes may be newly formed. The lymphatic channels leading to and from these nodes are greatly distended with clear and sometimes blood-stained fluid. The mesenteric nodes are not especially enlarged; they are much less pigmented, but there is a quite distinct coloration of the cortical layer.

The *pancreas* weighs 150 gm. It is large, firm, and measures about 19 by 4 by 2 cm. It is somewhat bound by adhesions to the surrounding tissues, in which there are many large pigmented lymph-nodes. The ducts are quite patent and contain no calculi. Their walls are about normal in thickness. The gland, on

section, is curiously mottled, portions of it quite irregular in size and firm, having a bright brown color, while the remaining intervening parts have the appearance of very solid fat. This is much firmer than ordinary fat and there is evidently a considerable amount of fibrous tissue running through it. The yellow fatty areas form a much greater proportion of the head of the pancreas than of the tail. In the brown areas, especially in the tail, the lobulation is extremely indistinct. Pale grayish dots are frequently seen which suggest the islands of Langerhans. From the surface the mottled appearance of the pancreas, with its alternating brown and yellow tissue, is very striking.

The *liver* weighs 2,400 gm. and measures 27 by 24 by 11 cm. It is very hard and its surface is extremely rough and knobbed. At the gall-bladder, the contractions of the liver in deep furrows is especially marked. On section the cut surface has a bright brown color. There is a very distinct lobulation, the nodular

Fig. 7.—Pancreas (Case 3). Low power. Hematoxylin and eosin. Note the especially heavy accumulations of pigment around the three islands of Langerhans.

masses being separated by retracted, translucent bands. The lobules are fairly homogeneous in size, although there are some areas in which the scar tissue contains numerous very minute lobules. In many areas this lobulation seems to follow the normal very closely, the peripheral part of each lobule being paler, while the central part appears as a reddish dot. But there is great irregularity, and in other places the masses of liver tissue 2 mm. in diameter probably correspond in no way to the normal liver lobule, although they stand out sharply. At one point there is a distinctly outlined, round, yellow mass about 1 mm. in diameter.

The *peritracheal lymph-nodes* are not markedly pigmented, although the larger ones do show some abnormal coloration.

There are four *parathyroids* which are normal in appearance.

The *thyroid* is firm, possibly slightly browner than normal, and contains a considerable amount of colloid.

The mucosa of the *esophagus* seems normal. There are no varicose veins.

The mucosa of the trachea and of the epiglottis seem distinctly pigmented, mottled with yellowish patches.

The *aorta* is, in general, smooth, but shows some yellowish plaques of sclerosis. These are of small size and perhaps more abundant in the carotid arteries.

The *adrenals* are alike. The cortex is distinctly marked out, pale and sprinkled with opaque, yellow dots. The medulla is grayish brown.

The *kidneys* weigh 430 gm. The right kidney measures 14 by 7.5 by 5 cm. The two kidneys are quite similar in appearance. The capsule strips easily, leaving a surface somewhat roughened by depressions and on which the stellate veins are conspicuous. The surface is pale, pinkish-white, with a yellow cast. Minute dots of yellow opacity can be seen everywhere on the surface and minute hemorrhages are abundant. There are also translucent points. On section

Fig. 8.—Pancreas (Case 3). High power. The iron-pigment had been converted into Prussian blue and the section counterstained with carmin. Note the granules of hemofuscin (yellow in the preparation) in the wall of the vessel and the dark hemosiderin in the surrounding pancreatic cells.

the cortex averages 7 mm. in thickness. The striations are quite straight. The central translucent rays are very distinct. The labyrinthian portion is very opaque and pale, the glomeruli standing out conspicuously as bright red points.

The solitary nodules are conspicuous throughout the large *intestine* and the mucosa is markedly edematous. Peyer's patches in the small intestines are swollen. The mucosa is not distinctly pigmented; in fact, it is quite pink. The upper portion of the small intestine is very much congested and of a purplish color. The edema persists throughout.

MICROSCOPIC EXAMINATION

Liver.—There is a well-marked cirrhosis, with rather dense bands of connective tissue intersecting and dividing the liver into lobules. These lobules are

very much larger than the normal liver unit and, in them, the cells are not arranged with the same regularity. Only rarely can a central vein be made out among the liver cells; almost always they are surrounded by connective tissue, and in this respect do not differ from those of the portal system.

The pigment is abundant and very striking in certain places. As a general rule, the liver cells contain about the same amount, closely grouped in golden-brown granules around or at one side of the nucleus and of such quantity as to occupy about one-third or one-fourth of the cell. This is the general rule, but in many places, scattered throughout the tissue, are seen areas having a diameter of ten or a dozen liver cells, which are filled with pigment, contained, for the most part, within the cells, but also found in smaller amounts free in the tissue. These cells are alike in resembling masses of pigment surrounded by a limiting cell membrane, but differ in the appearance of the nucleus, which in a few is the large round vesicular nucleus of the liver cell, and in the majority an

Fig. 9.—Heart muscle (Case 3). High power. The iron-containing pigment appears black in the figure. Stained by Nishimura's method.

elongated or oval, darker staining variety resembling more nearly that of a connective tissue cell or polyblast. There are also seen here and there smaller masses of pigment between the liver cells contained in the cells of von Kupfer or polyblasts inside the blood capillaries. The larger lobules of liver parenchyma are singularly free from any invasion of fibrous tissue. The cells, for the most part, except their pigment content, seem very nearly normal. The protoplasm shows no other evidence of degeneration and the nuclei, as a rule, stain well, though here and there may be seen a very faintly staining one or one undergoing karyorrhexis.

The connective tissue trabeculae would be prominent of themselves and are rendered much more so by the abundance of the pigment lying free between the bundles and contained also in cells. The pigment here is concentrated into narrower space and occurs in larger granules and globules.

The formation of new bile-ducts is certainly not a striking feature of the liver, but wherever found the lining epithelium contains a share of the pigment usually in fine yellow grains.

Section through the atrophied portion near the gall-bladder shows a few strands and groups of pigmented liver cells embedded in rather dense connective tissue, which is infiltrated also with small, round cells. Occasionally, there occur larger islands of liver tissue, resembling in every respect that found in other parts of the organ. In the very fibrous connective tissue surrounding the large vessels and bile-ducts in this region, there are evenly and rather sparsely scattered connective tissue cells loaded with pigment in very fine, even grains and of a slightly darker brown color than that of the prevailing type elsewhere. This

Fig. 10.—Thyroid (Case 3). Low power. Stained by Nishimura's method. Iron-containing pigment appears black in the figure and is situated in the lining cells of the alveoli.

variety (hemofuscin) is found in the adventitial cells of vessels in other sections, and has not reacted positively in any test for iron which I have used. The pigment of the liver cells, bile-ducts, connective tissue, generally, with the single exception noted, gives a positive reaction for iron.

In comparing sections treated by Perl's or Nishimura's methods the difference in distribution of the pigment in Case 1 and in Case 2 is well shown. As noted in the description of that case, the pigment in the former is present in striking amount in the liver cells in the periphery of the lobule or cell mass while those in

the center are comparatively little affected, as are also the connective tissue septa. In Case 2, the connective tissue contains the pigment in most conspicuous amounts, while that in the liver lobules is very evenly distributed.

Pancreas.—On an average about one-third of the sectional area is occupied by pancreatic tissue, the remainder of the space being filled with fat. The parenchyma is grouped together, usually, in one end of the section in lobules or islands which are separated from each other by fatty septa of varying widths. The glandular tissue itself is further subdivided into numerous small collections of acini by fibrous tissue strands which are very rich in pigment-bearing cells, a fact which renders the false lobulation especially distinct. There are in places fine strands of connective tissue between the individual acini, but this is not striking. The acini, as a rule, are poorly formed and irregular but the cells stain fairly well. Almost all the cells are pigmented with yellowish-brown granules, varying in extent from a few granules to a mass of globules. The connective tissue strands are rich in pigment occurring to some extent free in the interstitial spaces and to much greater extent contained in heavily loaded mononuclear cells. These cells are massed here and there in clumps between

Fig. 11.—Adrenal (Case 3). Low power. Hematoxylin and eosin. Figure shows capsule, heavily pigmented glomerular layer, and part of the fasciculate layer, in which may be seen a few clumps of pigment.

the groups of persisting acini and the fibrous bands, lying in small groups in the meshes of a coarse connective tissue reticulum.

In certain small, rather sharply circumscribed areas, the pancreatic cells, while still arranged in acini, have lost their usual appearance and staining reactions. The protoplasm appears swollen, vacuolated and foamy, staining hardly at all. The nucleus does not depart greatly from the usual type, but occasionally is smaller and darker. These cells, as a rule, contain a great many pigment granules gathered especially about the nucleus, but they are not by any means always the most heavily pigmented cells (Fig. 3).

The lining cells of many of the ducts contain the pigment, even those of large size. In the walls of the vessels, too, are occasional pigment-bearing cells.

The islands of Langerhans appear in all sections, but vary greatly in number and somewhat in appearance. In the head of the organ, they are quite numerous

and appear as sharply circumscribed areas of the usual size containing rather small cells in thick columns arranged in a single, simple or branching coil. Only a very few of these cells are pigmented. In the body the islands are less numerous but of typical appearance in every way and having a somewhat more complex arrangement than in the head. Here the pigmentation is more noticeable, most of the islands containing at least one or two pigmented cells. The tail of the organ is greatly scarred. There is loss of all appearance of acini; the cells are arranged in rows and small masses surrounded by strands of fibrous tissue. Hence the differentiation of the islands from the rest of the parenchyma is not so easily accomplished. Almost all of the cell areas might be taken for atypical islands of Langerhans, but scattered fairly frequently throughout the section are typical appearing circumscribed strands of cells which closely resemble those seen in other portions of the organ and like them are distinctly less pigmented than the surrounding parenchyma.

By using Nishimura's method with tissues fixed in methyl-alcohol, practically all the pigment gives an iron reaction, becoming either of a pure Berlin blue color or a dirty, yellowish-green, with many intermediate stages. Even in the best preparations, however, a little of the pigment retains its original golden character. This iron-reacting pigment does not seem to have any special predilection for any particular cell or situation, but is found now in the ducts, now in the acini, or again in the connective tissue cells. By using Perl's method, the modification recommended by Dr. Maud Abbott,⁵ a greater amount of the pigment remains unchanged, more is of the yellowish-green hue and there is less showing of the clear blue. In these sections, the islands of Langerhans are rendered conspicuous by the absence of the blue color and its presence in abundance in the surrounding tissues.

Spleen.—The connective tissue is diffusely increased.

The interstitial pulp is greatly decreased in amount. The Malpighian bodies are small and circumscribed. The golden-brown pigment is noticeably increased in all of the sections, but does not begin to compare in amount with that in the liver, pancreas, and other organs. Most of it is deposited, finely divided, in the walls of the sinuses with here and there larger intracellular masses. Occasionally granules and small globules are found in the venous spaces, either free or in large phagocytic cells. Many of the fine grains in the walls of the sinuses seem to be contained within the endothelial lining cells. None is found in the Malpighian bodies. The pigment reacts readily to iron tests.

Lung.—The sections show a moderate grade of emphysema. There is a moderate amount of black coal pigment. In addition to this there may be seen also yellowish-brown pigment in very much less amount, but easily distinguished from it. As a rule the hemosiderin occurs in small masses (10 or more millimeters), far apart, in the alveolar walls, either loose in the interstices of the tissues or contained in large mononuclear cells. It may be seen, also, forming small casts of the capillaries for a short distance. The small lymph-nodes within the lung substance, while containing an abundance of coal pigment, show no hemosiderin. The bronchi are slightly dilated within the lung substance and the cells of their lining membrane are definitely, though slightly, pigmented with golden-brown grains lying, usually, deeply placed in the cell, away from the lumen and in close proximity to the nucleus. The cells lining the pleura are not pigmented.

The mucosa of the higher air passages (the larger bronchi, trachea and larynx) is quite free from the pigment, which here occurs in the acini, ducts and interstitial tissue of the serous glands beneath the mucosa and in the cells of the cartilages. In the glands, the pigment is moderate in amount and finely divided into yellowish grains. Often only a few of the cells of each acinus may show the pigment; occasionally the lumen of an acinus or a duct may show a mass of the fine granules. There are mononuclear cells loaded with pigment in the interstitial tissue between the acini and in some of the glands there appears

5. Abbott, Maud: Pigmentation of Cirrhosis of the Liver in a Case of Hemochromatosis. *Jour. Path. and Bacteriol.*, 1901, vii, 55.

to be an increased round-celled infiltration. The mucosa of the ducts shows pigmentation up to within a short distance of the lining membrane of the air passage. The cells of the cartilages, many of them, are large and well staining; many others are small and shrunken away from the matrix. Cells of both orders may contain finely divided pigment granules clustered about the nucleus or filling the whole cell. Perhaps a third or a half of the cartilage cells are thus pigmented. In the small peritracheal lymph-nodes the dilated sinuses are filled with mononuclear leukocytes which, in turn, are loaded with brownish pigment. There is no increase in fibrous tissue. The pigment in the serous glands reacts poorly to iron tests; some of it becomes blue, most of it a dirty greenish color, but never does all the pigment in a single section react positively. This is true to even greater extent of that in the cartilage cells. Most of this remains unaffected, even when treated by Nishimura's method, but the granules in some of the cells become dark green and in a very few the bluish color appears. However, that in the sinuses of the adjacent lymph-nodes reacts promptly. In these nodes the iron pigment is the only one present. There is no coal pigment.

Kidney.—There is no increase in connective tissue. The capsule of Bowman contains a granular detritus. The cells of the convoluted tubules are cloudy, swollen and much vacuolated, especially toward the basement membranes. Their lumina are irregular in shape and contain the characteristic "foamy" material of epithelial degeneration.

The pigment in the kidney is sharply limited to a certain class of tubules, having low, regular, slightly cloudy epithelium and a relatively wide empty lumen, apparently connecting or collecting tubules. The pigmented tubules are situated, principally, in the outer half of the cortex in the convoluted areas, between the medullary rays of Ferrein, though occasionally, they are found within the rays themselves. The cells often carry an excessive load of the pigment, while the immediately adjacent tubules are entirely free. The pigment reacts quite readily with the usual iron reagents. In the basal portion of the medullary pyramids, there are many tubules whose cells are much swollen and quite clear and translucent, the nucleus staining fairly well. These cells contain a few large grains of yellow pigment which does not give the iron reaction.

Thyroid.—The vesicles are normal in size and contain the usual amount of colloid. The interstitial connective tissue is not increased. Pigmentation is slight. Many of the cells lining the vesicle contain fine golden-brown pigment granules; some of them are distended with pigment, most of the cells contain little or none. On the whole, the pigmentation of the thyroid is not striking. The interstitial thyroid cells occurring in nests and strands (undifferentiated, embryonic cells?) are the most heavily affected. The pigment reacts readily for iron. There are also fine yellow grains in the coats of some of the smaller arteries.

The *parathyroids* show excessive pigmentation within the essential cells and this is rendered all the more striking by the close anatomical relationships existing between the cells. Each cell seems to contain great numbers of fine yellowish granules but with the oil immersion lens unpigmented cells can be distinguished scattered here and there. Cells with clear protoplasm are more numerous than those staining with eosin and these clear cells are the most heavily pigmented. Many of the nuclei are pyknotic and have irregular serrated edges. In the connective tissue along the larger blood-vessels are seen pigmented spindle-shaped cells. There are no masses of pigment in the interstitial tissue, which, moreover, is very inconspicuous. The pigment is iron containing (Fig. 4).

Skin.—In the basal cells of the Malpighian layer of the epidermis there is a finely granular yellowish or greenish-brown pigment in a section stained by Perl's method. There are only a few pigment grains in the underlying connective tissue of the corium.

Testes.—The epithelium of the tubules is retracted away from the capsule. Many mitotic figures occur but no spermatozoa. The tubules are far apart, the

interstitial tissue is quite loosely arranged and seems edematous. The interstitial cells of Leydig are large and prominent by reason of an excess of diffuse brownish pigment throughout their protoplasm. In the walls of many of the blood-vessels, and especially of their lining endothelium, there is a golden-brown pigment of different morphology, occurring in grains and small masses as seen in the other organs. This latter pigment reacts quite readily for iron, but that in the interstitial cells is always negative.

Bone Marrow.—The marrow of the femur appears slightly edematous. There is a peculiar fibrin-like stroma between the fat spaces. Fat greatly preponderates over the cellular elements, which are grouped in small islands throughout the sections and are present in about the normal relative proportions. There is no special alteration in the marrow, except a slight increase in the amount of pigment, most of which lies free in the spaces between the marrow cells. Much of this pigment reacts positively for iron (Fig. 5).

Adrenal.—The cells of the glomerular layer are heavily pigmented with small golden-brown granules quite regular in size and appearance. As a rule the pigmentation is sharply limited to the cells of this layer, but here and there it may be seen dipping down into the fasciculate layer to a distance of from ten to fifteen cells. Many of the cells are degenerating and some have already disintegrated, their pigment being deposited in small masses. What is evidently the track, along which this pigment is removed, may be traced up through the capsule where small granules and masses of pigment may be found in rows in the interstitial spaces or occasionally in a mononuclear cell. Even in the surrounding fat little rows of pigment resembling tiny strings of beads may be seen in the spaces between the fat cells.

The zona fasciculata is composed of quite normal-looking cells, which, however, contain less than the usual amount of lipoid material. The cells of the zona reticulata contain the usual diffuse brownish pigment in increased amounts. This is darker than the chromaffin substance of the medulla and is found in sections regardless of the method of fixation. With high powers it is slightly granular with granules very much smaller than those of the iron pigment. It is not affected in the least by the reagents used in the demonstration of iron.

In tissue fixed in formaldehyd solution and potassium chromate the yellow chromaffin substance of the medulla is found in normal, or slightly increased amounts. It occurs in similar proportions also in the cells of two sympathetic ganglia examined. In addition, there is found in the medulla another pigment occurring sparsely scattered in the medulla cells and interstitial tissue, in tissues fixed in alcohol or otherwise, and morphologically resembling the pigment found in the glomerular layer.

In sections stained by Nishimura's method, the glomerular pigment becomes blue readily. The outer third of the cortex takes a diffuse light bluish stain, which under high powers is seen to be especially localized in the spaces between the cell columns. The medulla, too, takes a diffuse light-blue stain, marking it off sharply from the reticulate zone. The pigment in the medulla, however, though resembling closely the glomerular pigment morphologically, for the most part remains unaffected by this test, only a few clusters of the granules taking on a dirty, dark green color.

Heart Muscle.—The muscle cells are moderately increased in size. The pigmentation is striking. The pigment occurs not only in the undifferentiated sarcoplasm at the nuclear poles, but also throughout the entire muscle cell in fine yellowish-brown granules arranged in longitudinal rows between the fibril bundles. The picture reminds one strikingly of that seen in a section of a fatty heart muscle stained with Sudan III. The cross striations of the muscle cells are easily seen, except where obscured by closely packed masses of pigment. The nuclei, as a rule, stain rather faintly. In cross section of the muscle cells, the pigment granules are seen to be more numerous in the center, diminishing in numbers toward the periphery in inverse arrangement to that of the fibril

bundles. Here and there, especially in sections stained for iron, may be seen a mass of granules, perhaps with a clear space in the center, and surrounded by a narrow zone of the muscle cell showing the characteristic fibril bundles. Other cells with large amounts of pigment in the center may show many small clear vacuoles in their periphery. Pigment is also found in the interstitial tissue, usually, in mononuclear cells but occasionally free. Where this pigment is greatest in amount, the fibrous tissue is increased and many small areas of fibrosis may be found containing heavily pigmented mononuclears and one or two pigmented and atrophying muscle cells. In sections stained by Nishimura's method all the pigment takes on a beautiful blue color. No unaffected granules can be found, even immediately at the nuclear poles after careful search. Perl's method leaves some unchanged pigment at the nuclear poles.

Gastro-Intestinal Tract.—The mucosa and submucosa of the stomach are slightly edematous. The glands at the base of the mucosa are deeply pigmented. The pigmentation becomes less toward the surface and none is found in the superficial half of the mucosa. An occasional pigmented mononuclear cell is found in the interstitial tissue. There is no pigmentation of the muscle.

Sections of the small intestine show marked edema, especially of the mucosa and submucosa. There is no pigmentation.

Lymph-Nodes.—Sections of all lymph-nodes examined, with the exception of the bronchial, contain a larger or smaller amount of iron pigment, both intracellular and free in the interstitial tissue. There is also in many cases a chronic lymphadenitis with marked increase in fibrous tissue. Where the pigmentation is slight, it is found especially in the periphery of the sinuses along the lymph-nodes and cords, but in more extreme cases may be found in dense masses in the nodes and cords.

CASE 3.—*Medical No. 24,059. Hemochromatosis with Diabetes Mellitus. Onset with Diabetic Symptoms Five Months Before Death.*

R. D., white, farmer, 44 years old, was admitted to the Johns Hopkins Hospital May 20, and died June 5, 1909.

Complaint.—General weakness.

Family History.—Excellent. The patient has been married twenty-three years. His wife is in excellent health. There are three children.

Past History.—In general the patient has been a hearty man and enjoyed excellent health. He had the usual diseases of childhood.

Lesions of the head, cardiac, respiratory, neuro-muscular and genito-urinary systems were carefully inquired for with negative results.

Gastro-Intestinal.—He has always been a hearty eater, thinks he often ate too much, but suffered very little from indigestion. He occasionally had indefinite pains after eating. Bowels have been constipated since the present illness began, but before this for fifteen years were loose with two or three passages each day. Twelve years ago he had dysentery which lasted for one year. It was characterized by bloody, mucous stools and great abdominal tenderness. He was never jaundiced and has not noticed the presence of hemorrhoids.

Habits.—He is a hard worker and has undergone considerable exposure. Formerly he was a heavy smoker, but has used no tobacco since last fall. He has never been addicted to alcohol. Since the onset of the present illness he has taken a little whiskey twice a day. Before that he drank very little whiskey and no beer.

His average weight is 175 pounds. He has lost 40 pounds.

Present Illness.—This began about six months ago with an attack of indigestion for which his physician advised drinking of quantities of water. Following this advice the patient gradually developed a marked polydipsia and polyuria with the urine of pale color. He noticed that he ate more than formerly, but nevertheless lost weight rapidly. His bowels became constipated until a cathartic was always required. The patient grew weaker and weaker, finally becoming

exhausted after the slightest effort. In February, three months ago, he spoke to a physician, who on examination found sugar in his urine. Sexual power is much diminished.

Physical Examination.—The patient is a large man, has evidently lost weight. The pigmentation over the dorsum of the hands, over the face and neck, and over the legs and feet, from the knees down, is of an extreme brown color. The nipples, axillæ, genitalia, and skin of the body show no increase. There is scattered freckling over the chest and arms. The skin is dry and scaly. There is very little beard and the hair elsewhere is small in amount.

The pupils are unequal and react to light and accommodation.

The tongue is cracked and coated; the throat injected; and there is pus in the posterior pharynx. There is no general glandular enlargement.

The heart and lungs are negative.

The spleen is not felt.

The liver is enlarged, the dulness beginning on the fourth rib, flatness on the sixth rib and extending to the costal margin. The edge is felt at a point 3.5 cm. below the costal margin in the right mammillary line, is smooth, firm and resistant. It crosses the midline 7.5 cm. below the ensiform.

Descent of the liver with respiration is visible. There is also seen some lazy peristalsis about the umbilicus. Just to the right of the umbilicus there is a tender prominence corresponding to the head of the pancreas. No definite mass can be felt.

Genitalia.—The right testicle is atrophied.

Extremities.—There are no scars nor edema. The deep reflexes are sluggish. The left knee-jerk is obtained on reinforcement; the right knee-jerk not at all, nor are any of the other deep reflexes.

Urine on admission was pale yellow, acid, of specific gravity 1,030 and contained 48.6 gm. of sugar in the twenty-four-hour specimen, or 5.4 per cent. There was no albumin nor any casts. Acetone and diacetic acid were present. For subsequent examinations see the table.

Examination of the Blood.—Red cells, 5,320,000. Leukocytes, 7,080. Hemoglobin, 94 per cent. (Sahli). In a preparation of the fresh blood the red cells seem normal in size, shape and color. The blood smears readily with tendency to rouleaux formation. There are no evidences of degeneration. The leukocytes are not increased; no eosinophils are seen. Numerous fat droplets are seen in fresh smears. The diagnosis is verified by staining with Sudan III.

On June 3, the temperature, which had been about normal, rose to 100.2, and a red, tense, boggy swelling was discovered in the region of the left parotid gland. For the past few days the patient had been growing weaker and more toxic with a marked, fruity odor to the breath. The skin was very dry and desquamating on the hands where the pigmentation was definitely increased.

Over the abdomen there were many small reddish-purple spots, the color of which did not fade on pressure.

The liver seemed slightly larger than before and somewhat tender.

Blood Examination, June 4.—In a fresh smear the red cells show moderate distortion with tendency to a raquette shape. The color appears normal. No fat droplets are seen. Leukocytes number 3,000 per c.mm.

Later the right parotid became swollen and tender and the coma deepened in spite of increased carbohydrate in the food and intravenous infusions of alkali.

On June 1 the Wassermann reaction of the blood serum was negative.

The temperature remained about normal until June 2, when coincident with the parotid infection, it rose to 100.2. It continued to rise gradually until on the evening of June 4 it was 107, and the pulse 130. Just after the death of the patient some blood was aspirated from the right auricle. This showed great numbers of *Staphylococcus aureus* in pure culture.

Clinical diagnosis: Hemochromatosis; diabetes; double parotitis; septicemia.

Autopsy No. 3,228. (Autopsy one hour after death.)

Anatomical Diagnosis: Hemochromatosis (bronze diabetes); extreme pigmentation of liver, pancreas, lymph-nodes, heart, gastric mucosa, thyroid, submaxillary gland, Brunner's glands, adrenals, parathyroids, prostate, testes, seminal vesicles, and skin of face, hands, and lower legs; cirrhosis of liver, pancreas and lymph nodes; chronic splenic tumor; ascites; purulent tonsillitis; acute diphtheritic enteritis; edema and atelectasis of lungs; chronic adhesive pericarditis; chronic gastritis; chronic orchitis with atrophy; chronic fibrous myocarditis; chronic prostatitis.

TABLE 2.—URINARY FINDINGS, CASE 3

Date.	Total Amount of Urine in c.c.	Specific Gravity.	—Sugar— Per Cent.	—Per Gm.	Acetone.	Diacetic Acid.	Diet.
5/21	900	1.030	5.4	48.6	+	+	Ward diet.
5/22	2,930	1.032	4.4	129.	+	+	Ward diet.
5/23	2,600	1.033	6.	156.	+	+	Ward diet.
5/24	2,230	1.035	5.	113.	+	+	Carbohydrate-free, plus white bread, G. 25
5/25	2,300	1.035	2.5	32.	+	+	Carbohydrate-free, plus white bread, G. 25
5/26	2,920	1.035	2.3	67.	+	+	Carbohydrate-free, plus white bread G. 25
5/27	3,850	1.035	1.	38.	+	+	Carbohydrate-free, plus white bread G. 25
5/28	2,600	1.025	1.1	28.	+	+	Carbohydrate-free, plus white bread G. 25
5/29	2,600	1.025	0.5	13.	+	+	Patient refuses bread.
5/30	2,530	1.022	0.5	11.6	+	+	Patient refuses bread.
5/31	1,610	1.025	0.7	11.2	+	+	Patient refuses bread.
6/ 1	1,300	1.025	*		+	+	Glucose G. 10, q. h. ad. G. 50
6/ 2	1,750	1.021	1.5	27.	+	+	Glucose G. 10, p. h. ad. G. 50
6/ 3	2,680	1.025	3.9	104.	+	+	Glucose G. 10, q. h. ad. G. 50
6/ 4	2,710	1.025	3.7	97.	+	+	
6/ 5			..		..	..	

*Levorotatory.

The body is that of a large, fairly well-nourished, white man, 175 cm. long. The face and hands are deeply "sun-burned." The lower legs show a definite pigmentation, but not nearly as marked as the hands. The skin over the arms, legs and thorax is dry and scaly. The hair is slight in amount over the pubis and absent over the chest. There is a very small amount of subcutaneous fat. The abdominal cavity contains 1.5 liters of a clear, lemon-yellow fluid.

The *heart* is cut and torn considerably on removal through an abdominal incision. The pericardium is loosely adherent by old fibrous adhesions. The cavity is completely obliterated. The heart is small and the right side dilated. The left auricle is slightly dilated but not thickened, while the left ventricle is tightly contracted. The wall of the left ventricle measures 12 to 14 mm. The valve segments are all quite delicate with the exception of the aortic valves which though slightly thickened were evidently competent. The tricuspid ring measures 12 cm.; the mitral 9.5 cm. The myocardium of the right ventricle is a bright yellowish-brown color. Tangential section of the muscular wall of the left ventricle shows a very diffuse translucent tissue replacing a good deal of the heart muscle which is of a deep bronze color. The root of the aorta shows a considerable amount of sclerosis, especially around the orifices of the coronary arteries. The coronary arteries are diffusely thickened.

The *lungs* were edematous and showed some hypostatic congestion.

The *spleen* weighs 250 gm. and measures 13 by 8 by 4.5 cm. The capsule is thick, irregular and granular. On section the organ is of a uniform, dull red

color. The trabeculae are prominent, while the Malpighian bodies are very small and hard to see. The pulp is elastic and does not scrape off on the knife.

The *stomach* is dilated with fluid and gas. Its mucosa has a definite, rusty, pinkish color. On section this is seen to be due to a pigmentation of the mucosa. The muscle coats appear pale and normal. The rugae are considerably obliterated and the mucosa has a stamped leather appearance.

The *pylorus*, *duodenum*, and *bile-ducts* are normal.

The *gall-bladder* contains a very thick, slimy, black bile.

The *pancreas* is of a deep bronze color. It is definitely enlarged and somewhat increased in consistence. On section the lobules seem glued together.

The *liver* weighs 3,200 gm. The surface is slightly irregular and the edges rounded. The organ is very tough and elastic, and is deep bronze-red in color. Section is quite uniform in appearance, and has a rough, granular appearance. The lobules are greatly distorted; some of them being larger than normal. There are numerous depressed scars all over the cut surface. The vessels stand wide open. The liver tissue everywhere has this deep bronze tint. The connective tissue in some places is slightly translucent and grayish looking.

The right *kidney* measures 11.5 by 5.5 by 4 cm. and weighs 200 gm. Both kidneys show some congestion and cloudy smelling.

The *adrenals* are of normal size and general appearance except for a slight increase in pigmentation. The cortex is 2 mm. in thickness and the striae yellowish brown. The inner zone is deep brown, while the medulla is inconspicuous.

The *lymph-nodes* about the head of the pancreas, at the hila of the liver and kidneys and along the aorta are greatly enlarged, some of them measuring 2 or 3 cm. in long diameter. They are firm, but not very tough; in some instances indeed rather soft. They are deeply pigmented, the color showing distinctly through the capsule. On section this pigment is found to be of bright yellow or deep brown color. Some of the nodes below the thyroid are enlarged and show a definite rusty pigment, as well as black coal pigment. Their bronze tint, however, is not so marked as that of the nodes about the liver. Those just above the bifurcation of the trachea measure from 1.5 to 2 cm. in diameter, and are of a uniform, deep, rusty-brown color. They are conspicuous also just above the base of the heart. Here the pigmentation is as deep in some instances as the nodes in the neighborhood of the liver. Other nodes are scattered all through the fat of the anterior and posterior mediastinal regions. Many of them are of the size of small peas, and all are more or less pigmented. The nodes posterior to the hilum of the lung form a rather conspicuous chain along the esophagus. These, too, show a deep yellowish-brown, uniform pigmentation with a rather conspicuous amount of connective tissue. They are of all sizes and in places somewhat matted together. The bronchial nodes are enlarged and more or less pigmented, but contain more black than yellow pigment. The nodes become more conspicuous as we go along the aorta downwards, and their architecture is very sharply marked out by the bright yellow pigment. They are very numerous and large along the lumbar region, where they form a continuous chain beside the aorta. The nodes in the pelvis do not show this pigmentation, but are of a uniform, pinkish-gray color. They are somewhat larger than normal and rather juicy. The mesenteric nodes are examined with considerable care. They are slightly increased in size, some of them measuring 1 or 1.5 cm. in long diameter. They are uniform, pinkish-gray and on section moist and juicy, but have no abnormal pigmentation. They contrast very conspicuously with the nodes around the duodenum and pancreas close beside them which contain large amounts of the ochre pigment, together with a certain amount of black pigment in some instances.

The *aorta* shows a moderate grade of sclerosis throughout its entire course.

The *submaxillary gland* has a regular architecture. The gland acini are pigmented a deep yellowish-brown, much the color of maple sugar.

The left *tonsil* is large, hemorrhagic in appearance and contains some pockets of creamy material. *Pharynx*, *larynx* and *esophagus* are quite normal.

The *thyroid* is rather small and fibrous, and on section the connective tissue is rather conspicuous. The colloid substance is easily seen. The tissue has a deep bronze tint.

The *parathyroids* are of normal size and appearance, but the pigmentation is darker than that of the normal gland.

The *jejunum* is considerably dilated, and its mucosa swollen, pink and velvety-looking. No ulcerations are present. The same condition is found in the upper part of the *ileum*, but 60 cm. from the ileocecal valve the mucosa becomes very deep red, swollen and edematous and in places has a charred, granular appearance. The surface is covered with little knobs and tags of granular fibrin. The surface throughout is granular and dry. The same condition is found in the *cecum*. There are thickly scattered ecchymotic patches. In the *sigmoid* a considerable membrane of rather uniform appearance is found.

The *bone-marrow* of the femur is somewhat cellular, but soft and flabby. About two-thirds of the tissue is evidently fat, the remaining third being pale and cellular.

The right *testicle* is very small, most of the gland tissue being replaced by radiating scars. The tubules, where remaining, have a very pale, brown color. The left testicle, in general, has the same appearance, but is larger, measuring about 2 cm. in long diameter. There are scars in this testicle as well. The tubular structure has a deeper brown color.

MICROSCOPIC EXAMINATION

Liver.—The liver is divided into lobules, varying greatly in size and shape, by rather broad and cellular bands of fibrous tissue.

The pigmentation is very heavy, more marked than in the other cases, and very evenly distributed throughout the section. Each cell is almost filled with the mass of very fine brown granules. The nuclei are often eccentric and rather small. In every lobule there are one or more areas in which the cells are broken down and the pigment lies free in heavy masses or contained in phagocytic cells. These areas resemble those already fully described in the preceding cases. There are other places in which the cell relations are preserved but the cells excessively vacuolated and poorly stained in addition to being heavily pigmented. A prominent feature of this liver is the presence of nodes of liver cells, larger than their fellows, with large deeper staining nuclei and containing very little or no pigment. They vary from 0.25 to 0.5 mm. in diameter, are round or oval and are evidently masses of newly formed liver cells. In the recently regenerated areas the cells of Kupfer are heavily pigmented as elsewhere in the sections, while the parenchymatous cells contain only a fine, dust-like pigmentation, and occasionally even this is not evident (Fig. 6). There are other more or less circumscribed areas in which the cells are more definitely pigmented, but much less so than in most of the surrounding tissue, representing probably regenerated tissue of longer standing.

The connective tissue septa are rather cellular and contain numerous fine capillaries. Pseudo-bile-ducts are fairly numerous, also, and always pigmented. Pigment is found scattered throughout the connective tissue in the interstitial spaces, in wandering cells, in the pseudo-bile-ducts, and in the endothelium and walls of blood-vessels, but is less abundant, compared with that of the liver cells, than in the other cases of this series.

The so-called pseudo-bile-ducts are among the most heavily pigmented structures found. Transitions may readily be traced from the ordinarily pigmented liver cell, through those more heavily pigmented which are arranged in two parallel rows and the very heavily pigmented pseudo-bile-ducts, to those in which many of the cells have been entirely destroyed leaving an elongated mass of free pigment. Many such elongated structures are seen, one end consisting of heavily pigmented cells and the other of free pigment masses.

The connective tissue tends to be annular in arrangement, and does not invade the masses of liver cells to any extent. The lobulation is quite evidently a false

one. Sometimes a central vein may be located near the periphery of a mass of cells. Often the most degenerated areas and the newly regenerated portions seem to be in close relation to a central vein, and occasionally, a new growth of pigmented connective tissue can be found about what is apparently a small radicle of the hepatic vein. The endothelial cells of the blood-vessels, large and small, are frequently loaded with pigment, but the true bile-ducts contain little, and often are entirely free.

Hemofuscin is present in considerable amount in the walls of blood-vessels. Elsewhere the pigment reacts positively in iron tests.

Pancreas.—Sections of pancreas show a distinct lobulation, and there is practically no infiltration of fat between the lobules. The gland is heavily but not uniformly pigmented. Every cell seems to contain at least a few grains and the majority are but slightly pigmented, but there are several patches in each lobule in which the cells are loaded with pigment or have already disintegrated with the deposit of masses of pigment and the ingrowth of granulation tissue. These areas resemble those in the liver, but the focal nature of the pigmentary degeneration is much more pronounced in the pancreas.

In the least affected regions, the acinar arrangement, while not so clear and distinct as normal, is quite evident. The cells are slightly or moderately pigmented, their nuclei stain well, in some cells pyknotic, but the borders of the cells are very indistinct and they fade into each other. Some sections show edema where the acini are forced apart and rendered more distinct. In others they are close together and at times resemble cords or masses of nuclei with little distinct acinar arrangement. There is only the slightest amount of connective tissue growth in these less affected zones.

On the other extreme there are found masses of pigment, which, on close examination, reveal themselves to be excessively pigmented cells sometimes barely preserving some trace of their former arrangement. Into such areas, the connective tissue is growing and the later stage of deeply pigmented scar tissue may also be seen.

There are intermediate stages, also, in which heavily pigmented cells preserve their glandular arrangement.

The interlobular connective tissue is slightly increased in amount and shows scattered pigmented wandering cells. The mucosa of the larger ducts is free from pigment, but their walls and those of the blood-vessels show pigment bearing cells.

All the islands of Langerhans are either in the center of or in close association with one of the areas of excessive pigmentation and degeneration (Fig. 7). The islands vary in size; some few are large and well formed, more are quite small and contain only a few cells which are, however, well preserved. They are not numerous, however, and the degenerated areas are seen frequently without an island of Langerhans. As a rule, each island contains a few pigment granules.

The pigment reacts positively to tests for iron, except that in the walls of the large blood-vessels. This pigment is very finely divided and of a lighter color (Fig. 8).

Heart Muscle.—The heart muscle is deeply pigmented and here, too, the pigmentation is patchy. There are groups of cells heavily loaded throughout their protoplasm with the golden-brown grains alternating with other areas, in which the pigmentation is much lighter and more nearly confined to the undifferentiated sarcoplasm at the nuclear poles. The pigmentation is heavier than in the preceding case, but there has been little destruction of cells. A few very small areas may be found, in which occur mononuclear wandering cells loaded with pigment, but for the most part the pigment is confined to the intact muscle cells. The nuclei, as a rule, stain well; in some cells, the chromatin is small in amount. The appearance of the individual cells conforms to the description given for the preceding case. Almost all the pigment reacts positively to iron tests. It is only by rather careful search that some unchanged pigment is seen (Fig. 9).

Spleen.—There is a diffuse increase in the connective tissue. Many large pyknotic and karyorrhetic nuclei are seen, the protoplasm of their cells staining deeply with eosin. There is a slight increase in pigmentation. The pigment is found in golden-brown granules or small masses, scattered here and there throughout the splenic pulp. It is found, usually, in phagocytic cells with very irregular vesicular nuclei and often occurs free in the interstitial tissues or in the sinuses. Occasionally, heavily loaded phagocytes show marked signs of degeneration with nucleus undergoing karyorrhexis or karyolysis, but the majority of the large pink cells with pyknotic nuclei are free from pigment. No pigment is found in the large lymphocytes nor in the endothelial cells lining the sinuses. The pigment reacts positively to iron tests.

Submaxillary Gland.—There is a definite increase in the amount of connective tissue which is principally interlobular in distribution and to a less extent inter-acinar. In structure, it is loose and cellular with many mononuclear cells in its interstices.

The parenchymatous cells are practically all pigmented with finely divided golden-brown granules. Some of the tubules are heavily loaded, their structure almost lost or obscured by the surrounding granulation tissue; others are only slightly affected with a few scattered grains. Serous tubules greatly predominate, but there occur, scattered through the gland, groups of tubules which have a very well-defined basement membrane and practically clear non-staining cells with small, dark misshapen nuclei lying close to the basement membrane. A very fine, delicate network can be made out in some of the cell bodies and fine lines marking the cell boundaries. Most of these cells show scattered pigment in fine grains. There are crescent-shaped serous cells around the ends of these tubules—the demilunes of Giannuzzi—which, like the other serous glands, are much more heavily pigmented. Usually the demilunes are distended with the golden-brown granules and stand out very prominently in contrast with the clear mucous cell-bodies. In some of the mucous tubules the cells cannot be distinguished or nuclei found. There is simply a foamy, almost clear, space surrounded by the crescents and basement membrane. The serous tubules are the more heavily pigmented. Many of them are filled with granules, while others are only slightly affected. In most cases, the tubular arrangement is well preserved, but in places this is quite indistinguishable, and the cells are found apparently in columns and strands.

The connective tissue is almost free from pigment. A few pigment-bearing cells are found on careful examination. Some of the small interlobular arterioles have in their walls a pigment which very closely resembles that of the gland cells, but is perhaps of a lighter color. This pigment does not give a positive reaction with Nishimura's method, while the great majority of that in the secreting cells does. There is some brownish pigment remaining unaffected by the ammonium sulphid and other reagents. It is scattered rather diffusely throughout the sections and seems to belong especially to the serous glands. Unlike most reported observations, the pigment of the demilunes of Giannuzzi reacts positively as a rule, though in a few places there was no change in color. I could determine no difference in the cells or pigment which showed positive tests from those reacting negatively. Sometimes the same cell contains both blue and brown granules. The protoplasm of the mucous cell stains a diffuse light blue by Nishimura's method while their pigment grains stand out as much darker blue.

Stomach.—The section of tissue preserved shows both stomach and duodenum with the usual gastric glands and at the other end of the section typical Brünner's glands. The secretory cells are heavily pigmented with a very finely and evenly divided golden-brown pigment. This pigmentation involves those glands near the muscularis mucosæ especially and extends to a point about half-way to the free surface of the mucosa.

There is no pigment in the superficial half of the mucosa. All the cells of Brünner's glands are also heavily pigmented and the cells of their ducts contain granules to a point about half way through the mucosa. Some of the cells have

excessive deposits and show poor staining and distortion of the nuclei. As a rule, however, each cell contains only a moderately heavy accumulation in the end of the cell toward the lumen of the tubule. Only rarely is there found a pigment granule in the interglandular tissue. All this pigment gives a positive reaction for iron.

There is another pigment in the section which differs from the foregoing, especially in its failure to respond to iron tests and in its characteristic location. It has, too, a somewhat different morphology, being in finer particles and having a slightly lighter color. This is found in the walls of the blood-vessels. Here and there a muscle cell of the muscularis mucosæ or of the main muscular coats may show these fine yellowish-brown grains. It is found in the vessel walls characteristically in rather blunt, spindle-shaped cells with light blue oval nuclei, but here and there the pigment seems to be in the muscle-cells themselves. Occasionally the containing cell is quite large, irregular in shape with a large round or somewhat distorted, vesicular nucleus. This pigment is absolutely unaffected, apparently, by the various reagents used microchemically to detect iron, and corresponds to von Reeklinghausen's "hemofuscin." According to many writers, in sections stained with methylene blue, the hemofuscin takes a darker color, blue or violet, and is thus rendered much more distinct. I have been unable to substantiate this in this case. The pigment remained unaffected by Gruber's polychrome-methylene blue. Perhaps there is some difference in the stains used.

Sections from the duodenum lower down show a similar picture. In sections from the lower portion of the jejunum no iron reacting pigment (hemosiderin) is found and but little of the non-reacting hemofuscin in the walls of the vessels. There is, however, a striking inflammatory condition of the common type. Hemofuscin is present in the walls of the larger vessels but much less conspicuously than in the duodenum.

Lower down in the ileum the inflammatory process is further advanced and more hemorrhagic. The prominence of the acute condition overshadows any possible chronic enteritis which may have been present. The mucosa seems decidedly thin, however, and somewhat distorted. Hemofuscin is found in the vessel walls in amounts similar to that in the jejunum, and in addition, it may be quite plainly seen here and there in the fibers of the longitudinal muscle coat. Perhaps the excessive edema causes it to stand out more prominently. The pigment is not noted in the circular layer whose fibers are cut transversely.

Section of the colon shows dilatation of blood-vessels, some edema of the submucosa with the presence of large mononuclear cells in slightly increased amounts, and a few polymorphonuclear leukocytes. The mucosa and the muscle coats seem quite normal. In the interglandular spaces and very near the surface of the mucosa there is found a goodly number of large cells, mononuclear, which are distended with a very fine brownish pigment. They seem to be in close association with the capillaries. They are not observed in the depths of the mucosa, and no other pigment is seen in the section. It does not give a positive iron reaction.

Thyroid.—The cells lining the alveoli are heavily pigmented, even more so, perhaps, than those in Case 1 and much more than Case 2. The alveoli, themselves, are larger and contain more colloid material than in the first case, giving a slightly different picture, but except for the pigmentation neither gland is far from normal.

As in the first case, it is in the smaller, more irregularly shaped alveoli, containing a small amount of colloid, that the cells are most heavily loaded. This may be stated as a general rule, but is by no means invariable. The nuclei may be obscured or pushed aside, but where plainly seen are always apparently in good condition. Pigment is not found in the interstitial tissue. Instead of the usual hemofuscin, the larger darker granules of hemosiderin are present in quite noticeable amounts in spindle-shaped connective tissue cells in the walls of the blood-vessels.

The pigment in the glandular cells reacts positively to iron tests, but not so readily as that in some other situations (Fig. 10).

Kidney.—The capillaries, including those of the glomeruli, are congested and the convoluted tubules show a mild grade of parenchymatous degeneration with swollen granular cells surrounding a lumen, which contains a small amount of detritus. There is very little pigmentation. This occurs in small groups of three or four collecting tubules either in or close to the pyramids of Ferrein. Sometimes Henle's loop is involved. Those tubules which are affected at all have a heavy deposit. The pigment occurs in rather coarse grains, which not infrequently distend the cell, pushing the nucleus aside. It reacts quickly with iron reagents. No hemofuscin is found.

The amount and distribution of the pigment in the kidneys of the three cases are practically the same.

Testicles.—Section from the right testicle shows a greatly thickened capsule and complete atrophy of the seminiferous tubules. The basement membranes are intact and thickened, but only traces of cells remain in the tubules which are shrunken and massed together in lobules separated by bands of thick, fibrous tissue. The interstitial tissue between the tubules is loosely arranged and shows the presence of large mononuclear cells and thick walled, hyaline capillaries but no pigmented cells of Leydig are seen. Both the capillaries and larger vessels have greatly thickened walls, many of them showing advanced hyaline degeneration. There is a striking pigmentation of the endothelial lining cells of these vessels and of their walls, as well. The pigment is in rather coarse golden-brown grains and gives the iron reaction readily, even that part of it in the muscular walls of the vessels, the characteristic site of hemofuscin.

With Perl's and Nishimura's tests the whole vessel wall sometimes takes a diffuse light blue color, but the dark blue granules stand out quite distinctly. This pigmentation extends to the vessels of the thickened capsule and is quite prominent there. The pigment in the testis in Case 2 shows the same distribution, but in much less conspicuous amounts. In no other organ have I found such extensive pigmentation of the endothelial and muscle walls of the vessels, and in this organ and the thyroid only have I found pigment in the muscle wall of an artery reacting positively in tests for iron. No pigment whatever is seen which is not iron-reacting.

Sections from the left testicle show apparently the same process in a less advanced stage. The capsule is thick as are also the basement membranes of the tubules. The germ cells are in various stages of degeneration. Mitotic figures are seen, but no spermatozoa. Running through the section are broad bands of loosely arranged fibrous tissue containing shrunken atrophic tubules like those in the other gland. The interstitial tissue is edematous and contains mononuclear cells and capillaries similar to those in the right testis. In addition, there are also present the cells of Leydig with an excess of very fine, diffuse, yellow pigment throughout their protoplasm. This pigment is negative in all iron tests. In the tubules themselves there occurs a diffuse yellow pigment quite similar, if not identical, with that in the interstitial cells. As a rule it is found in degenerated cells with a vacuolated protoplasm often showing no nucleus, and may be seen in small or moderate amounts in all the living tubules, but rarely in those completely atrophied. It occurs occasionally in large yellow masses in the lumen of an atrophic tubule either free or within a cell, the nucleus of which is still visible. This pigment does not give the iron reaction.

The amount of the iron-reacting pigment in sections of the left testicle is very much greater than in those of the right, and there is a very distinct difference in the same section between the pigment content of those vessels in the atrophic, scarred areas and that of the vessels in the better-preserved portions. In the latter, the endothelial cells seem to be almost completely filled with pigment, while in the scars between the atrophied tubules the vascular endothelium is only moderately or slightly pigmented and occasionally not at all. This may be partially explained by the fact that the vessels in the scarred areas have

smaller lumina, thicker walls and more frequently show hyaline degeneration. But this is not sufficient, since vessels of the same size and similar appearance, otherwise, show great difference in their pigment content, dependent, apparently, in some way, on their relation to the seminiferous tubules.

Prostate.—Judging from the piece of tissue cut for microscopical examination, the prostate was about normal in size and configuration. Histologically there is a moderately advanced chronic prostatitis. The gland acini seem slightly compressed by the somewhat hypertrophied stroma.

The cells of the acini are fairly heavily pigmented with golden-brown grains which react positively in iron tests. The endothelium of the capillaries in close association with the basement membrane is frequently heavily loaded with hemosiderin which in some fields is more conspicuous than that in the epithelial cells. The prostatic ducts contain less pigment. As a rule, the nearer the urethra, the less pigment, and the greater proportion of granules which do not react positively for iron. A few cells in the lining membrane of the urethra contain clusters of hemosiderin granules about the nucleus. These are quite sparsely scattered and are found only in the deeper layers near the basement membrane.

The capillaries and small vessels of the stroma are pigmented in a way quite similar to that in the testicles, but to much less extent.

The epithelial cells of the vas deferens in the ampulla and of the seminal vesicles, both the stratified cylindrical lining membrane and the columnar and cubical cells of the glands, are heavily pigmented with a light yellow granular material which does not give the iron reaction. This same pigment is found in large quantities in the vas deferens further back and in the epididymis as well. In color it closely resembles that in the testis.

Hemosiderin is present in small amounts in the capillaries of the interstitial tissue in all sections of the sexual apparatus.

Adrenal.—The sections of this organ very closely resemble those of the preceding case. There is the same heavy pigmentation of the glomerular zone with hemosiderin (Fig. 11), the increase in the normal pigment of the reticular zone and the presence of a few pigment granules in the medulla, some reacting positively for iron, others negatively. There are, however, minor differences. In this case the pigmentation of the outer layer is more general, involves more cells and more frequently dips down into the fasciculate layer as a long continuous or interrupted strand of pigmented cells. The pigment grains are even and regular in size and are gathered around the nucleus in the less heavily affected cells. Many cells are filled with pigment, but the nucleus, as a rule, stains well and only occasionally is there evidence of cell destruction with the deposition of small masses of pigment. There is really more pigment present in this adrenal than in that of the preceding case, but one gets the impression that the process is much more recent and less complete. There is little cell destruction, little pigment has been set free in the interspaces and the lymphatics cannot be traced by their injection with pigment as in Case 2. In both cases there seems to be less of the lipid substances in the cortex than normally. The pigment in the zona reticulata is unaffected in the iron tests.

In tissue fixed in formaldehyd solution and potassium chromate, the medullary cells show a very diffuse yellow pigment. In addition there are many granules closely resembling those of hemosiderin in size and appearance, but they are not present in tissue fixed by other methods. The chromaffin tissue of the adrenal seems to be about normal or slightly increased in amount; certainly not diminished. The sympathetic ganglia and other tissues, in which this material occurs, were not examined.

Skin.—The cylindrical basal cells of the Malpighian layer of the epidermis contain a fine yellowish-brown pigment which is not affected by ammonium sulphid or by potassium ferrocyanid and hydrochloric acid. This pigment varies in amount in different cells from that scarcely perceptible to a very considerable pigmentation.

In the corium and subcutaneous tissues there are many scattered, branched, spindle-shaped or elongated cells containing the familiar iron-reacting, golden-brown granules of hemosiderin morphologically and microchemically identical with that in the internal organs. The nuclei of these cells are round, oval or irregular and elongated with a very distinct nuclear membrane and no definite nucleolus. The cells are in general scattered evenly between the connective tissue fibers, except about the blood-vessels, where they are collected in large numbers. The endothelium of the vessels is entirely free. In the muscle coats iron-free hemofuscin granules are occasionally seen. No sweat glands are seen in the sections of tissue available.

This picture is quite different from that presented in Case 2, where the pigmentation is limited to the basal cells of the epidermis and does not react positively in iron tests. Both kinds of skin pigmentation are reported and quite naturally have led the observers to different views concerning the clinical value of a microscopic examination of the skin in a suspected case of hemochromatosis.

Bone-Marrow.—The bone-marrow of the femur shows some edema, slight hyperplasia and a small increase in pigmentation. The general appearance is like that in the preceding case. There is the same preponderance of fat and the edematous areas between the fat cells containing the fibrin-like stroma, but the blood-forming cellular areas are slightly more numerous than in Case 2. The hematopoietic islands are not uniformly scattered among the fat cells, but many are grouped in one corner of a section while there are many fields in another portion of the same section which show nothing but fat. Cells of the myelocytic series greatly predominate, but here and there are found groups of megaloblasts and normoblasts. The hyperplasia is, however, very slight and in no way suggests any active new blood formation. The pigment is rather less than in Case 2 and for the most part is intracellular and iron-reacting. The endothelial lining of the blood-vessels contains no pigment.

Lymph-Nodes.—In the bronchial nodes, hemofuscin is present in considerable amounts in the walls and adventitia of blood-vessels and in the capsules of the nodes. In one, especially, in which the hemosiderin is relatively small in amount compared with that in other nodes, and in which there is even less coal pigment, the black pigment is more nearly in the center of the lymph-cords, while the iron-containing pigment is confined to the sinuses or their immediate neighborhood.

Those from the anterior and posterior mediastinum contain a moderate supply of hemofuscin and hemosiderin along with more or less coal pigment and show the presence of more fibrous tissue. In the tissue between separate nodes there are many connective tissue cells containing hemofuscin and small strings of hemosiderin suggesting its passage through lymph-spaces and vessels.

There are sections of several mesenteric nodes which in gross were not pigmented. In one there is a rich accumulation of hemofuscin in the walls of the vessels and in the capsule and a small amount of hemosiderin in the sinuses and cords, which might be easily overlooked with low powers of the microscope. On more careful examination, it is found to be present in amounts about equal to or somewhat greater than, for example, that in the spleen. Another smaller node shows conspicuous hemofuscin in the vessel walls, but not so much in the capsule and connective tissue generally. The hemosiderin is present in very small amount, collected here and there into large, tightly packed mononuclear cells.

A pelvic node shows a great deal of hemofuscin in its capsule and vessel walls and only an occasional small granule of hemosiderin. The sinuses in one portion are widely dilated and contain a few very large mononuclear cells. There is no trace of the fine yellow pigment seen in the sexual organs.

The most heavily pigmented are the portal nodes and those immediately surrounding the splenic artery. Some of these seem to have been converted into masses of pigment which completely obscure the structure of the larger part of the node. The trabeculae are increased in thickness, and are quite prominent. In the connective tissue cells of the capsule, trabeculae and walls of the vessels,

hemofuscin is present in large amounts. It is especially conspicuous at the hilum of the nodes where the connective tissue trabeculae are most prominent and the many cells are filled with the fine yellowish-brown granules which do not show microchemically the presence of iron.

The walls of the *splenic artery* are not pigmented or at most show a few granules here and there in the adventitia. The intima is not thickened nor is any abnormality observed. In the surrounding connective tissue, hemosiderin is found in the endothelium of the capillaries including those of the nerves, but neither here nor in any other sections has pigmentation of the nerve tissue itself been noted. Hemofuscin is found in the walls of the smaller vessels.

Sections of the *aorta* show a slight irregularity in the thickness of the intima which is not, however, anywhere markedly thickened. The media seems normal, and there is no pigmentation noted in any of the coats nor in those of the *vasa vasorum*. The cells of the peritoneum above the aorta stain deeply, but no definite granular pigmentation is observed. In the connective tissue immediately below the serous layer there are a few scattered cells containing a yellowish-brown pigment resembling hemofuscin.

MICROCHEMICAL REACTIONS OF PIGMENTS

In studying the effect of various reagents on the different pigments found in hemochromatosis, my experience has been that of most other workers on the subject.

Hemosiderin, for example, is dissolved or disappears from sections treated with 10 per cent. sulphuric acid or with hydrochloric acid alcohol, while the hemofuscin in such sections is not appreciably affected. It may be stated, as a general rule, that the pigment which reacts readily for iron is more quickly dissolved by acids than that in which it is more difficult to demonstrate iron. The pigment in the liver cell shows its iron readily when treated with the usual reagents, and quickly disappears under the influence of acids. On the other hand, the pigment found in the thyroid responds more slowly and less certainly to the reagents used in the demonstration of iron, and sections of the thyroid show a considerable amount of undissolved pigment after treatment with acids for a time sufficient to dissolve all the pigment from a liver section of equal thickness. If the section is then treated with ammonium sulphid or potassium ferrocyanid and hydrochloric acid the undissolved residue does not give an iron reaction.

Hemofuscin entirely disappears under the influence of alkalis, sodium or ammonium hydroxid, and hydrogen peroxid, none of which has any effect on hemosiderin. Anschütz⁶ states that ammonium sulphid dissolves hemofuscin. I have not found this to be the case, for in sections tested for iron by Nishimura's combination method, which involves exposure to ammonium sulphid for at least one hour the amount of hemofuscin has not been appreciably altered. Even in sections treated for one week with a mixture of ammonium sulphid and glycerin at a temperature of 60 C. (Macallum⁷), a considerable amount of the yellow pigment remains unchanged. The iron in the nuclei is well shown in sections so treated, and the fact that much, if not all, of the hemofuscin remains yellow clearly indicates that, if this pigment does contain iron, the metal exists in firmer combination than that in the chromatin of the nuclei.

Other pigments, that found in the basal cells of the epidermis and the yellow pigment in the genital organs, resemble hemofuscin in their behavior toward the solvents above mentioned. The yellow genital pigment takes on a deeper reddish tint when treated with Sudan III, but is not dissolved by alcohol, ether or chloroform.

6. Anschütz: Ueber den Diabetes mit Bronzefärbung der Haut, zugleich ein Beitrag zur Lehre von der allgemeinen Hämochromatose und der Pankreasschrumpfung, *Deutsch. Arch. f. klin. Med.*, Leipsic, 1899, lxxii, 411.

7. Macallum: The Distribution of Assimilated Iron Compounds, Other than Hemoglobin and Hematin, in Animal and Vegetable Cells, *Quart. Jour. Micr. Sc.* London, 1895-6, xxxviii, 175.

INCIDENCE

In 1899 Anschütz collected twenty-four cases of hemochromatosis with diabetes. In 1907, Fitcher⁸ counted thirty-five and Heller⁹ forty-two, including several doubtful cases. Including those of this series and several cases recently reported very briefly by Simmonds,¹⁰ I have been able to find about fifty cases with diabetes and thirteen which did not show diabetic symptoms.

The patients' ages are between 30 and 65 years. The patient in Heller's doubtful case died at 73.

The only well-authenticated case occurring in a woman is that reported by Abbott. Murri¹¹ and Berg¹² each report one other clinically, but in neither case can the diagnosis be accepted without question.

Concerning the predisposing causes little is known. Occupation seems to play no rôle. Many authors emphasize alcoholism; six exclude it as a possibility. Abbott and Beattie¹³ are inclined to think that gastrointestinal disturbances play an important part in the etiology. Many different diseased conditions appear, of course, in the former histories of this number of cases, but such is their number and diversity that no importance can be attached to any one of them.

CLINICAL FEATURES

In most cases of hemochromatosis, in the form termed *diabète bronzé* by the French, we have clinically the picture of a severe diabetes with large amounts of sugar in the urine, polyphagia, polydipsia and polyuria; extreme weakness and cachexia, generalized pigmentation of the skin, but not of the mucous membranes; and an enlarged, often tender, liver.

This association of symptoms occurs with remarkable regularity but not with absolute constancy.

Abnormal pigmentation of the skin was noted in fourteen of the twenty-four cases of which reports were collected by Anschütz, and in practically all of the authentic cases reports of which have been published since the appearance of his paper. It varies in intensity from icteric, light sun-burn, to earth-color, lead-color and the extreme bluish-black shade described by Letulle. It occurs characteristically on the face, neck, dorsum of hands and feet, extensor surfaces of forearms and legs, around

8. Fitcher: Hemochromatosis with Diabetes Mellitus, Am. Jour. Med. Sc., 1907, cxxxiii, 78.

9. Heller: Ueber Hautveränderungen beim Diabète bronzé, Deutsch. med. Wehnschr., 1907, No. 30, p. 1216.

10. Simmonds: Ueber Bronze Diabetes und Pigment-Cirrhose, Berl. klin. Wehnschr., 1909, xlv, 531.

11. Murri: Ueber Bronzediabetes, Wien. klin. Rundschau, 1901, xv, 345, 361, 406, 424.

12. Berg: Diabète Bronzé and the Pathogenesis of Diabetes Mellitus: Illustrated by a Case of Bronze Diabetes Preceded by Basedow's Disease, Med. Rec., New York, 1899, lvi, 881.

13. Beattie: Hemochromatosis with Diabetes Mellitus, Jour. Path. and Bacteriol., 1903-4, ix, 117.

the nipples and genitalia, and occasionally in the axillæ. In these regions there is an intensification of the abnormal coloring which covers diffusely and less conspicuously the skin of the whole body.

In only two of Anschütz' cases were there pigment flecks on the buccal mucosa. These appeared in Ridder's¹⁴ recent case and in the very doubtful one reported by Murri. This is of some importance in differentiating the pigmentation of hemochromatosis from that of Addison's disease.

Anschütz states that in fourteen of seventeen cases in which this organ was described, the enlargement of the liver was clinically demonstrated. More recently Abbott and Simmonds have each reported a case in which the liver was smaller than normal (Abbott's case was one of simple hemochromatosis, without diabetes). The left lobe is sometimes enlarged disproportionately and the abdomen has been explored under the impression that the mass was that of an hepatic neoplasm.

The diabetes, as already stated, is severe and usually rapidly fatal. The history of the disease frequently begins with the symptoms of hyperglycemia; and after the onset of glycosuria, death, as a rule, occurs within a year, though cases have been reported in which diabetes was present for two or even four years (Letulle¹⁵) before death.

Heller in his analysis of forty-two cases gives the average amount of sugar as 4.7 per cent. and the average amount of urine excreted daily as 4.5 liters.

Several authors have noted the diminution of the glycosuria under treatment or on a mixed diet just before the exitus. In six of Anschütz' series, this symptom disappeared entirely for varying periods, as long as two months in one of Letulle's cases.

This brings us to a consideration of the cases of so-called simple hemochromatosis which do not show diabetic symptoms, such as Case 1 of this report and those of Opie, Abbott, Brault¹⁶ (two cases), Letulle, Foa,¹⁶ Gilbert and Grenet,¹⁶ Richardière,¹⁶ Osler¹⁷ and Simmonds (three cases). With the important exception of the diabetes these cases bear a close resemblance clinically to those of true bronzed diabetes, and pathologically the two cannot be distinguished from each other.

The interpretations of these cases in the literature varies with the writers' views concerning the etiology of the affection. It is suggested by some authors that these patients may have been observed during a latent period of the diabetes, yet some of the cases were followed for

14. Ridder: *Bronze Diabetes*, Deutsche. med. Wehnschr., 1910, xxxvi, 36.

15. Letulle: Deux cas de cirrhose pigmentaire du foie dans le diabète sucré, Bull. et. mém. Soc. méd. d. hôp. de Paris, 1885, 406; Letulle: Note sur trois observations de cirrhose hypertrophique pigmentaire non diabétique, Bull. et mém. Soc. méd. d. hôp. de Paris, 1897, p. 205.

16. Quoted from Hess and Zurhelle (see note 4).

17. Osler: Clinical Remarks; Hypertrophic Cirrhosis of the Liver with Bronzing of the Skin: Hemochromatosis, Brit. Med. Jour., London, 1899, ii, 1595.

fairly long periods and gave no history of polyuria, polydipsia nor polyphagia. Simple hemochromatosis is now generally held as the earlier or less severe form of the disease. Those who assume that the diabetes is due to changes in some one organ, as the liver or pancreas, hold that the lesion in that organ has not reached a sufficiently advanced grade to cause a marked disturbance in carbohydrate metabolism. Others, who believe that the three so-called cardinal symptoms, general pigmentation, cirrhosis of the liver and diabetes, are entirely independent of each other and are equally due to the same primary cause, give no explanation of these cases except that they are less severe forms.

Comparatively few records of clinical blood examinations are available, but those of the cases here reported, as well as those cited by Jeanselme,¹⁸ Osler, Fletcher and Ridder, show the red cells normal or slightly reduced in number and of normal size and general appearance. The leukocytes are not increased, nor, according to the differential counts in Osler's and Fletcher's cases, do they vary especially from their usual relative proportions. The slight leukocytosis as well as the moderate anemia in Case 1 might well be accounted for by the tuberculous peritonitis.

The hemoglobin does not depart far from the normal. Margain,¹⁹ in a rather incomplete examination without cell counts or hemoglobin estimation, noted "a certain number of red blood-cells completely deprived of hemoglobin," and in the plasma masses of free pigment. The latter observation is a very rare one. Marie²⁰ reports a similar one in a smear studied by Auscher, but thinks the pigment may have come from the skin, pricked to obtain the drop of blood.

The spectroscopic examination reported by Fletcher showed only the usual absorption bands of oxyhemoglobin.

Lipemia is occasionally demonstrable, notably in Case 3 of this report and in Case 2 of Hess and Zurhelle.

Evidence of pancreatic disease in examination of the stools is very rare. In Anschütz' case it was quite pronounced.

Microscopic examination of a piece of skin is of great diagnostic value when one finds iron-containing pigment in the cells of the cutis and sweat glands; but no reliance can be placed on a negative report, since in some cases there is present only the iron-free pigment in the basal cells of the Malpighian layer of the epidermis as in Case 2, in Abbott's case and others.

18. Jeanselme: *Hématologie et Pathogénie du diabète bronzé*, Bull. et mém. Soc. méd. d. hôp. de Paris, 1897, xiv, 179; Presse méd., 1897, p. 58.

19. Margain: Contribution à l'étude de l'origine du pigment dans le diabète bronzé et de la pathogénie de cette affection, Rev. de méd., 1905, xxv, 378; Progrès méd., 1904, xx, 472.

20. Marie: Sur un cas de diabète bronzé suivi d'autopsie, Semaine méd., 1895, xv, 229.

The three cases of this report are typical of the two well-recognized forms of the disease; Case 1 of simple hemochromatosis, Cases 2 and 3 of hemochromatosis with diabetes.

The onset in each case was rather abrupt. The patients had not noted the abnormal coloring of the skin or signs of liver disease before the appearance of subjective symptoms.

In Case 2 the pigmentation was not noted until the patient had been under observation about three weeks. In all three cases the pigmentation became progressively more intense. Heller says that in only three cases, those of Anschütz, Ubert²¹ and Hyss,²¹ did the appearance of the abnormal skin coloring antedate the diabetes. In one of Fletcher's cases the skin had been dark-colored for several years. In Anschütz' case and in Case 1 of Hess and Zurhelle, swelling of the liver had been noticed before the diabetic symptoms. Ridder recently had a remarkable opportunity to observe the development of a case. Four years before death, pigmentation of the extremities was noted; one year later, disease of the liver was diagnosed; diabetes was present during the terminal four months.

A history of moderate alcoholism was given by each of the three patients. In Cases 1 and 3, there had been marked gastro-intestinal disturbances and dysentery ten or twelve years before.

The complication of tuberculous peritonitis, present in Case 1, has been reported several times and is perhaps second in frequency only to pulmonary tuberculosis. It is noteworthy that in both the other cases there occurred an acute parotitis. In Case 3 septicemia was demonstrated by blood-culture. In Ridder's case, also, there was an otitis media with staphylococcus septicemia and bilateral purulent parotitis. At autopsy both glands were heavily pigmented and showed numerous abscesses. These were found also in the kidneys and heart muscle.

The Wassermann reaction was negative in the two cases in which this test was applied.

PATHOLOGY

At autopsy the most striking finding is the ochre or bronze color of many of the glandular organs, the heart muscle and lymph-nodes. In addition, there is usually a much enlarged and cirrhotic liver and more or less pronounced changes in the pancreas, either a small atrophic organ or an enlarged, fat infiltrated and fibrous gland, as the case may be.

The spleen is large and firm, but varies little from its normal color. Ascites may or may not be present.

The liver is a prominent feature in any post-mortem examination; small wonder then, that, with the peculiar abnormal coloring, the cirrhotic changes, the large size and often irregular contour, this "pigmen-

21. Quoted from Heller (see note 9).

tary cirrhosis" has been regarded as of commanding importance in the pathology of the disease. Yet the changes in this organ differ but slightly from those in the pancreas. The lymph-nodes, the spleen, the heart muscle and, occasionally, other organs show fibrous change of much less degree.

Only those lymph-nodes which lie in the path of the lymph-stream from some heavily pigmented organ show in gross the ochre coloring, indicating clearly that they play but a passive part in this process.

Microscopically, the pigment remains the center of attraction. This is of several different kinds: hemosiderin or iron-reacting pigment; hemofuscin or non-iron-reacting pigment; the increased amount of those pigments physiologically present in certain situations, as that of the adrenal reticular zone, the substantia nigra, the epidermis, the interstitial cells of the testis, and the finely divided yellow pigment (described in Case 3) in the tubules of the testis, epididymis, vas deferens and seminal vesicles.

Hemosiderin is the predominant and much the most conspicuous of the pigments. The wide extent of its distribution may be seen in the report of these three cases. The characteristic sites of its deposit are in the parenchyma cells of the glandular organs and in the muscle cells of the heart. It is found in the connective tissue of an organ only when the essential cells of that organ have disintegrated and deposited it in the intercellular spaces. It occurs, also, in the lymph-nodes, in the spleen, in certain cells of the cutis, and in the endothelial cells of blood-vessels in certain situations.

Hemosiderin has no definite and fixed chemical composition, but all the granules contain iron, since this is the criterion by which they are recognized as hemosiderin. Its iron content may be readily demonstrated in gross and microscopically by the use of the usual reagents, ammonium sulphid, or potassium ferrocyanid and hydrochloric acid.

The amount of iron in certain organs, notably the liver, pancreas and lymph-nodes, is enormous. As Auscher expressed it, the lymph-nodes had been converted into veritable mines of iron.

Table 3 shows the iron content as determined in the cases of Anschütz and Hess and Zurhelle in comparison with the normal and an extreme case of siderosis in pernicious anemia by Quincke.

TABLE 3.—IRON CONTENT OF LIVER, PANCREAS AND LYMPH-NODE

	—Liver—		—Pancreas—		—Lymph-Node—	
	Dried	Ash.	Dried	Ash.	Dried	Ash.
	Substance. ...	Substance. ...	Substance. ...	Substance. ...	Substance. ...	Substance. ...
	%	%	%	%	%	%
Normal (Oidtman).....	0.08					
Pernicious Anemia (Quincke)....	2.1					
Hemochromatosis (Anschütz).....	7.62	55.7	5.0	44.0	14.6	58.36
Hess and Zurhelle.....	7.1					

In these cases the liver contained almost 100 times the normal amount of iron. Hess and Zurhelle estimate the total iron in the liver of their case at 38.7 gm., about fifteen times that contained in the hemoglobin of a normal person and at least ten times that in the whole body of the average human being.

Hemofuscin varies greatly in the cases of this series. In Case 1 in the tissues examined it is hardly appreciable, scarcely more so in Case 2, while in the third case it approaches the relatively large quantity described by several other writers. The morphology differs somewhat from that of hemosiderin. It occurs in finer grains more evenly distributed throughout the cell protoplasm and has a lighter yellow color, rarely a darker brownish tint. It is found characteristically in connective tissue cells, in the capsules and trabeculæ of lymph-nodes and in the walls of blood-vessels, especially those of the lymph-nodes and gastro-intestinal canal, though von Recklinghausen describes it in the vessel walls widely distributed throughout the whole body. Chemically this pigment does not give a reaction for iron, is much more resistant to acids than is hemosiderin, and, according to Buss,²² takes a dark stain with methylene-blue. It disappears from sections treated with dilute alkalies or with hydrogen peroxid, neither of which affects hemosiderin.

Some confusion exists regarding just what pigments should be designated as hemofuscin. Certain authors seem to consider all pigments not giving an iron reaction as hemofuscin. Since the different microchemical methods vary considerably in their efficiency in the demonstration of iron, the Perl method showing less than the ammonium sulphid or the combination of the two suggested by Nishimura, it follows that the relative amounts of hemosiderin and hemofuscin in the same tissue would vary with the method used or in the hands of different observers. It is the contention of Hess and Zurhelle, as well as of others, that no sharp distinction can be drawn between them since they find both pigments side by side in the same cell. The non-iron-reacting pigment in the glandular cells resembles hemosiderin morphologically and is quite different in appearance from that found in the trabeculæ and capsules of lymph-nodes. The two extremes at any rate have very distinct differences which have already been indicated.

Several workers have noted increased pigment in the organs where it is found under physiological conditions and say, further, that in these regions an iron-containing pigment is never deposited. This, however, does not hold true for the heart muscle cell.

So far as I know, the striking accumulation of fine yellow pigment in the sexual organs in Case 3 is unique. Hess and Zurhelle mention in the gross description of one of their cases an intense pigmentation of the

22. Buss: Ein Fall von Diabetes Mellitus mit Lebercirrhose, Pankreas-atrophie und allgemeiner Hemochromatose, Inaug. Dissertation, Göttingen, 1894.

seminal vesicles, but no microscopical examination was made, nor is there any report of other portions of the genital tract. The seminal vesicles frequently show a similar pigmentation of much less degree, especially in old or functionally inactive cases.

Increase in connective tissue is found in those organs, notably the liver and pancreas, in which great numbers of the cells have degenerated and deposited pigment in the interstitial spaces; and in the lymph-nodes and spleen, to which the pigment has been conveyed through the lymph- and blood-streams.

Individual cases of hemochromatosis present distinct differences in the distribution and extent of the lesions in various organs as may be seen by a comparison of the three cases here reported. Some idea of the grade of the process may perhaps be obtained by attention to certain details. In Case 2, for instance, the even distribution of the pigment in the liver cells with relatively few and small focal areas of extreme pigmentation; the fibrous nature of the connective tissue strands in the liver; the old sclerotic lesions in the pancreas; the degeneration of the cells in the glomerulation zone of the adrenal with pigment deposition in the interstitial tissue and the presence of this pigment in the lymphatics of the capsule; all these points together suggest to me a comparatively sluggish process of long standing. On the other hand, in Case 3, an acute and rapidly progressing lesion is suggested by the larger and more numerous focal areas of pigmentary degeneration in the liver and pancreas; by the more cellular nature of the connective tissue strands in these organs; and by the heavy pigmentation of the adrenal glomerular zone and of the cardiac fibers with little or no cellular disintegration and, consequently, no increased connective tissue.

PATHOGENESIS

Some idea of the development of the subject is of value in the discussion of its pathogenesis. These two phases will, therefore, be considered together in this section as briefly as is consistent with a clear expression of the authors' opinions.

According to Osler, Troisier, in 1871, was the first to mention bronzed cachexia occurring in diabetes.

The first description to attract attention, however, is that of Hanot and Chauffard,²³ in 1882. They described two cases of diabetes associated with pigmentation of the skin and enlarged liver. At autopsy there was cirrhosis of the liver and fibrosis of the pancreas with a great accumulation of brown pigment, both in the gland cells and in the interstitial tissue.

Quincke,²⁴ Tillmanns²⁴ and Hindenlang²⁴ had already described cases with rather extensive pigmentation throughout the body, associating the

23. Hanot and Chauffard: *Cirrhose hypertrophique pigmentaire dans le diabète sucré*, *Rev. de méd.*, 1882, ii, 385.

24. Quoted from Opie (see note 3) et. al.

pigment deposits with severe anemias or with large extravasations of blood.

Hanot and Chauffard consider the diabetes primary and the other manifestations merely complications induced by diabetic endarteritis. Under the conditions of diminished circulation thus brought about, the liver cells produce more coloring matter than normally, leading to a deposit of pigment, a pigmentary hypergenesis. From the liver, the pigment is distributed to other organs, probably, in an embolic manner.

Letulle, in 1885, reported two similar cases and was impressed by the extreme pigmentation of the heart muscle cells and by the appearance of degeneration in the pigmented liver cells rather than any indication of activity as might be expected in a hypergenesis. The pigmentation of the cardiac fibers with little or no pigment in the interstitial tissue, he considers an unanswerable argument against the embolic distribution of the pigment from the liver. He thinks the diabetic condition induces an alteration of the blood, the pigmentation being due to a reduction *in situ* of the hemoglobin whether it be of the blood or of the different tissues.

Brault and Galliard, in 1888, follow Letulle in believing the pigment deposited wherever found in the gland cells, but emphasize the liver more strongly. They consider the diabetes and liver cirrhosis concomitant factors in the production of pigmentation. The altered blood-pigment furnished daily to the degenerated liver cells cannot be well elaborated and utilized by them and thus accumulates in the blood and in the other organs.

In 1889, von Recklinghausen, under the name of "general hemochromatosis," described a wide-spread accumulation in the organs of an iron-containing and an iron-free pigment, both of which he regarded as derived from hemoglobin, together with cirrhotic changes in the liver and pancreas. Most of the glands of the body were of a deep brown color and in their secreting cells he found the iron-containing pigment, which he called hemosiderin.

He named the iron-free pigment occurring in finer granules, hemofuscin, and describes it in smooth muscle cells of the stomach and intestine, of the blood-vessels and lymphatics, and rarely in those of the urinary bladder, ureters and vas deferens; also in certain connective tissue cells, those of Glisson's capsule, the splenic trabeculae, and the sheaths of blood-vessels. He described these peculiar findings as occurring in the course of different diseases, and gives diabetes as the cause in one case.

Buss, in an inaugural dissertation published in 1894, reported a case having clinically the characteristics of the *diabète bronzé* described by the French, and which in its pathology was the exact counterpart of the hemochromatosis of von Recklinghausen. He discussed the then recent work on pancreas extirpation and its relation to glycosuria and considered the hemochromatosis, caused by some primary blood disease of

unknown origin, to be the predisposing cause of diabetes, which was induced by the severe pancreatic changes.

Marie, in 1895, says we must look to the hemoglobin of the blood and tissues for such quantities of iron. In his opinion, the following is the sequence of events: dissolution of the hemoglobin of the blood under the influence of a general primary cause; transformation of the hemoglobin into pigment granules by the action of the cellular protoplasm of the organs; exaggerated elimination and accumulation of the pigment in the organs by irritation and consequent atrophy of the cells; supplementary elimination of the pigment by way of the lymph-channels with irritation of these channels and the production of a sclerosis of the interstitial connective tissue; and overloading of the lymph-nodes with pigment.

He deplored the prevailing opinion that bronzed diabetes was an accident or termination of ordinary diabetes. Some of his predecessors had spoken of it as pigmentary cirrhosis in diabetes, others as bronze cachexia in diabetes. To him, it was a true clinical and pathological entity, as distinct from the ordinary form as was, in his opinion, pancreatic diabetes, with which he would classify it rather than with any other form.

Acard,²⁴ Dutournier²⁴ and Jeanselme in their publications on the subject during the next two years, agree in all essentials with Marie.

Rendu and de Massary²⁵ in 1897, laying emphasis on cases of malaria reported with extensive pigmentation, conclude that the iron pigment, due to the particular action by the cellular protoplasm of a glandular organ or of a cardiac cell on hemoglobin, is the result of a general cachexia of the organism, and that this cachexia may be caused by diabetes complicated with a cirrhosis of the liver, by an hepatic disease alone, or by the malarial intoxication.

In 1899, Anschütz published his very comprehensive paper in which he reports a case of his own with thorough pathological investigations, and collects and tabulates in an interesting way the twenty-four cases of bronzed diabetes published up to that time. In his opinion, the blood destruction of unknown origin is primary. At the same time, there is a diseased condition of the cells of the glandular organs, which, under some mysterious influence, take up and hold tremendous amounts of the products of blood destruction and undergo a gradual degeneration. In those organs in which the degeneration is most pronounced, there occurs a reactive inflammation resulting in connective tissue growth. The diabetes is a result of the extensive lesions thus induced in the pancreas.

He remarks that "it is certain that a definite and significant blood dissolution occurs," that without very active new blood formation such large amounts of pigment could not be deposited. He is naturally surprised and somewhat nonplussed that there is never any sign of anemia,

25. Rendu and de Massary: *Cirrhose et Diabète Bronzé*, Presse méd., 1897, p. 58.

that the brain and kidneys contain almost no pigment and the spleen relatively little.

In the same year, Opie, after an admirable presentation of the subject, closes his paper with the following conclusions:

"1. There exists a distinct morbid entity—hemochromatosis—characterized by the wide-spread deposition of an iron-containing pigment in certain cells and an associated formation of iron-free pigments in a variety of localities in which pigment is found in moderate amount under physiological conditions.

"2. With the pigment accumulation there is degeneration and death of the containing cells and consequent interstitial inflammation, notably of the liver and pancreas, which become the seat of inflammatory changes accompanied by hypertrophy of the organ.

"3. When chronic interstitial pancreatitis has reached a certain grade of intensity, diabetes ensues and is the terminal event in the disease."

He thinks it possible that some primitive alteration of the blood, associated with intravascular destruction of red corpuscles, precedes the pigment deposition.

Two years later, Abbott reported the first case of hemochromatosis occurring in a woman. Diabetes was not present. Pathologically, the descriptions are typical of the disease except that the pigmentation was perhaps a little less widely distributed than usual (none in suprarenals or kidneys, little in skin) and the liver was slightly smaller than normal.

Her view is that some unknown primitive agency is at work on the red blood cells and on the parenchymatous cells of certain organs leading to the pigmentation. The cirrhosis results as a chronic interstitial inflammation from the presence in the tissues of the pigment set free by destruction of the parenchymatous cells. She is inclined to emphasize gastro-intestinal disturbances in the etiology and is rather doubtful as to the propriety of accepting this disease as a pathological entity.

During this year, Murri reported, from a clinical standpoint, a unique case. The patient, a woman 59 years old, for fifteen months had experienced symptoms of diabetes and a darkening of the skin. The urine contained from 2 to 4.5 per cent. of sugar; the skin pigmentation seemed fairly typical of bronzed diabetes with the exception of pigment flecks on the buccal mucosa, but there was no sign or symptom of lesions in the liver, pancreas or spleen. At different times there were abdominal pain and pains in the groin radiating to the abdomen and leg. There was no fever. A piece of skin with subjacent lymph-node was removed and found to contain a considerable amount of pigment, some of which gave a doubtful iron reaction. After the administration of epinephrin tablets was begun, the urine became sugar-free. The patient's condition remained the same for a while, then rapidly improved and, within less than a year after entering the hospital, she was discharged with most of

her weight regained, her urine physiological, and the abnormal skin pigmentation entirely gone. In 1901, she had remained well for two years.

As Murri says, this is undoubtedly a case of bronzed diabetes if we accept the name literally, but very unlike the disease as usually described. It is the only case on record of recovery, but its relation to hemochromatosis is not sufficiently established to give it a place among the well-authenticated cases of this disease.

Murri holds that the frequent occurrence together of the three phenomena, diabetes, hemochromatosis and cirrhosis of the liver, is not to be explained on the assumption that the process from which they arise is an individual and peculiar one; but that the disease processes, which are peculiar to each of the phenomena, possess certain characteristics which are partly biological (age), partly hygienic, partly toxic, and partly of diseased nature, but all of which reach a climax in an injury to the household of general metabolism.

The rational conclusion is on the hypothesis, he says, of a general dystrophy of the cell elements of the body, brought about by long continued action of the causes which are at work injuring their metabolism. This dystrophy goes along with the diminution of a normal capacity, namely, the oxidation of carbohydrates, and with the occurrence of two abnormal properties. The one consists in the capacity of the tissues to absorb and fix blood coloring material, the other in the increase of connective tissue, which, especially in the liver, assumes the character of an interstitial inflammation.

In common with Murri, most of the other German writers in the past decade, Hess and Zurhelle, Aschoff and Hess,²⁶ and recently Simmonds and Martineck,²⁷ hold that the three striking phenomena of the disease, diabetes, hemochromatosis and cirrhosis of the liver, are independent of each other and are each referable to the same common cause. They emphasize alcohol as a most important etiological factor.

Ridder considers a disease of the liver cells primary and follows Anschütz in his opinion as to the causes of the cirrhosis and diabetes.

Beattie, writing in 1903, is impressed by the possibility of a toxic agent derived from the intestine by way of the portal circulation, causing a degeneration of the cells of the liver, pancreas and other glands and at the same time a cirrhosis of these organs. The pigment, he thinks, is due partly to the inability of the degenerated cells to perform properly their metabolic processes and partly to transportation from the liver and pancreas. While the cirrhosis is, in his opinion, due, chiefly, to the toxic degeneration of parenchyma cells, it may, in part, be due to irritation produced by the pigment in the tissues.

26. Aschoff and Hess: Clinical Presentation of a Case of Hemochromatosis, *Berl. klin. Wchnschr.*, 1894, xli, 1231.

27. Martineck: Zur Pathogenese und klinischen Diagnose der Diabète bronzé, *Charité Ann. Berl.*, 1909, xxxiii, 40.

Parker,²⁸ reporting a case in the same year, is the only author who in so many words questions the occurrence of an abnormal blood destruction and suggests that the chief factor may be a loss in the power of elimination of certain iron-containing products rather than any abnormal amount of hemolysis.

He, in common with Heller and Martineck, mentions a supposition concerning the pathogenesis of hemochromatosis which they attribute to Croftan. I have been unable to find it in the elaborate form given by them in any of the references to Croftan's²⁹ work which they cite.

Croftan finds a ferment in liver extracts which he considers identical with trypsin, which has the power of breaking up both sugar and hemoglobin in the presence of each other, but is without effect on either, should the other be absent. According to the authors just named, the supposition is that some toxic agent produces a degeneration and chronic inflammatory changes in the liver and pancreas. When the pancreatic changes have reached such a grade of intensity that there is an insufficiency of the ferment trypsin, hyperglycemia results. Under the influence of this increased sugar content of the blood and in the absence of trypsin, the hemoglobin is changed into hematochromatin. The normal liver has the power of removing this abnormal pigment, but the cirrhotic liver lacks this ability; hence occurs the accumulation of iron-containing pigment in the liver and other tissues.

Rössle,³⁰ in 1906, reported a remarkable case of a patient 29 years old, who died in coma a few hours after entering the hospital. Pathologically, there were found cirrhosis of the liver, wide-spread lesions of the capillaries with many small hemorrhages, and, most remarkably, phagocytosis of the red blood corpuscles by the parenchyma cells. These unique findings were observed in the liver and kidney especially, to less extent in the pancreas, and not at all in the other organs. The pneumococcus was isolated from the tissues at autopsy.

He reviews the literature of hemochromatosis and considers his case as the earliest form yet observed. The primitive cause, in his opinion, is an acute infection with wide-spread "capillaritis" resulting in hemorrhagic edema and phagocytosis of the blood corpuscles by the parenchyma cells. The subsequent change of the hemoglobin to hemosiderin produces the siderosis of those cells. There is a rapid regeneration of the blood to normal after the acute stage has subsided. The cirrhosis may be due

28. Parker: Case of Bronze Diabetes, *Brit. Med. Jour.*, 1903, ii, 1052.

29. Croftan: Some Experiments on the Formation of Bile Pigment and Bile Acids; A Contribution to Our Knowledge of Icterus, *Phil. Med. Jour.*, 1902, ix, 75, 143; An Experimental Investigation into the Causes and Treatment of Diabetes Mellitus, *Am. Jour. Med. Sc.*, 1902, cxxiii, 662.

30. Rössle: Ueber Phagocytose von Blutkörperchen durch Parenchymzellen und ihre Beziehung zum haemorrhagischen Oeden und zur Haemochromatose, *Beitr. z. path. Anat. u. z. allg. Path.*, 1907, xli, 181.

partly to the "capillaritis" and partly to the destruction of the parenchyma. Diabetes is a terminal symptom.

This view is of interest because it departs so strikingly from those usually given and is founded on an observation which is interesting of itself. But he fails entirely, it seems to me, to substantiate his claim that there is any connection whatever between his unique case and those of hemochromatosis.

Several attempts have been made to reproduce the disease in animals experimentally by injecting blood into the peritoneal cavity and by producing intravascular hemolysis by the use of poisons such as toluylen-diamin; but such procedures have failed to produce anything but the faintest analogies to hemochromatosis.

Auscher and Lapique³¹ and, later, Meunier,³² produced, in this way, heavy pigmentation of the spleen, bone-marrow and abdominal lymph-nodes with less pigment in the liver. By increasing the amount of blood injected the disparity between the spleen and liver pigment deposits was diminished and they considered that they had reproduced the disease. There was never any connective tissue growth nor any reaction whatever, even when the spleen contained a much greater amount of iron than is found in any case of hemochromatosis.

The most thorough work of this kind, of which I know, is that of Biondi,³³ who poisoned dogs, cats and guinea-pigs with toluylendiamin administered by mouth and subcutaneously. In this way he produced anemia, jaundice and, sometimes, hemoglobinuria. Pigment deposits were found in the liver, spleen, bone-marrow and lymph-nodes, and occasionally in the kidney and other organs. He thinks that in any case of hemoglobinemia the liver cells increase their secretion of bile (*Pleiochromie*) and perhaps to such an extent as to cause jaundice without any obstruction of the bile-passages. The other part of the hemoglobin molecule, the iron-containing portion, is taken up by the leukocytes, giving rise to what he calls *siderifere Zellen*, and in this way to siderosis. These cells are carried through the blood to the spleen, bone-marrow, lymph-nodes and other organs. He says, therefore, that siderosis is the anatomical exponent of bile secretion. (He found a certain amount of iron pigment normally present.) Only when the liver cells were degenerated and manifestly abnormal did he find hemosiderin within their protoplasm. He thinks the diseased cell has lost its normal capacity of ridding itself of the iron-containing portion of the hemoglobin molecule. The

31. Auscher and Lapique: Quelques recherches chimiques sur un cas de diabète pigmentaire, *Compt. rend. Soc. de biol.*, 1895, 402, 510; 1898, 185.

32. Meunier: Contribution expér. à l'étude pathogénique de la cirrhose pigmentaire, Thèse, Paris, 1898.

33. Biondi: Experimentelle Untersuchungen über die Ablagerung von eisenhaltigem Pigment in den Organen infolge von Häematolyse, *Beitr. z. path. Anat. u. z. allg. Path.*, 1895, xviii, 176.

siderosis of the liver cells, then, he says, is an expression of their alteration. Anschütz thought this a particularly happy statement of the case and apparently considered this portion of Biondi's work applicable to the parenchymatous siderosis in hemochromatosis. The picture which he draws of the general siderosis in these experiments has little or no resemblance to that of hemochromatosis.

Most authors who express an opinion on this point think that, in addition to some injury to the parenchyma cells, an increased blood destruction is an essential factor in the production of such heavy pigment deposits. Others seem tacitly to accept this prevailing opinion.

Meltzer³⁴ has suggested that by some fault in the mechanism of iron metabolism by which this metal is excreted, the organs and cells of the body retain all or most of the iron which reaches them on its normal round through the body. Parker, also, asks the question whether it may not be possible that the essential factor is a failure in the elimination of iron rather than an increased hemolysis.

Since the work of M. B. Schmitt,³⁵ it has been generally accepted that all iron-reacting pigments have their origin from hemoglobin, and that, although a pigment fails to respond to the available microchemical tests, it cannot be said that it does not contain iron. According to Schmitt, the iron-reacting pigment represents only a stage in the continual development which these pigments are undergoing. Hemofuscin, then, is really the older state of the same pigment, which at an earlier period would have given a positive reaction for iron. By a further elaboration of this conception, Schmitt comes to the conclusion that melanin is a still later stage beyond the hemosiderin and hemofuscin period.

Many workers in metabolism experiments have proved that iron salts, whether organic or inorganic, administered by mouth, subcutaneously or intravenously, cause a deposit of an iron-reacting pigment in various parts of the body without any evidence of blood destruction. Wells,³⁶ in speaking of melanin, after mentioning the fact that the amount of iron present is usually extremely small, and often is entirely absent, says: "Furthermore, the presence of iron is no proof that the pigment is derived from hemoglobin, since many other proteids contain iron."

That iron forms an integral part of various tissues of the body and exists in firmly bound organic form is certain. Especially in nuclein substances iron has been found in relatively large amounts. According to Hammarsten,³⁷ a large amount of proteid substance of the liver exists

34. Meltzer: *The Pathogenesis of General Hemochromatosis (Diabète Bronzé)*, Med. Rec., New York, 1900, lvii, 43.

35. Schmitt: *Ueber die Verwandtschaft der hämatogenen und autochthonen Pigmente und deren Stellung zum sog. Hämosiderin*, Virchows Arch. f. Path. Anat., 1889, cxv, 397.

36. Wells: *Chemical Pathology*, W. B. Saunders Co., 1907, p. 393.

37. Quoted from Meinertz: *Ueber den Eisenstoffwechsel*, Zentralbl. f. d. ges. Physiol. u. Path. d. Stoffwechs., 1907 (new series), ii, 652, 689.

as an iron-containing nucleoprotein. Zaleski³⁷ isolated such a combination and called it "hepatin." Schmiedeberg³⁷ described an iron-containing nucleoprotein, rich in nucleic acid, "ferratin," which on hydrolysis yields a nuclein base, a pentose and levorotatory xylose.

What function tissue-iron serves is not well known. Schmiedeberg considers ferratin as reserve iron for blood formation. He gets a diminution in the ferratin content of the liver by the administration of iron-poor food and by repeated blood letting. It is known that in many chemical reactions iron is very potent as a catalyzer and in recent times catalytic reactions in the organism are being looked on with increasing favor. Spitzer³⁷ ascribes to iron a definite rôle as an oxygen-carrier. The oxidizing power of water extracts of the liver is due, according to him, to an iron-containing nucleoprotein, which is contained also in other organs. Iron is found in both split products, the nucleic acid and the protein; both are oxygen carriers. In further splitting, the oxidizing power persists as long as the split products contain iron.

Dastre and Floresco³⁸ accept the oxygen-carrying capacity as an important function of liver iron. They show that in certain invertebrates, the cephalopodia for example, the liver, so important in metabolic combustion, is rich in iron. The blood, on the contrary, does not contain iron. In these animals, certainly, the tissue iron is entirely independent of blood destruction, whatever may be said of the higher species.

A. B. Macallum,³⁹ basing his results on microchemical tests devised by himself, states that iron is found in all nuclei in amounts proportional to the chromatin. Assimilated iron is present in the cytoplasm of cells in higher animal forms in yoke-holding ova, the cells of yoke-holding embryos, and the hematoblasts of vertebrates. In addition, iron occurs in the cytoplasm of *ferment-forming gland cells of all descriptions*.

It is precisely in these ferment-forming gland cells that hemosiderin is deposited most characteristically in hemochromatosis.

In the majority of the cases in the literature, no blood examinations are reported. There are, however, some available figures. An average of four red cell counts made in the three cases of this series shows 4,709,000 cells per cubic millimeter. The hemoglobin estimations by different methods never fell below 90 per cent. Jeanselme gives as the first count in his case 3,379,000 red cells with 90 per cent. hemoglobin. Eight days later, one week before death, the count was 3,308,000. He noted no abnormal nucleated cells. The coagulability of the blood and retraction of the clot were normal. Spectroscopic examination of the blood serum showed no hemoglobin.

38. Dastre and Floresco: *Function martiale du foie chez tous les animaux en général*, Arch. de physiol. norm. et path., 1898, x, 176.

39. Macallum: See note 7; also *Die Methoden und Ergebnisse der Microchemie in der biologischen Forschung*, Ergebn. d. Physiol, 1898, vii, 552.

In Fletcher's two cases, the counts were 4,800,000 and 5,304,000 with the hemoglobin 87 per cent. and 95 per cent. respectively. Spectroscopic examination of the blood in the second case, made by Dr. Walter Jones, showed only the normal absorption bands of oxyhemoglobin. In this case the urine was examined for hematoporphyrin as a test of blood destruction, but this substance could not be demonstrated.

In Osler's case, the red cell count was above normal and no nucleated reds were seen in a differential count of 200 white cells. Ridder gives two red counts in his case, 4,800,000 and 4,400,000, with normal proportions in two differential counts of the leukocytes.

The bone-marrow of the femur was examined in two of my cases. Case 2 shows practically a normal picture. The marrow in Case 3 is slightly hyperplastic, but there is nothing to indicate any unusually active hematopoiesis.

Insignificant purpura has been described in a few cases, epistaxis in others and hemoglobinuria in one. These findings have been emphasized by different writers. The majority of the reported cases, however, show no such lesions.

Having failed to find in the clinical examination of the blood and in the histological appearance of the hematopoietic organs any reason to believe that there is a primary pathological blood destruction or blood alteration of any kind, we will direct our attention to the iron compounds of the tissues.

That increased blood destruction is not necessary to account for the large amounts of iron found in the tissues in this disease may be shown by a simple calculation, provided we assume a marked retention of the iron which is normally excreted. Iron excretion occurs, principally, by way of the alimentary tract and, to less extent, in the urine in cases of blood or tissue destruction. The question of its retention should be thoroughly investigated by chemical studies during the clinical observation of the patient.

Hess and Zurhelle estimated the total iron in the liver of their case, which contained a very high percentage, at 38.7 gm. Twelve additional grams for the pancreas, heart, lymphatic tissue, etc., is certainly a liberal allowance. This gives us 50 gm. of iron accumulated in the body. The whole amount of blood in the body of a man is said to contain about 2.4 gm. of iron. It is estimated that one-tenth of the red blood corpuscles undergo dissolution physiologically each day, thus liberating 0.24 gm. of iron. At this rate, if little is lost by excretion, 50 gm. of iron might be accumulated in less than seven months, which would be a very short course for the disease judged even by the period of active symptoms.

We may suppose that under some unknown influence there is a change in the equilibrium of the different cell constituents and under the changed conditions certain chromogenic groups of the proteid molecule can no

longer be held in the usual chemical and physical form and are deposited in the shape of pigment granules. The important cell constituent thus lost is being constantly replenished in the case of the glandular organs from the iron-containing portion of the hemoglobin molecule derived from the physiological dissolution of red cells in the circulating blood. Different cells are affected in varying degree. Newly regenerated cells may resist for a while, but on becoming physiologically active, they, too, suffer from the same pathological process, but of course, show much less pigment than their fellows (Fig. 6).

There is, then, a cell degeneration leading in the most active cells to dissolution, due both to the pigmentary degeneration of the protoplasm and to the accumulated pigment acting as a foreign body within the cell membrane. Differences in the chemistry of different cells could account for the chemical differences in the pigment found.

I would regard the sclerosis in different organs as a result of chronic interstitial inflammation produced, as Opie says, by two factors: pigmentary degeneration of the parenchyma cells and irritation produced by the presence of the pigment in the interstitial tissue. The latter factor is well illustrated in the lymph-nodes where the amount of fibrous tissue is roughly proportional to the pigmentation, and where in some cases (Case 1) foreign body giant cells are easily found.

The pathology of diabetes is a much mooted question. Naunyn and a few others have ascribed the diabetes in this affection as well as in common diabetes to cirrhosis of the liver. Palma, Von Noorden and others hold that diabetes and liver cirrhosis occur together so rarely that they cannot be considered more than the coincidence of two common diseases.

The relationship of chronic interstitial pancreatitis to diabetes has been fairly well established, and according to Opie, Anschütz, Marie and others, it is in the pancreas that one must look for the cause of the diabetes of hemochromatosis. In practically all the cases which are reported in sufficient detail, there have been more or less pronounced lesions in the pancreas. Yet what pathologist would be willing to say from an examination of that organ whether or not diabetes had been present? It is true that in Case 1 (simple hemochromatosis) the pancreas is more nearly normal than those of the other two cases, yet, it is in the islands of Langerhans in Case 1 that the pigment in the pancreas is most marked (Fig. 2). These structures seem relatively unaffected in the other cases which showed diabetes. Again, the pancreas from Case 3, a marked diabetic, seems less severely diseased than that of Opie's case, sections of which I have had the opportunity of examining. In his case, sugar was never present in the urine, nor were there ever any other symptoms of diabetes.

Emphasis has already been laid on the probable function of the iron-containing tissue compounds as catalytic agents, especially in oxidizing processes. There are, apparently, serious disturbances among these compounds in hemochromatosis. It is not surprising, therefore, that we find a diminution of the capacity of the organism to oxidize carbohydrates, without looking for further gross changes in any one organ. This view, it seems to me, would harmonize better with those cases in which the sugar disappears from the urine for varying periods. The process as outlined above, need not necessarily be continually active, nor present always in the same degree. It may well be that fluctuations occur and that the periods of lessened activity in the morbid process become manifest in a diminution or disappearance of the glycosuria and perhaps improvement in other ways. Similarly, in those cases in which the process is sluggish from the beginning, glycosuria may never appear and abnormal metabolism along other lines produce the weakness and cachexia of these cases of simple hemochromatosis.

Concerning the unknown influence which inaugurates the pigmentary changes and the mechanism of the pigment deposit, very little can be said at present.

Opinions concerning the origin and deposition of melanins, the most extensively studied pigments, have undergone radical changes in recent years. They are no longer regarded as being composed of the products of hemoglobin, but rather as having an origin in albumin, especially in those products with an aromatic or heterocyclic nucleus (tyrosin, etc.). It is thought that from these, under the action of various oxidizing ferments, the melanotic pigments are formed. Gortner⁴⁰ has recently demonstrated that the pigmentation of the integuments of the larva of *Tenebrio molitor* is the result of the interaction of an oxydase and a chromogen. The process of coloration is capable of preceding in the absence of life, but ceases entirely if the enzyme is killed by heat or if oxygen is withdrawn.

Brown⁴¹ has recently shown that hemosiderin may be produced outside of the animal body and is increased in the liver of rabbits during autolysis for a period of from twenty-four to forty-eight hours. Since this post-mortem hemosiderin formation is most marked in parts of livers exposed to the air, he concludes that the hemosiderin is an oxidation product of hemoglobin due to enzyme action.

In view of the fact that physiological pigment deposits may be due largely to enzyme action, we may consider this as a possibility, at least, in the abnormal pigmentation of hemochromatosis. Gortner says of the

40. Gortner: Origin of the Brown Pigment in the Integuments of the Larva of *Tenebrio Molitor*, Jour. Biol. Chem., 1910, vii, 365.

41. Brown: Changes in the Hemosiderin Content of the Rabbit's Liver During Autolysis, Jour. Exper. Med., 1910, xii, 623.

larva of *Tenebrio molitor* that the chromogen is present in very small quantity and is probably secreted only as needed for the pigmentation, while the oxydase is constantly present in relatively large amounts. Perhaps the relations are much the same in human tissues and perhaps excessive chromogen is formed as an intermediary product in the peculiar degeneration of the cell protoplasm; and, acted on by the ever-present oxydase, is precipitated as pigment granules. This may be one explanation of the changes seen in the rabbit's liver during autolysis and some experimental work now in progress indicates that such may be the case.

SUMMARY

Hemochromatosis is a distinct morbid entity characterized by the wide-spread deposition of an iron-containing pigment, especially in the secretory cells of glandular organs, by the presence of an iron-free abnormal pigment in certain situations, and by an increase in the pigmentation of those localities which contain pigment under physiological conditions.

There is no evidence of any abnormal blood destruction or of reparatory activity of the hematopoietic organs.

Hemochromatosis may be considered a metabolic disease implicating many of the body tissues and manifested, especially, by a change in the chromogenic groups of the proteid molecule with the deposition of pigments.

As a result of the pigmentary degeneration and the accumulation of pigment in the interstitial tissue, there occurs a reactive inflammation with a fibrosis of various organs.

The diabetes may be referable to the diminished oxidative powers of the body due to the pronounced disturbance in the iron-containing constituents of the tissues, assuming these iron compounds to be catalytic agents in oxidizing processes.

There is no constant predisposing cause.

In conclusion, I wish to express my appreciation to Professors Barker and Thayer for the privilege of using the clinical histories, and especially to Dr. G. H. Whipple for his unfailing interest during the course of this study and for valued suggestions in the preparation of this paper.

Johns Hopkins Hospital.