The Archives of Internal Medicine

Vol. XIII

JANUARY, 1914

No. 1

STUDIES ON THE CIRCULATION IN MAN

X. THE BLOOD-FLOW IN THE HANDS IN DISEASES OF THE HEART *

G. N. STEWART, M.D.
CLEVELAND, O.

GENERAL REMARKS

It has been mentioned in the introduction to the last paper¹ that when proper precautions are taken there is a very fair degree of constancy in the results obtained in successive observations on different days on one and the same patient, provided gross changes in his clinical condition are excluded. Accordingly, a marked change in the flow not accounted for by changes in the external conditions, especially in the external temperature, always suggests some decided alteration in the clinical state of the patient. On the other hand, in certain cases, information may be obtained as to the state of the patient; for example, in regard to the lability of his vasomotor mechanism or the power of adaptation of his heart, by purposely changing the external conditions or by taking advantage of changes that have spontaneously occurred, in order to determine their effect on the flow in the hands or feet.

In comparing the flows in the whole series of cases investigated with those in normal persons, one is struck by the fact that while the flow in persons suffering from well characterized pathological changes (compensated valvular lesions, for example) may be of the normal magnitude, having regard to the age of the patient, the room temperature, etc., in a much larger number of cases it is below the normal, often greatly below it. Sometimes there is good evidence that this diminution in the hand or foot flow is the expression of an increased excitation of the vasoconstrictor mechanism as a whole or at least of the portion of it which presides over the cutaneous vessels not necessarily related to any material diminution in the driving power or the output of the heart. In other cases there is good evidence that the diminished flow in the hand (or foot) is in the main the expression of diminished functional capacity on the part of the heart, associated either with engorgement of the veins

* From the H. K. Cushing Laboratory of Experimental Medicine, Western Reserve University.

1. Paper IX of this series.

110

and the right side of the heart or with diminution in the filling of the arteries without increased venous pressure. In still other cases evidence may be obtained of a diminished flow in one or both hands or feet, which is solely or mainly a local condition. A local increase in the flow may accompany such conditions as inflammation or vasomotor paralysis.

The question whether the flow in the hands or feet gives any indication of the functional capacity of the heart would be best discussed on the basis of an experimental comparison of the heart output, as determined gasometrically, with the hand flow in normal and pathological cases. Even in the absence of such a comparison, however, it would appear, that where the hand (or foot) flow is not less than normal it is only in exceptional conditions (local vasomotor paralysis of hands and feet?) that any gross error will be made in assuming that the heart is discharging an amount of blood not inferior to the normal. When, however, the flow in the extremities is permanently and decidedly less than normal, several points have to be considered. In many of these cases in which the normal ratio of the hand flow to the foot flow is preserved² in spite of the diminution in the flow and in which no anemia is present, there is often good evidence that the lessened driving power of the heart is primarily responsible for the diminished flow. On the other hand, in the anemias³ the diminution of the flow in the hand and in the skin generally seems to be a compensating mechanism which permits an even increased quantity of blood to be sent by the heart through the lungs.

A marked disturbance of the normal ratio between the hand and foot flow, of course, gives no information as to the functional capacity of the heart, but indicates in the absence of local anatomical changes which might account for the difference (arteritis, paresis, etc.), a change in vasomotor excitability. Usually, this is manifested in a vasoconstriction of the feet, and a foot flow smaller than the normal in comparison to the hand flow. Since the feet are normally but little exposed in comparison with the hands, local vasoconstriction connected with the technic of the measurements is more likely to occur in the feet than in the hands. In various pathological conditions, for instance, fever,⁴ where the vasoconstrictor mechanism of the skin is abnormally excitable the tendency of vasoconstriction in the feet on exposure is intensified. For this reason, as well as on account of the inferior flow per unit of volume in the feet, measurements of the hand flow are always to be preferred to measurements of the foot flow when the object is to obtain general information concerning the circulation and not to study local circulatory conditions. A disturbance of the normal ratio between the hand and

2. See Paper VII of this series, *Jour. Exper. Med.*, 1913, xviii, No. 1.

3. Paper VI of this series, *Jour. Exper. Med.*, 1913, xviii, No. 2.

4. See Paper VIII of this series, *Jour. of Exper. Med.*, 1913, xviii, No. 4.

foot flow, due not to vasomotor, but to anatomical causes (arteritis obliterans, for example), would be revealed by the fact that the ratio remains permanently abnormal even when the external conditions, the room temperature especially, are changed so as to abolish vasomotor differences of the lower and the upper extremities, or when this is done by the administration of vasodilator drugs.

A few words are necessary in regard to the classification of the cases dealt with in this paper. Some overlapping between the groups is unavoidable and the attempt has been made to assign each case presenting a complication of lesions to the group in which the factor seemingly predominant in the clinical picture would place it. For example, in the group of myocardial affections, cases are necessarily included in which there were definite and demonstrable valvular lesions. Thus certain cases with mitral stenosis (and insufficiency) are placed in the myocardial group because there is evidence that auricular fibrillation was also present, and that probably this condition with the consequent marked disturbance of the action of the ventricle was at the time of the investigation a more important factor in the clinical situation than the valvular lesion, or at least a factor which differentiated the case from one with the valvular lesion alone, and gave to it an added gravity. Among the few cases classed as arteriosclerosis, a group which might, of course, have been greatly swollen at the expense of the other groups had attention been paid solely to this condition, is one⁵ which might also have gone with the valvular lesions. It is included with the cases of arteriosclerosis simply because that condition was extremely marked in the accessible arteries.

I. CONDITIONS IN WHICH SIGNS OF INVOLVEMENT OF THE MYOCARDIUM PREDOMINATED

There is probably no pathological condition of the heart in which such small hand flows are encountered as in this group. While the myocardium is acting well, grave valvular defects may, as is well known, be overcome, and the output remain for long periods quite at the normal level. The results of the blood-flow measurements in many cases of valvular lesions with good compensation are perfectly normal. With a heart whose myocardium is seriously crippled, even when the valves are anatomically and probably functionally intact, the hand flow may sink to astonishingly small figures. Nor is this difficult to understand. A pump with somewhat leaky valves may still deliver a good stream when the strokes are strong and regular. No matter how tight the valves are the output will be small if the stroke is feeble, uncertain and irregular.

5. Fred. S., Paper XI of this series.

Thus in Stephen F., a man 31 years old, the flow in the hands was only 0.2 gram per 100 c.c. of hand per minute with a room temperature of 23.5 C. increased to 0.5 gram in the right hand during immersion of the left in warm water, or not more than one-twentieth of the normal flow. The protocol of this case has already been published.⁶

The man, a contractor in a small way for brick and carpenter work, was examined in the dispensary and was admitted the same day to Lakeside Hospital where he died a few days later. He stated that between three and four years previous to admission he became ill after a heavy lift and had never been well since. All the signs of cardiac failure were present. To the left the cardiac dulness extended almost to the anterior axillary line. There was marked arrhythmia with numerous premature contractions. For some time past he had been aware of the irregular action of his heart and stated that it sometimes stopped for a beat. The pulse-rate was only 60 (counted at the apex beat with the stethoscope). The neck veins were much engorged in the supine position. There was edema of the feet and ankles. The hands and feet were habitually cold although the weather was mild. The whole surface was pale and cool. There was no indication of valvular lesions. No murmurs were detected during the time the patient lived. The lungs were clear. The clinical diagnosis was myocarditis without valvular lesion. The immediate cause of death was pontine hemorrhage. At autopsy the findings were fibrous myocarditis with relative insufficiency but no valvular lesion, pericarditis, perihepatitis and atrophic cirrhosis of the liver, and acute pulmonary edema.

In Mrs. A., a woman aged 72, who came to the dispensary complaining of palpitation of the heart and irregular pulse, the blood-flow in the left hand came out 0.78 gram per 100 c.c. per minute, and in the right 0.45 gram, with room temperature 22.5 C. It must be taken into account that the hands in this case, being the typical hands of old age, consisted largely of bone, which has a small blood-flow.

The irregularity of the heart has been noticed for the past six years. The heart sounds were clear and no murmurs could be detected, but there was marked arrhythmia. The pulse rate was 92 in sitting position. There was no edema, no cough. There was a history of rheumatism at 50 years of age, and again the winter before admission. The hands were very thin and rather cool, although she wore gloves and said they were as warm as usual. A diagnosis of senile heart with chronic myocarditis, arrhythmia and arteriosclerosis was made. She continued to come periodically to the dispensary for over four months, still complaining of palpitation although the pulse became more regular.

The difference in the flow in the two hands (it must of course be noted that the accuracy of the measurements is less with these small flows than with normal flows) suggests that the smallness of the flow may be partly due to local vasoconstriction, which is often temporarily unequal for the two hands, and not altogether to the small output of the heart. The question may indeed be raised whether as in the anemias⁷ a peripheral constriction may not occur in these cases in the interest of an increased flow through the lungs (and through the brain and other

6. *Cleveland Med. Jour.*, 1911, x, 385.

7. *Jour. Exper. Med.*, 1913, xviii, No. 2.

vital organs). In some ways the question is merely an academic one, since it is well known that in marked anemia the heart muscle may be impaired. It is clear that the percentage content and the total absolute amount of hemoglobin in the blood being normal, there may still be oxygen deficiency if the weakened heart is unable to propel the normal amount of blood from the left ventricle through the systemic vessels to the right side of the heart. On the supposition that the right ventricle could deal with a larger quantity of blood than is actually brought to it, a diminution in the average circulation time of the blood through the systemic path brought about by vasoconstriction confined to or predominating in the periphery might increase the minute volume of the heart. In this case the small hand flow, where the myocardium is seriously damaged, would be due to two things: First, diminished driving power of the heart; and second, constriction of the vessels of the peripheral parts in excess of that which would correspond to the diminished filling of the vessels due to the deficient action of the heart alone. Although the output in such cases as Stephen F. is diminished, it is not conceivable that it is diminished in the same proportion as the hand flow.

Jacob W., a house carpenter aged 59, height 5 feet, 1 inch, weight 173 pounds, was admitted at the dispensary April 4, 1911, suffering from myocarditis with anginal attacks. Six months previously while going to church he was suddenly seized with substernal pain, with great dyspnea. He had to sit down and could not speak. Since that time he has had a spell every morning, lasting two to five minutes and it is getting worse. If the pain is severe the left hand and forearm get dead. Since yesterday he has had palpitation. His general health was good until six months ago. He had rheumatism eighteen years ago. He used to drink beer and whisky moderately and to smoke moderately. He denies lues. Physical examination, cyanosis, some icterus; emphysema; heart generally enlarged; no murmurs. The heart sounds are weak; blood-pressure high; some sclerosis of the arteries. Urine negative.

He came periodically to the dispensary after this. April 24, 1911, his systolic blood-pressure was 190, diastolic 110. April 27, 1911, he had five attacks during the night. May 1, 1911, he had a mild attack while being examined in the dispensary. The pain was felt right beneath the sternum and radiated down the inner side of the left arm and up the neck to the chin. The pulse becomes somewhat smaller during the attack but the rate does not change. The heart is enlarged 4 cm. to the left but not to the right. He is still stout. He is being treated with potassium iodid and sodium nitrite. The blood flow in the hands was examined May 1, 1911. Mouth temperature 37.1 C. Pulse (sitting) 100, and rather small. Hands in bath at 2:44 p. m., in calorimeters at 2:55.* At 3:10 p. m. left hand put into water at 42.9 C. (Table 1.)

Cooling of calorimeters, R., 0.11 degrees in fourteen and one-half minutes; Left 0.14 degrees in twenty-one and one-half minutes. Volume of right hand, 508 c.c., of left hand 486 c.c. The hands were fat.

The flow in the right hand works out at 2.46 grams per 100 c.c. per minute, and in the left hand at 1.96 grams. When the left hand was

* Three thousand and fifteen c.c. of water were put into each calorimeter. In the protocols, unless otherwise mentioned, this was the quantity always used for the hand calorimeters.

immersed in warm water the flow in the right rose to 3.86 grams, a fair reflex vasodilatation. Subsequent immersion of the left hand in cold water caused little change in the flow in the right.

TABLE 1.—CALORIMETRIC MEASUREMENTS IN CASE OF JACOB W.

Time	Right	Left	Room Temp. C.	Time	Right	Notes
2:54	29.83	29.91	24.1	3:14	30.12	Room 25.0 C.
2:56	29.81	29.86		3:15	30.14	
2:57	29.82	29.86		3:16	30.18	
2:58	29.84	29.87		3:17	30.20	
2:59	29.87	29.88	24.9	3:18	30.23	
3:00	29.87	29.89		3:19	30.26	At 3:20 left hand put in water at 11.5 C. Room 25.1 C.
3:01	29.88	29.91		3:20	30.29	
3:02	29.89	29.92		3:21	30.32	
3:03	29.89	29.92		3:22	30.34	
3:04	29.91	29.93	25.0	3:23	30.36	
3:05	29.92	29.94		3:24	30.39	Left hand dried and wrapped. Room 25.1 C.
3:06	29.93	29.94		3:25	30.41	
3:07	29.95	29.95		3:26	30.44	
3:08	29.98	29.97		3:27	30.47	
3:09	29.99	29.98		3:28	30.49	
3:10	30.02	30.00		3:29	30.51	Hand out of calorimeter at 3:30. Temp. of L. is now 29.86 C. Temp. of L. is now 29.76 C.
3:11	30.03			3:30	30.54	
3:12	30.05			3:31½		
3:13	30.08		25.2	3:44½	30.43	

In James M., a man 60 years old, with a very irregular pulse, enlarged heart and a history of edema of the feet and legs, the flow was 1.99 grams per 100 c.c. per minute in the right hand, and 1.35 grams in the left (for the last ten minutes in the calorimeters). Anemia was probably not an important factor in the small blood-flow, since the hemoglobin content of the blood was 80 per cent. three days before the examination. The calculated flows for the last twenty minutes in the calorimeters are 1.56 grams per 100 c.c. per minute and 1.06 grams for the right and left hands respectively, the ratio between the flows being precisely the same as for the last ten minutes (1.47:1).

James M., a laborer aged 60, was admitted to the City Hospital June 16, 1912. Three weeks before when in another hospital his feet and legs swelled and pitted on pressure. On admission to the City Hospital petechiae were present on both legs but there was no edema. The right side of the chest lags in inspiration and shows diminished resonance beginning at the sixth interspace in the axillary line, and also posteriorly. Fifteen months ago he had dyspneic attacks and the right pleural cavity was tapped. June 21, 1912, the right chest was aspirated but no fluid was obtained. The left border of heart dullness is outside the nipple line, the right border at the right margin of the sternum. The first sound is roughened. Pulse irregular, arrhythmic, fair volume, sclerosis marked. The rectal temperature since his admission has varied from 100.4 to 98.2 F.

Examination of the blood-flow in the hands was made June 27, 1912, on which day he was discharged. Hands in bath at 1:34½ p. m.; in calorimeters at 1:43¾ p. m., and out of calorimeters at 2:15.

TABLE 2.—CALORIMETRIC MEASUREMENTS IN CASE OF JAMES M.

Time	Right	Left	Room Temp. C.	Time	Right	Left	Room Temp. C.
1.43	31.02	31.12		2.01	30.93	30.97	
1.45	30.99	31.09		2.02	30.93	30.965	
1.46	30.98	31.08	22.9	2.03	30.93	30.955	23.0
1.47	30.98	31.07		2.04	30.925	30.95	
1.48	30.975	31.06	22.9	2.05	30.925	30.945	
1.49	30.97	31.05		2.06	30.93	30.94	23.1
1.50	30.965	31.04	23.0	2.07	30.925	30.935	
1.51	30.96	31.03		2.08	30.925	30.93	
1.52	30.96	31.03		2.09	30.93	30.93	23.1
1.53	30.955	31.025	23.1	2.10	30.925	30.93	
1.54	30.95	31.02		2.11	30.93	30.925	
1.55	30.95	31.015		2.12	30.925	30.925	
1.56	30.94	31.005	23.1	2.13	30.93	30.925	23.0
1.57	30.94	31.00		2.14	30.93	30.92	
1.58	30.94	30.995		2.15	30.925	30.915	
1.59	30.935	30.99		2.27	30.795	30.78	
2.00	30.935	30.98	23.2				

Cooling of calorimeters in twelve minutes, R., 0.13 degrees; L., 0.135 degrees. Volume of right hand 368 c.c., of left hand 356 c.c. He is right handed,* wrists very thin, hands skinny. Pulse 87 (sitting). Very irregular. Rectal temperature 37.65 C. Water equivalent of calorimeters with contents, R., 3,389; L., 3,380. Blood-pressure, systolic 149 (palpation), 152 (stethoscope), 72 (sound gone). Another observation, systolic 144 (palpation).

A case diagnosed as auricular fibrillation (in N. P) is included under this heading, although there was a valvular lesion (mitral stenosis) present as well. The case did not come under my observation until the patient had improved considerably under treatment (digitalis), yet even then the flow in the hands was quite small for her age (1.97 grams per 100 c.c. per minute for the right hand, and 1.59 grams for the left for the last eleven minutes in the calorimeters). The small flow agreed with the persistent coldness of the hands and feet. It was not associated with anemia.

N. P., aged 27, was admitted for her present illness to the City Hospital March 12, 1913. (She was first admitted to the hospital April 4, 1910, suffering from rheumatism.) Heart: Apex beat visible in fifth interspace, one finger's breadth outside the nipple line. A presystolic thrill can be felt here. Left border two finger's breadth outside the nipple line; right border at the midsternal line, upper border at third rib. A rough presystolic murmur is heard over a small area at the apex. All over the precordia and transmitted to the axilla and back

* It can always be assumed that a patient was righthanded unless the contrary is stated in the protocols.

is heard a soft systolic murmur. The pulse shows marked arrhythmia. Considerable edema. The urine, March 13, 1913, contained neither albumin nor casts. On May 12, 1913, there was some albumin but no casts. Under treatment (including digitalis) the pulse rate diminished. The arrhythmia became less marked although it did not disappear, and the edema cleared up. Her appetite continues poor, but she sleeps well and is not troubled with dyspnea in any position. Her hands and feet are always cold. She says she is only really warm when surrounded by hot water bottles. Polygraph tracings from the jugular and radial taken on May 16, 1913, showed the ventricular form of venous pulse, "a" waves being absent from the jugular tracing. The rate of the heart in the semi-reclining position was 80 as counted at the apex with the stethoscope, and precisely the same at the wrist. The variation in the duration of successive beats is great. Twelve successive beats were measured in each of two portions of the tracing. In the first series the durations were 1.1, 0.8, 0.9, 0.8, 0.7, 0.65, 0.65, 0.5, 0.6, 0.9, 0.75, 0.9 second, the average being 0.77 second corresponding to 78 beats per minute. In the second series the durations were 0.85, 0.65, 0.75, 0.7, 0.7, 0.55, 0.75, 0.85, 0.9, 0.85, 0.95, 0.85 second, the average being 0.78 second corresponding to 77 beats per minute.

The blood-flow was examined May 12, 1913. Hands in bath at 2:43 p. m.; in calorimeters at 2:54½, and out of calorimeters at 3:16. Mouth temperature 36.3 C. (the thermometer was in five minutes and special care was taken).

TABLE 3.—CALORIMETRIC MEASUREMENTS IN CASE OF N. P.

Time	Right	Left	Room Temp. C.	Time	Right	Left	Room Temp. C.
2.54	31.56	31.41		3.06	31.455	31.28	23.9
2.55	31.53	31.37		3.07	31.45	31.27	
2.56	31.51	31.36	24.2	3.08	31.45	31.27	
2.57	31.50	31.355		3.09	31.44	31.26	
2.58	31.50	31.35	24.1	3.10	31.43	31.255	23.8
2.59	31.49	31.34		3.11	31.43	31.25	
3.00	31.49	31.33		3.12	31.42	31.24	
3.01	31.48	31.31		3.13	31.42	31.235	
3.02	31.46	31.30		3.14	31.41	31.23	
3.03	31.46	31.29	23.6	3.15	31.41	31.225	
3.04	31.46	31.285		3.16	31.405	31.215	
3.05	31.455	31.28	23.9	3.35	31.19	30.995	

Cooling of calorimeters in 19 minutes, R., 0.215 degrees; L., 0.22 degrees. Volume of right hand in calorimeter 298 c.c., of left hand 276 c.c. Pulse (sitting) 75 in one count, 66 in another, irregular and weak. It is not easy to count the pulse at the wrist because of the variability in the beat. Systolic blood-pressure in right arm 95. Difficult to determine the blood-pressure with the stethoscope.

A man 44 years old (Jacob S.), with mitral stenosis and insufficiency, an irregular heart, and weak and irregular pulse, from a study of whose case a diagnosis of auricular fibrillation was made, had a flow of only 1.97 grams per 100 c.c. per minute in the right hand, and 0.96 gram in the left, with room temperature 22.2 C. The disproportion in the flow in the two hands is associated with old hemiplegia of the left side. Immersion of the right hand in warm water caused a great percentile increase in the flow in the left hand (to 1.54 grams per 100 c.c. per minute for

the first five minutes and 3.86 grams for the next four minutes), although the maximum was still absolutely a small flow. The fair crossed vasomotor reflex from the normal to the formerly paralyzed hand, which had largely recovered, contrasts with the absence of reflex in a case of old hemiplegia where there had been no recovery of the hand.⁸

Jacob S., a tailor aged 44, height 5 feet, 3½ inches, weight 140 pounds, first presented himself at the dispensary Oct. 27, 1910. Three years before he had rheumatism and had "heart trouble" at that time and also pneumonia. Eleven years ago when sitting in a chair he had a left-sided paralysis which came on him suddenly. He lost consciousness. He also had aphasia. He could understand what was said but could not speak. He recovered his speech in two months. The arm and leg recovered slowly. There was no disturbance of sensation at the time of the stroke, but two months later he had loss of sensation so that a pin could be stuck into his arm or leg without his knowing it. He burnt his left elbow severely without feeling it by leaning on a stove. The lines on the left side of the face are not so deep as on the right. He drags the left foot somewhat. The tongue is protruded in the median line. The reflexes are much increased in both upper and lower extremity. Babinski's sign can be obtained on the left side. Touch and pain are normally appreciated on the left side but heat cannot be distinguished over the left arm. The physical signs of chronic bronchitis and emphysema are present. Heart slightly enlarged both to right and left. Pulmonic second sound accentuated. The heart sounds are very irregular both in force and rhythm. Pulse rate 80.

Nov. 5, 1910, a marked presystolic murmur and thrill can be made out.

Dec. 10, 1910, he is very dyspneic and cannot sleep. Ears, fingers, tip of nose and lips are cyanotic. Positive venous pulse in the neck. The heart dullness extends two fingers outside the nipple line on the left side and two fingers outside the sternal margin on the right. There is a sharp first sound preceded by a faint presystolic murmur and followed by a systolic murmur. The pulse is small, irregular and of very poor volume and tension. The left radial pulse is smaller than the right. The liver is increased in size to percussion. He was received into the hospital to-day where he improved. Blood-pressure, systolic 162, diastolic 120. Blood: leukocytes, 10,500; hemoglobin, 90 per cent.

April 11, 1911, he returned to the dispensary in about the same condition as before he entered the hospital. Pulse rapid and irregular.

April 25, 1911, enlargement of the heart and arrhythmia is still marked. Knee-jerk much exaggerated on the left side. When he gets out of breath he feels the left hand getting stiff. The difference in size of the two hands is not conspicuous. The great difference in the two arms is in the girth of the shoulders, the left shoulder being much smaller than the right. He cannot completely extend the left forearm on the elbow owing to contracture. Sometimes when he is grasping an object in his left hand the fingers remain closed around it involuntarily and he has to release the object with the other hand. He lost sexual power at the time of the stroke and has not regained it. He has been constipated since the stroke and has had to use cathartics constantly.

The blood-flow in the hands was measured April 25, 1911. Pulse (sitting) 104. Mouth temperature 36.25 C. Hands in bath at 2:31 p. m., in calorimeters at 2:42.

At 2:55 p. m. the right hand was put into water at 43.3 C.

Calorimeter R. was constantly stirred until 3:10 p. m. After this R. and L. were only stirred occasionally. Cooling of R. from 2:55 to 3:10½ is 0.18 degree (in fifteen one-half minutes.) Cooling of R. from 3:10½ to 3:27 (with little stirring) is 0.15 degree (in sixteen one-half minutes.) Cooling of L. from 3:10½ to 3:28 (with little stirring) is 0.16 degree (in seventeen one-half min-

8. Heart, 1911, iii, 81.

utes.) Accordingly with little stirring L. cooled at the same rate as R. We can therefore assume that with constant stirring this would also be true and that the amount of cooling for L. during the observations is 0.18 degree in fifteen and one-half minutes. Volume of right hand 390 c.c., of left hand 378 c.c.

TABLE 4.—CALORIMETRIC MEASUREMENTS IN CASE OF JACOB S.

Time	Right	Left	Temp. Room C.	Time	Left	Notes
2:41	29.73	29.72		2:58	29.60	Room 22.7 C.
2:43	29.68	29.65		2:59	29.60	
2:44	29.68	29.65	22.0	3:00	29.60	
2:45	29.68	29.64		3:01	29.59	
2:46	29.68	29.64		3:02	29.61	
2:47	29.67	29.63		3:03	29.62	At 3:05 right hand dried and wrapped.
2:48	29.67	29.63		3:04	29.63	
2:49	29.67	29.62		3:05	29.64	
2:50	29.68	29.63		3:06	29.64	At 3:10 hand out of calorimeter. Temp. of R. is now 29.52 C. Temp. of R. is now 29.37 C.
2:51	29.68	29.62		3:07	29.65	
2:52	29.68	29.61	22.4	3:08	29.65	
2:53	29.69	29.61		3:09	29.67	
2:54	29.69	29.60		3:10	29.68	
2:55	29.70					
2:56		29.60	22.6	3:10½		
2:57		29.59		3:27		
				3:28	29.52	

In a third case diagnosed as one of auricular fibrillation (T. F. K.), a man 57 years old, the flow was 4.45 grams per 100 c.c. per minute for the right hand, over an eighteen-minute period, and 5.23 grams for the left hand (ratio of the flows in the two hands 1:1.17). For the last nine minutes of this period, the flows were 6.21 grams and 7.06 grams, respectively, for the right and left hands (ratio of the flows in the two hands 1:1.13). These are fair flows for the man's age, indicating that the heart in spite of the irregular action of the ventricle is maintaining a fair output. This corresponds entirely with the clinical condition at the time of the blood-flow examination. The edema had completely disappeared. There was no dyspnea. The appetite was good and the patient was no longer confined to his bed. Many of the pulse beats at the wrist were of good amplitude, although others were small (Fig. 1), while in the case of N. P. all of the beats were small.

T. F. K., a bricklayer, aged 57, height 5 feet, 9½ inches, weight 140 pounds, was admitted at the City Hospital May 1, 1913. He had typhoid about thirty years ago and rheumatism sixteen or eighteen years ago. His present illness began last November. His ankles became swollen and also his legs, nearly up to the hips, and he lost strength and weight rapidly. He noticed at the same time that his pulse was irregular. He had pain in the right side of the chest. The swelling in the legs had diminished somewhat before he entered the hospital, but on admission edema and cyanosis were still to be observed. His chest was barrel-shaped with broad costal angle and poor expansion. No rales or dulness over the lungs, but there was a friction rub over the anterior surface of the right

lung. The cardiac impulse was diffuse, and extended considerably beyond the nipple line on the left side. Cardiac dullness enlarged to the left. The liver was somewhat enlarged and pulsation evident in the epigastrium. The veins in the neck were distended. A systolic murmur was heard over the apex. His maximum temperature (on May 9) was 103.6 F., minimum 97.4 on the same day. The maximum temperature on May 16 was 102; May 17, 102; May 19, 101.4 F. The pulse-rate on May 1 was 120; May 19, 92. The pulse-rates were usually above 90. The radial pulse is markedly irregular in amplitude and rhythm, and the irregularity is not affected when respiration is suspended. Polygraph tracings from the distended external jugular vein and the radial artery showed that the venous pulse was of the ventricular character (Fig. 1). Diagnosis, auricular fibrillation. The blood-flow in the hands was examined May 19, 1913, at which time the swelling had entirely disappeared from the lower extremities and the patient was allowed to sit up for a portion of the day. Hands in bath at 3:14 p. m.; in calorimeters at 3:26 $\frac{3}{4}$, and out of calorimeters at 3:47.

Fig. 1.—Upper part of tracing, simultaneous record of apex beat and radial pulse. The respiratory movements interfered with the apex trace except towards the end of expiration. At the right is a portion of the apex tracing which allows a sufficiently good comparison to show that some of the apex beats are only feebly represented on the radial trace. The lower portion of the figure gives simultaneous tracings from the external jugular and the radial. As in all tracings, time is marked in fifths of a second. (All tracings reduced to about one-half size of originals.)

TABLE 5.—CALORIMETRIC MEASUREMENTS IN CASE OF T. F. K.

Time	Right	Left	Room Temp. C.	Time	Right	Left	Room Temp. C.
3.26	31.59	31.48		3.38	31.63	31.59	24.0
3.28	31.56	31.46	23.5	3.39	31.66	31.625	
3.29	31.565	31.47		3.40	31.695	31.665	23.95
3.30	31.57	31.475		3.41	31.72	31.685	
3.31	31.57	31.48		3.42	31.74	31.73	
3.32	31.57	31.485		3.43	31.78	31.76	24.0
3.33	31.58	31.50	24.2	3.44	31.79	31.77	
3.34	31.58	31.50		3.45	31.82	31.82	24.1
3.35	31.59	31.525		3.46	31.84	31.86	
3.36	31.60	31.55		3.47	31.88	31.90	
3.37	31.62	31.56		4.16	31.52	31.555	

Cooling of calorimeters in 29 minutes, R., 0.26 degrees; L., 0.245 degrees. Volume of right hand in calorimeter 444 c.c., of left hand 458 c.c. He is left handed. Water equivalent of calorimeters with contents, R., 3,450; L., 3,461.

Rectal temperature 37.35 degrees. Blood-pressure in left brachial, maximum systolic (when the larger pulsations begin to be heard) 111, 58 (sound gone). Another observation, maximum systolic 114, 58 (sound gone).

In two cases in which acute symptoms diagnosed as due to myocarditis had been present, but which had markedly improved under treatment so that the patients were able to come without trouble to the dispensary, a fair hand flow was found. Both were men 59 years old. In one of these patients (J. S. W.) who had been in the hospital two years before and had been near death, the flow was 6.82 grams per 100 c.c. per minute in the right hand and 6.58 grams in the left, with room temperature at 21.9 C. Simultaneous tracings from the jugular bulb, the radial artery and the apex showed that the "c" wave was notably large; the a wave was also well marked. All the ventricular beats were recognizable on the radial trace.

J. S. W., a fireman, colored, height 5 feet, 10 inches, weight 160 pounds, born in 1851, was first admitted to the dispensary Feb. 21, 1905, complaining of pain in chest and cough. The heart sounds were faint but clear. No murmurs. Heart dulness enlarged to the left well beyond the mammary line. The pulse was irregular. A diagnosis of myocarditis was made. April 13, 1908, he was again admitted to the dispensary. Heart enlarged to the left two fingers breadth beyond the mammary line. Pulse rapid and irregular. Some albumin in the urine. Nov. 4, 1908, he was admitted to Lakeside Hospital and again on Feb.

TABLE 6.—CALORIMETRIC MEASUREMENTS IN CASE OF J. S. W.

Time	Right	Left	Room Temp. C.	Time	Right	Left	Room Temp. C.
3.10	29.40	29.29		3.22	29.96	29.86	
3.14	29.47	29.38		3.23	30.00	29.91	
3.15	29.53	29.43		3.24	30.08	29.97	
3.16	29.59	29.50		3.26	30.20	30.10	
3.17	29.66	29.55		3.27	30.23	30.12	
3.18	29.73	29.61		3.34	30.16	30.06	
3.19	29.78	29.67		3.39	30.09	30.00	
3.20	29.83	29.71		3.51	29.96	29.88	
3.21	29.89	29.80					

20, 1909. He had attacks of syncope with fluttering in the chest and came near death. The pulse-rate was 120, systolic blood-pressure 160. Pulse regular and all the beats reached the wrist. Under treatment the pulse rate diminished and as it slowed the arrhythmia returned (premature contractions). March 19, 1909, the rate was 68, systolic blood-pressure 160. Tracings from the bulbus venosus showed well marked a, c and v waves of which the c wave was very large. Pre-systolic, systolic and diastolic pulsations were visible over the jugular bulb. He was discharged from the hospital May 5, 1909, improved. Oct. 5, 1909, he reported at the dispensary that he had gained 20 pounds since leaving the hospital. Oct. 4, 1910, he was feeling better, but had occasional smothering sensations. There was now no arrhythmia and no albumin in the urine.

The flow in the hands was measured Jan. 10, 1911. He feels much improved. Systolic pressure still very high; when last taken 210. Hands in bath at 3:00 p. m.; in calorimeters at 3:11½, and out of calorimeters at 3:27; 3,050 c.c. of water in each calorimeter. Room temperature 21.7 C. (Table 6.)

Volume of right hand 570 c.c., of left hand 570 c.c. His hands are very large. Pulse 60 (sitting). Mouth temperature 36.5 C. Rectal temperature 37.6 C. Room temperature at end of observations 22.2 C.

Since this time he has continued to report at the dispensary and has been more than once in the hospital with the signs of cardiac breakdown. On Dec. 7, 1912, "the left border of the cardiac dulness is at the anterior axillary line, and the right one finger's breadth beyond the right sternal border, heart thumping and somewhat irregular. Feb. 11, 1913, spitting blood and short of breath. Right border of cardiac dulness two fingers breadth to right of sternal margin. Sounds distant and irregular. No murmurs. Rales throughout both lungs." In June, 1913, eight years from the beginning of his illness, he was still coming to the dispensary and had periods of temporary improvement.

The other patient (Peter B.) was twice examined. At the first examination he had a flow of 6.44 grams per 100 c.c. of hand per minute in the right hand and 5.99 grams in the left, with room temperature 24.5 C. This is by no means a large flow for the temperature, and while fair reflex vasodilatation was obtained by the application of warmth to the contralateral hand, cold did not cause vasoconstriction sufficient to reduce the flow to the initial amount. This agrees well with the fact that arteriosclerosis was distinct, for reflex vasoconstriction is not well obtained in this condition. Almost a month later his hand flow was again measured. He had meanwhile much improved. The flow was 9.95 grams per 100 c.c. per minute for the right and 7.46 grams for the left hand. However, the room temperature was relatively high (26.6 C.), and this probably accounted for a portion of the increase in the flow. The vasodilatation consequent on the high external temperature was considerable and permitted a small reflex vasoconstriction when the contralateral hand was immersed in cold water.

Peter B., a janitor, aged 59, height 5 feet, 2 inches, weight 126 pounds, was under observation at the dispensary with symptoms diagnosed as due to myocarditis. For many years he did heavy work in a gas works. Though his work is now light he has had to give it up for the present. He suffered greatly from rheumatism till about fifteen years ago, since which time he has not been troubled. His joints would swell and he could not walk. He is married and has seven children, all in good health. His present illness dates from somewhat less than two years ago, when he suddenly became unconscious while mowing the lawn. He remained unconscious for five minutes. Three months later he had another attack. He came to the dispensary a number of times, and the records state that he had a slow pulse (60 a minute). The intervals between attacks became gradually shorter, and they occurred about once in six weeks. He falls down and "goes stiff" (extensor spasm) and now he sometimes has two attacks in a week. He seems particularly liable to become unconscious in bed. When he arises after bending down he is blind for about two minutes. Pupil reactions to light and accommodation are normal. He has long suffered from chronic bronchitis. Since the present trouble began his feet are always cold. Previously they used to be very hot and perspired greatly. He does not feel that his hands are cold. Physical examination negative for lungs. Heart dulness two finger's breadth to left of nipple. Sounds clear, distant, the first a little shortened. No accentuation of aortic or pulmonary. Arteries somewhat thickened. Pulse slow, fair volume, well sustained, easily compressible; rate 62. Marked slowing when the vagi are compressed in the neck. There is a venous pulse in the neck.

The flow in the hands was first examined April 21, 1911. Pulse (sitting) 80. Mouth temperature 36.45 C. Room temperature 24.8 C. Hands in bath at 2:31½ p. m., in calorimeters at 2:42. At 2:53 left hand put into water at 42.9 C.

TABLE 7.—CALORIMETRIC MEASUREMENTS IN CASE OF PETER B.

Time	Right	Left	Room Temp. C.	Time	Right	Notes
2:40½	30.04	30.03		3:00	30.57	At 3:03 put left hand in water at 8 C. Room 24.3 C.
2:43	30.01	30.02		3:01	30.60	
2:44	30.03	30.03	24.7	3:02	30.63	
2:45	30.06	30.06		3:03	30.69	
2:46	30.09	30.09		3:04	30.71	
2:47	30.11	30.10		3:05	30.77	At 3:11 dried left hand and exposed it for inspection. Then wrapped it in warm cloth.
2:48	30.14	30.12		3:06	30.79	
2:49	30.17	30.14		3:07	30.81	
2:50	30.20	30.18		3:08	30.83	
2:51	30.24	30.21	24.3	3:09	30.87	
2:52	30.28	30.23		3:10	30.89	
2:53	30.30	30.25		3:11	30.91	
2:54	30.32			3:12	30.93	
2:55	30.36		24.4	3:13	30.97	
2:56	30.40			3:14	31.00	
2:57	30.44			3:15	31.04	Room 24.2 C.
2:58	30.49		24.2	3:30	30.89	Hand out of calorimeter at 3:15.
2:59	30.52					At 3:30 temp. L. 29.95 C.

Cooling of calorimeters, R., 0.15 degree in fifteen minutes; L., 0.30 degree in thirty-seven minutes. Volume of right hand 360 c.c., of left 320 c.c. Water equivalent of calorimeters with contents, R., 3,383; L., 3,351.

Second examination May 17, 1911. He is feeling much better than when last examined. He was being treated with potassium iodid and nitroglycerin. Although he has had slight attacks of dizziness and "feeling stupid," he has not fallen or "gone stiff." There is no arhythmia now, nor can any murmurs be detected. Pressure on the vagus in the neck readily inhibits the heart causing the pulse at the wrist to disappear for a beat or two. He feels dizzy during the compression, and he can be made unconscious at any time by pressing on the vagus.

Hands in bath at 2:33 p. m., in calorimeters at 2:43½; mouth temperature 36.5 C.; pulse (sitting) 84. At 2:55 p. m. left hand put into water at 8 C.

Cooling of calorimeters, R., 0.07 degree in ten minutes; L., 0.17 degree in twenty-nine minutes. Volume of right hand in calorimeter 410 c.c., of left hand 394 c.c. The marks were put higher up than usual as his hands were short. The lower mark was about the middle instead of the lower border of the styloid process of the ulna. He seemed gradually to improve and reported at his last visit to the dispensary, June 7, 1911, that he was better. Pulse 84. (See Table 8.)

II. VALVULAR LESIONS

The flow in these cases may be normal or much less than normal, depending on the degree of compensation. Since anemia may be associated with a valvular lesion, it may be a factor in diminishing the hand

flow in the way already discussed in a previous paper.³ Since, further, the degree of involvement of the myocardium will differ in different cases with one and the same valvular lesion, it is not to be expected that a relatively large flow should always be associated with one particular type of valvular defect and a small flow with another. Nevertheless, so far as can be judged from the series of cases examined, certain of the valve lesions are more apt to be associated with the normal flow than others or even with a flow somewhat above the normal. Mitral insufficiency, for instance, it would seem, is less likely to be associated with a diminished

TABLE 8.—CALORIMETRIC MEASUREMENTS IN CASE OF PETER B.
SECOND TEST

Time	Right	Left	Room Temp. C.	Time	Right	Notes
2:43	30.29	30.18		3:00	31.125	At 3:05 left hand put in water at 43 C.
2:45	30.35	30.22	26.4	3:01	31.16	
2:46	30.42	30.27		3:02	31.195	
2:47	30.51	30.32		3:03	31.23	
2:48	30.57	30.38	26.5	3:04	31.29	
2:49	30.63	30.425		3:05	31.325	Room 26.7 C.
2:50	30.68	30.455		3:06	31.355	
2:51	30.73	30.50		3:07	31.38	
2:52	30.79	30.535		3:08	31.41	
2:53	30.835	30.57		3:09	31.44	
2:54	30.88	30.60		3:10	31.475	Room 26.6 C. Hand out of calorimeter at 3:14. At 3:24 temp. L. 30.455 C.
2:55	30.915	30.625		3:11	31.505	
2:56	30.94			3:12	31.53	
2:57	30.99		26.8	3:13	31.565	
2:58	31.035			3:14	31.59	
2:59	31.08			3:24	31.52	

hand flow than aortic insufficiency. In the only case in which a diagnosis of pure aortic stenosis seemed justified, the hand flow was normal, and this agreed with the clinical condition of the patient, who indeed did not enter the hospital for his heart lesion, but for a surgical cause. To emphasize the fact, however, that the mere naming of the valvular lesion is no indication of the amount of the hand flow, we shall begin the description of this group of cases with one of mitral incompetence in which the flow was very markedly diminished.

John T., contractor, aged 28, height 5 feet 6 inches, was admitted to the City Hospital March 31, 1912, with edema of the legs and hands and considerable cyanosis. A systolic murmur can be heard at the apex of the heart. The apex beat is visible and palpable with a heaving impulse in the sixth interspace in the nipple line. There is pulsation in the suprasternal notch. The cardiac dullness in the sixth left interspace is three fingers breadth outside the nipple line. The radial pulse is not easy to feel on account of the edema. It is small in volume and tension but has a regular rhythm. The urine contains albumin. He

was treated with digitalis. April 9, 1912, the blood-flow in the hands was examined. Hands in bath at 2:21 p. m., in calorimeters at 2:31. He had some cough during the observations. Systolic blood-pressure 122. Pulse 70. At 2:47 p. m. left hand put into water at 42.7 C.

TABLE 9.—CALORIMETRIC MEASUREMENTS IN CASE OF JOHN T.

Time	Right	Left	Room Temp. C.	Time	Right	Notes
2:30½	31.40	31.49		2:48	31.385	Room 24.3 C.
2:32	31.35	31.44	24.3	2:49	31.385	
2:33	31.35	31.44		2:50	31.39	
2:34	31.34	31.43		2:51	31.39	
2:35	31.34	31.43	24.1	2:52	31.395	
2:36	31.335	31.425		2:53	31.395	Hand out of calorimeter. Room 24.3 C.
2:37	31.335	31.425		2:54	31.395	
2:38	31.34	31.43		2:55	31.40	
2:39	31.345	31.435		3:11	31.23	At 3:11 temp. L. 31.225 C.
2:40	31.35	31.435				
2:41	31.35	31.435	24.3			
2:42	31.36	31.435				
2:43	31.365	31.44				
2:44	31.37	31.445				
2:45	31.37	31.45				
2:46	31.38	31.455	24.6			
2:47	31.38	31.455				

Cooling of calorimeters, R., 0.17 degree in sixteen minutes; L., 0.23 degree in twenty-four minutes. Volume of right hand 450 c.c., of left hand 438 c.c. Water equivalent of calorimeters with contents, R., 3,455; L., 3,445. Rectal temperature 37.95 C.

April 24, 1912, he was discharged "improved."

In this case the flow was only 2.09 grams per 100 c.c. of hand per minute in the right hand, and 1.67 grams in the left, with room temperature 24.4 C., a flow very much too small for his age.

In a case of mitral stenosis, mitral and tricuspid insufficiency in a man 63 years old (James H.), the flow, measured on an exceedingly hot day with a room temperature of nearly 31 C., was 7.31 grams per 100 c.c. of hand per minute for the right hand and 6.41 grams for the left. Probably not more than one-third to one-half of the flow to be expected in a healthy young adult for that room temperature. Measurements made at such high room temperatures do not lend themselves readily to comparisons. The examination was made on this day purposely to observe the effect of maximal vasodilatation on the flow in this case, since at two previous examinations the hand flow seemed to be greatly diminished by persistent local vasoconstriction which the patient attributed to the low temperature of the wards. He said he had felt chilly all the time, but to-day he felt warm. The pulse showed arrhythmia (Fig. 2), partly respiratory.

James H., a hatter, aged 63, admitted to the City Hospital May 2, 1913, suffering from chronic endocarditis with mitral stenosis and insufficiency and tricuspid insufficiency. No history of rheumatism. He complains of weakness and shortness of breath for the last three or four weeks. His feet swell at times. Ears, lips, nails and mucous membranes were markedly cyanotic on admission. There was slight edema of the legs with great varicosity of the leg veins. Marked

Fig. 2.—Tracings from James H.

engorgement and pulsation of the veins of the neck. Lungs revealed nothing special on physical examination. Heart enlarged. Left border of cardiac dullness at the anterior axillary line in the sixth interspace; right border 3 cm. to right of sternum. A presystolic thrill followed by impact and systolic thrill palpable over apex. Liver pulsatile and the lower edge palpable 8 cm. below the costal margin. Radial pulse markedly irregular and arrhythmic with fair volume and tension, fairly well sustained. He was discharged May 17 improved. He went to work but became ill again and returned to the hospital May 23, 1913.

TABLE 10.—CALORIMETRIC MEASUREMENTS IN CASE OF JAMES H.

Time	Right	Left	Room Temp. C.	Time	Right	Left	Room Temp. C.
2.00	31.30	31.24		2.17	31.07	30.98	23.3
2.02	31.25	31.17	23.1	2.18	31.055	30.97	
2.03	31.23	31.16		2.19	31.04	30.96	
2.04	31.22	31.15		2.20	31.03	30.955	23.4
2.05	31.20	31.13	23.3	2.21	31.02	30.95	
2.06	31.19	31.12		2.22	31.01	30.945	
2.07	31.18	31.105	23.3	2.23	31.00	30.94	23.4
2.08	31.17	31.09		2.24	30.995	30.93	
2.09	31.15	31.08	23.2	2.25	30.995	30.925	
2.10	31.13	31.06		2.26	30.99	30.915	23.5
2.11	31.125	31.055		2.27	30.98	30.90	
2.12	31.11	31.05	23.2	2.28	30.975	30.895	
2.13	31.10	31.04		2.29	30.97	30.89	
2.14	31.09	31.03	23.2	2.30	30.96	30.88	23.6
2.15	31.08	31.01		2.31	30.955	30.84	
2.16	31.07	30.99					

The first examination of the hand flow was made on May 29, 1913. His hands were cold when he was brought into the room, having been exposed for some time near an open window. The day, however, was not especially cool. He was allowed to keep the hands wrapped up for twenty minutes in his gown before the observations were begun. Pulse 70 (sitting). The volume of the pulse seemed fair and was thought to indicate a much better flow than was actually shown by the examination, but the veins on the back of the hands, although prominent, filled but slowly when emptied of blood. Hands in bath at 1:50 p. m.; in calorimeters at 2:00¹/₂, and out of calorimeters at 2:30.

Cooling of calorimeters in fifty-one minutes, R., 0.565 degree; L., 0.54 degree. Volume of right hand 418 c.c., of left hand 396 c.c. The hands do not feel cold to the touch on coming out of the calorimeters. They are fairly red. The veins on the back of the hand and in the forearm are large, but fill with marked slowness on being emptied. Blood-pressure, left arm, systolic 136, 63 (sound gone). Another observation 134, 68. There was no sensible interval between the change in sound and its disappearance. A few of the larger beats were heard a little before the readings given as systolic pressure. The number of waves coming through increased as the pressure fell.

The flow in the hands in this examination works out at only 0.80 gram per 100 c.c. per minute for the right hand, and 0.64 gram for the left, for the last ten minutes of immersion in the calorimeters. If the whole period in the calorimeters is taken, the calculated flow is still less, as is always the case with a very small flow, the period required for the attainment of the steady state where the thermometers are changing uniformly being long.

A second examination on June 2, 1913, gave an even smaller calculated flow (0.40 gram per 100 c.c. per minute) for the last ten minutes in the calorimeters. The temperature readings followed precisely the same course as at the first examination. The question arises how such small flows can be reconciled with the fairly good volume of the arterial pulse at the wrist. The explanation is found partly in the fact that the venous return was hindered as indicated by the appearance of the veins. The increase in venous pressure while it did not hinder the conduction of the arterial pulse wave, apparently diminished the rate of translation of the blood. A second factor in the small flow was apparently a persistent local vasoconstriction in the hands. That this second factor was concerned is shown by the great improvement in the hand flow at the third examination, although the pulse-rate, far from being accelerated, was even slower than at the previous examinations.

Third examination June 16, 1913. He is feeling well, only weak. No dyspnea and no cough whatever. The day is very warm and the room temperature the highest in the whole series of observations. The veins on his hands and arms fill much more quickly after being emptied than on the two previous occasions. Amplitude of the pulse at the wrist is fair, but not strikingly greater than at the other examinations. Pulse 57 (sitting), with some irregularity but not so marked as before. Hands in bath at 4:06½ p. m.; in calorimeters at 4:15½, and out of calorimeters at 4:27.

Cooling of calorimeters in ten minutes, R., 0.045 degree; L., 0.04 degree. Volume of right hand in calorimeter 408 c.c., of left hand 392 c.c. Water equivalent of calorimeters with contents, R., 3,421; L., 3,409. Blood-pressure in left brachial, systolic 110, another observation 106, 80 (change in sound). Rectal temperature 37.6 C. (See Table 11.)

The general cutaneous vasodilatation at the third examination was evident and is doubtless related to the decidedly diminished systolic pressure. The hands shared in this general vasodilatation, and it is

probable for this reason that the hand flow gives an index to the heart output. Since the hand flow with practically maximal vasodilatation is not half that of a young adult at this room temperature, the heart output is probably at most 50 per cent. of that of a healthy young adult. The three observations together show that in this man of 63 the range of the vasomotor adaptation of the hand blood-vessels is still considerable, a confirmation of his statement that the hot weather suited him.

TABLE 11.—CALORIMETRIC MEASUREMENTS IN THIRD EXAMINATION OF JAMES H.

Time	Right	Left	Room Temp. C.	Time	Right	Left	Room Temp. C.
4.15	31.45	31.50		4.22	31.71	31.72	30.9
4.16	31.46	31.52		4.23	31.765	31.76	
4.17	31.49	31.54		4.24	31.795	31.83	
4.18	31.54	31.58	30.8	4.25	31.86	31.84	30.9
4.19	31.58	31.62		4.26	31.89	31.87	
4.20	31.62	31.655	30.9	4.27	31.925	31.90	
4.21	31.67	31.69		4.37	31.88	31.86	

From the standpoint of the technic of the method it ought still to be mentioned that a local vasoconstriction so persistent as not to be overcome during the usual period of an examination (the disappearance of it being shown by a rather abrupt and henceforth steady rise of the thermometers) is rare in hospital cases, although in hands long exposed to cold it may be frequently met with. Where it exists the results cannot, of course, be used as average flows for the patient in question, although the intensity and persistence of the vasoconstriction may be of interest in relation to the patient's condition.

In a case of badly broken compensation in a boy (Joseph Z.), 16 years old, suffering from chronic endocarditis with aortic insufficiency and stenosis and mitral insufficiency, the flow in the right hand was 6.28 grams per 100 c.c. per minute, and in the left hand 4.79 grams. This is much too small for the age of the patient and the very high room temperature (28.1 C.). He was obviously in bad condition. Immersion of the left hand in cold water was followed by but a small diminution of the flow in the right, and immersion of the left hand in warm water by no increase at all, either because the dyspneic condition of the blood caused increased stimulation of the vasoconstrictors of the hand which the vasodilator effect of warmth to the contralateral hand could not overcome, or because the vasoconstrictor tone was already much diminished under the influence of the warm room, although, owing to the handicap of the heart, this was not shown in a large flow. Another possibility is that the dyspneic condition of the blood may have rendered the vasomotor centers less excitable to this form of afferent stimulation.

Joseph Z., boy aged 16, was admitted to the dispensary Nov. 11, 1910, suffering from aortic insufficiency and stenosis with attacks of acute pulmonary edema. He has had rheumatism for ten years. He was in the hospital three weeks for heart trouble which started four years ago. Has dyspnea and weakness. His heart is enlarged 16 cm. to the left, and 5 cm. to the right of the median line. Pulsation in the arteries of the neck. Apex beat heaving. Systolic impulse over apex. A systolic thrill can be felt to the right of the sternum. A double murmur over the apex and louder over the aortic region. Pulse 95. Moderate tension, rather slow rise and quick fall. Pistol shot in femorals. Capillary pulse.

Nov. 15, 1910, he was admitted to the hospital where he improved and after his discharge he remained well, keeping in bed, however, most of the time.

Feb. 2, 1911. When in the dispensary he had an attack of palpitation during which the pulse rate rose to 160, with great throbbing in the carotids and general cyanosis. He began to spit up pinkish frothy liquid. The attack lasted forty-five minutes. He was again admitted to the house, where the diagnosis of chronic endocarditis, mitral insufficiency and aortic insufficiency, was made. He suffered from occasional attacks of tachycardia and shortness of breath. The blood-examination showed hemoglobin 90 per cent. He was discharged July 5, 1911, "improved."

July 14, 1911, he had an attack in the dispensary. There was great pain referred to the right infrascapular region. He received morphin (grain $\frac{1}{8}$) and atropin sulphate (gr. 1/300) hypodermically to relieve and calm him, and fifteen minutes later the blood flow in his hands was measured. Pulse (sitting) 140. Hands in bath at 3:01 p. m., in calorimeters at 3:11½. Mouth temperature 38.15 C. At 3:21 p. m. left hand was put into water at 8 C.

TABLE 12.—CALORIMETRIC MEASUREMENTS IN CASE OF JOSEPH Z.

Time	Right	Left	Room Temp. C.	Time	Right	Notes
3:11	29.87	29.905		3:27	30.59	
3:12	29.88	29.915		3:28	30.60	Pulse 132.
3:13	29.91	29.93		3:29	30.635	
3:14	29.98	29.97		3:30	30.695	At 3:30 put left hand in water at 43 C.
3:15	30.07	30.04	28.1	3:31	30.735	
3:16	30.13	30.10		3:32	30.775	Room 28.4 C.
3:17	30.21	30.13		3:33	30.81	
3:18	30.22	30.15		3:34	30.87	
3:19	30.27	30.20		3:35	30.895	
3:20	30.29	30.22		3:36	30.93	
3:21	30.32	30.235		3:37	30.975	Room 28.3 C.
3:22	30.34			3:38	31.01	3:38 hand out calorimeter.
3:23	30.37	30.22		3:38½		Temp. of L. 30.16 C.
3:24	30.40			3:48	30.97	Pulse 120. Mouth temp. 38.45 C.
3:25	30.49		28.1			
3:26	30.54					

Cooling of calorimeters, R., 0.04 C. in ten minutes; L., 0.06 C. in fifteen one-half minutes. Volume of right hand 406 c.c., of left 412 c.c. He is right handed. Water equivalent of calorimeters with contents, R., 3,420; L., 3,425.

He attended from time to time at the dispensary feeling pretty well for awhile, but compensation broke down again and on March 3, 1913, he had again to seek admission to Lakeside Hospital with the cardiac dulness enlarged to the left to the mid-axillary line.

In another case of aortic stenosis and insufficiency and mitral insufficiency with fair compensation at the time of the blood-flow examination in a man (Matthew K.), 42 years old, the flow was 7.20 grams per 100 c.c. per minute in the right, and 5.49 grams in the left, with room temperature 22.5 C., a flow somewhat below the normal for his age and the external temperature, although by no means remarkably small. He had no edema and no dyspnea, but he complained of pain over the heart and was getting weaker.

Mat. K., aged 42, height 5 feet, 7 inches, weight 165 pounds, a worker in a railway shop, was admitted at the dispensary Dec. 2, 1910, suffering from aortic stenosis and insufficiency and mitral insufficiency. His illness began two months before and he has been getting worse ever since. He feels chilly at times. His weakness is increasing each day and he has had to give up work. He suffers from pain over the heart, palpitation and inability to sleep. There is a probable history of rheumatic fever seven years ago. The left border of the cardiac dulness is 12½ cm. to the left of the mid-sternal line, the right border 4 cm. to the right of it, upper border at third rib. No capillary pulse, no edema, urine normal.

The blood-flow in the hands was measured Jan. 24, 1911. He has no shortness of breath. The systolic blood-pressure is 170. Mouth temperature 36.6 C.; pulse (sitting) 80. Hands in bath at 3:16 p. m.; in calorimeters at 3:27, out of calorimeters at 3:42. There was 3,050 c.c. of water in each calorimeter. Room temperature 22.0 C. before the observations were begun.

TABLE 13.—CALORIMETER MEASUREMENTS IN CASE OF MAT. K.

Time	Right	Left	Room Temp. C.	Time	Right	Left	Room Temp. C.
3.26	29.42	29.19		3.37	29.73	29.43	
3.28	29.40	29.18		3.38	29.78	29.46	
3.29	29.42	29.18		3.39	29.82	29.48	
3.30	29.44	29.20		3.40	29.86	29.51	
3.31	29.47	29.21		3.41	29.90	29.54	
3.33	29.54	29.29		3.42	29.96	29.58	23.1
3.34	29.60	29.33		3.59	29.76	29.42	
3.35	29.64	29.37		4.04	29.70	29.37	
3.36	29.70	29.40					

Cooling of calorimeters in twenty-two minutes, R., 0.26 degree; L., 0.21 degree. Volume of right hand in calorimeter 440 c.c., of left 435 c.c. He is right handed.

He continued to come to the dispensary. On Feb. 2, 1911, he complained of pain over the precordia and down the shoulder and arm. His last visit was on April 2, 1913, when his condition was not essentially altered.

In a case of mitral and tricuspid insufficiency (P. B.) with venous engorgement, edema of the ankles and fluid in the peritoneal cavity, a perfectly good flow was seen in the hands (12.80 grams per 100 c.c. per minute for the right and 13.46 grams for the left, with room temperature 25.5 C.). A fair reflex vasoconstriction was obtained by the application of cold to the contralateral hand, but the application of warmth failed to increase the flow, even to the initial level, owing no doubt to the fact

that the vessels were already so much dilated. There was no dyspnea during the experiment, and the driving power of the heart with its obviously powerful stroke was adequate to maintain a normal flow through the periphery. The vasomotor centers seem to have retained their normal excitability. The cord, therefore, in spite of the venous engorgement, was not suffering appreciably at the time.

P. B., section hand on a railway, aged 31, height 5 feet, 11 inches, was admitted to the dispensary May 12, 1911, with mitral and tricuspid insufficiency. He has been ill for eighteen or nineteen months, during which time he has been unable to work. He had acute articular rheumatism nineteen months before admission, had swollen feet, enlargement of the abdomen and palpitation of the heart and was in bed six weeks. The left border of the cardiac dullness is at the mid-axillary line. There is a large diffuse area of cardiac pulsation visible (positive systolic) and the framework of the chest undulates with each beat. Positive venous pulsation in both jugulars. Liver much enlarged with marked systolic pulsation. There is venous engorgement, edema of the ankles and fluid in the peritoneal cavity.

The blood-flow in the hands was measured May 12, 1912. There was no dyspnea during the examination. Pulse (sitting) 100. Mouth temperature 36.9 C. Hands put into bath at 2:29 p. m., into calorimeters at 2:39. At 2:50 p. m. left hand put into water at 8 C.

TABLE 14.—CALORIMETRIC MEASUREMENTS IN CASE OF P. B.

Time	Right	Left	Room Temp. C.	Time	Right	Notes
2:38	30.21	30.11		2:57	31.67	Room 25.7 C. At 3:01 put left hand in water at 43.1 C.
2:40	30.28	30.19		2:58	31.71	
2:41	30.375	30.28		2:59	31.77	
2:42	30.48	30.40		3:00	31.82	
2:43	30.59	30.495		3:01	31.875	
2:44	30.68	30.61	25.5	3:02	31.91	Room 25.5 C.
2:45	30.79	30.71		3:03	31.97	
2:46	30.88	30.80		3:04	32.02	
2:47	30.97	30.875		3:05	32.085	
2:48	31.06	30.96		3:06	32.135	
2:49	31.14	31.06	25.6	3:07	32.195	3:10 hand out calorimeter. Temp. of L. is now 30.98. Room 25.7 C.
2:50	31.23	31.16		3:08	32.26	
2:51	31.295			3:09	32.32	
2:52	31.345			3:10	32.38	
2:53	31.395			3:10½		
2:54	31.475			3:22½	32.25	Temp. L. is now 30.88 C.
2:55	31.545		25.8	3:23½		
2:56	31.605					

Cooling of calorimeters, R., 0.13 degree in twelve one-half minutes; L., 0.28 degree in thirty-two minutes. Volume of right hand 510 c.c.; of left hand 480 c.c.

He entered Lakeside Hospital August 21, 1911. He had then neither cyanosis nor edema, but there was still fluid in the abdomen. The apparent incongruity between the distention of the abdomen, the pulsation of the liver and the great comfort of the patient was remarked. Heart dullness extends on the right to 7 cm.

from the mid-sternal line in the third interspace, on the left to the anterior axillary line in the sixth interspace. The blood examination showed 90 per cent. hemoglobin. He again entered Lakeside Hospital Nov. 2, 1911, and was discharged "improved" Feb. 20, 1912.

The large flow in P. B. contrasts with the small flow in Joseph Z., in whom the delivery of blood to the systemic vessels was seriously interfered with by the insufficiency of the aortic valve.

In Ernest M., another case of mitral insufficiency with chronic bronchitis, in a boy 14 years old, the flow was also good (12.74 grams per 100 c.c. per minute for the right hand and 12.45 grams for the left hand, with room temperature 23.4 C.). Compensation was so far established that he was able to come to the dispensary and had no dyspnea during the examination.

Ernest M., boy, aged 14, height 5 feet, 1 $\frac{1}{4}$ inches, was admitted at the dispensary Jan. 26, 1911, suffering from mitral insufficiency with chronic bronchitis. The fingers of both hands show well marked clubbing. He has had a cough ever since he was two years old. He still has the cough and is getting thinner. His arms are thin for his age. He gives a history of heart trouble. He has never been very strong and is not well enough to go to school. His appetite is good. Up to the present time there has never been any blood in the sputum, but on Feb. 7, 1911, he stated that since his last visit he spat some blood. The sputum examination was always negative as regards tubercle bacilli.

The hand flow was measured Jan. 26, 1911. Mouth temperature 38.2 C. Rectal temperature 38.5 C. Hands in bath at 2:56 p. m.; in calorimeters at 3:07, and out of calorimeters at 3:23. There was 3,150 c.c. of water in each calorimeter. Pulse (sitting) 116. The weather was mild with rain. Room 23.4 C.

TABLE 15.—CALORIMETRIC MEASUREMENTS IN CASE OF ERNEST M.

Time	Right	Left	Room Temp. C.	Time	Right	Left	Room Temp. C.
3.06	29.71	29.68		3.17	30.18	30.10	
3.09	29.73	29.68		3.18	30.24	30.16	23.4
3.10	29.77	29.72		3.20	30.34	30.27	
3.12	29.89	29.82		3.21	30.40	30.33	
3.13	29.96	29.89		3.22	30.45	30.39	
3.14	30.00	29.93		3.23	30.50	30.42	23.1
3.15	30.06	29.99		3.42	30.30	30.27	22.7
3.16	30.12	30.05					

Cooling of calorimeters in nineteen minutes, R., 0.20 degree; L., 0.15 degree. L. was on the side of the room next to the radiator, though at a good distance from it. A small centrifuge had been started at 3:23 at the side of the room nearest to R. The difference in the rate of cooling of the calorimeters, which is decidedly greater than usual, may be thus accounted for. Volume of right hand 255 c.c., of left 240 c.c.

In another boy (Frank C.), 17 years old, with mitral stenosis and insufficiency, the flow was also quite normal in amount (11.25 grams for the right and 11.58 grams for the left hand, with room temperature 22.4

C.). Compensation was fairly established. The protocol has already been published.⁶ Later he was admitted several times to Lakeside Hospital suffering from broken compensation and was discharged for the last time on March 4, 1912. He afterward died.

One of the most interesting cases was that of Albert H., since the heart underwent rapid deterioration within the period of the observations and the patient died soon after the last examination, when an autopsy was obtained. The patient was a man 31 years of age with a clinical diagnosis of mitral and tricuspid insufficiency, confirmed by the autopsy, which in addition showed pericarditis with large effusion into the pericardial sac. There was no doubt that the myocardium had suffered. The patient suffered much from dyspnea if he attempted to lie down. The flow in the right hand was 4.08 grams per 100 c.c. per minute, and in the left 4.54 grams, with room temperature 24.7 C., a distinctly subnormal flow. Reflex vasoconstriction of the hand was easy to obtain on immersion of the contralateral hand in cold water, but no reflex vasodilatation could be obtained, which agreed well with the obviously poor filling of the vessels of the hand. Ten days later he was again examined. The abdomen was now full of liquid, the legs and feet greatly swollen, the heart-sounds weak and the pulse at the wrists so weak that the heart-rate was counted with the stethoscope. The hands also had increased in size. The blood-flow in the hands was very small and what was considered significant at the time, less than half its amount at the last examination (2.0 grams for the right and 1.88 grams for the left). This was for a period of seven minutes, beginning fifteen minutes after insertion of the hands in the calorimeters. For the first fifteen minutes the flow was still less. The patient said he felt better, but the great decline in the blood-flow was remarked as an ominous symptom. He died in the hospital ten days after the last examination.

Albert H., colored hotel porter, aged 31, was admitted to the City Hospital April 3, 1912, complaining of dyspnea, cough and swelling of the extremities. He had rheumatism three weeks before admission and the legs were very painful. The edema and dyspnea began one week before admission. He has grown worse, coughs most of the time and is very tired on any exertion. The physical signs of fluid in the right pleural cavity are present. Heart dullness, upper border third rib, right border parasternal line, left border midway between nipple and anterior axillary line. The rate of the heart is rapid, the sounds feeble. There is a soft mitral systolic murmur with reduplication of the second sound. Radial pulse small and easily compressible. No fibrosis. Great edema of scrotum and prepuce, and edema of lower extremities. Albumin in urine. April 5 and April 7 the dyspnea and edema were less, the mitral murmur not so pronounced, the pulse soft and not well sustained. April 9, on which day the hand flow was first measured, the signs of an increase in the fluid in the right pleural cavity were present. The heart boundaries were the same as before and the edema was still considerable. Seven hundred and fifty c.c. of liquid was aspirated from the right thorax before the blood-flow examination. His maximum temperature on April 3 was 100.6 F. It rose gradually till April 9 when it reached 104.5 F. The systolic blood-pressure on April 5 was 152; April 7, 160; April 9, 160.

Hands in bath at 1:09 p. m., in calorimeters at 1:19. Mouth temperature 37.8 C. At 1:31 p. m. left hand put into water at 7.9 C.

TABLE 16.—CALORIMETRIC MEASUREMENTS IN CASE OF ALBERT H.

Time	Right	Left	Room Temp. C.	Time	Right	Notes
1:18	30.82	30.78		1:35	31.13	
1:20	30.81	30.75		1:36	31.14	
1:21	30.84	30.76	24.6	1:37	31.16	Room 24.7 C.
1:22	30.87	30.79		1:39	31.19	
1:23	30.89	30.82		1:40	31.195	At 1:40 left hand put in water at 43 C.
1:24	30.90	30.85	24.9	1:41	31.21	
1:25	30.96	30.90		1:42	31.22	Room 24.8 C.
1:26	31.03	30.96		1:43	31.23	
1:27	31.07	31.00		1:44	31.245	
1:28	31.08	31.02	24.6	1:45	31.26	
1:29	31.085	31.04		1:46	31.28	
1:30	31.095	31.07		1:47	31.295	
1:31	31.09	31.09		1:48	31.305	
1:32	31.10			1:49	31.325	
1:33	31.105		24.5	1:50	31.33	1:50 hand out calorimeter.
1:34	31.115			1:59	31.23	Temp. of L. is 30.81 C. at 1:59.

Cooling of calorimeters, R., 0.10 degree in nine minutes; L., 0.28 degree in twenty-eight minutes. Volume of right hand 447 c.c., of left hand 450 c.c. He says he does most of his work left handed. Water equivalent of calorimeters with contents, R., 3,452; L., 3,455. Pulse (lying down) 120. Rectal temperature 39.1 C.

April 10 his maximum temperature was 103.8 F.; April 14, 103.3 F. April 17, "the dulness in the right thorax begins as high as the fifth rib posteriorly. Heart boundary is the same." Systolic blood-pressure April 16, 130; April 18, 140; April 19, 130.

Second examination of the hand flow, April 19, 1912. The abdomen now contains much liquid. The legs and feet are greatly swollen and he cannot sleep because he is unable to lie down on account of dyspnea. The heart-sounds are weak and the pulse at the wrist barely discernible. He says he feels better than before the tapping on April 9, but he seems worse than when examined after the tapping on that date. The arms and hands, according to his statement, are not swollen, but simply fat and there is no pitting. Yet measurement of the volume of the hands demonstrates an increase in their size, although the edema is not great enough to be detected by pitting. He has no fever now.

Hands in bath at 1:58 p. m.; in calorimeters at 2:09 p. m., and out of calorimeters at 2:45. At 2:35 the right hand was put into water at 42.7 C.

Cooling of calorimeters, R., 0.295 degree in twenty-eight and one-half minutes; L., 0.17 degree in eighteen and one-half minutes. Pulse 102 (counted with stethoscope). Rectal temperature 37.75 C. It was noticed that he now has piles. He never had them before. Volume of right hand in calorimeter 475 c.c., of left hand 505 c.c. Water equivalent of calorimeters with contents, R., 3,475; L., 3,499. (See Table 17.)

April 26, 1912, edema was general in the legs, arms and face. Maximum temperature 102.4 F. He died April 29. Results of necropsy: Pericarditis; 600 c.c. fluid in the pericardial sac. Mitral valve, edges thickened, insufficient. Tricuspid valve, insufficient. One thousand c.c. fluid in the abdomen. Chronic passive congestion of the liver, spleen and kidneys.

Whether the mere mechanical effect of the liquid in the pericardial sac contributed to the small flow in Albert H. at the second examination cannot of course be determined, although it is likely enough that this was a factor in addition to the weakening of the myocardium and the insufficiency of the valves. However, if the clinical diagnosis in the next case (Roman B.) be accepted, liquid may be present in considerable amount in the pericardial sac without reducing the hand flow very markedly. This patient was a man aged 22, in whom the diagnosis of mitral insufficiency and pericardial effusion was made. Blood-flow measurements in the hands were obtained on three occasions, the first on May 29, 1912, when the heart dulness was much diminished and the

TABLE 17.—CALORIMETRIC MEASUREMENTS IN SECOND EXAMINATION OF ALBERT H.

Time	Right	Left	Room Temp. C.	Time	Right	Left	Room Temp. C.
2.08	29.81	29.72		2.28	29.66	29.59	
2.10	29.74	29.67		2.29	29.675	29.61	
2.11	29.72	29.65	22.7	2.30*	29.685	29.62	
2.12	29.705	29.635		2.31	29.68	29.61	
2.13	29.695	29.625		2.32	29.69	29.625	
2.14	29.69	29.62		2.33	29.695	29.63	
2.15	29.68	29.61		2.34	29.70	29.63	
2.16	29.67	29.60	22.5	2.35	29.705	29.635	
2.17	29.665	29.595		2.36		29.63	
2.18	29.65	29.58		2.37		29.635	23.5
2.19	29.64	29.575		2.38		29.635	
2.20	29.635	29.565	22.6	2.39		29.645	
2.21	29.63	29.56		2.40		29.65	
2.22	29.625	29.555		2.41		29.655	
2.23	29.62	29.55		2.42		29.655	
2.24	29.62	29.55		2.43		29.66	
2.25	29.63	29.56		2.44		29.665	
2.26	29.64	29.57	22.7	2.45		29.67	
2.27	29.65	29.58		3.03½	29.41	29.49	

* Here the bringing in of the warm water attracted his attention. In excitable patients this may cause a change in the flow.

swelling of the legs present on admission had disappeared. The flow came out 7.61 grams per 100 c.c. per minute in the right hand and 10.59 grams in the left hand, with room temperature 22 C. He seemed fairly comfortable and had no dyspnea. On May 31, when the weather was warmer, the room temperature being 26.3 C., the flow was 8.76 grams for the right hand and 11.05 grams for the left. The average for the two hands is normal, or nearly so, and greater of course at the second examination, in accordance with the increased room temperature. He left the hospital, and after working in a foundry for two days, his legs became swollen and he had to return to the hospital on June 11 with the same

symptoms of broken compensation which he had at first, dyspnea, cough, greatly increased area of cardiac dulness and the edema already mentioned. On June 13 his hand flow was again examined, and came out only 5.52 grams per 100 c.c. per minute for the right hand and 8.74 grams for the left, with room temperature 21.8 C., a considerable smaller flow than at the first examination, although the external temperature was practically the same. The flow, then, concomitantly with the cardiac breakdown, has been decidedly diminished. The marked inequality in the flow in the two hands persists throughout, although the relative difference is greater at the third than at the other two examinations. No local difference in the hands or arms could be discovered to account for the difference in the flows. It is possible that it is connected with intrathoracic changes not taken account of in the diagnosis. The question of differences of flow in the two hands associated with unilateral changes in the lungs and pleurae will be taken up in a future paper.

Roman B., a laborer, aged 22, was admitted to the City Hospital May 17, 1912. He had rheumatism for the first time three years before admission. He has a typical cardiorenal facies. He seems fairly comfortable, with no dyspnea. The heart presents a pear-shaped area of dulness, left border in the mid-axillary line, right border one to three fingers' breadth to the right of the mid-sternal line, making an obtuse cardiohepatic angle. The cardiac dulness is increased when he leans forward. There is a loud systolic murmur at the apex. The second sound in the pulmonic area is split. The sounds are distant. Pulse regular in rhythm, of fair volume, rather rapid rise, and well sustained. Liver three fingers' breadth below costal margin; hard, sharp edge, not tender. Edema of legs. Urine negative.

TABLE 18.—CALORIMETRIC MEASUREMENTS IN CASE OF ROMAN B.

Time	Right	Left	Room Temp. C.	Time	Right	Left	Room Temp. C.
3.21	31.03	31.05	21.4	3.32	31.52	31.68	
3.23	31.08	31.09		3.33	31.56	31.74	22.2
3.24	31.12	31.15	21.5	3.34	31.58	31.77	
3.25	31.19	31.23		3.35	31.60	31.825	22.6
3.26	31.23	31.315		3.36	31.64	31.89	
3.27	31.28	31.36		3.37	31.68	31.94	22.8
3.28	31.30	31.42		3.38	31.705	31.98	
3.29	31.36	31.47	21.0	3.39	31.735	32.025	22.5
3.30	31.42	31.56		3.52	31.555		
3.31	31.48	31.64	21.5	3.52½		31.83	

Cooling of calorimeters, R., 0.18 degree in thirteen minutes; L., 0.195 degree in thirteen one-half minutes. Volume of right hand 454 c.c. of left hand 439 c.c. He is right handed. Water equivalent of calorimeters and contents, R., 3,458; L., 3,446. Pulse (sitting) 96. Rectal temperature 37.95 C.

May 25, 1912, left cardiac border just outside the nipple line. Right border one finger's breadth outside of sternal margin. Cardio-hepatic angle less obtuse. Liver one finger's breadth below costal border. Heart sounds louder and more distinct.

May 29, 1912, the flow in his hands was measured. Hands in bath at 3:12 p. m.; in calorimeters at 3:21¼, and out of calorimeters at 3:39. Mouth temperature 37.25 C.

Second examination May 31, 1912. He says his right ankle pains him now. It was painful three years ago when he had rheumatism for the first time. Hands in bath at 1:51 p. m; in calorimeters at 2:01½, and out of calorimeters at 2:20.

The weather was considerably warmer than at the last examination.

TABLE 19.—CALORIMETRIC MEASUREMENTS AT SECOND EXAMINATION OF ROMAN B.

Time	Right	Left	Room Temp. C.	Time	Right	Left	Room Temp. C.
2.01	31.04	31.03	26.2	2.12	31.52	31.65	26.2
2.03	31.04	31.06		2.13	31.57	31.71	
2.04	31.08	31.125	26.3	2.14	31.61	31.77	26.3
2.05	31.125	31.17		2.15	31.67	31.85	
2.06	31.20	31.25	26.4	2.16	31.72	31.925	26.3
2.07	31.27	31.34		2.17	31.76	31.97	
2.08	31.32	31.42	26.4	2.18	31.795	32.025	26.2
2.09	31.38	31.48		2.19	31.83	32.08	
2.10	31.42	31.54	26.3	2.20	31.88	32.13	26.2
2.11	31.48	31.60		2.32	31.76	32.02	

Cooling of calorimeters in twelve minutes, R., 0.12 degree; L., 0.11 degree. Volume of right hand 451 c.c., of left hand 445 c.c. Water equivalent of calorimeters with contents, R., 3,456; L., 3,451. Pulse (sitting) 87. Rectal temperature, 37.65 degrees.

June 12, 1912, he was readmitted to the hospital, complaining of precordial pain, dyspnea, and cough. The right leg pains him more than before. Lips cyanosed. Finger tips bluish. Edema in legs. There is a diffuse wave during systole not prominent in any one region. Limits of cardiac dulness were as follows:

Interspace	Supine Posture Distance in Cm. from Mid-sternal Line		Sitting Posture Distance in Cm. from Mid-sternal Line	
	Right Border	Left Border	Right Border	Left Border
6	9.5	16	12	17.5
5	8.0	15	9.5	15.5
4	6.5	10	6	11.0
3	4.0	7	..	

Auscultation, with moderate pressure of stethoscope, rough systolic murmur at apex. Pulmonic second accentuated and split. Aortic second, weak. With pressure of stethoscope a soft blowing murmur is heard at apex during diastole. The sounds are reduced in intensity. Free fluid in abdomen.

June 13, 1912, third examination of the hand flow. Hands in bath at 1:58½ p. m.; in calorimeters at 2:09½, and out of calorimeters at 2:25.

TABLE 20.—CALORIMETER MEASUREMENTS IN THIRD EXAMINATION OF ROMAN B.

Time	Right	Left	Room Temp. C.	Time	Right	Left	Room Temp. C.
2.09	31.145	31.06	21.4	2.20	31.345	31.47	22.0
2.11	31.13	31.07		2.21	31.37	31.51	
2.12	31.14	31.095		2.22	31.385	31.545	
2.13	31.165	31.13	21.8	2.23	31.395	31.59	22.1
2.14	31.18	31.17		2.24	31.405	31.625	
2.15	31.20	31.225	21.9	2.25	31.415	31.635	
2.16	31.22	31.26		2.34	31.28	31.51	
2.17	31.26	31.325	22.0				
2.18	31.29	31.375					
2.19	31.315	31.42					

Cooling of calorimeters in nine minutes, R., 0.135 degree; L., 0.125 degree. Volume of right hand 451 c.c., of left 443 c.c. Rectal temperature 37.3 C. Water equivalent of calorimeters with contents, R., 3,456; L., 3,449.

In a case (Robert T.) diagnosed as pure aortic stenosis with perfect compensation in a man, aged 22 years, the average flow in the hands was at least equal to normal (10.45 grams per 100 c.c. per minute for the right hand and 13.97 grams for the left, with room temperature 20.3 C.). He entered the hospital not for his heart condition, but for surgical treatment.

Robert T., a railroad worker 22 years old, was admitted to the City Hospital Jan. 23, 1913, for surgical treatment (crushed toes on left foot). He had typhoid fourteen years ago; also rheumatism in the neck for several weeks when a child. Denies venereal diseases. For four years after typhoid had shortness of breath and still gets short of breath if he has to hurry up, especially if he changes the

TABLE 21.—CALORIMETRIC MEASUREMENTS IN CASE OF ROBERT T.

Time	Right	Left	Room Temp. C.	Time	Right	Left	Room Temp. C.
2.52	31.22	31.13		3.00	31.38	31.42	20.3
2.53	31.19	31.10		3.01	31.42	31.48	
2.54	31.205	31.12	20.1	3.02	31.47	31.54	20.4
2.55	31.215	31.15	20.1	3.03	31.515	31.59	
2.56	31.23	31.18		3.04	31.56	31.65	
2.57	31.27	31.24	20.2	3.05	31.595	31.71	20.4
2.58	31.30	31.29	20.3	3.06	31.64	31.77	
2.59	31.35	31.36		3.27	31.28	31.41	

manner of his work, but not when he has become accustomed to a job and can work steadily without spurts. He is well-developed and well-nourished, no cyanosis. Boundaries of cardiac dulness one and a half finger's breadth to left of nipple line, right sternal border, third interspace. A blowing systolic murmur is heard over the entire area, but best in the aortic area where it is exceedingly

loud. From time to time the murmur disappears for one beat, the apex beat becomes weak and the radial pulse vanishes, the murmur corresponding to the next beat being very loud, and the apex beat very strong. The radial sphygmogram is the typical trace of aortic stenosis, with flat top and sloping up-stroke. The heart is slow. The radial pulse shows fair volume, but to appreciate this the artery must be in a particular position as the tension is low and the arm fat. The physical examination reveals nothing abnormal elsewhere. The flow in the hands were measured Jan. 31, 1913. Hands in bath at 2:41½ p. m.; in calorimeters at 2:52½, and out of calorimeters at 3:06. He sat during the whole examination with the injured left foot lying on a stool, the left leg horizontal and the right foot down. (Table 21.)

Cooling of calorimeters in twenty-one minutes, 0.36 degree for R. and L. Volume of right hand 458 c.c., of left 452 c.c. He is mostly right handed, he says. Water equivalent of calorimeters with contents, R., 3,461; L., 3,456. Blood-pressure, left arm, systolic 105 (palpation), 103 (stethoscope), 91 (sound sharply diminished), 87 (sound gone). Rectal temperature 36.6 C.

The last cases to be described are two of mitral stenosis and insufficiency, one with good (Mary S.) and the other with only slightly broken compensation (Frank W.) at the time of the examination. In the first case, a woman, 26 years old, the flow works out at 15.74 grams per 100 c.c. per minute for the right hand and 14.26 grams for the left. Even taking into account the relatively high room temperature (26.7 C.) and the youth of the patient, these flows are, if anything, above the normal.

Fig. 3.—Tracings from Mary S.

When the blood-flow in the hands is not below the normal we are usually justified in assuming that the output of the ventricles is not less than normal. The large flow in this patient agrees well with the obviously exaggerated force of the ventricular stroke, and the absence of venous engorgement, of edema and of cyanosis. It confirms the conclusion that the increased respiratory rate at the time of the examination is a nervous affair not depending on deficient exchange in the lungs. Specimens of polygraph tracings are given in Figure 3. The *a* wave is prominent.

Mrs. Mary S., married woman, aged 26, a Hebrew, was admitted to the City Hospital May 29, 1913, complaining of pain in the chest, weakness and shortness of breath. Seven months before admission she had an attack of acute articular rheumatism and during this attack she began to have pain in her breast. Since then there have been several short periods during which she got short of breath. There seems, however, to be a strong neurotic element in the case, and she does not look particularly ill nor does she show any marked dyspnea. She is fat and well-nourished. No cyanosis, and little, if any, edema of the legs. No venous engorgement, no pulsation in the neck. The thorax is resonant throughout. The breathing everywhere vesicular. The apex beat is in the fifth interspace below the nipple; upper border of cardiac dulness in third interspace, right border, at right sternal margin; left border two finger's breadth out. Auscultation showed the signs of mitral stenosis and insufficiency. The

rather rapid breathing appears to depend largely on nervous excitement. It is certainly not associated with any signs of deficient oxygen intake. The lips are red and she is even more comfortable lying down with her head on a single pillow than sitting up.

The blood-flow in the hands was estimated June 13, 1913. The day was rather warm (external temperature about 80 F). Hands in bath at 2:28 p. m., in calorimeters at 2:40 and out of calorimeters at 2:52. Mouth temperature (two observations) 36.6 C. Pulse (sitting) 84.

TABLE 22.—CALORIMETRIC MEASUREMENTS IN CASE OF MARY S.

Time	Right	Left	Room Temp. C.	Time	Right	Left	Temp. Room C.
2.39	31.51	31.52		2.47	31.86	31.81	26.7
2.41	31.55	31.57	26.4	2.48	31.90	31.84	
2.42	31.60	31.60		2.49	31.93	31.87	26.8
2.43	31.65	31.64		2.50	31.97	31.90	
2.44	31.71	31.68	26.5	2.51	32.01	31.94	
2.45	31.76	31.73		2.52	32.06	31.99	26.8
2.46	31.80	31.76		3.07	31.89	31.83	

Cooling of calorimeters in fifteen minutes, R., 0.17 degree; L., 0.16 degree. Volume of right hand 280 c.c., of left 259 c.c. She is right-handed. Blood-pressure, left arm, systolic 114, 95 (second phase), 62 (fourth phase), which was with practically no further fall in the pressure succeeded by the fifth phase (cessation of sound). Water equivalent of calorimeters with contents, R., 3,319; L., 3,302.

The last case is that of Frank W., a man 34 years old, with mitral stenosis and insufficiency with somewhat broken compensation, which was rapidly restored. Three examinations of the hand flow were made. The first on the day of admission, before any treatment had been given, showed a flow of 15.20 grams per 100 c.c. per minute for the right hand and 15.29 grams for the left, with the unusually high room temperature of 30.9 C. At the second examination, three days later, when the patient's clinical condition had improved, the flows were 11.95 grams for the right and 11.55 grams for the left hand, with the lower but still high room temperature of 28.5 C. The third examination, made six days after the second, with a room temperature (30.8 C.) the same as that during the first examination, showed a flow in the right hand of 11.44 grams per 100 c.c. per minute, and a flow in the left of 11.20 grams. The patient was feeling well. The re-establishment of compensation in this case was associated, then, with a diminution in the hand flow of 25 per cent. This diminution in the hand flow coincided with a reduction in the pulse-rate, quieter action of the heart and an improvement of the patient's clinical condition. It must be noted that the loss of compensation when the patient was admitted was not great, that his heart is permanently large and that there are no signs that the myocardium is, as yet, greatly

impaired. The irregularity of the heart noted in the protocol is associated with premature auricular beats (Fig. 4). This condition is not included among the graver cardiac irregularities by those who have specially studied the clinical disorders of the mechanism of the heart beat. Loss of cardiac, or to speak more accurately, of cardiovascular compensation as expressed in an abnormal distribution of the blood, need not be always associated with a diminished flow through the periphery. Doubtless the abnormal distribution of the blood, resulting in engorgement of the venous side of the vascular system and a rise of venous pressure, is harmful, and is going eventually to cripple the machine. Yet it is not difficult to see that for a while the rate of the blood-flow through a part like a hand, far from being diminished, may be increased provided the driving power of the left ventricle, the aortic valves being intact, does not fail. For the rise of venous and therefore of capillary pressure may, up to a certain point, by increasing the caliber of the small vessels, favor the flow of blood. In order that this may result in increased flow through the part it is obvious that the arterial pressure must also be increased so that

Fig. 4.—Tracings from Frank W. The left hand portion of the figure shows simultaneous tracings from the jugular bulb (upper trace) and the radial artery (lower trace). In the right hand portion of the figure simultaneous tracings from the apex (the large elevations) and the radial are given.

a sufficient slope may be maintained. And as a matter of fact, in Frank W., both the systolic and diastolic pressures in the brachial were materially higher at the first than at the second and third examinations. Corroborative evidence for the conclusion that some loss of compensation when the myocardium is not seriously affected is not incompatible for a time with a fully normal hand flow, is found in the observations on Mary S. and P. B., already referred to.

Frank W., an electrician aged 34, was admitted Feb. 9, 1912, to Lakeside Hospital. He says he never had rheumatism. His illness began six weeks before admission with pain all over, especially in the lumbar region and then in the epigastrium. In about a week shortness of breath came on and palpitation of the heart and he became blue in the face. A few days before admission edema was observed in the lower eyelids and the feet. The physical examination showed distended jugular veins with a systolic pulse. Heart: Apex beat in anterior axillary line in fifth interspace. Here there is a systolic impulse and thrill followed by a diastolic impact. A systolic impulse over the *conus arteriosus*. Left border of cardiac dulness is beyond the anterior axillary line, 18 cm. from the mid-sternal line, and the right border 9 cm. to right of midsternal line. The

auscultatory signs are those of mitral stenosis and insufficiency. The pulse is irregular and arrhythmic. Systolic blood-pressure 145, diastolic 108. Systolic pulsation of the liver. Liver dullness extends to umbilicus in right nipple line. Some edema of lower part of the back. Legs and feet edematous. When dependent, hands and feet are cyanotic. No clubbing of fingers. After a few days' rest the most prominent murmur became the presystolic rumble, heard best at the extreme apex, and indistinct elsewhere. The systolic sound was followed by a short blowing murmur, heard best to right of apex. The size and shape of the heart did not change materially. The edema rapidly disappeared. On March 6 the patient walked out of the hospital feeling well and with good cardiac compensation. The pulse rate which was 160 on admission gradually fell, and varied from 70 to 80 at the time of discharge. The minimum was 58. He broke down again after a time and was admitted to the City Hospital with the same symptoms and physical signs and was discharged improved. He was readmitted June 16, 1913. He worked for three days before he came in but was losing appetite and becoming short of breath. Some edema of the feet and ankles, but not a great deal. Slight cyanosis and dyspnea only on exertion. The point of the nose is red. The boundaries of the cardiac dullness are practically the same as those given in the physical examination at Lakeside Hospital. Pulsation in the neck veins.

The hand flow was measured June 16, 1913, a few hours after his admission and before he had had any drug treatment. The day was very warm. Pulse (sitting) 72, irregular. Hands in bath at 3:11½ p. m.; in calorimeters at 3:22¼, and out of calorimeters at 3:33.

TABLE 23.—CALORIMETRIC MEASUREMENTS IN CASE OF FRANK W.

Time	Right	Left	Room Temp. C.	Time	Right	Left	Room Temp. C.
3.22	31.49	31.49		3.29	32.09	32.08	
3.23	31.56	31.55	30.7	3.30	32.18	32.17	
3.24	31.64	31.64		3.31	32.27	32.255	
3.25	31.72	31.72	30.8	3.32	32.355	32.34	31.0
3.26	31.82	31.82		3.33	32.44	32.43	
3.27	31.90	31.90	30.9	3.49	32.37	32.36	
3.28	31.995	31.99					

Cooling of calorimeters in sixteen minutes, 0.07 degrees.

Volume of right hand in calorimeter 456 c.c., of left hand 452 c.c. Water equivalent of calorimeter and contents, R., 3,460; L., 3,456. Blood-pressure in left brachial 133 (systolic), 103 (change in sound). Another observation 133 (systolic), 133 (change in sound). The pulse was slow and the sound diminished very rapidly to extinction after the change was heard. Rectal temperature 37.55 C.

Second examination June 19, 1913. He says he feels better. He has been receiving digitalis. There is no edema of the ankles, no dyspnea, no distention of the neck veins, no redness of the nose. The day was hot, but not so hot as that of the first examination. Hands in bath at 1:55 p. m.; in calorimeters at 2:10¼, and out of calorimeters at 2:24.

Cooling of calorimeters in twenty-three minutes, R., 0.18 degree; L., 0.16 degree. Radial pulse (sitting) 51. Volume of right hand in calorimeter 450 c.c., of left 439 c.c. Water equivalent of calorimeters with contents, R., 3,455;

L., 3446. Rectal temperature 37.75 C. Blood-pressure left arm, systolic 104, 84 (change in sound); the sound disappeared very soon after this reading. Another observation, systolic 108, 88 (change in sound followed almost at once by its complete disappearance). Pulse (sitting) 60, counted with stethoscope over apex beat. There is still arrhythmia. On holding his breath the heart-rate suddenly increased considerably.

TABLE 23.—CALORIMETRIC MEASUREMENTS ON SECOND EXAMINATION OF FRANK W.

Time	Right	Left	Room Temp. C.	Time	Right	Left	Room Temp. C.
2.10	31.56	31.47		2.18	32.09	32.02	28.3
2.11	31.61	31.53		2.19	32.165	32.08	
2.12	31.685	31.60	28.7	2.20	32.215	32.135	28.4
2.13	31.76	31.68		2.21	32.295	32.20	
2.14	31.83	31.745	28.6	2.22	32.33	32.255	28.4
2.15	31.895	31.82		2.23	32.39	32.305	
2.16	31.97	31.89	28.5	2.24	32.465	32.36	
2.17	32.02	31.955		2.47	32.285	32.20	

Third examination, June 25, 1913. He feels well. Appetite fair, but does not sleep well. Since last examination the pulse-rate has varied between 60 and 72. The respiratory rate has rarely exceeded 20. The temperature has not exceeded 98.4 F. The day was very hot, about the same as the day of the first examination. He says his heart thumps much less than when he first came in although it gives an occasional thump. Hands in bath at 1:53 p. m.; in calorimeters at 2:04¼, and out of calorimeters at 2:19.

TABLE 24.—CALORIMETRIC MEASUREMENTS ON THIRD EXAMINATION OF FRANK W.

Time	Right	Left	Room Temp. C.	Time	Right	Left	Room Temp. C.
2.04	31.77	31.73		2.13	32.32	32.26	30.85
2.05	31.78	31.74		2.14	32.39	32.33	
2.06	31.83	31.80	30.6	2.15	32.44	32.38	
2.07	31.89	31.85		2.16	32.50	32.44	30.85
2.08	31.975	31.935	30.8	2.17	32.565	32.49	
2.09	32.04	31.995		2.18	32.62	32.555	30.85
2.10	32.12	32.06	30.85	2.19	32.70	32.63	
2.11	32.19	32.14		2.35	32.62	32.555	
2.12	32.26	32.20					

Cooling of calorimeters in sixteen minutes, R., 0.08 degree; L., 0.075 degree. Volume of right hand in calorimeter 443 c.c., of left 432 c.c. Water equivalent of calorimeters with contents, R., 3,449; L., 3,440. Blood-pressure, left arm, systolic 104, 82 (change of sound followed very soon by its disappearance). Another observation. Systolic 109, 86 (change of sound). Pulse (sitting) 60, counted with stethoscope over apex. Respiration (sitting) 18. Rectal temperature 38.0 C.

SUMMARY

1. The smallest flows observed in the series of cases dealt with in this paper have been associated with marked irregularity of the heart, indicating involvement of the mechanism of the heart beat. In some of these cases the cardiac irregularity was associated with valvular lesions, but in those with the very poorest hand flows there was no evidence that the valves were involved.

2. Three cases were diagnosed as auricular fibrillation, all associated with mitral stenosis (and insufficiency). While the flow in all was below the normal it was decidedly greater in one of the patients than in the other two, although he was much the oldest of the three. This corresponded well with his clinical condition, which was good at the time of the examination, better than that of the other two patients. The ventricle in spite of its irregular action was on the whole beating fairly strongly, and was obviously delivering on the average of its beats, a fair amount of blood to the periphery.

3. In a case of mitral stenosis and insufficiency and tricuspid insufficiency, in which the heart showed marked arrhythmia in part associated with the respiration (sinus arrhythmia) the hand flow was also much below the normal.

4. Abnormally small flows were seen in a number of cases diagnosed as myocarditis with regular action of the heart, but feeble heart sounds not accompanied by murmurs.

5. In some cases diagnosed as myocarditis without valvular lesion, in which a serious clinical condition had previously existed, very fair hand flows, not much inferior to the normal, were found when the clinical condition had markedly improved.

6. In several cases of mitral insufficiency with good compensation and no signs of impairment of the myocardium, the hand flow was found to be quite up to the normal amount. In one case of mitral stenosis and insufficiency the flow appeared to be even somewhat greater than normal as if the big, well-compensated heart were discharging more than the usual quantity of blood.

7. In a case of mitral insufficiency with broken compensation, venous engorgement and edema, but no dyspnea or cyanosis, the hand flow was quite up to normal. In another case with the same lesions where there was little edema, no shortness of breath except on exertion, and none at the time of examination, the flow was also little if at all inferior to the normal flow, and the re-establishment of compensation was associated with an actual moderate diminution in the hand flow.

8. In a case of mitral insufficiency with physical signs of pericardial effusion a flow not much inferior to the normal was found after the patient had been some time in the hospital and had regained compensation, and when the heart dulness had decidedly diminished. When the patient, two days after leaving the hospital, broke down again and

TABLE 25.—DATA CONCERNING

Case	Age	Date	Pulse-Rate	Blood-Pressure	Temperature (C.) of				Volume of Hands in c.c.	
					Room	Art. Blood	Calorimeters		Right	Left
							Right	Left		
Jacob W..	59	5/ 1/11	100	190, (110)	25.0	37.1	29.92	29.93	508	486
					25.1		30.16		...	...
					25.1		30.38		...	...
James M..	60	6/27/12	87	152, 72	23.0	37.15	30.93	30.93	368	356
N. P.	27	5/12/13	70	95	23.8	36.3	31.44	31.33	298	276
Jacob S. .	44	4/25/11	104		22.2	36.25	29.69	29.62	390	378
					22.6			29.76	...	
					22.7			29.78	...	
					22.7			29.83	...	
T. F. K....	57	5/19/13	120	112, 58	24.0	36.85	31.76	31.75	444	458
Peter B....	59	4/21/11	80		24.5	36.45	30.17	30.14	360	320
					24.3		30.50		...	
					24.3		30.79		...	
Peter B....		5/17/11	84		26.6	36.5	30.63	30.43	410†	394†
					26.8		30.80		...	
					26.8		31.12		...	
					26.7		31.46		...	
John T. . .	28	4/ 9/12	100	122	24.4	37.45	31.36	31.44	450	438
					24.3		31.39		...	
James H. .	63	5/29/13	70	134, 68	23.4	36.1	31.00	30.92	418	396
		6/16/13	57	110, (80)	30.9	37.1	31.69	31.74	408	392
Joseph Z..	16	7/14/11	140		28.1	38.3	30.10	30.10	406	412
					28.2		30.51		...	
					28.3		30.85		...	
Mat K....	42	1/24/11	80	170	22.8	36.6	29.72	29.40	440	435
P. B.	31	5/12/11	100		25.5	36.9	30.76	30.68	510	480
					25.8		31.55		...	
					25.5		32.13		...	
E. M.	14	1/26/11	116		23.4	38.0	30.14	30.07	255	240
Frank C..	17	12/19/10	116		22.4	37.3	30.19	30.10	275	250
Albert H..	31	4/ 9/12	120	160	24.7	38.6	30.95	30.92	447	450
Albert H..		4/19/12	102	130	22.6	37.2	29.65	29.58	475	505
Albert H..			...		23.8			29.65	...	
Roman B..	22	5/29/12	96		22.0	37.45	31.41	31.52	454	439
Roman B..		5/31/12	87		26.3	37.15	31.46	31.60	451	445
Roman B..		6/13/12	...		21.8	36.8	31.28	31.36	451	443
Robert T..	22	1/31/13	70	103, (91), 87	20.3	36.1	31.44	31.48	458	452
Mary S....	26	6/13/13	84	114	26.7	36.6	31.81	31.78	280	259
Frank W..	34	6/16/13	72	133, (108)	30.9	37.05	32.00	31.99	456	452
Frank W..		6/19/13	60	106, (86)	28.5	37.25	32.04	31.95	450	439
Frank W..		6/25/13	60	106, (84)	30.85	37.5	32.24	32.19	443	432

* Good compensation.

† As his hands were short the marks were put higher up on the wrists than at the previous

‡ Badly broken compensation. The results on Stephen F., Mrs. A. and J. S. W. are sum-

BLOOD-FLOW IN AUTHOR'S CASES

Heat Given Off in Gram-Calories			Blood-Flow in Gm. Per Min.		Flow Per 100 c.c. of Hand Per Min.		Notes
Right	Left	In Mins.	Right	Left	Right	Left	
1,050	801	13	12.50	9.55	2.46	1.96	Myocarditis.
1,225		10	19.62		3.86		Left hand in water at 42.9 C.
805		7	19.03		3.75		Left hand in water at 11.5 C.
366	270	10	7.32	4.82	1.99	1.35	Senile heart. Arrhythmia.
283	216	11	5.88	4.39	1.97	1.59	Auricular fibrillation.
545	238	11	7.69	3.63	1.97	0.96	Auricular fibrillation.
....	170	5		5.82		1.54	Right hand in water at 43.3 C.
....	340	4		14.59		3.86	Right hand in water at 43.3 C.
....	340	5		11.76		3.11	Right hand dried and wrapped.
1,138	1,336	9	27.60	32.34	6.21	7.06	Auricular fibrillation.*
1,180	980	9	23.19	19.17	6.44	5.99	Myocarditis.
1,636		10	30.51		8.47		Left hand in water at 42.9 C.
913		7	25.62		7.12		Left hand in water at 8 C.
2,156	1,605	10	40.81	29.38	9.95	7.46	
924		5	36.03		8.79		Five min. just preceding vasomotor test.
1,643		10	33.93		8.28		Left hand in water at 8 C.
1,130		9	27.69		6.75		Left hand in water at 43 C.
516	396	10	9.41	7.32	2.09	1.67	Mitral insufficiency.‡
362		8	8.29		1.84		Left hand in water at 42.7.
154	119	10	3.35	2.55	0.80	0.64	Mitral stenosis, insufficiency; tricusp. insufficiency; arrhythmia.
1,454	1,214	10	29.86	25.16	7.31	6.41	
1,505	1,165	8	25.50	19.75	6.28	4.79	Aortic insufficiency and stenosis.‡
1,402		9	22.22		5.47		Left hand in water at 8 C.
1,197		..	22.31		5.49		Left hand in water at 43 C.
2,159	1,704	11	31.70	23.90	7.20	5.49	Aortic insufficiency and stenosis.
3,608	3,618	10	65.29	64.63	12.80	13.46	Mitral and tricusp. insufficiency.
2,627		11	49.60		9.72		Left hand in water at 8 C.
2,102		9	54.40		10.67		Left hand in water at 43.1 C.
2,988	2,772	13	32.49	29.88	12.74	12.45	Mitral insufficiency.
2,378	2,331	12	30.96	29.97	11.25	11.58	Mitral stenosis and insufficiency.
1,380	1,555	11	18.22	20.45	4.08	4.54	Mitral and tricusp. insufficiency.‡
452	455	7	9.50	9.48	2.00	1.88	
....	472	10		6.94		1.37	Right hand in water at 42.7 C.
3,008	3,970	16	34.58	46.49	7.61	10.59	Mitral insuff. with pericardial effusion.
3,440	4,176	17	39.5	49.18	8.76	11.05	
1,610	2,466	13	24.92	38.74	5.52	8.74	
2,007	2,626	10	47.85	63.15	10.45	13.97	Aortic stenosis.
2,091	1,763	11	44.09	36.94	15.74	14.26	Mitral stenosis and insufficiency.
3,152	3,148	10	69.35	69.11	15.20	15.29	Mitral stenosis and insufficiency.
3,280	3,146	13	53.80	50.73	11.95	11.55	
3,359	3,237	14	50.68	48.38	11.44	11.20	

examination.

marized in a previous paper (Heart, 1911, iii, 84).

returned with some dyspnea, cough and swelling of the legs, the hand flow was found distinctly below its previous value.

9. In two cases of mitral insufficiency with badly broken compensation, with venous engorgement, edema, cyanosis and dyspnea, the hand flow was much below the normal. In one of these cases a striking diminution occurred a few days before death.

10. In a case of apparently pure aortic stenosis with perfect compensation, the hand flow was normal in amount.

11. In a case of aortic stenosis and insufficiency, with mitral insufficiency, while there were no signs of broken compensation, the hand flow was somewhat below the normal.

12. In a case of aortic insufficiency and stenosis, with badly broken compensation (great dyspnea, pulmonary edema), the hand flow was greatly diminished.

While it is obvious that some of the groups of cases are much better represented in the observations than others, as is easily understood, since they had to be collected as chance offered, certain general conclusions which, although of course at present only provisional, are yet entitled to some weight, seem to emerge with sufficient clearness. It is scarcely necessary to say that in some points these conclusions simply confirm what is found by palpation of the radial pulse. But it is by no means true that even the qualitative result of the blood-flow measurement can always be predicted from feeling the pulse.

CONCLUSIONS

1. The hand flow is more apt to be markedly deficient in cases in which there is evidence of serious impairment of the myocardium, or more comprehensively and probably more accurately, of the mechanisms on which the force and rhythm of the heart's action depend, even when valvular lesions are absent, than in cases in which gross valvular lesions exist while the heart's action is strong and orderly. This holds even when in valvular diseases clinical signs of loss of compensation are present. In aortic insufficiency the hand flow appears to be less easily maintained at the normal level and more readily to fall below it when compensation is lost than in mitral insufficiency.

2. Even where there is considerable venous engorgement in cardiac cases the flow in the hands may be little if at all diminished, provided that the myocardium is not impaired. When in a case of broken compensation the flow in the hand remains good the reestablishment of compensation may be associated with a moderate decline.

3. When in a cardiac case with broken compensation the hand flow is normal or not much diminished, the indication is that the myocardium has not as yet suffered serious impairment. A decided diminution in the course of the case may be significant of impending failure of the myocardium.