The Archives of Internal Medicine

Vol. XIV

OCTOBER, 1914

No. 4

THE RELATION OF METHODS OF DISPOSAL OF SEWAGE TO THE SPREAD OF PELLAGRA *

J. F. SILER, M.D.
Capt. Med. Corps, U. S. Army.

P. E. GARRISON, M.D.
Passed Asst. Surg., U. S. Navy.

AND

W. J. MacNEAL, M.D.
NEW YORK

INTRODUCTION

In analyzing the data collected in 1912 we were struck by the inequalities in the distribution of pellagra in the townships of the county in which our studies were undertaken. This rate ranged from no cases to 71 cases per ten thousand of population. A further analysis, based on the distribution of the population in rural, urban and mill-village communities showed that in rural communities the incidence rate was 16 per ten thousand of population; in the city of Spartanburg, exclusive of the mill-village section, it was 29 per ten thousand, while in the mill-village communities it averaged 104 per ten thousand.

One of the features of our epidemiological work in 1912 was the investigation of methods of disposal of human wastes. In certain sections of the city of Spartanburg a water-carriage system was in use. In the rural districts the white farmers usually had the unscreened surface privy. Privies were but seldom found on the premises of negro tenants on the farms. In the cotton-mill villages, about twenty-eight in number, in which pellagra prevailed much more extensively than elsewhere, two types of privy were in use. The usual type was an unscreened wooden outhouse where the excreta were deposited on the ground. In a few mill-villages the privy type was an unscreened wooden outhouse provided with metal containers for the reception of fecal matter. In some of the villages scavengers cleaned the privies at weekly intervals; in others at less frequent inter-

* Submitted for publication, April 17, 1914.

* This paper forms a part of the Second Progress Report of the Thompson-McFadden Pellagra Commission of the New York Post-Graduate Medical School and Hospital. It was presented in part before Section K of the American Association of Science, at Atlanta, Ga., Jan. 2, 1914.

vals or irregularly. These villages were laid out in streets, the dwelling-houses were but short distances apart and the privies necessarily in close proximity to the dwelling-houses. No effort was made to screen these privies; flies and other insects had free access to the collections of excreta.

In a series of 243 cases of pellagra investigated in 1912, the following methods of disposal were found to be in use.

In 5 per cent. (11 of the cases) a modern water-carriage system was used.

In 20 per cent. (51 of the cases) the unscreened wooden outhouses were equipped with metal receptacles.

In 60 per cent. (152 of the cases) the privy was an unscreened wooden outhouse, the feces being deposited on the ground.

In 12 per cent. (29 of the cases) there were no privies.

The average sanitary index of the privies used by these 243 pellagrins was 16 per cent., computed in accordance with the methods adopted by the Rockefeller Sanitary Commission. The existing sanitary conditions were summarized as follows:

The most insanitary condition found in the county is the absence of properly constructed privies. Outside of a part of the city of Spartanburg, which is supplied by a water-carriage sewage system, there is no effective provision in the county for the proper disposal of human excreta. A second striking insanitary condition is the almost complete absence of effective screening of dwellings. These two conditions present a situation highly favorable to the transmission of disease organisms eliminated in the excreta, both by direct contamination of food and persons and by insects. This situation is naturally aggravated in the mill villages and small towns by the greater congestion of houses.

As a result of these, as well as other observations, the members of the commission were impressed with the possibility that insanitary methods of disposal of human wastes might be an important factor in the spread of pellagra.

In the course of our investigations during 1913 it was determined that this feature should receive special attention. The principal point which we endeavored to ascertain was whether the endemic foci of the disease were more common in those communities and in those sections of cities in which the methods of disposal of human wastes were insanitary. In order to ascertain these facts we made a careful study of the exact distribution of the cases of pellagra, including a history of its distribution in the past. By personal visits to pellagrins and their relatives it has been possible for us to secure definite information as to the exact residence of pellagrins. We have endeavored to locate precisely all houses occupied by them from the inception of the disease up to the present time. The information relative to residence, as well as to other pertinent facts, has been transferred to our pellagra "house card," a copy of which we show. These "house cards"

have been filed by locality and from them we have constructed spot-maps showing the exact location of the cases of pellagra of which we have record.

HOUSE CARD

House No. 190. Township, *Spartanburg*. *Whitney*.
 Location, *Whitney Mills No. 101*.
 Pellagrous History Date, *July, 1913*.
 No. First Erythema (Place, Date)
 86. Here from *Jan. 22, 1912, to early in 1913*. *July, 1911. Clifton No. 2.*
H. No. 191.
 87. Here from *Jan. 22, 1912, to early in 1913*. *June, 1912. Here.*
 88. Here from *Jan. 22, 1912, to early in 1913*. *June, 1912. Here.*
 89. Here from *Jan. 22, 1912, to early in 1913*. *June, 1912. Here.*
 Notes—86 = *mother*; 87 = *son*; 88 = *daughter*; 89 = *daughter*.
 Apparent house of origin for Cases Nos. 87, 88, 89.
 House: Construction—*Frame*. Screened, *no*.
 Sanitary Condition—Good, poor, *fair*.
 Water Supply—Well: *dug, with bucket. Deep.* Surface contamination, *no*.
 Disposal of Excreta—Privy type: A, B, C, D, E, F. Distance from house,
twenty yards.
 Method of Ultimate Disposal—*Scavenger*.
 General Sanitary Condition—*Poor*.

In all the mill-villages in Spartanburg County unscreened surface privies were in use and, in order that possible contrasts in the prevalence of pellagra might be shown, for the purpose of this study, it was necessary to extend our investigations to industrial communities elsewhere in the state in which water-carriage systems of sewage disposal had been installed. Through the state health authorities we learned that four cotton-mill villages in the state had adopted such methods of disposal. In two of these villages, one situated in Greenville, S. C., and the other in Union, S. C., the water-carriage sewer systems had been installed less than a year. The third was the Newry mill-village in Oconee County in the western part of the state, in which a water-carriage system of sewage disposal was installed about twenty years ago. The fourth village was the Republic mill-village in Chester County, situated in the northern part of the state, in which a water-carriage system was installed about three years ago. We have investigated the sanitary condition and prevalence of pellagra in the Newry mill-village and the Republic mill-village. As the water-carriage system of disposal had been in use in the other two villages for only a few months, their investigation was left to the future.

This study has enabled us, first, to compare the prevalence of pellagra in sewered and unsewered portions of the city of Spartanburg, and second, to make a similar comparison of two entirely sewered villages with a large number of villages where unscreened surface privies were in use.

THE SPREAD OF PELLAGRA IN THE CITY OF SPARTANBURG AND ITS
RELATION TO VARIOUS METHODS OF DISPOSAL
OF HUMAN WASTES

We have record of a total of 268 cases of pellagra in the city of Spartanburg (204 white and 64 colored). This includes not only the pellagrins living there in 1913 but also all known cases which have occurred. Of these 268 cases, 168 are known to have originated in the city.

Fig. 1.—Map of Spartanburg, S. C., showing plan of city.

Figure 1 is a reproduction of the plan of the city, the black portion in the center indicating the location of the business section. It will be noted that there are three mill-villages within and two just beyond the city limits. The largest cotton-mill is situated in the north-western quadrant, the next largest in the northeastern, and the smallest in the southwestern quadrant.

Figure 2 shows the geographical distribution of pellagra in the city of Spartanburg together with the distribution of the sewer mains. It is clearly evident from this map that practically all the active foci of the disease were in the sections of the city in which unscreened surface privies were used. Two active foci of the disease existed in the southeastern quadrant of the city. One of these foci was in a sewer district but none of the pellagrous houses had sewer connections. The other was in a negro settlement and all the privies in this vicinity were of the unscreened surface type.

Fig. 2.—Map of Spartanburg, S. C., showing distribution of cases of pellagra in the city and distribution of sewer mains. Note that 168 of these cases originated in the city and that the largest number of cases occurred in the northwestern quadrant.

Of the total 268 cases of which we have record in the city of Spartanburg, we have investigated methods of disposal of excreta in 241 instances. In 230 cases, or 95 per cent. of the total investigated, the method of sewage disposal was the unscreened surface privy. In eleven instances, or 5 per cent. of the total, a water-carriage system of disposal was in use. A detailed examination of the records of these eleven cases has shown the following facts:

Patient 12 was a white woman, aged 40, married, whose residence had not changed since 1907. She contracted pellagra in 1911 and had mild recurrences in 1912 and 1913. She lived on one of the better residential streets; her circumstances were comfortable, and her only occupation was housework. The house was not screened. She gave no history of association with pellagrins, except visits to two persons suffering with pellagra. Our records, however, show that in a house immediately in rear of hers, Patient 496, a colored female, died of pellagra, July 12, 1910, and that during 1911, Patient 379, a colored female, lived directly across the street from her. The houses of these two colored pellagrins had the usual unscreened surface privies.

Patient 19, a white man, aged 20, contracted pellagra in the spring of 1911. His residence remained the same from 1905 to the fall of 1912. His home was in the best residential section of the city and the house was effectively screened. He was a college student and his circumstances were comfortable. His general health was good until the summer of 1909, when he contracted malaria in a saw-mill camp in South Carolina, near Augusta, Ga. Since that time his health has not been good. In 1910 he again spent five months (January to June) in a lumber-camp near the Georgia state line in South Carolina. This patient had severe recurrences during 1912 and 1913. No cases of pellagra have been located in the vicinity of his home in Spartanburg and there is no history of association with pellagrins.

Patient 24, a white man, aged 61, contracted pellagra in May, 1912. His residence had remained the same for several years. His home was situated in a good residential section but was not screened. He had been engaged in clerical work since 1881; his health had been good, and his circumstances appeared to be comfortable. He gave no history of association with pellagrins and we were not able to locate any cases in the vicinity of his residence. The disease recurred in 1913 and he died in the fall of 1913.

Patient 40, a white woman, aged 63, contracted pellagra in February, 1912. She had not changed residence for two years except for prolonged visits to relatives in South Carolina. Her home was situated in the business section of the city and, though it was screened, the screening was not effective. Two visits were made to this home by members of the commission. We found the screens raised, swarms of flies present and the general sanitary conditions very unsatisfactory. The patient's economic circumstances were poor and her occupation was housework. Patient 342, a white woman, who was a chronic pellagrin, lived with her during August, September and October, 1911. This patient (No. 40) died of pellagra, Aug. 10, 1912, having first shown symptoms in February, 1912.

Patient 57, a white woman, aged 43, first showed symptoms of pellagra on June 27, 1912. She lived in the best residential section of the city and had been living there since 1910. Her home was effectively screened and her husband well-to-do. She apparently had had some gastric disorder for a number of years. Her only occupation was the supervision of her housework. There was no history of association with pellagrins and no pellagrins lived in the immediate vicinity. There was no definite recurrence during 1913.

Patient 59, a white woman, aged 38, contracted pellagra in March, 1909, three months after confinement. Her home was in one of the best residential sections of the city and she had been living there since 1907. Her house was screened in part and her circumstances were those of comfort. There was no history of pellagra in her family and she gave no history of association with pellagrins. She had slight recurrences in 1910, 1911, 1912 and 1913.

Patient 208, a white man, aged 59, developed pellagra in December, 1909, and severe symptoms persisted until September, 1910. He had completely recovered from the attack in November, 1910, since which time his health had been good. The house of origin for this case was situated in one of the better residential sections and he lived there from 1907 to early in 1912. His circumstances were those of comfort and his health prior to the development of

pellagra had been good. No pellagrins have been located as living in the immediate vicinity of the house of origin for this case. For several years it had been the patient's custom to make visits from time to time to relatives in certain country districts in the county. One of his sisters died of pellagra in 1907 and another died of pellagra in 1910. Both were living in Spartanburg county and were well-to-do. His health has been excellent since November, 1910.

Patient 233, a white woman, married, contracted pellagra in May, 1910, in a house located in the best residential section of the city. She lived in this house from August, 1908, to March, 1911, and it was not screened. Her husband was well-to-do and she had enjoyed good health prior to the development of pellagra. Symptoms recurred in 1911, but there was no recurrence in 1912 and 1913. We have no record of pellagrins living in her immediate vicinity during 1909 and 1910. Her father-in-law (Case 321) died of pellagra in September, 1910, at his home in an unsewered portion of the city of Spartanburg and she had visited him there frequently during his illness.

Patient 426, a white woman, single, died of pellagra in 1902. Her father was well-to-do. Prior to the development of pellagra her health is said to have been good. She had been living in the same house for many years. As the condition was not recognized as pellagra until several years after her death the history is very incomplete. Evidence of association with other cases of pellagra is wholly lacking.

Patient 541, a white woman, aged 21, married, first showed erythema in April, 1913, though she gave history of ill health since November, 1912. She had lived since 1909 in a good residential section and the house was screened in part. Her circumstances were those of comfort and her health had been good prior to November, 1912. The nearest case of pellagra was a white man, Patient 5, who contracted pellagra in 1911 and died of it in 1912 in a house directly across the street. His house was provided with an unscreened surface privy. This woman, Patient 541, made a visit of three weeks' duration, during the summer of 1912, to relatives in a southern state on the gulf coast where surface privies were in use. There was no history of association with pellagrins.

Patient 755, a white man, aged 32, druggist, contracted pellagra in the spring of 1913 and died of it in August, 1913. During 1913 he lived in the best residential section of the city and was well-to-do. The history of this case is very incomplete and we have no data concerning association with pellagrins.

Of the eleven patients, three (Cases 12, 233 and 541) gave a history of living or visiting in the same house or next door to pellagrins using surface privies; another (Case 40) lived under very insanitary conditions in the same house with a pellagrin; and a sixth (Case 208) had two sisters who were pellagrins, one dying before he developed the disease and the other dying the following year. Two, Patients 24 and 755, were men engaged in occupations bringing them constantly into contact with the general public. In the other four cases, (Nos. 19, 57, 59 and 426) we have been unable to learn of any relationship whatever to antecedent pellagra.

THE SPREAD OF PELLAGRA IN UNSEWERED MILL-VILLAGES IN SPARTANBURG COUNTY

Our records show that pellagra occurred and has originated in all the mill-villages in Spartanburg County. In all these villages unscreened privies were in use. In some, the fecal material was col-

lected in pails; in the majority, however, it was deposited on the ground. Many of these mills were situated but a few miles distant from one another and for a number of reasons there was a tendency for certain elements of their population to move, with great frequency, from one mill to another. This element of the population—generally termed “floating”—varied from about 15 per cent. to even more than 33 per cent. Cases of pellagra commonly occurred in this floating element and frequently it was necessary for us to trace pellagrins from mill-village to mill-village before finally locating them.

Six of these villages were selected for intensive studies and by house-to-house visitations all existing cases of pellagra were accurately located. The incidence rate for 1912 and 1913 was higher in

Fig. 3.—Distribution of pellagra in mill-village “W,” Spartanburg County. Unscreened surface privies are shown as small blocks in the rear of dwelling-houses. Note that twenty cases of pellagra have originated in this village.

some villages than in others, but incident cases occurred in every village during 1912 and 1913. It was observed that in several of these villages incident cases of pellagra would be quite numerous one season and relatively few the next, or *vice versa*.

Spot-maps have been constructed of these mill-villages showing the distribution of pellagra. Figure 3 is a reproduction of the plan of village “W” showing the distribution of the disease, and this is a fair example of conditions in all these villages. It will be noted from this map that at least twenty cases of pellagra have originated in this village.

HISTORY AND SPREAD OF PELLAGRA IN MILL-VILLAGES IN WHICH WATER-CARRIAGE SYSTEMS OF DISPOSAL OF SEWAGE
HAVE BEEN INSTALLED

Newry Mill-Village

This village is located in Oconee County in the northwestern part of South Carolina. It was built about twenty years ago, its first president being Captain Courtenay of Charleston. Captain Courtenay installed a water-carriage system of sewage disposal before the mill-village dwellings were occupied and this system has been continuously in use to the present time. The closets have been grouped (from four to eight in each group) in detached houses convenient to a group of dwelling-houses, rather than as individual closets in each dwelling-house. One of these privies has been assigned to each dwelling and the tenant held responsible for its proper care. The annual cost for repairs and upkeep has amounted to but little. The total population was 750. The operatives employed in the mill were recruited from the same sections and from the same elements of the population, as were operatives in mills elsewhere in the state. They were accustomed to eat the same foods as were eaten by the inhabitants of other mill-villages throughout the state. The company store was selling about a bushel of shipped (western) corn-meal to each family per month. The only respect in which their dietary habits possibly differed from those of some of the mill-villages in the state, lay in the fact that fresh meat could be obtained throughout the year from a local market. In some of the mill-villages in Spartanburg County fresh meat was not on sale in the village during the hot summer months. In others, however, where pellagra prevailed extensively, fresh meat could be obtained throughout the year from the mill-village market.

The Newry village is almost surrounded by permanent streams, and the eggs and larvae of *Simulium* were observed in abundance. The only demonstrable difference between this mill-village and (with three exceptions) those elsewhere in the state lay in the fact that a water-carriage system of sewage disposal was used there, while the method employed elsewhere was the unscreened surface or pail privy. With these facts in mind it is of interest to study the history of pellagra in this village.

In August, 1911, a woman showing active pellagrous skin lesions came into the village from Georgia. She departed the same day on the advice of the local physician and has since died.

In the spring of 1908, a family which had been resident in the village for several years, purchased a home near the railway (at Courtenay Station) about one mile away. They lived there until December, 1911. A woman with pellagra resided next door during the period of their residence at Courtenay Station. In June, 1909, the mother (Case 2064) contracted pellagra. In December, 1911,

they returned to the Newry mill-village and were still there in October, 1913. The family consisted of six adults and one child. No other case of pellagra was observed in the family and this one case had recurrences in 1910, 1911, 1912 and 1913.

Another woman (Case 2060), one of the four patients now resident in the village, gave a rather indefinite history of pellagrous symptoms as far back as 1910. The family had been extremely migratory and it was difficult to determine with certainty exactly where and when the disease developed. She first lived in the village for two months in the winter of 1910 and it is possible that she was at that time pellagrous. The first positive history of the disease began in the spring of 1912, when a frank erythematous rash appeared. At that time she was living on a farm some ten miles north of the village. She returned to Newry in November, 1912, and lived there until the last of February, 1913. She was then away until June, 1913, when she returned to the village and was still there in September, 1913.

Fig. 4.—Distribution of cases of pellagra in the Newry mill-village having a water-carriage system for disposal of sewage. Note that all patients with one possible exception contracted pellagra before moving into the village.

Patient 2061, who had been a pellagrin since 1910 and had had a recurrence each succeeding year, first moved to the village in November, 1912, and still lived there in September, 1913. Her 10-year-old son (Case 2062) had pellagra, but he likewise developed the disease early in 1912 before coming to the village.

Case 2063, that of a young woman who died of pellagra in Newry, Aug. 17, 1913, offers the only instance of possible origin in this mill-village, inasmuch as she moved there late in the fall of 1912 (exact date not known) and apparently did not have any erythema prior to Aug. 3, 1913, two weeks before death. She was under treatment by the local physician for persistent diarrhea from June 6, 1913, and gave a history of stomatitis, diarrhea and general poor health the previous year, while living at another village in a house within a few doors of at least two other cases of pellagra. It seems highly probable that she was a pellagrin in 1912 before coming to Newry.

The history of pellagra in this village may be summarized by saying that five pellagrous patients have been observed, all of whom, with one possible exception, were suffering with pellagra on arrival in the village. No case of pellagra clearly originating in the village has been found. Figure 4 indicates the distribution of pellagra at the Newry mill-village.

As a control study we investigated the prevalence of pellagra in a mill-village on the outskirts of Seneca, S. C., about four miles distant from Newry mill. The methods of disposal of human wastes in this village were exactly similar to those observed in mill-villages in Spartanburg County (unscreened surface privies). We were shown active cases of pellagra originating in this village. The history of pellagra in this village as given us by the local practitioners indicated that it had prevailed here to much the same extent as in the mill-villages in Spartanburg County.

Oconee County had not so large a number of cotton-mills as Spartanburg County, nor were the mills in such close proximity one to another. For this reason the "floating" or transient population was appreciably smaller.

Republic Mill

This mill is situated at Great Falls, Chester County, in the north central part of South Carolina. It was built in 1910 and was put into operation in June of that year. There were ninety-two houses in the village and its population was between 475 and 500. Before building the village, Mr. R. S. Mebane, president of the mill, made a special study of the problem of the proper sanitation of mill-villages, especially with reference to the water-supply and sewerage. In building the village he constructed a complete sewer system, connected with every house, and placed in each house a kitchen sink and flush closet. The system differed from that at Newry Mill in that the closets were placed in the houses instead of being grouped outside. This mill-village was visited for two days in September, 1913, and as complete a study made of the pellagra situation as was possible in that time. Dr. J. B. McKeown, the resident physician, kindly gave practically the entire two days to assist in the inquiry. Notes were secured of six cases of pellagra concerned in the history of the village during the three years of its existence.

The first patient, a woman, aged 30, had lived in the village only four months. She was at about term with her fourth child. She had no erythema when seen and said she had never noticed any. When seen by us she had a very bad stomatitis and glossitis and had suffered with persistent diarrhea for the past nine months. She had lost considerable weight and had suffered recently with severe burning of feet. While, in the absence of any skin manifestation, it was not considered possible to make an absolutely positive diagnosis, we agreed

with Dr. McKeown in considering hers a very probable case of pellagra. There was no question, however, but that her symptoms antedated her coming to Republic Mill.

A second case, not now in the village, was reported by Dr. McKeown as having moved there in January, 1913. The patient first came to his attention in April, 1913, when she was confined. About three weeks later he was again called and found her with all the symptoms of an active attack of pellagra—diarrhea, nervous symptoms, bad stomatitis, and a symmetrical erythema on the backs of her hands. She left the village the following June. In November we received a letter from Dr. L. S. Hay of Rock Hill, a former physician of the case, saying that she had been a pellagrin under his care the previous year before coming to Republic Mill.

The third patient was a woman who had been a pellagrin since the spring of 1909 and had lived in Republic Mill for the past two years. She had had no active symptoms of the disease during the period of her residence in the village.

The fourth case was that of a boy aged 8, who came to the village in March, 1913. When seen he showed no evidence of the disease except a symmetrical scaliness and pigmentation of the backs of the hands. In the spring of 1913 he had an erythema of hands and feet and an attack of sore mouth. He was seen by Dr. McKeown at that time and the case diagnosed as pellagra. According to his father's statement he had suffered an exactly similar attack, though rather more severe, in March of the previous year (1912) before coming to Republic.

The fifth case was that of a boy, aged 6, who, according to Dr. McKeown's statement, presented a typical erythematous rash of both hands and feet and was suffering from diarrhea and sore mouth and tongue during the summer of 1912. The erythema had appeared in June, 1912, and he had come to the village the preceding January. His mother stated that he had a somewhat similar rash the previous year (1911) before coming to Republic, which she thought was "poison oak." She said that she "broke out" the same way on her hands, face and neck about six years ago and that her mouth got sore. She also stated that her father was said to have died of pellagra. The woman was of a very low order of intelligence and her statements were not considered very reliable. A diagnosis in her case was not possible and the time of appearance of the boy's first symptoms could not be determined positively. The case must be considered as possibly originating at Republic Mill, but there was a fair probability that the boy was pellagrous prior to coming to the village.

The sixth patient, a colored woman, aged 35, did not live in the village proper, but in a cabin about one fourth of a mile distant. This was a typical case, but the patient had been living there only a short while and her symptoms clearly antedated her present residence.

Of the total six cases which could be discovered in the three years history of the village, only one could possibly have arisen there and in that case the evidence seemed to point to probable origin elsewhere. Figure 5 shows the distribution of pellagra in this village. Two of the cases cited above are not shown on this map because the patients lived beyond the limits of the village.

One significant fact in these two villages was the difficulty encountered in finding cases of pellagra even by careful search, whereas in all other villages studied, a few moments' inquiry was all that was necessary to locate many cases of pellagra, well known to the population at large.

The second striking feature, of course, was that, with two doubtful exceptions, each of the eleven patients affected with the disease who could be found in these two water-sewered villages had been pellagrins prior to coming to the respective villages. It appears, therefore, that while cases of pellagra had been introduced into these villages from time to time during their history of twenty and of three years, respectively, no case of the disease could be shown definitely to have originated in these villages and only two could possibly have done so.

With reference to Republic Mill, the point arises whether three years is not perhaps too short a time for pellagra to show itself in a newly constructed village. That this is not true is demonstrated in the case of a mill-village in Spartanburg County which was built in 1911 and went into operation in April, 1912. During the following year five known cases of pellagra moved into the village. Two chil-

Fig. 5.—Distribution of pellagra in the Republic mill-village. Note that with one possible exception all pellagrins contracted the disease elsewhere and moved into the village with pellagra.

dren (Cases 546 and 585) who in July, 1912, moved into the house next door to one of these imported cases, developed typical pellagrous symptoms in May and June, respectively, the following year (1913). There were very clear histories of the absence of any previous manifestations of the disease. A third patient (No. 745), a boy of 7 years, developed pellagra in March, 1913. There was no history of a previous attack. The immediate connection with a previous case of pellagra appeared to be the boy's grandfather, who was a pellagrin and had visited the family during the previous winter. The development of these three cases within little over a year after the newly built village was first occupied would argue strongly that the apparent freedom of Republic mill-village from pellagra cannot be explained

by the assumption that a period of three years is not sufficient time for the disease to manifest itself in a new locality.

Each of the villages here cited is still under observation, and it is our intention to follow them for at least another year, both in order to determine the presence or absence of pellagra and to ascertain, if possible, any factors other than the presence of water-carriage systems of sewage disposal which may differentiate them from those villages showing a high prevalence of pellagra.

THE SPREAD OF PELLAGRA IN UNSEWERED RURAL COMMUNITIES

We have referred to inequalities in the distribution of pellagra in Spartanburg County and have cited the fact that our records showed an incidence rate of sixteen per ten thousand of population in rural communities, against 104 per ten thousand in mill-village communities.

So far as disposal of excreta is concerned the only important difference between these two elements of the population was the fact that unscreened surface privies were always found in mill-village communities, while in rural communities, not infrequently, there was no visible method of disposal. The absence of privies applied more particularly to the colored tenant farmers. Where privies were found in rural communities they were practically always of the unscreened surface type.

The most important contrast between rural and mill-village communities appears to be density and congestion of population. There were isolated instances in rural communities in Spartanburg County where foci of pellagra appeared to exist and we made an extensive study of one such district. In this farming district covering an area 2 miles square, there was a total population of approximately 260 persons, 100 white and 160 colored. We secured records of a total of eleven cases of pellagra occurring in this population since 1906, four cases in the white population and seven cases in the colored. The incidence rate for pellagra in this rural community was 423 per ten thousand of population while the incidence rate for the rural population of the county as a whole was only 16 per ten thousand of population. In this district a history of close association between previously existing and new cases was demonstrable. There was a general tendency for active foci of pellagra in rural communities to occur in townships in which mill-villages were located and the focus mentioned above occurred within about 2 miles of a mill-village community where pellagra was endemic.

The methods of disposal of human wastes in the county in which our studies were undertaken are typical of those in use throughout the southern states where pellagra prevails so extensively. Recent sta-

tistical reports of the Rockefeller Sanitary Commission¹ indicate quite clearly the sanitary conditions existing in the Southern states at the present time. Of 189,510 homes inspected in rural communities it was found that 95,735 had no privy. The sanitary index for the total number investigated was only 5.8 per cent.

THE PREVALENCE OF PELLAGRA IN INSTITUTIONS WITH WATER-CARRIAGE METHODS OF DISPOSAL OF SEWAGE

The prevalence of pellagra in institutions for the insane, having water-carriage systems of disposal of sewage, is of particular importance in the discussion of any possible relationship between the spread of pellagra and methods of disposal of human excreta. We have been unable to obtain any definite information relative to the actual incidence rate of pellagra in asylums in the southern states. The disease is endemic in these states and cases of pellagrous insanity are being admitted to these institutions at frequent intervals. Very frequently symptoms of the disease are observed within a short time after admission and even though there be no suspicion or definite evidence of the disease on admission, it is impossible, from the histories obtainable, to arrive at any definite conclusion as to where the person contracted pellagra. It is a fact, however, that, not infrequently, patients show active symptoms of the disease for the first time, several years after commitment.

So far as we have been able to ascertain, there is probably only one institution for the insane in this country in which a severe epidemic of pellagra has occurred and in which there appeared to be no question that practically all the patients contracted the disease in the institution. We refer to an epidemic of pellagra at the Peoria State Hospital at Peoria, Ill.² Two of the members of this commission took part in an investigation of this epidemic and it was studied most thoroughly by the Pellagra Commission of the State of Illinois. Dr. George A. Zeller, superintendent of this institution, recalls very distinctly that cases of pellagra occurred in the Peoria Hospital as early as 1907 and has records of such cases, though they were not, at that time, recognized as such. In 1909, however, there was a sharp outbreak of pellagra in the institution involving more than 8 per cent. of the inmates (177 cases). Table 1 shows the total number of incident cases for each year from 1909 to 1913 inclusive. All cases for 1909 are tabulated as incident cases because accurate records were not available prior to that time. The mortality rates in

1. Fourth Annual Report, The Rockefeller Sanitary Commission.

2. Report of the Pellagra Commission of the State of Illinois, November, 1911, Springfield, Ill., 1912.

this epidemic were very high, particularly so during the year 1909 (50 per cent.). The severe outbreak of pellagra in 1909, with the sharp decline in succeeding years, is most interesting.

TABLE 1.—INCIDENT CASES OF PELLAGRA AT THE PEORIA STATE HOSPITAL, PEORIA, ILL.

Year	1909	1910	1911	1912	1913	1914 (to March 15)	•
Cases	177	67	14	0	0	0	

The institution had a most excellent water-carriage system of disposal of sewage and it is not possible to attribute the 1909 epidemic

TABLE 2.—PELLAGRA ARISING IN THE WARDS OF THE PEORIA STATE HOSPITAL

Ward	Type of Patient	Habits*	Sex †	Population	No. of Cases of Pellagra	Incidence Per Cent.
Infirmary A	Infirm chronic...	±	♀	60	10	16.6
1 A	Staff house.....	±				
2 A	Chronic	±	♀	50	7	14.0
3 A	Demented	—	♀	62	19	30.6
4 A	Epileptic	+	♀	60	7	11.7
5 A	Acute and chronic	+	♀	52	2	4.0
Infirmary B	Infirm chronic...	±	♂	60	3	5.0
1 B	Acute and chronic	+	♂	52	2	4.0
2 B	Demented	—	♂	52	13	25.0
3 B	Demented	—	♂	62	4	6.4
4 B	Epileptic	+	♂	58	8	13.8
1 C	Demented workers	+	♂	110	8	7.3
2 C	Demented	±	♂	110	9	8.2
3 C	Demented	±	♂	110	8	7.3
4 C	Irritable dements	+	♂	110	7	6.4
5 C	Irritable dements	±	♀	110	14	12.8
6 C	Irritable dements	+	♀	110	7	6.4
7 C	Feeble dements...	±	♀	110	6	5.5
8 C	Infirm old	+	♀	110	23	20.8
South dormitory..	Quiet workers...	+	♀	...	2	...
Tent Col. B.....	Tuberculous	±	♀	57	12	21.0
Tent Col. C.....	Tuberculous	±	♂	57	5	8.8
Supply ward	Workers	+	♂	140	4	2.9
Domestic ward....	Workers	+	♀	64	1	1.5

* The sign + indicates that patients are all cleanly in habits; ± indicates that some are cleanly but many are not; — indicates that all patients are uncleanly. In this column ♂ stands for male and ♀ for female.

to improper methods of disposal of human excreta. It is to be remembered, however, that in many infectious diseases of intestinal origin, dependent mainly for their spread on faulty methods of disposal of human excreta, intimate association and contact may also be important factors. With this in mind, there are some suggestive facts which may be cited in connection with this epidemic.

The wards of this institution were constructed on the pavilion plan. Twelve of the pavilions had capacity for 60 patients. Eight pavilions housed 110 patients each; the supply ward had 140 patients and the domestic ward, 64. In assigning patients to wards an effort was made to distribute them according to cleanliness of habits. Through the courtesy of Dr. H. D. Singer, director of the Illinois Psychopathic Institute, Dr. George A. Zeller, member of the State Board of Administration and Dr. E. Z. Levitin of the Medical Staff of the Peoria State Hospital, we have secured information³ covering the distribution of 181 cases of pellagra arising in the early stage of the epidemic at Peoria. Table 2 shows the distribution of cases in the different wards and each case has been charted in the ward of apparent origin.

It will be noted from this table that, except for Wards 8 C and 3 B, there was a distinctly greater tendency for pellagra to spread in wards housing untidy patients (3 A and 2 B) than in those housing patients of cleanly habits. It is also evident that the patients in 8 C, which shows a high rate of incidence, were of advanced age and infirm,

Omitting the two cases which occurred in the south dormitory, for which population figures are not at hand, there remain 179 pellagrins. The distribution of these cases according to cleanliness of habits is shown in Table 3.

TABLE 3.—DISTRIBUTION OF PELLAGRA IN WARDS OF THE PEORIA STATE HOSPITAL ACCORDING TO CLEANLINESS OF HABITS

Character of Inmates	Population	No. of Pellagrins	Incidence Per Cent.
All cleanly habits	866	69	8.0
Mixed: some cleanly habits and many uncleanly.....	724	74	10.2
All uncleanly habits.....	176	36	20.5
Total	1,766	179	10.1

The incidence rate of pellagra in the untidy wards was 20.5 per cent., two and a half times greater than in wards housing patients with cleanly habits, in which it was 8 per cent.

The food for the inmates was distributed from a central kitchen. One of the early and severe cases in the institution was that of a woman employed in the central kitchen. She had a very severe attack of pellagra during the summer of 1908 while working in the kitchen and her symptoms became so severe as to necessitate her admission to the hospital.

3. Personal communication, March, 1914.

In August, 1909, Dr. Zeller, the superintendent of the institution, decided to establish in the hospital proper, male and female pellagra wards. In the fall of 1909 practically all the pellagrins in the institution were transferred to these wards, a graduate nurse was assigned to each ward and daily notes were made on the cases. During the summer of 1910, the pellagrins were still segregated and as new cases of the disease arose they were transferred to hospital. Dr. E. Z. Levitin of the hospital staff informed us recently (March, 1914)⁴ that segregation was practiced until the fall of 1911, when the patients were so few that the practice was discontinued. He states further that, except for one chronic pellagrin, there are at the present time (March, 1914) no active cases of pellagra in the institution. This segregation and practical isolation of pellagrins may possibly have been a factor in the remarkably rapid disappearance of pellagra from this institution. The death-rate among pellagrins in this institution was also remarkably high, thus diminishing the number of hypothetical sources of infection.

DISCUSSION

This study shows inequalities in the geographical distribution of pellagra in Spartanburg County. Furthermore, in those communities and elements of the population in which methods of disposal of human excreta were similar, its distribution was still unequal. In the section in which our studies were undertaken, congestion of population, general sanitary conditions being equal, appeared to be an important factor in producing higher incidence rates. Thus we found that, where the population was congested in Spartanburg County, as was the case in mill-village communities, the pellagra morbidity per ten thousand of population was six and a half times greater than in rural communities where the population was sparse. (Rural rate per ten thousand was 16, mill-village rate per ten thousand was 104.)

In the city of Spartanburg there was a distinct tendency for the endemic foci of the disease to be confined to those sections of the city in which unscreened surface privies were in use. Information given us by physicians practicing elsewhere in the South indicates that the distribution of the disease, in at least some other southern cities, coincides, so far as methods of sewage disposal are concerned, with its distribution in the city of Spartanburg. It is our intention, during 1914, to study the distribution of the disease in a number of cities and industrial communities, with particular reference to methods of disposal of sewage.

4. Personal communication.

Fig. 6.—Type of unscreened trough privy formerly in use in Spartan mill-village.

Fig. 7.—General view of unscreened trough privies formerly in use in Spartan mill-village.

Pellagra prevailed and originated in all the mill-village communities in Spartanburg County and in all these communities the methods of disposal of excreta were unscreened surface privies. We investigated the history of pellagra in two mill villages elsewhere in the state in which the only striking contrast was the fact that a complete water-carriage system of disposal of sewage had been installed. In these

Fig. 8.—Type of water-closet recently installed in Spartan mill-village.

villages we found that, notwithstanding the fact that pellagrins moved in from time to time, thus far, there had been no tendency for pellagra to originate or to spread in these communities. These observations appear to us to be somewhat striking and strongly suggest the possibility of a direct relationship between the spread of pellagra and faulty methods of disposal of human excreta.

There is a wide-spread movement under way at the present time in the southern states toward the improvement of methods of disposal of human excreta. Many towns and cities as well as industrial communities in which pellagra is endemic are either improving their types of surface privies or putting in modern water-carriage systems of disposal. It is in such communities that most important information can be obtained, either confirming or refuting these observations. Superficial studies and general impressions will not suffice. If any information of value is to result, the local health officers and physicians in communities, in which improved methods of disposal are being installed, must necessarily know the history of the prevalence and distribution of pellagra prior to the installation of improved methods.

Fig. 9.—Rear view of a row of houses showing type of construction adopted in installation of water-carriage system of disposal of sewage in Spartan mill-village. The closet is on the back porch.

They must also necessarily keep accurate records, for at least three or four years, of the prevalence of the disease subsequent to the installation of improved sanitary methods.

One community exists in Spartanburg in which a water-carriage system of disposal has been substituted for unscreened surface privies. Reference to Figure 2 (the distribution of pellagra in the city of Spartanburg and its relation to methods of disposal of sewage) shows that the disease has prevailed most extensively in the northwestern quadrant of the city. In this quadrant is located one of the largest cotton-mills in the county. About 300 of the dwelling houses in this section are the property of this mill, and during the summer of 1913 the president of the mill, Mr. W. S. Montgomery, decided to install individual water-closets in each dwelling. Up to the present, pel-

lagra has been excessively prevalent in this section and the method of disposal of human wastes has been the unscreened trough system (Figs 6 and 7). The installation of a water-carriage system was begun in the fall of 1913 and is being pushed to completion. On Feb. 1, 1914, 123 installations had been completed and connected with the sewer mains. The type of construction adopted is shown in Figures 8 and 9. A special non-freezing type of closet has been installed, as the fixtures are exposed to sudden and sharp changes in temperature.

We have studied the prevalence of pellagra in this village during 1912 and 1913 and each case has been carefully located. We know that at least eighty cases of pellagra have originated in the north-western section of the city during the past six years. It is our intention to observe carefully the behavior of the disease with respect to the development of new cases during 1914 and for two or three years thereafter.

In presenting these facts at this time, we hope that similar studies may be undertaken in communities elsewhere in the Southern states where pellagra is endemic and where contrasts in methods of disposal of human excreta present themselves.

SUMMARY

1. Pellagra morbidity was higher in congested communities using surface privies than in more sparsely settled districts in which similar methods for the disposal of excreta were employed.

2. In the city of Spartanburg the endemic foci of pellagra were located in the districts in which surface privies were in use.

3. In cotton-mill villages equipped with surface privies pellagra was found to be endemic and new cases of the disease arose there year after year.

4. In two cotton-mill villages completely equipped with water-carriage systems of sewage disposal it was impossible to find cases of pellagra which had certainly originated there, although some cases which had originated elsewhere were present.

5. There is some evidence that pellagra spreads in hospitals for the insane more readily in the wards housing untidy patients.

6. This study indicates that methods of disposal of human wastes may prove to be a determining factor in the spread of pellagra in certain communities and it suggests a possible method of prophylaxis, which is now being tested in a practical way.

Twentieth Street and Second Avenue.