PROGRESSIVE LIPODYSTROPHY *

CHARLES HERRMAN, M.D.
Attending Pediatricist to the Lebanon Hospital
NEW YORK

A few years ago I had under observation at different times two girls aged 10 and 12 years, respectively, in whom I was struck by the marked contrast between the emaciated face with hollow cheeks and the good development of the rest of the body. These children were brought to the clinic for slight ailments and were not affected with any constitutional disease. I realized that I was dealing with a rare condition, but was at a loss to know where to search for a description of it in the literature. A paper recently published by Feer¹ on "Lipodystrophia Progressiva" immediately cleared up the diagnosis. Unfortunately no detailed history, photographs or measurements were taken at the time, and the patients are no longer under observation.

I had the good fortune recently to discover a typical example of this condition in the person of a mother who brought her children for treatment. The history is briefly as follows:

REPORT OF CASE

History.—Mrs. K., aged 32, Hebrew, has been married for ten years to her cousin. She had three brothers and two sisters. One sister died in childhood of pneumonia. One brother and the sister are married and have healthy children. No other member of the family has had a similar condition; neither the mother of the patient nor the other female members were unusually fat or thin, nor did any one have an increased amount of adipose tissue in the lower part of the body. There is no family history of syphilis, tuberculosis, alcoholism, diabetes or nervous diseases. The birth was normal, the patient was breast fed, and her physical and mental development during childhood was normal. At the age of 3 she had measles without complications. Since that time, with the exception of occasional headaches, her health has always been good. The change in her face was first noticed at the age of 6. It began insidiously, without any fever, pain or discomfort, and gradually became more marked, so that at 11 the fat of the face had almost entirely disappeared. Later the dystrophy spread to the neck, upper part of the chest and arms, so that these parts became distinctly thinner. When she was 11 years old the family emigrated from Russia to this country. The patient was taken to a well known New York physician, who after a careful examination told the mother that he hoped her other children were as healthy. The patient began to menstruate at 13 years. She married at 22 years; her weight at that time was 44.5 kg. (98 pounds). After her marriage she began to notice that the lower part of her body was growing stouter, and this continued, so that at the age of 28 she weighed 53.5 kg. (117 pounds) and at 31 years 55.5 kg. (121 pounds). Except for the disfiguring appearance of her face, she has not suffered in any way, her appetite is good, her

* Submitted for publication Jan. 27, 1916.

1. Feer: *Lipodystrophia Progressiva*, Jahrb. f. Kinderh., 1915, lxxxii, 1.

bowels regular, she has no disturbances of sleep, no palpitation or nervous symptoms. She has two children, a girl aged $8\frac{1}{2}$ years, who is perfectly healthy, and a boy aged 7 years, who has chronic endocarditis following an attack of rheumatism. She has had no miscarriages or stillbirths, and is now pregnant for the third time. Both children were breast fed for one year.

Physical Examination.—Measurements: weight, before pregnant, 55.5 kg.; in seventh month of pregnancy, 59 kg.; height, 150 cm.; circumference of the head, 53 cm.; neck, 31.5 cm.; chest, 80 cm.; arm, 22 cm.; elbow, 21.5 cm.; forearm, 21.5 cm.; thigh (at level of great trochanter), 62 cm.; at 15 cm. above patella, 53.5 cm.; knee, 36.5 cm.; leg, 37.5 cm.; ankle, 23 cm.; thickness of the raised fold of fat and skin—cheek, 5 mm.; breast (over nipple), 6 mm. axilla (level of nipple), 6 mm., interscapular region, 8 mm.; arm, 5 mm.; forearm, 5 mm.; neck, 7 mm.; thigh, 45 mm.; leg, 27 mm.

One is immediately struck by the disproportion between the upper and lower parts of the body. The face is haggard, and has a cadaverous expression (Fig. 1). It is hollowed in the temporal region, the malar bones are prominent, and below them there are two hollows, an upper small one, and a lower large one

Fig. 1.—Progressive lipodystrophy in a woman of 32 years. Author's case

separated by a thin bridge, corresponding to the zygomatic muscle. This renders the masseter more distinct, and gives a fullness around the corners of the mouth. There are also distinct grooves under the lower lip and on either side of the chin. When the patient smiles, folds form on both sides of the mouth. The neck shows a marked reduction of the subcutaneous fat; also the arms and upper part of the trunk. There is no disturbance of function or sensation in these parts. Examination of the hair, eyes, ears, nose, throat and teeth shows nothing abnormal. The patient does not complain of flushing, profuse perspiration or chilliness. The breasts are well developed, and on account of the small amount of subcutaneous fat are unusually hard. Examination of the heart and lungs shows normal conditions. From the crests of the ilia downward there is a very marked increase in adipose tissue; this is most distinct in the gluteal region and on the outer side of the thighs. It is less distinct on the leg, and the ankle is not at all enlarged. The function of the muscles, sensation and the tendon reflexes are normal. Examination of the blood and urine shows nothing abnormal. The roentgenologic examination of the arm and of the skull (sella turcica) shows nothing pathological.

Number of Cases, Race, Sex.—Up to the present, fifteen cases of progressive lipodystrophy have been reported by the following authors: Barraquer,² Campbell,³ Holländer,⁴ Simons,⁵ Laignel-Lavastine and Viard,⁶ Cohn,⁷ Lewandowsky,⁸ Weber,⁹ Christiansen,¹⁰ Feer,¹ and Pic and Gardère.¹¹ In most of these reports a detailed history and measurements are not given. Three cases in males which are somewhat doubtful are reported by Shaw,¹² Hertz and Johnson,¹³ Husler.¹⁴ The condition is probably not as rare, however, as the figures would seem to indicate, for like other uncommon conditions, it is sometimes not recognized because not known. A number of the patients were Hebrew, though there does not seem to be any marked predisposition of the race to this condition. All of the undoubted cases have been in females; but it is not at all certain that males may not be occasionally affected.

Etiology and Pathogenesis.—In those few instances in which the family history is given, there is no indication of any inherited tendency, either to the same condition or to a disturbance of the fat metabolism. Syphilis, tuberculosis and alcoholism apparently play no part in the causation of this condition. In those patients in whom the Wassermann test was made the result was uniformly negative. The patients may of course be affected with intercurrent diseases, but these apparently bear no relation to the underlying condition, which appears to be a form of trophoneurosis. One naturally thinks of some disturbance in the function of one or more of the ductless glands, but symptoms of hyperthyroidism such as palpitation, tremor, tachycardia, those of hypothyroidism, such as mental and physical sluggishness, subnormal temperature, constipation, are absent. Symptoms pointing to a disturbed function of the pituitary, suprarenals or ovaries are likewise absent. The condition occurs in married and in unmarried women; they present no unusual disturbances of menstruation, pregnancy or lactation. In addition, none of the patients has received any benefit from the administration of the extracts of the ductless glands. It is not unlikely that

2. Barraquer: Abstr. Neurol. Centralbl., 1907, p. 1072.

3. Campbell: Tr. Clin. Soc. London, 1907, xl, 272; Proc. Roy. Soc. Med., London, 1913, vi, 71.

4. Holländer: Ztschr. f. d. ges. Neurol. u. Psychiat., 1911, v, 633.

5. Simons: Ztschr. f. d. ges. Neurol. u. Psychiat., 1911, v, 29; *Ibid.*, 1913, xix, 377; Ztschr. f. Kinderh., 1915, xi.

6. Laignel-Lavastine and Viard: Nouv. iconog. de la Salpêtrière, 1912, xxv, 473.

7. Cohn: Abstr. Neurol. Centralbl., 1913, xxxii, 779.

8. Neurol. Centralbl., 1913, xxxii, 866.

9. Weber: Brit. Med. Jour., 1913, p. 1154.

10. Christiansen: Mentioned by Feer, Note 1.

11. Pic and Gardère: Lyon méd., 1909, cxiii, 61.

12. Shaw: Tr. Clin. Soc. London., 1905, xxxviii, 222.

13. Hertz and Johnson: Proc. Roy. Soc. Med., London, 1913, vi, 92.

14. Husler: Ztschr. f. Kinderh., 1914, x, 116.

something in the circulating blood is the causative factor. The symmetrical, bilateral character would seem to indicate this. The fact that only certain parts of the body are affected, would not be a strong argument against such an assumption, for it is well known that certain organisms, toxins and other substances have a special affinity for certain tissues; and it is not impossible to conceive of a substance that would cause a diminution or disappearance of the fat in one part of the body and an increase in another part. By injecting the blood of patients in

Fig. 2.—Progressive lipodystrophy in a woman of 21 years. Simons' case.

the *active progressive stage* of the disease into monkeys it might be possible to produce somewhat similar changes.

Symptoms and Course.—The loss of the subcutaneous fat of the face begins insidiously without fever, pain or other symptoms, and gradually becomes more marked; it is bilateral and symmetrical. In six of the patients, including my own, the disease began between the fifth and the seventh year; in the remainder, about the time of puberty. In the majority of the cases the loss of subcutaneous fat then extended

to the neck, arms and upper part of the trunk. The orbital fat is little if at all affected and as no postmortem examinations have been made it is still unknown whether the internal fat is diminished. The increase of adipose tissue in the lower part of the body begins or becomes more marked at the time of puberty. This increase is most distinct in the gluteal region and on the outer side of the thigh, but also involves the rest of the leg. As a rule, both sides are equally affected; occasionally the increase is greater on one side. The amount of subcutaneous

Fig. 3.—Microscopic sections. A. Progressive lipodystrophy. Skin removed to the fascia; almost complete absence of fat; traces in the sebaceous glands and at the roots of the hair. B. Fat content of the normal skin of the head in man 34 years old. Both specimens stained with hematoxylin-Sudan (Simons').

fat gradually increases, but after a certain number of years probably remains stationary, so that, strictly speaking, the disease can no longer be considered progressive. The examination of the parts affected shows no impairment of function of the muscles, no change in the electrical reactions of the muscles or nerves, or in the tendon reflexes, no

disturbances of sensation, no vasomotor disturbances. Roentgenologic examination of the extremities and the skull (sella turcica) shows nothing abnormal. The measurements in the seven cases in which they are given are as shown in Table 1.

TABLE 1.—MEASUREMENTS OF SEVEN PATIENTS WITH LIPODYSTROPHY

Author	Circumference						
	Age, Years	Weight, Kg.	Height, Cm.	Arm, Cm.	Forearm, Cm.	Thigh, Cm. * †	Leg, Cm.
Simons.....	21	46.8	158			56	
Simons.....	28	56.0	159			61 51.5	37.5
Laignel-Lavastine and Viard.....	33	53.0	150	22	20	56 46.0	36.0
Cohn.....	17					57 53.5	39.0
Feer.....	9½	26.9	130	16.5	15.5	.. 36.0	26.5
Boissonas.....	7½	22.1	119	17	18.0	.. 34.5	26.5
Herrman.....	32	55.5	150	22	21.5	62 53.5	37.5

* Measured at the level of the great trochanter.

† Measured 15 cm. above the patella.

The measurements of the thickness of the fold of raised skin and subcutaneous fat,¹⁵ as first suggested by Oeder,¹⁶ and carried out on children by Batkin,¹⁷ are given by only three authors as set forth in Table 2.

TABLE 2.—MEASUREMENTS OF THICKNESS OF FOLD OF SKIN AND SUBCUTANEOUS FAT

Author	Age, Years	Cheek, Cm.	Breast, Cm.	Axilla, Cm.	Arm, Cm.	Fore-arm, Cm.	Thigh, Cm.	Leg, Cm.
Laignel-Lavastine...	39	..	..	...	...	...	64	59
Feer.....	9½	4	3	3.5	3.5	2.5	30	18
Herrman.....	32	4	6	6.0	5.0	5.0	45	27

In none of the patients were changes found in the internal organs which could be attributed to the disease. The examination of the blood

15. The measurement of the thickness of the layer of subcutaneous fat is made as follows: A fold of skin with the subcutaneous fat is raised from the underlying muscle and held by an assistant between the thumb and index fingers of both hands, the thumbs being separated 3 or 4 cm. This double layer of skin and fat is then measured with a pair of calipers; not the kind which are used for measuring the diameters of the head or pelvis, but the kind used by mechanics. This touches the fold along a larger surface and the thickness can be more accurately determined.

16. Oeder: *Med. Klin.*, 1910, p. 17; *Fortschr. d. Med.*, 1911, p. 41.

17. Batkin: *Jahrb. f. Kinderh.*, 1915, lxxxii, 103.

and the urine, and the roentgenologic examination have shown no characteristic changes. In those cases which were tested no disturbance of metabolism and no vasomotor or trophic disturbances were noted. Three hundred grams of grape sugar have been given without producing an alimentary glucosuria; no increase of lipase has been found in the blood; after the ingestion of 200 gm. of a 20 per cent. cream there was no lipemia; the blood serum obtained by puncture of a vein was clear. When large quantities of cream and butter were taken the feces

Fig. 4.—Boissonas' case, a girl aged 10. Disappearance of subcutaneous fat in the face, arms and chest; marked increase in the gluteal region, abdomen, hips and calves.

did not show any marked change. Injections of pilocarpin produced moderate sweating, less distinct on the face; the pulse, respiration and intestinal peristalsis were not affected.

Macroscopically a section of the skin of the affected part shows a marked diminution or a complete absence of fat; microscopically (Simons' Fig. 3) almost a complete loss of the subcutaneous fat; only in the sebaceous glands and at the roots of the hair can traces of fat be seen. The epithelium and the corium present no changes.

Diagnosis and Differential Diagnosis.—If we bear in mind the characteristic features of this condition—bilateral, symmetrical loss of fat in the upper part of the body, with increase of adipose tissue in the lower part, and the absence of other trophic and constitutional disturbances—the diagnosis is usually easy. The cadaverous expression of the face in an otherwise healthy individual is almost pathognomonic. The good general health immediately excludes tuberculosis, malignant

Fig. 5.—Pulmonary tuberculosis in a girl 9 years old (Schlossmann).

tumors and diabetes. The retention of the normal function of the muscles excludes the various forms of muscular atrophy and dystrophy; the normal sexual characteristics and the absence of ocular disturbances separate it from dyspituitarism of the Froelich type.

In passing, it is interesting to note that in this disease the subcutaneous fat is lost in a location in which in infantile atrophy, even when extremely marked, it is retained, namely, as the fat or sucking pad of the cheek. On the other hand, in some cases of advanced pulmonary tuberculosis in children, the fulness of the face may persist after the rest of the body is greatly emaciated (Fig. 5). It is not uncommon

to see girls and young women in whom there is a striking contrast between the thinness of the face and the normal or more than normal development of the lower part of the body; and although I do not believe that such cases should be included in the group of progressive lipodystrophy, they may possibly bear some relation to that condition.

Prognosis.—Although the disappearance of the subcutaneous fat of the face is progressive, after it has all been lost the process must of necessity come to a standstill. A spontaneous reappearance of the fat is extremely unlikely. The increase in adipose tissue in the lower part of the body is also progressive, but after a certain number of years apparently reaches its maximum. The disease appears to interfere in no way with the general health, and it is doubtful if life is shortened.

Treatment.—Thyroid, pituitary, and ovarian extract have been tried without benefit. Feer proposes to use the milk of a thyroidectomized goat, with which he obtained good results in a case of Graves' disease. The rationale of this method of treatment is not clear, since no similar disturbances have been noted in this disease. It would seem more logical to inject the blood of a patient in the active progressive stage of the disease into monkeys with the idea of stimulating the production of antibodies and later to inject the serum of such animals into the patient.

The patients are all female and naturally complain, chiefly, of the disfigurement, especially if they are still unmarried. For cosmetic reasons human fat and mutton suet have been injected into the hollows of the face, but after a time this is absorbed. Transplantation of the excess fat from the buttocks to the face has been suggested, but it is doubtful whether that will remain unabsorbed. Paraffin injections have been objected to because they interfere somewhat with the play of the features; but in view of the great deformity this does not appear a strong objection, and this is the method which I intend to employ in my patient.

250 West Eighty-Eighth Street.