HEART-BLOCK

I. TWO CASES OF COMPLETE HEART-BLOCK SHOWING
UNUSUAL FEATURES *FRANK N. WILSON, M.D., AND G. CANBY ROBINSON, M.D.
ST. LOUIS

The following two cases of complete heart-block are considered worthy of report because of several unusual features shown by electrocardiograms.

REPORT OF CASES

CASE 1.—Mrs. E. P., an American housewife, aged 39, had been under observation in the Washington University Dispensary for two years. She first came to the dispensary because of various gynecologic complaints associated with retroversion of the uterus. At that time it was found that she had an unusually slow pulse rate, and electrocardiograms revealed the presence of complete heart-block. This condition had been constantly present since, but in spite of this fact she had had no cardiac symptoms aside from slight dyspnea on exertion. She had never had a syncopal attack. Physical examination of her heart had constantly shown moderate enlargement of the cardiac dulness, the presence of an apical systolic murmur transmitted to the axilla, and a faint aortic systolic murmur poorly transmitted. The patient's past history revealed no infections which might account for the cardiac condition aside from frequent attacks of tonsillitis during childhood. The Wassermann reaction was negative. When she was 19 years old, she was told by a physician that her pulse rate was only about one half the normal, so that it seems likely that the heart-block had been present for at least twenty years.

CASE 2.—Mr. M. R., a German cigarmaker, had been under observation in the Barnes Hospital and the Washington University Dispensary since March 31, 1917. He presented himself on that date complaining of syncopal attacks, vertigo, and dyspnea on exertion. During the previous seven weeks he had had five syncopal attacks of short duration, all of which came on without warning and without any apparent exciting factor. None of these attacks followed exertion. Between attacks he was troubled by vertigo on turning quickly and by dyspnea on exertion. Physical examination revealed moderate enlargement of the cardiac dulness and a faint systolic murmur was heard over the whole precordium. The heart was beating regularly at the rate of 36 per minute. There was a moderate arteriosclerosis. Electrocardiograms showed the presence of complete heart-block. The past history and the laboratory findings, including the Wassermann reaction, were negative.

DISTURBANCES OF AURICULAR ACTIVITY

The electrocardiograms revealed a tendency in both patients for those interauricular periods during which a ventricular systole occurred to be shorter than those which followed them (Figs. 1 and 2). This

* Submitted for publication Sept. 1, 1917.

* From the Department of Internal Medicine, Washington University Medical School.

Fig. 1 (Case 1, Lead II).—Complete heart-block. Atropin produced an acceleration of both the auricular and ventricular rates and caused the auricular arrhythmia to disappear.

Fig. 2 (Case 2).—The usual cardiac mechanism. Complete heart-block; auricular rate 66.8, ventricular rate 33.4. Those auricular periods which contain a ventricular contraction are shorter than the others.

tendency was observed only when the auricular rate was comparatively slow; it was not definitely present when the auricular rate was rapid, as after exercise or after the administration of atropin (Fig. 1). This type of auricular arrhythmia has been previously described by Hecht,¹ who observed it in a case of complete heart-block in a child. This author offers no explanation for its occurrence. Erlanger and Blackman² have described a similar auricular arrhythmia in experimental heart-block. They found that each ventricular systole produced a marked slowing of the auricular rate. In marked instances this slowing was so great that only one auricular beat occurred during each interventricular period; when several auricular beats occurred between two consecutive ventricular beats, the first interauricular interval was long and the following auricular periods became gradually shorter. Erlanger and Blackman ascribed this type of auricular arrhythmia to

Fig. 3 (Case 2).—Ectopic auricular beats represented by inverted auricular complexes occur when an auricular contraction is due shortly after the beginning of ventricular systole.

rhythmical “variations in vagal tone, vagus inhibition increasing with each arterial pulse and diminishing again between pulses.” It is quite possible that the auricular arrhythmia observed in our patients was the result of a similar mechanism. The absence of the arrhythmia when the heart rate was fast after exercise and after atropin (Fig. 1) lends support to the view that it was dependent on vagus activity.

The occurrence of another type of auricular disturbance in one of our patients leads us to believe, however, that another explanation of this arrhythmia is possible. In Case 2 a peculiar type of ectopic beats frequently occurred. These beats are represented in the electrocardiograms (Fig. 3) by inverted auricular complexes (in Leads II and III)

1. Hecht, A. F.: *Ergebn. d. inn. Med. u. Kinderh.*, 1913, **11**, 324.
2. Erlanger and Blackman: *Heart*, 1909-1910, **1**, 177.

occurring from 0.20 to 0.34 second after the beginning of the QRS group of the ventricular complex. These abnormal auricular complexes appear only when a normal P wave is almost due at this point. They bear, therefore, a fairly constant relationship to both auricular and ventricular systoles. They differ from ordinary extrasystoles in this respect, and also because they occur so very late in auricular diastole. On the other hand, it is impossible to believe that they represent retrograde auricular contractions, which they somewhat resemble, because of the presence of complete heart-block. Similar ectopic beats have been observed in a case of complete and incomplete heart-block by Cohn and Fraser,³ and these authors mention their occurrence in a case of complete heart-block studied by Parkinson. They attribute their occurrence to the mechanical effect of ventricular systole on some ectopic auricular center. In our case, as well as in that described by Cohn and Fraser, the form of the abnormal complexes indicates that this center was situated in each case, in the lower auricular or upper junctional tissues, and it seems likely that it was located in the specialized tissues to be found in that region. We must assume that ventricular systole hastened the discharge of the stimulus in this center, and it seems possible that a similar effect of ventricular systole on the ordinary auricular pacemaker may account for the auricular arrhythmia discussed in the foregoing. A careful study of the electrocardiograms has not enabled us to decide with confidence between this explanation and the one advanced by Erlanger and Blackman, which was mentioned previously.

The effect of ventricular systole on the auricular period may explain another interesting phenomenon observed in both patients. It sometimes happened — frequently and for long periods in Case 2, less frequently and for shorter periods in Case 1 — that the auricular rate was exactly twice the ventricular. Whenever this occurred, the relationship of the auricular contractions was always nearly the same, and was of such a nature that every alternate auricular complex fell on the T wave. It can be easily understood that this relationship might be brought about by a tendency for ventricular systole to bring about an auricular contraction at this point. This relationship between the positions of auricular and ventricular contractions, when the auricular rate is an exact multiple of the ventricular, may lead to the diagnosis of partial block when complete block is really present. This fact has also been noted by Erlanger.⁴ It is only when the ratio of auricular rate to ventricular rate is disturbed by exertion, or some other factor, that the true cardiac mechanism becomes evident.

3. Cohn and Fraser: *Heart*, 1914, 5, 141.

4. Erlanger: Personal communication.

In Case 1 an entirely different type of auricular disturbance was observed. This disturbance consisted in a very marked change in the form of the P wave in the third lead. The usual form of the P wave in this lead is shown in the first six cycles of Figure 4. In the seventh cycle of this figure the P wave becomes very much larger and broader. This change in the form of the P wave was observed in this patient on two occasions. Each time it was accompanied by a definite increase in the auricular rate. We have observed similar changes in the form of the P wave in other patients and in normal individuals; they are usually accompanied by a change in rate and are usually most marked in Lead III. We believe that they are due to changes in the location of the pacemaker within the sino-auricular node, similar to the change in location of the pacemaker within this node which has been shown to occur in the experimental animal under the influence of vagus stimulation.⁵

Fig. 4 (Case 1).—A spontaneous change in the form of the P wave accompanied by an increase in auricular rate.

THE VAGI AND THE IDIOVENTRICULAR RATE

There has been considerable discussion in the past as to whether the vagi exert any chronotropic influence on the independently beating mammalian ventricles. The question has been repeatedly investigated both experimentally and clinically. As a result of the clinical investigations, it has been well established that in the majority of the cases of complete heart-block, pressure on the vagus nerves in the neck and the administration of atropin do not produce any change in the ventricular rate. In a few cases, however, the administration of atropin has produced a definite ventricular acceleration. Such cases have been reported by Hecht¹ and by Hart.⁶

Vagus pressure and atropin experiments were carried out on each of our patients. Pressure on the vagi in the neck failed to affect the ventricular rate in either. The subcutaneous administration of one-

5. Lewis, Meakins and White: *Phil. Tr. Roy. Soc., London, Series B*, 1914, **205**, 375.

6. Hart, T. S.: *Am. Jour. Med. Sc.*, 1915, **149**, 62.

fiftieth grain of atropin in Case 1 increased the auricular rate from 89.9 to 124.4 per minute and the ventricular rate from 43.4 to 60.0 per minute (Fig. 1). In Case 2, atropin increased the auricular rate from 74 to 96 beats per minute, but failed to affect the ventricular rate.

Two general hypotheses have been advanced to explain such findings. According to the first of these, the influence of the vagi on the idioventricular rate is dependent on the site of the lesion producing the block, and consequently on the level within the junctional tissues at which the ventricular pacemaker is located as described by Zander.⁷ If this is true, in the majority of the cases of complete heart-block the lesion, and consequently the ventricular pacemaker, is situated relatively low down in a region not subject to vagus control, but in an occasional case the lesion and ventricular pacemaker are located at a higher level in a region which is normally under the control of the vagi. Organic structural lesions may occur in any portion of the His bundle, but the recent experimental work of Lewis, Meakins and White⁸ indicates that functional block usually has its site in the region of the A-V node. It seems likely, therefore, that, if the influence of the vagi on the ventricular rate is dependent on the site of the lesion, the chronotropic influence of the vagi on the ventricles would be more marked in functional than in organic heart-block. There is not sufficient data at hand, however, to determine this point. The greatest objection to this whole hypothesis is the fact that the idioventricular electric complexes, which occur when complete block is brought about in animals by vagus stimulation, are usually markedly abnormal and distinctly different from those seen when complete block is produced by other means (Kahn⁹), and also different from the ventricular complexes which usually occur in clinical complete heart-block. This unquestionably indicates that the vagi do exert an influence on that region within the junctional tissues which lie above the bifurcation of the His bundle, a region which, judging from the usual form of the ventricular complexes in complete heart-block, contains the idioventricular pacemaker.

According to the second hypothesis which has been advanced to explain the established facts with regard to the vagi and the idioventricular rate, the region which contains the idioventricular pacemaker is normally subject to vagal influence, but this influence is destroyed by the same lesion which produces the block (Lewis¹⁰).

7. Zander, E.: *Nord. med. Ark.*, 1915, **2**, No. 6, Part 2. Reviewed in *Zentralbl. f. Herz. u. Gefässkrank.*, 1915, **19**, 1914.

8. Lewis, Meakins and White: *Heart*, 1914, **5**, 289.

9. Kahn, R. H.: *Ergebn. d. Physiol.*, 1914, **14**, 1.

10. Lewis, T.: *Lectures on the Heart*, New York, Paul B. Hoeber Company, 1915.

This might easily occur if, as is probable, the majority of the vagus fibers reach the ventricles by way of the A-V bundle. Advocates of this theory have, so far as we know, not attempted to explain the occurrence of cases, such as one of those which we are reporting, in which, in spite of the presence of complete block, the ventricular rate is subject to vagal influence. The occurrence of such cases is, however, not an insuperable objection to the theory, for it has been shown by Garrey¹¹ that the passage of the excitation wave through a layer of muscle may be interrupted by compression without preventing the passage of nerve impulses through the same area. It is possible also that a lesion may be so located as to interrupt the muscular without interrupting the nervous connection between auricles and ventricles.

Fig. 5 (Case 2).—The effect of exercise on the cardiac mechanism. The auricular rate is markedly intensified, and the ventricular rate only slightly increased.

We also investigated the effect of exercise on the cardiac rate in one of our cases (Case 2, Fig. 5). The exercise consisted in doing 3,300 foot-pounds of work in thirty seconds by climbing stairs. Electrocardiograms were taken immediately before and after the exercise. The auricular rate was increased from 75 to 120 per minute and the ventricular rate from 34 to 40 per minute. According to Gasser and Meek¹² exercise may increase the heart rate in four distinct ways: (1) by an inhibition of vagus activity; (2) by an increase of accelerator activity; (3) by an increase in the secretion of epinephrin, and (4) by an increase in the temperature of the blood. They found that the first of these mechanisms was by far the most important. Since

11. Garrey: *Am. Jour. Physiol.*, 1911, **28**, 249.

12. Gasser and Meek: *Am. Jour. Physiol.*, 1914, **34**, 48.

the ventricular pacemaker in our patient was apparently not under the control of the vagi, the increase in the ventricular rate observed after exercise must have been due to the last three factors mentioned. Erlanger and Hirschfelder¹³ have shown that the idioventricular rate in complete heart-block is subject to accelerator influence, and the other two factors concerned would probably act without reference to the location of stimulus production. The slight increase in the ventricular rate as compared with the marked auricular acceleration indicates the great disadvantage from which patients with complete heart-block suffer during exercise as a result of the loss of vagus control of the ventricular rate.

DISTURBANCES OF VENTRICULAR ACTIVITY

In Case 2, two distinct disturbances of ventricular activity were observed. On one occasion there was a spontaneous change in the form of the ventricular complex (Fig. 6). This was not accompanied by any definite change in rate, and two possibilities offer themselves for its explanation: there was either a change in the location of the ventricular pacemaker, or a change in intraventricular conduction at this time. This phenomenon may have been caused by a massive dose of the tincture of digitalis which had been administered twenty-three hours before it was observed.

The second disturbance of ventricular activity shown by this patient was a marked tendency to multiple ventricular extrasystoles after exertion. Multiple extrasystoles could be produced almost at will by having the patient rapidly perform about 3,000 foot-pounds of work by climbing stairs. They were first observed about twenty-three hours after the administration of a massive dose of digitalis, but could be produced as long as the patient remained under observation. The extrasystoles were not always of the same type and they occurred at a very rapid rate (Fig. 7). Single extrasystoles were observed during the same period and the number occurring in succession varied from two to eight. These extrasystoles did not occur immediately after the cessation of effort, but about two to three minutes later, and there was usually considerable variation in the form of the ventricular complex at the time of their appearance. At first it was thought that there was some relationship between the tendency to multiple extrasystoles after exertion and the administration of the digitalis, but the fact that this tendency continued unabated for a period of sixteen days after the digitalis had been given makes this relationship doubtful.

It is also interesting to speculate on the relationship of this tendency to multiple extrasystoles and the syncopal attacks which the patient

13. Erlanger and Hirschfelder: *Am. Jour. Physiol.*, 1905-1906, **15**, 153.

Fig. 6 (Case 2).—A spontaneous change in the form of the ventricular complex.

Fig. 7 (Case 2).—Multiple extrasystoles after exercise.

had. Ventricular tachycardia might conceivably produce such attacks, if sufficiently long continued, and, moreover, it is possible that the ordinary pacemaker might fail to take up the rhythm promptly after its interruption by such a series of ectopic beats as in a case observed by Lewis.¹⁰ As the patient had no syncopal attacks while under observation, no direct evidence could be obtained on this point. Against the existence of any causal relationship between the multiple extrasystoles and the syncopal attacks is the fact that the latter never occurred after effort, while the former occurred at no other time during the period that the patient was under observation. The constant appearance of multiple extrasystoles after exercise in this case suggests that exertion may have influenced the nutrition of the heart unfavorably and thus led to their production. This idea is borne out by observations on a patient with angina pectoris who developed an idioventricular tachycardia on exertion.

SUMMARY

Two cases of complete heart-block showing unusual features are reported. The following phenomena were observed:

1. Auricular arrhythmia dependent on ventricular activity.
2. Ectopic auricular beats dependent on stimuli received from the contracting ventricles.
3. Increase in the idioventricular rate after the administration of atropin.
4. Multiple ventricular extrasystoles after exertion.
5. Spontaneous changes in the form of the auricular and of the ventricular complexes.

The explanation of these phenomena is discussed.