PULMONARY BOTRYOMYCOSIS

REPORT OF A CASE

F. A. McJUNKIN, M.D.

ST. LOUIS

Botryomycosis is known to veterinarians as a pedunculated fibrous granuloma of horses especially. The growth usually arises in connection with skin surfaces, and often in the cut spermatic cord following castration. However, in the first description of the condition by Bollinger¹ in 1870 an internal organ, the lung of a horse, was the seat of the disease. In 1897, Poncet and Dor² identified the infection in a woman in a small growth on the palmar surface of the hand. Although human cases have been reported from France, Italy and Switzerland, it has been observed most commonly in northern Africa and in Morocco where it is said to occur as commonly as actinomycosis.

The gross lesions consist of abundant chronic granulation tissue which is not especially characteristic but which on microscopic examination reveals granules made up of coccus-like organisms embedded in and surrounded by a hyaline matrix. In the chronic inflammatory lesions described as granuloma pyogenicum or pseudobotryomycosis, these characteristic "botryomycotic granules" are not present.

In 1913, Opie³ studied a case of hepatic botryomycosis in a girl, 11 years of age, which is the first human case with involvement of an internal organ, and appears to be the only instance of the disease in the United States. In this remarkable case a large part of the liver had undergone necrosis with extensive scarring in some portions and with formation of abscesses containing the characteristic "botryomycotic granules" in others. According to Opie, veterinarians in Missouri and nearby states have seen the disease in horses, while reports indicate that it is uncommon on the Atlantic coast.

In 1914, an article on botryomycosis with a review of the literature to that date was published by Margrou,⁴ who was able to reproduce in the testicle of guinea-pigs spherical masses of staphylococci analogous to the "botryomycotic granules" by injection of measured quantities of this organism. The conclusion reached by him is that the zooglea-like hyaline matrix in which the organisms are embedded, and the peculiar

1. Bollinger: Virchows Arch. f. path. Anat. **49**:583, 1870.

2. Poncet and Dor: Lyon méd. **43**:213, 1897.

3. Opie: Arch. Int. Med. **11**:425 (March) 1913.

4. Margrou, quoted by Masson.

cellular reaction, do not signify a specific organism but a certain relationship between the resistance of the host and the virulence of the usual staphylococcus.

Masson,⁵ in 1918, observed a case of botryomycosis in a gunshot wound with fracture of the femur. The granulation tissue in which the typical granules are present was curetted from the sinus tract six months after the injury. The wound, after being irrigated with surgical solution of chlorinated soda for two weeks, was curetted and healing followed. He concurs in the opinion expressed by Margrou in regard to the staphylococcus nature of the organism, and places emphasis on nonencapsulated masses of the organisms along the Haversian canals of necrotic sequestrums.

Tussan⁶ recently observed four cases of botryomycosis of the skin in Morocco; he discusses them only from a clinical standpoint.

The botryomycotic granules made up of mulberry-like masses of gram-positive coccus-like organisms surrounded by hyaline material are fully as characteristic as the "sulphur" granules of ray-fungus infection, and the histologic diagnosis of botryomycosis is certain and easy. Two views are held in regard to the organism, many regarding it as the usual *Staphylococcus pyogenes aureus*, while others express the opinion that it is a specific coccus. Cultures made from the lesions by the different observers on the ordinary media have usually given a growth quite like that yielded by the staphylococcus.

Especial interest attaches to the case here reported because of its occurrence in the same community from which a previous one with localization in an internal organ was reported.

REPORT OF CASE ⁷

A male infant, four months of age, was admitted to the St. Louis Children's Hospital, Sept. 15, 1920. The father is a painter. The father, mother and other members of the family have been well during the life of the child, and there is nothing in the family history to indicate the source of the infection.

The child was born at home and nursed the mother for three months, during which time it remained well. It has been on various artificial foods since being weaned and has had diarrhea at times.

The baby has always had a hacking cough which has been much worse during the past six weeks. About three weeks ago it had a high fever which continued for one week.

Clinical Examination.—The temperature was 99.2 F., respiration 50. The baby seemed quite lively and cried continually in a loud voice. There was a papular eruption over the shoulders which was probably miliariasis.

There was an impaired percussion note over the entire right side and the breath sounds were suppressed and accompanied by fine crackling rales, with

5. Masson: Lyon Chir. **15**:230, 1918.

6. Tussan: Lyon Chir. **15**:241, 1918.

7. The clinical history I have been permitted to use by Dr. Williams McKim Marriott into whose service the case was admitted.

suppressed breathing posteriorly over the left chest. Impression was that of a beginning bronchopneumonia or of an old process clearing up. Examination of heart, abdomen and extremities, negative.

Clinical Course.—September 18, the baby had a distressing cough bringing up a seromucous secretion. The temperature remained low. Leukocytes, 20,000 with 62 per cent. neutrophils. September 23, the lungs showed everywhere medium sized crackling râles. The low temperature is certainly peculiar with so much in the lungs.

October 6 it was evident that the patient was steadily growing worse. Glucose and saline solution were given intravenously. The temperature only occasionally reached 101 F. The lung process had been very peculiar all along—many signs with little temperature elevation. It had been considered of probable tuberculous origin by everyone. Von Pirquet test was repeatedly negative.

The child continued to grow worse and died October 8.

Fig. 1.—A low power photomicrograph to show the areas of bronchopneumonia in which are the dark-staining "botryomycotic granules."

REPORT OF NECROPSY

The body is that of a poorly nourished male infant weighing 2,610 gm. There is slight rigor mortis, no lividity and no demonstrable edema; necropsy three hours postmortem.

The peritoneal cavity contains 200 c.c. clear fluid. The stomach and upper portions of the small intestine are greatly distended with gas.

Gross examination of the organs other than the lungs shows nothing definitely abnormal. The myocardium has a slightly yellowish tinge suggesting some deposition of fat. The spleen is firm in consistency and the lymph nodules may be distinguished. The mucosa of the small and large intestines shows nothing unusual. The pancreas, kidneys, adrenals, genito-urinary organs and aorta appear normal.

Lungs: Each of the lungs weighs 40 gm. On the pleural surface of both lungs there are centimeter areas of a dark bluish color most of which are somewhat depressed below the surrounding pleural surface. These are present in all lobes but are most conspicuous in the posterior portions of the right lung. On section these irregular areas appear to be airless. The bronchi when cut across exude a mucopurulent material distinctly slimy in character. The peribronchial lymph nodes are not especially enlarged.

Fig. 2.—A medium power photomicrograph of a pneumonic focus containing two foreign-body giant cells, one incorporating organisms, the other without organisms.

Histologic Examination.—The lesions in numerous sections taken from various parts of the lungs are much alike and consist of areas in which the air sacs are collapsed (atelectasis), and of a bronchitis which in many places involves the entire bronchial wall and extends into the surrounding alveolar sacs (Fig. 1). Even with low magnification the latter lesion is quite different from the usual type of bronchopneumonia owing to the presence of giant cells of the type seen in tubercles and of peculiar dark masses which consist of gram-positive coccus-like organisms embedded in a hyaline eosin-staining matrix. In some instances both within the bronchi and outside giant cell forma-

tion (Fig. 2) has taken place about the groups of organisms so as completely to surround them. In Gram-Weigert stains, where the relation of organism and zooglea-like matrix appears to the best advantage, some of the groups are seen to be surrounded by a rather wide zone of hyaline substance the periphery of which is regular or forms small radial projections; others lie free between the cells with little indication of hyaline about them. However, a small amount of pinkish background is seen about some of the groups of a half dozen or more micro-organisms (Figs. 3, 4 and 5).

The exudate in the alveoli and bronchi consists of endothelial leukocytes and neutrophils in about equal numbers. Many of the former have abundant

Fig. 3.—Photomicrograph of "botryomycotic granules." Zeiss 3 mm. apochromat. objective and 8 $\times$ compensating ocular. Certain ones of the spherical organisms embedded in hyaline are in focus; about the entire mass leukocytes are fusing to form a giant cell.

cytoplasm with fat vacuoles, some have ingested neutrophils, and others have fused to form foreign body giant cells. In the bronchial exudate and in the alveolar sacs there are comparatively few cells of the lymphocytic series but the bronchial walls are the site of an extensive lymphocytic infiltration. In places the epithelium of the bronchi has undergone necrosis and disappeared with extension of the inflammatory exudate into the surrounding air cells. The submucosa of the bronchi is heavily infiltrated with lymphocytes and plasma cells. There are granular foci in the exudate within the bronchi in which most of the leukocytes have undergone necrosis. The alveolar epithelium in the pneumonic areas is intact even in alveoli greatly distended with exudate.

Fig. 4.—The capsule at one side of the mass of organisms stains intensely and presents a radial arrangement.

Fig. 5.—Although the homogeneous capsule which lies between the dark mass of organisms and the cells is not stained intensely, it is quite distinct.

The Gram stain of smears of the bronchial exudate made at the time of the necropsy shows further details in regard to the organism. The larger masses of the coccus-like organism are embedded in and frequently encircled by the hyaline matrix, but smaller groups of a half dozen or more often have none of the pink-staining substance about them. There is considerable variation in

Fig. 6.—Photomicrograph of the organisms in a Gram stain of a smear of bronchial exudate. Zeiss 2 mm. apochromat. objective and $8\times$ compensating ocular. Three small groups of the organism in one of which the hyaline has been smeared out on one side and stains positively.

Fig. 7.—A group of the organisms from a part of the smear in which there appears to be a delicate positively staining capsule about each unit of the group. Magnification and staining are the same as in Figure 6.

size and many are less than one-half the diameter of those embedded in the matrix. Groups of two, four, six and eight predominate, and frequently the pairs present a coffee-bean appearance. There are a few chains, the longest consisting of about twenty-four units. In some of the small groups the

hyaline is distinct and about the individual organisms in portions of the smear there are regular pink-staining capsules contrasting with the purplish and more heavily staining organism (Figs. 6 and 7).

Microscopic examination of the other organs reveals no lesions of evident importance. The heart, spleen, stomach and liver are normal. In the mucosa of the intestine there are small collections of plasma cells and an occasional epithelial cell is necrotic and invaded by leukocytes. There is a moderate edema of the kidney with hyaline casts in the tubules. In the intima of the aorta opposite the intercostals the collagen is pushed apart by a small amount of granular material and in this there are a few migrating endothelial leukocytes.

Cultures on blood agar plates made from the bronchial exudate gave a heavy growth of gram-positive cocci which presented the usual appearance of *staphylococcus albus*. Nothing unusual was suspected at the time of necropsy, and unfortunately the cultures taken had been discarded when the lesion was identified from the sections of lung. Smears and sections of the lung stained by the Ziehl-Neelsen method show no acid-fast bacilli.

SUMMARY

With the pulmonary symptoms predominant at the beginning of the disease and persisting for several weeks in such a manner as to suggest pulmonary tuberculosis, one is led to think that this case was a primary bronchopneumonia which terminated fatally. It is not perfectly clear, however, that this was the sequence of events.

The histologic lesion is that of a typical granuloma and is one not encountered in infections with pyogenic cocci commonly examined microscopically. The zooglea-like matrix likewise is unique. The appearance of the delicate circle of hyaline material about individual organisms which stains positively and is not a retraction zone suggests a true bacterial capsule and, if confirmed by further observations, especially in connection with animal inoculation, would indicate a distinct species of capsulated coccus.

In conclusion I wish to express my indebtedness to Dr. E. L. Opie who brought to my attention the lesion present in his case and to Dr. H. H. Bell for an examination of the cultures.