OSTEOMALACIA *

E. P. CORSON WHITE, M.D.

PHILADELPHIA

Osteomalacia is a generalized softening of adult bones which were at one time quite calcified and hard. It occurs in adult man and in some animals and birds. In man several types are recognized.

1. A mild form is found in pregnant, puerperal and lactating women, which recurs with each pregnancy. The skeletal changes become worse with each attack, but tend toward a spontaneous recovery when the drain of pregnancy and lactation is removed. A very similar group of cases arises in famine districts, or when the variety of diet is limited. The avidity of the tissues for calcium and phosphorus in these forms of osteomalacia is, however, undisturbed, and when these elements are added to the dietary in usable form, the patient recovers.

2. A second or senile type occurs in old people, in which the bone softening is, as a rule, confined to spine and pelvis. It generally progresses rapidly for a time, then ends in recovery with marked residual deformity.

3. A severe form occurs in pregnant, puerperal and multiparous women, in boys, men and castrated women, which is steadily progressive and shows no tendency toward recovery, ending in a marasmic state.

In lower animals practically all cases examined were of this last type. It occurs equally in male and female, mammal and bird. Up to the present time there have been forty-nine cases among mammals and thirty-four among birds. Considering the many species of animals exhibited, the disease is rather sharply limited to a very few groups.

TABLE 1.—CASES OF OSTEOMALACIA IN MAMMALS AND BIRDS

MAMMALS			
	Cases	Males	Females
Primates	26	19	7
Lemurs	4	0	4
Carnivora	6	4	2
Hyraces	1	0	1
Rodentia	5	3	2
Marsupialia	8	5	3
AVES			
	Cases		
Columbae	12		
Galli	9		
Psittici	6		
Passeres	2		

This table shows that 50 per cent. of all the mammals infected are primates, and in this order are limited almost exclusively to the Cebidae.

* From the Laboratory of the Philadelphia Zoological Society.

Of the birds 40 per cent. are doves (*Columbae*), 30 per cent. are pheasants (*Galli*), 12 per cent. parrots (*Psittici*). In the first two groups the disease is very extensive, involving practically every bone in the body, and the lesions are much more severe than those found in any of the other groups of animals or birds. Many other animals exposed to exactly the same environmental condition show a real or apparent immunity. Necropsy findings are practically the same in birds and mammals.

In mammals, the long bones of the extremities are early and seriously involved. They yield under the weight of the adult body, twist and bend, causing extraordinary and grotesque deformities. Very soon the animal is unable to stand (cage paralysis).

In birds, the more serious lesions occur in the bones of the thorax, ribs and sternum. Claws and beak are frequently affected, often so softened that the bird is unable to pick up seeds.

The gross appearance of the bones is again roughly similar. The skull may be reduced to the consistency of writing paper, the margins of the bony plates are often markedly thinned, while centers show irregular spots of thinning interspersed with areas of thickening. In one case, an adult Cebus, the skull was so completely deprived of calcareous matter that it could be twisted. The ribs are irregularly softened. Thin translucent areas are found associated with white opaque fusiform or nodular beadings, especially at the costosternal junction and at the epiphyses. These beadings are apparently efforts at splinting, occurring where the weakness from excessive bone absorption is the greatest. The whole rib may be soft, the consistency of soft rubber, and almost entirely deprived of the bone salts. It will cut with scarcely any crunch and bend easily in the fingers.

The long bones show the same irregular thickening closely associated with areas so thinned that they are almost translucent. They are usually distorted because of the body weight, generally a forward bowing. Fractures are extremely common; often more than one is found at necropsy. These injuries may show no attempt whatever at repair, or more often a fibrous or inadequate cartilaginous union. In all instances the deposition of bone salts is very inefficient. More rarely there may be an overactive effort at repair, but here also with only a scanty deposition of bone salts. In the immediately adjacent areas there may be marked subperiosteal reaction, but, as a rule, the periosteum shows no reaction at all.

In man, the small joints often become deformed early and in appearance suggest an osteo-arthritis. Large joints may ankylose. The spine and pelvis are involved sooner or later, ending often with a total loss of power over the legs.

Besides the bone lesions, there is constantly present in these animals a more or less severe secondary anemia, a lowered alkalinity of the blood and a fermentative type of intestinal indigestion. All show slight dilatation of the stomach, with some atrophy of walls. The small intestines are thin walled, and the accumulation of gas gives to it a sausage appearance. Lactic acid was found once in the urine. The kidneys show mild nephritis changes, and in some, fatty degeneration or merely pallor is present.

Microscopically, the bones show in every case a more or less pronounced absorption of the calcareous salts (osteoporosis) and with the abnormal bone absorption there is a marked tendency toward the deposition of newly formed osteoid tissue in which calcification never reaches the normal. Sections show osteoid margins along each lamella of bone, covering in the parts still calcified. Normal lamellar arrangement of bone is sometimes retained; usually, however, the bone cortex is very thin with coarse, broad trabeculae, wide marrow spaces and only the slightest tendency toward the formation of Haversian systems. Lacunae and osteoblasts may be normal but contain little calcareous matter. Osteoblasts and osteoclasts may be few, equal in number, or the osteoclasts may be unduly numerous. Osteophytes and Sharpey's fibers, similar to those found in osteitis deformans, are frequently seen. Cancellated tissue is often much decreased, especially at the ends of the long bones. In some cases it has practically disappeared.

The periosteum may be sharply marked off from the bone cortex; as a rule, it shows no activity in its osteogenetic layer. The subperiosteal bone may be very thin, almost like paper, but normal in appearance. At the junction of the bone and the cartilaginous rib there is often a thickening, seemingly from the periosteum. Endochondral bone formation at the epiphysis is usually abnormal, as shown by its irregularity of foci and extremely large intratrabecular spaces; osteoblasts are numerous. The marrow cavity is wide and filled with red more or less edematous marrow. The bone marrow is very much like that found in aplastic anemias. It is generally scanty and collected near the ends of the bones. Bone shows active removal of old trabeculae by means of numerous osteoclasts, and the production of fibroblasts, some of which differentiate into bone cells for forming new trabeculae in which only scanty amounts of lime are deposited. Rarely ever is normal bony architecture reproduced.

An analysis of the bones shows a marked diminution in the amount of calcium and an excess of sulphur and magnesium. Metabolic studies show a loss of calcium primarily in the feces, but also a marked increase in the calcium output in urine, with the retention of sulphur and magnesium. A salt metabolism of one monkey was attempted on four four-day periods, which showed a high calcium and phosphorus loss with a moderate magnesium and sulphur retention.

No hypothesis yet advanced satisfactorily explains the observed facts. The one most generally accepted is Fehling's¹ theory, based on Forchier's observation that osteomalacic women were improved or cured by the removal of the ovaries. His idea was that the disease was due to a hyperfunction of the internal secretion of the ovaries, which produced a sympathetic paralysis of the vasomotor system, which, in turn, caused a hyperemia of the bones and a consequent abstraction of calcium. Careful examination of our autopsy material failed to reveal any specific or distinct lesion in the ovaries, or in any of the ductless glands. All were, however, apparently smaller than glands usually found, with the exception of the adrenals, which were either normal in size or larger, and in two cases much congested. Minor changes were noted in the thyroid in a few instances. Ovaries and testicles were grossly normal in appearance. Microscopically, spermatazoa were found to be much reduced in numbers.

The second theory of origin was that in some manner the alkalinity of the blood was decreased, which permitted the lime salts to go into solution. This decreased alkalinity might be due to (1) a primary ductless gland disorder, especially a dysfunction of those glands which in some manner were concerned with bone metabolism; (2) a loss of alkali; (3) an excessive acidity, either the result of a predominantly acid forming diet, or due to an excessive production of acid in the body. These factors might act independently or in coalition, and might act directly or indirectly through their influence on the ductless glands. The absorption and deposition of salts are dependent on the chemical equilibrium of the cells, and slight shiftings of the acid base equilibrium of the cells in either direction may be sufficient to cause extensive lesions.

In favor of this view are many observations and experiments. Virchow² described deposits of calcium in the organs of people dying of osteosarcoma, osteomalacia and other disease states in which there is excessive bone absorption. These deposits were found in all organs, but especially those excreting an acid into their cavities—lungs, stomach and kidneys. In these organs there was no evidence of diffuse necrobiosis previous to the deposit of the lime salts. It seems probable, therefore, that the calcium was precipitated into the normal organ because the excretion of the acid in these parts left the tissue fluids relatively more alkaline and less able to hold the same amount of calcium in solution. Pawlow³ found a bone condition very similar

1. Fehling: Ueber Wesen und Behandlung der Osteomalazie, *Arch. f. Gynaek.* **28**:1890; **29**:1891.

2. Virchow: Virchows *Arch. f. path. Anat.* **197**:112, 1909.

3. Pawlow, cited by Babkin: *Ztschr. f. Stoffwechsel* **11**:561, 1910.

to osteomalacia in dogs, in which a pancreatic fistula was continued for long periods. Here there was a constant loss of alkali through the fistulas, and the body alkalis had to be utilized for neutralization of the acid production.

Diet has at various times been proposed as a factor, at least, in the production of this condition. This idea has gained considerable weight through the increase of the number of cases recorded from famine districts and parts of Central Europe where the dietary is restricted and unbalanced. Voit ⁴ fed pigeons for one year on diets very poor in calcium, with clinically no ill effects, but found at necropsy marked wasting of the sternum and skull.

In adults a poor calcium diet can be continued for long periods without any apparent ill effect, because the losses from the blood and soft tissues can be replaced from the bones. In the young, however, poor calcium diets very promptly cause weakness and flexibility of the bones.

During pregnancy the calcium requirements of the animal are markedly increased. Steenbock and Hart ⁵ found that the production of milk made heavy drains on the skeleton, unless the amount of calcium and phosphorus contained in the food was abundant. Forbes and Beegle ⁶ also noted that calcium was often lost when the animal was storing nitrogen. Steenbock ⁷ and his associates have produced osteomalacia in cattle by feeding them "shorts," a diet high in magnesium but otherwise well balanced. Sherman and Gettler ⁸ found, on a diet giving a high acid ash, as rice and potatoes, that 33 per cent. of the acidity could be accounted for by increased ammonia production and about 40 per cent. by the increased acidity of the urine, leaving a remainder which must be eliminated or neutralized by sodium or potassium carbonate of blood or other fixed alkali. Etienne ⁹ found that excess of magnesium in an otherwise well balanced food apparently caused a continual loss of calcium. Schiff and Strausky injected subcutaneously a magnesium salt and produced a marked increase in calcium output in urine and a diminution in calcium retention. In children after weaning, calcium disturbances are frequent and often are due to disorders of digestion and obscure irregularities of nutrition.

4. Voit, E.: Significance of Calcium in Animal Nutrition, *Ztschr. f. Biol.* **16**:55, 1880.

5. Steenbock and Hart: Influence of Function on the Lime Requirements of Animals, *J. Biol. Chem.* **3**:23, 1913.

6. Forbes and Beegle: Mineral Metabolism of the Milch Cow, Ohio Agriculture Experiment Stat. Bull. 295.

7. Steenbock and Hart, Development of Osteoporosis by Swine Induced by High Magnesium Intake, *J. Biol. Chem.* **3**:23, 1913.

8. Sherman and Gettler: The Balance of Acid Forming and Base Forming Elements in Foods and Its Relation to Ammonia Metabolism, *J. Biol. Chem.* **11**:323, 1912.

9. Etienne: *J. Physiol. et Path.* **14**:108, 1912.

From these observations and experiments, it seemed advisable, first, to examine carefully the diet of the Cebidae monkeys, in which family the disease was most prevalent and most severe. The food of these monkeys, two bananas or two apples or six small sweet potatoes, with a portion of boiled rice about the size of an egg; occasionally bread, peanuts and lettuce. This diet was studied first, for the amount of essential protein, fat, carbohydrate and ash; then, for the general character of the protein and ash; and third, for the ability of this entire diet and of each of its constituents to maintain normal growth, health and

The average growth curves are plotted for the entire time on which the rats were fed the separate constituents of this diet and they are compared with the curve of normal rats on a well balanced diet and with rats on the complete diet fed to these monkeys.

reproduction. A diet is considered normal when it maintains a normal weight curve, gives a normal immunity to disease, allows completion of a normal span of life, and allows normal breeding of healthy young that shall be vigorous when weaned.

Analysis of this diet computed from Atwater's tables and by actual analysis of the amounts of food consumed by the animal in four four-day periods gave: (1) Protein low in quantity and poor in quality;

especially low in phosphorus content. (2) Fat very low. (3) Carbohydrate very high, almost eleven times the value of all other ingredients. (4) Ash decidedly low and predominantly acid. Further analysis of this ash showed only a trace of calcium and phosphorus and iron; only a small amount of sodium and potassium; sulphur and magnesium were slightly higher. (5) Vitamins A, B and C were present in extremely small amounts. Vitamin A was exceptionally deficient and in the rations of some days was entirely lacking.

The accompanying graph shows the effects of the separate constituents of this diet and the diets as a whole on the growth and nutritive balance of rats. Rats were chosen for this study because they were susceptible, and the only animal for which we have normal standards for comparison. Lettuce, corn and apples did not support life; bananas gave little growth with no reproduction, but life continued for a considerable length of time. Eggs were extremely detrimental to the health of rats, causing diarrhea with many fatty acids in the stool, eczema and early death. Sweet potatoes caused distention of the abdomen, sore eyes, diarrhea with frothy stools, rough coarse fur, hump back, no breeding and considerable atrophy of testicles. Rice produced a loss of weight, constant diarrhea, hump back, swollen joints, sore eyes and no breeding. With bread the general condition was good; weight curve depressed and diarrhea frequent. Reproduction was much reduced. The young died or were eaten by parents before weaning. With peanuts the eyes were sore, a diarrhea frequent, but coat was fair and general condition good. The rats bred poorly and the young all died before weaning. In other words, not any constituent of the diet was able in itself to support life and health. The entire diet gave a generally good condition, with a moderate depression of weight, immunity and breeding. Frothy diarrhea was present in most of the rats at some time during the experiment, and in some was constant. Litters were vigorous at birth but were always eaten by the parents.

A second series of rats was subjected to the principal ingredients of this diet for one month and was then fed with additional food which, theoretically, should replenish the deficient diet and make it adequate, as shown by resumption of normal growth curve.

Table 2 shows the additions necessary for corrections of the separate ingredients of the diet.

The complete diet was rendered adequate by the addition of fresh, whole milk and leafy vegetables, or by butter fat, salt mixture and leafy vegetables.

In this monkey diet several factors are of importance: (1) Low vitamin content, especially vitamin A, factors which are essential for life and growth. (2) A high carbohydrate diet which on oxidation

TABLE 2.—CORRECTION OF DIET INGREDIENTS

DIET	CORRECTED BY
Rice	casein, salt mixture, carrots or lettuce, butter fat.
Bread	casein, butter fat, salt mixture, especially phosphorus.
Potato	salt mixture, especially sodium chlorid and calcium carbonate.
Raw peanuts ..	salt mixture.
Bananas	casein, yeast or carrots.
Corn	casein, tryptophan, lactalbumen.
Egg	detrimental to rats; no recovery.
Apple	casein or gelatin, butter fat.
Onion	casein or gelatin, butter fat.

yields an acid ash and which favors the growth of intestinal bacteria producing acid and gas. The acid from these two sources must be neutralized either by the alkalis derived from food or from the body storage. This diet, however, is abnormally low in ash, and especially in the alkaline salts of the ash, therefore making it an ideal diet for the production of osteomalacia.

The full significance of predominantly acid forming diets has only lately been fully recognized, especially as they affect protein storage, and, therefore, growth and recuperation by deflecting the nitrogen and draining the body alkalies. Goto shows that in experimental acidosis the carbonates are first called on, next, the alkali phosphates, particularly the potassium phosphate of muscles and calcium carbonate of bones.

The main service of milk is, probably due to its alkaline residue when oxidized in the body, and its plentiful supply of calcium, iron and vitamins.

This diet is deficient in those salts that were lacking in the bones of the monkey and which the animal was excreting in excess of its intake, possibly because of the excess of magnesium.

This ration at best would only keep the animal in calcium and phosphorus equilibrium, and any upset making even the slightest demand on its alkaline storage would be sufficient to set up a faulty metabolism.

CONCLUSIONS

It would seem that in this inefficient diet lies, if not the cause of osteomalacia, at least a very potent factor in its production.

1. The disturbance of the calcium and phosphorus metabolism may be due primarily to the deprivation of the alkaline salts from the diet (famine osteomalacia); or, second, to a drain from the alkaline storage of the body, associated with a deficient diet, as in the cases of osteomalacia of pregnancy and lactation; or, third, in the combined action of a diet faulty in more than its salt content, which, by the production of acid in its oxidation and by favoring the development of acid

forming bacteria, causes a drain of the body alkali for the neutralization of this acid; or, fourth, and this is more probable, especially in the severe form, it is due to the combination of all these factors acting upon the ductless glands and through them altering definitely the inorganic metabolism.

2. In severe osteomalacia the power to retain and absorb calcium and phosphorus is practically lost, while in the cases developing during famine, and in pregnancy and lactation, this power is retained. These subjects immediately recover when calcium and phosphorus are presented in usable form. This permanent loss of power to absorb or retain calcium and phosphorus can be explained only by some definite changes in the endocrine balance.

3. It is important, also, that while this disease is very common among the Cebidae it has never been found among our Macaques. This may be due to the fact that owing to the storage sacs in the mouth of the Macaques more food proportional to body weight is consumed; and to a probable essential difference in the basal metabolism of these families and individuals. All the factors enumerated do tax the metabolic resources of the body and depress the functions of the endocrine glands. Detailed and accurate quantitative studies of normal metabolism and their effect on the ductless glands will alone give a more definite answer to the problem. The starting point of treatment must be a diet adequate in vitamins and containing a proper inorganic balance.

.