SENSORY CHANGES IN INJURIES OF THE MUSCULO-SPIRAL NERVE *

ARTHUR S. HAMILTON, M.D.

MINNEAPOLIS

In view of the several very brilliant and equally exhaustive reports on the innervation of the skin which have appeared in the literature of the last few years, one must, with some temerity, approach the study of the sensory functions of any one of the peripheral nerves. And this is the more true when one considers that much of the work recently published has been done on patients well fitted both by education and by training to carefully analyze their sensations and by workers who have given to the task many months of painstaking examination, whereas the conditions obtaining in army life render a similar study wholly impossible, partly by lack of that preliminary education and training on the part of the patient which is essential to the best results in interpreting new and confusing sensations, and partly by reason of the lack of time which permits only a limited study in a routine examination. The difficulty which the latter factor offers is appreciated when one recalls the statement of Trotter and Davis¹ that it requires from forty-five to fifty hours to examine and record the sensibility to cold alone in a fairly large area. For these reasons, therefore, no attempt has been made to describe adequately and to measure all the different sensory disturbances that may appear in the region supplied by the musculospiral nerve. The excuse for this report must lie in the scarcity of literature dealing with the sensory field of this particular nerve and in its surprising areas of distribution found in numerous cases of nerve wounds among soldiers — surprising at least in view of the anatomic distribution of the nerve as commonly taught.

It is important to mention that, as all the cases here reported were studied at the Walter Reed General Hospital in Washington, D. C., it was not possible to determine the area of sensory change in the early stages of the injury and in several instances well marked evidences of healing had appeared before the study was begun. It was especially in those cases in which the loss of function of the musculospiral nerve was dependent on fracture of the humerus, with the development of scar tissue and which presented a cessation of function but not necessarily an anatomic loss of continuity, that the picture was at times

* Read before the meeting of the American Neurological Association, held in Atlantic City, N. J., June 16-18, 1919.

1. Trotter and Davis, *Brain* **38**:137 (Feb. 9) 1919.

greatly altered, for in from three to five or six months that had usually intervened before the study was made, marked improvement could, and often had occurred.

In many cases the amount of destruction of tissue, in addition to the actual section of the nerve, has seriously interfered with the work, often by the injury or section of nerves, especially of the musculo-cutaneous, in addition to the one desired to be studied, thereby confusing the sensory picture. So far as possible, however, cases were chosen for special study in which the section of the nerve appeared to have been at least physiologically complete, in which a sufficient period of time for healing to occur had not intervened, in which the wound was small and appeared not to have involved nerves other than the musculospiral and in which the intelligence of the patient was such as to permit a reasonably reliable study. It must be admitted, however, that when all these restrictions were applied, the number of cases suitable for study was so much reduced that the limits were more than once overstepped.

The total number of musculospiral cases studied was fifty-five, and in all of these the investigation was carried far enough to determine the area of sensibility to pressure, pin prick and cotton or camel's hair touch, but beyond these routine tests, the type of study made varied so much and in many of the cases is still so incomplete that no absolute percentages of the difference findings can be given.

THE SENSORY BRANCHES OF THE MUSCULOSPIRAL NERVE

There are three sensory branches of the musculospiral nerve, distributed in the average case as follows:

The internal cutaneous branch is given off in the axillary fossa and supplies the internal part of the posterior surface of the upper arm below the deltoid and as far down as the point of the elbow. On account of its high origin, this branch is affected in only a small number of cases, in which the musculospiral nerve is injured at a very high point. The external cutaneous branch is given off at a point where the musculospiral nerve passes around the external edge of the humerus and supplies the external part of the posterior surface of the arm, the lower part of the outer surface of the arm and a more or less narrow band of the posterior surface of the forearm lying between the musculocutaneous and the internal cutaneous nerves. Often there are two external cutaneous branches, and in such cases the area of numbness will be different if only one branch is involved. The superficial radial nerve has its origin at, or just below, the elbow joint, and after passing down the forearm is distributed to the skin of the external part of the dorsum of the hand, to a variable but

usually small part of the thenar eminence, to the dorsal surface of the thumb, to the dorsal surface of the first phalanx of the index finger and to the outer half of the dorsal surface of the first phalanx of the second finger (Fig. 1).

With so wide an anatomic distribution and with its well known extensive motor supply, one might expect a very widespread and complete loss of sensation in the various injuries to which the musculospiral nerve is subjected. As a matter of fact, quite the contrary is true, and the area of sensory loss is sometimes surprisingly small and is almost always incomplete.

Fig. 1.—Area of sensory distribution of the musculospiral nerve as given by Tinel in *Nerve Wounds*, p. 103.

One of the unexpected findings in all nerve injuries is the failure of the individual to recognize subjectively and with any degree of accuracy the area of sensory loss, especially of epicritic loss and particularly so in the area of distribution of the musculospiral nerve. Frequently patients with well marked paralysis of the extensors of the hand, following injury, if asked for their subjective sensory area will say they have none. If their attention is insistently brought to the matter, they will tell of a small area of numbness over the first interosseous space and over the dorsum of the thumb and perhaps over the dorsum of the first phalanx of the index finger, and they will be greatly surprised when a larger area of sensory loss can be demonstrated.

As already explained, the internal cutaneous branch rises so high that it is seldom involved in injury of the musculospiral nerve and

even the external cutaneous frequently rises above the area of injury, at least its supposed area is often found not involved even in injuries high up in the arm. It is impossible, however, to explain all the variations found in the area of numbness on the basis of the point of origin of the sensory branches, and other explanations are offered, chiefly that of the overlapping or anastomosis of the musculocutaneous, internal cutaneous, and median nerves involving the musculospiral field and giving varying areas of sensory loss when these nerves are injured. In the upper arm the intercostohumeral or the lesser internal cutaneous may supply areas ordinarily innervated by the internal cutaneous branch of the musculospiral and may also supply that part of the upper arm ordinarily supplied by the external cutaneous branch. I have also recently seen a case in which a lesion of the circumflex nerve caused a large area of numbness in that part of the upper arm ordinarily supplied by the external cutaneous branch of the musculospiral nerve.

As is so common after severance of any large nerve of the arm, the patient will usually complain of complete loss of feeling in all parts of the arm distal to the wound, directly following injury to the musculospiral nerve. During this preliminary period the arm is described as a very heavy appendage, or the statement is made that the arm feels as if wholly separated from the body and must be seen or felt by the other hand in order that the patient may appreciate its presence. After the lapse of a few hours or days, the patient discovers that there is feeling in the parts not supplied by the nerve immediately injured and the area really involved is more or less clearly recognized.

In the fifty-five cases that I studied there was only one instance in which the musculospiral nerve alone had been severed sufficiently high to produce a loss of sensation corresponding to all three branches.

REPORT OF CASES

CASE 1.—The previous history was not noteworthy. October 13, 1918, the patient while holding his left arm in the position of guard with the forearm horizontal, was struck below the clavicle by a shell fragment which punctured the lung and passed out the back. His left arm immediately became numb and fell to his side. There was no jerking or twitching. He felt as if the arm were shot off. His neck was stiff and he was unable to look down. He repeatedly asked those about him if his arm were not really gone. For some time after the injury he would seem to feel his arm in the position of guard and would reach up with his other arm to scratch it. At first he had excessive pain in the shoulder; later his whole arm ached a good deal, but he could not localize the ache as any worse in one part than in another. There was a slight causalgia for a time in the median. He was told that his musculospiral nerve was sutured directly after his injury, but the record is not clear on this point. At the examination, in February, 1919, he was conscious only of some numbness on the dorsum of the index finger, which he thought worse

in the first phalanx than in the second, and of some numbness on the back of the upper arm. An examination made with considerable difficulty, at this time, showed sensory changes as seen in Figure 2. Since February there has been some improvement in the sensory and motor condition and on May 31 even epicritic sensibility had returned over the dorsum of the hand with the exception of the area over the first metacarpal bone. The analgesic islands, outlined in the diagram, persisted. The dorsum of the first phalanx of the index finger was anesthetic and hypalgesic. Tinel's sign was elicited as far as the elbow and at this point stroking produced a tingling feeling in the fingers.

In a general way this case conforms fairly well to the typical distribution of the musculospiral nerve. The fact that sensibility is more disturbed in the upper part of the forearm than at the wrist was a

Fig. 2 (Case 1).—Area of sensory disturbance. In this and in the accompanying illustrations the meaning of the legends employed is given as a part of each illustration. When lines lie close together it is not intended that they should represent the relative limitations of the different sensory fields. They were not permitted to overlap merely as a matter of clearness of representation. In most cases also the work was not sufficiently carefully done to permit the accurate tracing of the relative sizes of the different fields.

frequent finding in our cases. The enlargement of the sensory area at the wrist and just above is atypical, though found in one other case, and on stroking touch the patient was able to identify an area of decreased sensibility even to the extreme ulnar side of the arm, just above the wrist joint.

In the fifty-five cases there were twenty-seven with a definite area of sensory involvement in the forearm as well as the hand, and one

instance of involvement of both musculospirals, each case involving the forearm. Seven of these were rejected on account of various complications. The subjective sensory phenomena were usually about as in Case 1, with an initial sensation much as if a heavy blow had struck the arm and a subsequent feeling of loss of the limb and a still later more clearly defined and localized loss of sensibility. The following is a fairly typical case except that there seems to have been less initial loss of sensibility than was ordinarily present.

CASE 2.—While standing the patient was struck by a rifle bullet in the upper portion of the left upper arm, Sept. 15, 1918. The arm was in a horizontal position, and he had the sensation of being struck a heavy blow. The arm rotated inward, bringing the palm downward. The fingers flexed like a claw

Fig. 3 (Case 2).—Area of sensory disturbance.

and remained so for some time. There was no pain except at the first blow and no feeling of the arm being gone. He did not recall noticing much numbness at any time. His record states that he was operated on the night of the 15th and his musculospiral nerve sutured.

When examined, May 7, 1919, a rather extensive area of sensory disturbance was found (Fig. 3). There was no connection between the area on the forearm, which extended well over into the supposed region of the musculocutaneous nerve, as did some of the others, and the area on the hand. There was a loss of sensibility to cold at 2 C. slightly wider than the area of touch loss on the forearm and especially extending considerably farther over on the palmar surface of the forearm than any other form of loss of sensibility. Two point discrimination was greatly decreased on the dorsum of the forearm (33 per cent. of correct answers at 110 mm. with 53 per cent. of correct answers at 50 mm. on the right forearm). On the dorsum of the hand two point sensibility could not be measured on account of the inability to perceive pressure.

Joint sensibility was normal in the fingers and thumb. Vibration sense was normal in the fingers and somewhat reduced over the thumb.

CASE 3.—While kneeling and digging in, the patient was struck by shrapnel in two places on the left upper arm and in six or seven places on the right upper arm, Oct. 5, 1918. He was also struck in the right knee and thigh. He fell and was unconscious at least a few minutes. On recovering consciousness, both hands and arms were wholly paralyzed for motion and sensation and remained so for one day. The left hand and forearm felt numb on the dorsum until the end of January and then began rapidly to improve. There was some numbness in the musculospiral area of the right hand and lower part of the forearm for two weeks with rapid improvement following.

When examined, May 9, 1919, there was a slight hypesthesia and hypalgesia in the right hand with a subjective area practically the same in extent (Fig. 4).

Fig. 4 (Case 3).—Area of sensory disturbance.

May 28, 1919, it was impossible to map any definite area of disturbed sensibility in the right hand or arm.

May 9, 1919, there was an extensive area (Fig. 5) of diminished touch and pain sensibility on the posterior surface of the left hand and arm, most marked on the forearm. May 28, 1919, the area was considerably diminished and could be outlined in anything like its original form only by stroking touch. There was no loss of appreciation of weight but two point discrimination was definitely lessened on the dorsum of the forearm and a small area of somewhat lessened appreciation of cold at 4 C. was found on the dorsum of the forearm a little below the elbow.

CASE 4.—The patient while advancing was struck by shrapnel in the left upper arm and left leg, Nov. 1, 1918. His arm felt dead but he could move it at the elbow directly after receiving the wound. He could not move the fingers; the arm did not jerk or twitch. The next day he noticed a definite numb area on the back of his left hand, forearm and arm. He could move the arm at the shoulder but not at the elbow or wrist. The fingers could be feebly flexed.

Fig. 5 (Case 3).—Condition, May 9, 1919.

Fig. 6 (Case 4).—Area of sensory disturbance.

When examined May 13, 1919, he had a large scar, 6 by 2½ inches, over the lower end of the outer side of the left upper arm, adherent to the bone. The area of sensory disturbance (Fig. 6) extended down the back of the forearm in a rather narrow line and in the hand there was loss or great diminution of touch and pain over the first phalanx of the thumb but the skin over the first metacarpal and over the first interosseous space responded normally. In this case the loss of sensibility was more complete in the areas affected in the hand than in the forearm and in the lower part of the forearm than in the upper.

A further examination, May 28, 1919, showed an area of complete loss of sensibility to temperature of 5 C. and to ice over the region of greatest sensory loss on the hand, but on the thumb similar temperatures were fully appreciated. Joint sensibility was normal. Pressure sense was greatly diminished on the hand (200 gm.) and much less diminished on the forearm (27 gm.). On the hand, in two point discrimination, he gave 47 per cent. of correct answers at

Fig. 7 (Case 5).—Area of sensory disturbance.

40 mm. as against 53 per cent. at 22½ mm. on the right hand. On the left forearm he gave 53 per cent. of correct answers at 80 mm. and on the right forearm 53 per cent. of correct answers at 47 mm.

In July he went to the beach and both forearms became much sunburnt. The left was a little more affected than the right and almost blistered, but the two felt equally sore and hot.

CASE 5.—Sept. 27, 1918, while walking, the patient was struck by a machine-gun bullet in the right leg and by a high explosive fragment in the left upper arm. The humerus was fractured. The left hand dropped immediately at the wrist and there was subsequently no movement at the wrist joint, though he could flex the fingers slightly. The hand and arm from the elbow down were numb on the dorsal surface, but he retained the sense of feeling on the palmar side and there was no pain except at the site of injury.

When examined, May 28, 1919, the ordinary line of numbness (Fig. 7) was found on the posterior surface of the forearm and thumb and the first and part of the second phalanges of the first finger were involved, but the skin

Fig. 8 (Case 5).—Condition, June 30, 1919.

Fig. 9 (Case 6).—Area of subjective numbness.

over the first metacarpal and a part of the first interosseous space was unaffected. Pressure sense was much more affected in the hand and wrist than in the middle and upper part of the forearm, whereas two point discrimination was much more deficient in the upper part of the left forearm than in the wrist or hand. The area of loss to cold was wider than that of loss of touch and pain. There was no diminution of vibration or joint sensibility.

On June 30, 1919, the loss of sensation was about the same on the dorsum of the hand and upper two thirds of the forearm, but had quite disappeared over the wrist and lower one third of the forearm (Fig. 8).

CASE 6.—Oct. 14, 1918, the patient, while standing, was struck by a fragment from a shell which exploded about 3 feet from him. He sustained a severe compound, comminuted fracture of the left humerus. He bled considerably and was unconscious for three or four days. His arm was heavily bandaged when he regained consciousness. His idea of the sensory loss was not exact,

Fig. 10 (Case 6).—Area of sensory disturbance.

but he was sure that at least he had had numbness of the back of the left thumb, of the proximal part of the left index finger and did have feeling somewhere toward the ulnar side of the hand. There was complete wrist drop.

When examined, May 5, 1919, the area of subjective numbness (Fig. 9) included a large portion of the dorsum of the left forearm, but he was not aware of any loss over the hand. The examination showed (Fig. 10) a definite area of anesthesia and analgesia of the dorsum of the thumb and of the wrist and anesthesia of practically the entire dorsal region of the forearm and of the lower part of the upper arm with a less wide area of analgesia. On the volar surface of the forearm, toward the radial side, there was hypesthesia and over the outer side of the thenar eminence and thumb there was anesthesia and analgesia. There were small areas of hyperalgesia over the distal phalanx of the thumb and over the knuckle of the index finger.

The involvement of the radial side of the forearm, in this case, suggests the possibility that the musculocutaneous was injured, but it may at least be said

Fig. 11 (Case 6).—Condition, June 1, 1919.

Fig. 12 (Case 7).—Area of sensory disturbance.

that the injury was not in such a position as to be likely to involve the musculocutaneous, unless it did so above its supply to the biceps and there was no evidence of any motor involvement to the biceps.

Examination, June 1, 1919, showed (Fig. 11) hyperesthesia and hyperalgesia and loss of cold at 5 C. over the area of the hand previously showing sensory loss, no evidence of any sensory involvement of the dorsum of the wrist and a decided diminution of the area on the forearm.

CASE 7.—The patient was struck by a machine-gun bullet, Sept. 29, 1918, causing compound, comminuted fracture of the left humerus. Following the injury there was extreme pain in the whole arm and considerable hemorrhage. He does not recall whether there was loss of feeling or motion following the injury. He was operated on thirty-six hours later, and is sure there was no paralysis of motion following the operation. The arm was placed in a Thomas splint with extension. Three weeks later he noticed that wrist drop was

Fig. 13 (Case 8).—Area of sensory disturbance.

developing and that the end of the thumb was growing numb. The loss of motion and sensation increased for a time, but he thinks that during the past few months there has been progressive improvement.

When examined, Aug. 19, 1919, a wide area of sensory change was found (Fig. 12). Sensibility was much less disturbed in a narrow line over the first interosseous space than in the areas to either side.

CASE 8.—Sept. 30, 1918, while standing, the patient was struck on the outer side of the right upper arm and the bullet came out on the inner side. Directly following the injury, he had no power of sensation or motion in the right arm or hand. Seven hours after the injury he was operated on and the musculospiral nerve sutured. Just before the operation, he found he could flex the right fingers feebly. After the operation there was numbness down the outer side of the right arm, on the dorsum of the forearm and on the radial side of the hand.

Fig. 14 (Case 9).—Area of sensory disturbance.

Fig. 15 (Case 10).—Area of sensory disturbance.

April 8, 1919, examination showed (Fig. 13) an extensive loss of sensation on the dorsum of the hand and forearm and on the outer side of the upper arm. Diminution of pain and touch was much more marked distally than proximally in the forearm and more marked in the hand than in the forearm or arm. Vibration and joint sensibility were normal but two point discrimination was diminished. The mental condition of the patient did not permit a very satisfactory examination.

CASE 9.—Sept. 29, 1918, the patient was struck by a piece of shrapnel on the dorsal surface of the right arm just below the elbow. The fragment passed out through the ulnar surface. Directly after the injury was received the entire extremity from the elbow down felt numb and "tingled as if a nerve had been struck." He was operated on the following day, but thinks no nerve was sutured. Subsequently he found about the same area of numbness on the dorsum of the arm and hand as at present, except that it was slightly larger. At an examination made May 26, 1919, it was found that the loss of sensibility (Fig. 14) was much more complete on the dorsum of the forearm than on the hand, and there was an area of normal sensibility over the first metacarpal and the first interosseous space. There was no perceptible loss of sensation on the index finger, but the dorsum of the first phalanx of the second finger showed hypesthesia. Vibration sense was practically normal.

CASE 10.—While lying down, Oct. 24, 1918, the patient was struck on the outside of the left upper arm by a machine-gun bullet. He felt "as if a hot poker were being run through it and the arm felt numb from the wound down." When dressed he could feel the bandages about the palm of the hand and the ulnar part of the wrist but not anywhere on the forearm. Feeling began to return in the forearm about the middle of November. When examined, June 4, 1919, there was a large area on the dorsum of the forearm (Fig. 15) of very marked sensory disturbance. Anesthesia was wider than analgesia. The margin of stroking outline lay outside both and the line of thermoanesthesia and of thermohypesthesia (8 C.) was still wider. The dorsum of the thumb showed anesthesia and analgesia, but the dorsum of the first metacarpal and of the first interosseous space showed hypesthesia and, over part of the former, hypalgesia. Vibration sense was reduced on the thumb. In two point discrimination he gave 53 per cent. of correct answers at 25 mm. in the right forearm and 53 per cent. of correct answers at 70 mm. in the left forearm.

CASE 11.—The patient was struck in the left upper arm Nov. 7, 1918, by a machine-gun bullet which caused a compound, comminuted fracture at the middle of the humerus. There was immediate and complete motor paralysis of the musculospiral nerve, but he is uncertain as to the sensory condition. Examination, April 24, 1919, revealed anesthesia and analgesia (Fig. 16) of the area on the second metacarpal and the second interspace and hypesthesia and hypalgesia on the dorsum of the thumb and the first metacarpal with a narrow band of normal sensibility between them. There was also hypesthesia and hypalgesia over a band of considerable width extending down the dorsum of the forearm. June 21, 1919, the area on the forearm was considerably reduced. The condition of the hand was not much altered except for hyperalgesia over the thumb and an increase in the size of the normal band over the first interosseous space. Aug. 16, 1919, no sensory loss could be discovered in the forearm, and the hand was greatly improved (Fig. 17).

CASE 12.—While the patient was asleep, Oct. 15, 1919, a bullet passed through the left forearm, just below the elbow. Directly the entire extremity was numb from the elbow down. Feeling later returned in all parts except in a line down

Fig. 16 (Case 11).—Area of sensory disturbance.

Figure 17

Figure 18

Fig. 17 (Case 11).—Area of sensory disturbance, Aug. 16, 1919.

Fig. 18 (Case 12).—Area of sensory disturbance.

the center of the back of the hand and forearm. Examination, July 10, 1919, showed hypesthesia and analgesia in the area shown in Figure 18 and a partial paralysis of the musculospiral motor supply.

CASE 13.—Nov. 1, 1918, the patient was struck on the inside of the left upper arm by a bullet which was removed on the upper side of the upper arm. So far as the patient knows, there was no motor or sensory loss following this operation, but about March 1, 1919, a sequestrum was removed from the upper arm and subsequently he had difficulty in extending the fingers and thumb. When examined, June 25, 1919, there was still no extension of the thumb or fingers of the hand, but extension of the wrist was fair. On the dorsum of the forearm was an area $1\frac{1}{4}$ by $1\frac{1}{2}$ inches (Fig. 19) of loss of sense of touch. In this area pin prick was described as "less sharp," but as "hurting more," with a feeling of being different from a normal pin prick. No other sensory disturbance could be found.

Fig. 19 (Case 13).—Area of sensory disturbance.

There were twenty-seven cases in which the sensory phenomena, due to injury of the musculospiral nerve, were limited to the hand or to a region extending slightly above the wrist. Of these, fourteen were selected for reporting.

CASE 14.—While riding a motorcycle, Nov. 9, 1918, the patient's left arm dropped suddenly to his side. He was not conscious of having been injured, and there was no pain or jerking of the hand or arm. An examination showed he had been struck in the left arm by a bullet. He could flex his fingers, and he is sure that the dorsum of the thumb and of the first finger and the space between were numb and that he could not appreciate pin prick in this region as pain.

When examined, April 17, 1919, only an area on the dorsum of the first phalanx of the thumb and a small area on the outer surface of the dorsum of the first phalanx of the index finger felt numb. Examination showed (Fig. 20) analgesia to 12 gm. of pressure over the knuckle of the index finger and to

Fig. 20 (Case 14).—Area of sensory disturbance.

Fig. 21 (Case 15).—Area of sensory disturbance.

10 gm. of pressure over a band extending back about half way to the wrist. The dorsum of the first and second phalanges of the thumb was analgesic to 12 gm. of pressure. Both areas were anesthetic to touch as was also an area extending proximally from the knuckle almost to the wrist. He gave two point discrimination in 53 per cent. of tests over the normal hand at 21¼ mm. and in 53 per cent. of tests over the affected area at 47 mm. On the area over the knuckle, he could barely discriminate 25 gm. of pressure. With the exception of this region, discrimination of pressure was not greatly reduced in the affected area.

CASE 15.—The patient was injured Oct. 3, 1918, by a machine-gun bullet entering the right arm at the elbow. He was sitting with his elbow on his knee. He felt as if a "log had struck him" and was knocked over but did not become unconscious. There was no twitching or jerking in the hand or arm. The dorsum of the thumb and of the back of the hand (with the exception of the extreme ulnar side) was numb. He cannot recall anything concerning the feeling in the back of the fingers, but does not remember any numbness, and he is positive there was no numbness in the forearm. There was some stinging feeling in the back of the hand for a time, but this had disappeared at the examination on April 18, 1919, when he showed an area of anesthesia and analgesia as outlined in the diagram (Fig. 21).

A second examination on June 10, 1919, showed some improvement in sensation. Vibration and joint sensibility at this time were normal. Cold at 5 C. was readily felt. In discrimination of two points he gave 53 per cent. of correct answers at 45 mm. in the right hand and 100 per cent. at 26 mm. in the left hand.

CASE 16.—Aug. 10, 1918, while lying flat on his face, the patient was struck by a fragment of shell which entered the left arm 2 inches below the elbow on the radial side and passed out through the posterior surface of the arm just above the elbow, fracturing the humerus. Directly after he was struck, the hand and arm drew up and the "fingers clenched," after which the entire arm became limp and numb. He felt as though the arm from the elbow down did not belong to him, and he could make no movement from the hand or forearm.

When examined, May 23, 1919, there was an area (Fig. 22) of anesthesia and hyperalgesia over the phalanges of the thumb and of anesthesia and analgesia over the second metacarpal bone and second interspace and over most of the first interspace. Over the first metacarpal bone there were hypesthesia and hypalgesia. Sensibility to cold at 2 C. was lost in all this area and over a considerable area over the thenar eminence. Cold at 22 C. was lost over a somewhat wider area. The threshold for two point discrimination was about one-half higher in the affected arm than on the unaffected side. Weight discrimination was very little changed. In the first, second and third fingers there was an area of uncertain loss to touch, probably due to local skin condition.

CASE 17.—July 16, 1918, a fragment of shell entered over the lower end of the left biceps and passed out a little above the outer condyle of the humerus. When examined, May 26, 1919, there was anesthesia (Fig. 23) over the entire dorsum of the thumb and over the radial side of the dorsum of the hand with the exception of a band about 1 inch wide extending down the first interspace. Pin prick was appreciated as such at from 4 to 8 gm. of pressure; two point and weight discrimination were both only slightly impaired in the same region. There was no loss of joint or vibration sensibility. An area of hyperalgesia was found over the web of the thumb.

Fig. 22 (Case 16).—Area of sensory disturbance.

Fig. 23 (Case 17).—Area of sensory disturbance.

CASE 18.—The patient was injured in the left arm, on the outer side, just above the elbow by a piece of shrapnel, Oct. 4, 1918. He was knocked down, and remained unconscious for a few minutes. On recovering, he felt as though the arm had been blown off. When he attempted to move it, it hung limply by his side, and he said it "felt as if it weighed a ton." The next morning there was numbness in the area shown in the diagram and over the dorsum of the first finger. He is positive there was no numbness in the arm. He noticed improvement first in the distal part of the forefinger and then in the region of the first metacarpal.

Fig. 24 (Case 18).—Area of sensory disturbance.

Fig. 25 (Case 19).—Area of sensory disturbance.

When examined, May 12, 1919, a roentgenogram showed some small fragments of bone still in the wound and in the region of the musculospiral nerve. There was an area of anesthesia to touch and analgesia to prick over the second and third metacarpals as shown in Figure 24 and outside of this there was an area of hypesthesia and hypalgesia. The dorsum of the phalanges of the thumb and of the distal part of the first metacarpal was anesthetic and analgesic. In the first interosseous space was an area of normal sensibility.

CASE 19.—Oct. 29, 1918, the patient was struck in the left arm just to the outer side of the elbow. The missile passed out at a slightly higher level on the posterior surface of the upper arm, fracturing the humerus. His arm was immediately paralyzed from the elbow down and subsequently there was an area of "numbness" (Fig. 25) involving the dorsum of the first phalanx of the thumb, the distal part of the first interosseous space, a band over the second metacarpal and the first phalanx of the forefinger. Over and just proximal to the knuckle of the second finger was a circular area of subjective hyperesthesia. Examination, May 16, 1919, showed (Fig. 26) hypesthesia and hypalgesia over the first phalanx of the index finger, analgesia and hypesthesia over the thumb and anesthesia and analgesia over the remainder of the area. The subjective hyperesthetic area gave normal response to touch and prick.

Fig. 26 (Case 19).—Condition, May 16, 1919.

CASE 20.—The patient was injured Oct. 4, 1918, by a machine-gun bullet. He was lying down, and the bullet entered at the lower part of the right deltoid on the outer surface, and came out on the inside of the arm just below the axilla. The whole arm felt numb subsequently, and motor power was completely lost. After a few hours the arm was less numb and began to feel "sore" but not in any special locality. At the same time the numbness largely disappeared from the upper extremity but for three weeks there was some loss of feeling in the forearm, not clearly localized. There has been no numbness since. After this there was not much change in feeling but marked improvement in motion.

When examined, May 13, 1919, he outlined an area of numbness on the back of the hand a little smaller than that found on examination and an area on the inside of the hand considerably greater than could actually be demonstrated. Examination showed (Fig. 27) anesthesia and analgesia over the dorsum of the phalanges of the thumb, over the first and part of the second

Fig. 27 (Case 20).—Area of sensory disturbance.

Fig. 28 (Case 21).—Area of sensory disturbance.

metacarpals, including the first interosseous space, with an area of hypesthesia and hypalgesia considerably wider. There was also hypalgesia over the first and second phalanges of the right index finger.

CASE 21.—The patient was hit by a machine-gun bullet, Oct. 16, 1918, in the upper part of the left arm, and the humerus was fractured. He felt no pain and did not know that he was hurt, until his arm, on which he was leaning, gave way. For one week the arm felt numb up to the shoulder and after three weeks there was no sensory loss of which the patient was aware other than the area shown on examination, June 5, 1919 (Fig. 28). The area of hypesthesia on the dorsum of the hand was wider than the area of hypalgesia. The dorsum of the second phalanx of the thumb was anesthetic and hypalgesic. The dorsum of the distal half of the first phalanx was anesthetic and of the proximal half hypesthetic, but he could feel pin prick in both places.

Fig. 29 (Case 22).—Area of sensory disturbance.

CASE 22.—The patient was injured, Nov. 1, 1918, in the right arm and forearm. The only part of the early history available, was that after the injury was received he had a wrist drop. When examined, May 2, 1919, he had slight extension of the wrist, a very slight extension of the fingers and none of the thumb. There was anesthesia and analgesia (Fig. 29) of the dorsum of the thumb, of the first phalanx of the index finger and of the radial side of the hand, extending upward a little distance above the wrist. Anesthesia was somewhat wider than analgesia. The area of stroking change was slightly larger than either and the subjective area was somewhat smaller. Sensation was preserved in a region along the edge of the web of the thumb.

June 9, 1919 (Fig. 30), sensation had improved decidedly. The dorsum of the thumb and of the second metacarpal remained anesthetic and analgesic but the area between was very slightly hypalgesic. Loss of sensation of cold at 8 C. and the stroking outline almost coincided. On the front of the right wrist was a region of increased sensibility to cold at 8 C.

Fig. 30 (Case 22).—Improvement in sensation, June 9, 1919.

Figure 31

Figure 32

Fig. 31 (Case 23).—Area of sensory disturbance.

Fig. 32 (Case 24).—Area of sensory disturbance.

CASE 23.—The patient was struck by shrapnel, Oct. 9, 1918, in the lower one-third of the right upper arm, receiving a compound, comminuted fracture of the humerus with considerable loss of bone. When examined, May 16, 1919, there was considerable loss of substance of the shaft of the humerus, a complete motor paralysis of the musculospiral supply and an area of anesthesia and analgesia on the dorsum of the hand (Fig. 31).

CASE 24.—Oct. 14, 1918, a bullet passed through the right arm, just below the elbow, resulting in musculospiral paralysis. The history of the early symptoms is lacking. When examined, May 22, 1919, there was (Fig. 32) hypesthesia and hypalgesia of the dorsum of the thumb and of the radial side of the hand. Extension at the wrist was poor and of the thumb and fingers was entirely lacking.

CASE 25.—The patient was shot through the left elbow, Oct. 12, 1918. Musculospiral paralysis resulted. The early history was not available. When examined, June 30, 1919, there was fair extension of the wrist, fingers and thumb, but there was a large area of anesthesia and analgesia on the dorsum of the hand and an area of hypesthesia and hypalgesia on the dorsum of the thumb (Fig. 33).

Fig. 33 (Case 25).—Area of sensory disturbance.

CASE 26.—The patient was struck on the outside of the right arm just above the elbow, Oct. 8, 1918, by a machine-gun bullet which came out on the posterior and inner surface at about the same level. Motion was lost at once in the musculospiral supply and for a time the arm felt numb, but he thinks he could distinguish touch at all times. He recognized no complete loss of sensation. When examined, June 5, 1919, there was a small area of anesthesia (Fig. 34) about the knuckle of the right index finger and on the dorsum of the second phalanx of the thumb with hyperalgesia in both areas and inability to extend the thumb. There was also rather weak extension of the fingers.

CASE 27.—The patient was injured Oct. 8, 1918, by high explosive fragments which entered the left upper arm and forearm. The arm at once became limp and he felt as though he had lost it. There was no sensation at any point on the arm until the following day, after which sensation returned except in the area shown in the diagram. When examined, April 24, 1919 (Fig. 35), the entire dorsum of the thumb was anesthetic and there was a small area of analgesia on the distal part of the dorsum of the first phalanx.

CASE 28.—Oct. 15, 1918, the patient was struck by high explosive fragments at various points, including the front of the right axillary fold. The entire right arm and hand were paralyzed for motion and sensation for some hours

Fig. 34 (Case 26).—Area of sensory disturbance.

Figure 35

Figure 36

Fig. 35 (Case 27).—Area of sensory disturbance.

Fig. 36 (Case 28).—Area of sensory disturbance.

with subsequent gradual improvement. After one week the only sensory loss was over an area on the thumb about as large as at present. A musculospiral motor paralysis persisted for three months but at present his only marked motor loss is his inability to extend the thumb and index finger.

When examined, June 30, 1919, there were analgesia of the dorsum of the first and second phalanges of the thumb and anesthesia of the dorsum of the second phalanx (Fig. 36). No other sensory loss could be elicited.

SENSORY PHENOMENA

The literature in existence in reference to the sensory phenomena in musculospiral injuries is surprisingly limited, but in a general way there appears to be an agreement that the musculospiral is chiefly a motor nerve and that the sensory changes resulting from its injury are limited in extent and unstable in their location. Head and Sherren² even state, "It has long been known that division of the musculospiral in the region of the spiral groove causes no definite loss of sensation over the thumb or back of the hand," and work on the hand and arm was done by Head after division of the external cutaneous nerve as well as of the superficial radial.

Borchardt³ reports seventeen cases of injury of the musculospiral nerve. In six instances there is no statement as to sensory deficiency. In three cases there was definite sensory change and in the other eight cases the sensory condition is dismissed with such statements as: "Nothing of consequence in the way of sensory loss," "no true sensory disorders," or "a dull sensation in the fingers."

Tinel,⁴ in speaking of the subject, says: "Anesthesia in musculospiral paralysis is not often found anywhere else than on the hand. Seldom does it reach the typical distribution of the musculo-spiral, far more seldom does it go beyond it. Most frequently it limits itself to a very restricted part comprising the dorsal region of the first and the second metacarpal. It is often very slight, sometimes even it can scarcely be seen; in any case, it is exceptional to find complete anesthesia in this region; at most we have more or less marked hypesthesia."

Byrne's case⁵ showed both epicritic and protopathic loss over the dorsum of the thumb, part of the thenar eminence, the radial half of the dorsum of the hand and the lower wrist.

My own observations lead me to believe that though a lesion of the musculospiral nerve at any point produces less sensory disturbance

2. Head and Sherren: *Brain* **28**:177, 1905.

3. Borchardt: *Bruns Beiträge* **97**:253.

4. Tinel: *Nerve Wounds*. 1918, p. 107.

5. Byrne: *J. Nerv. & Ment. Dis.* **43**:489, 1916.

than one would expect considering its anatomic distribution, it is rarely, if ever, that some definite area of sensory loss may not be found if carefully sought.

Sensory phenomena, as appreciated by the patient, are not pronounced and are limited in extent. Usually he will tell of an area of numbness over the dorsum of the thumb and especially of the first phalanx, over the first interosseous space and perhaps over the first phalanx of the index finger and more or less over the distal portion of the second metacarpal. The area on the back of the forearm is often ignored even when by examination a large area of well defined sensory change can be demonstrated. In the one instance of well marked sensory change on the back of the upper arm that came under my observation, this was recognized by the patient. A possible satisfactory explanation for the failure of the patient so many times to recognize his sensory loss may be that the musculospiral nerve, limited in sensory fibers, supplies only epicritic and protopathic sensibility to the integument and that deep sensibility is supplied by other nerves. The patient, depending largely on some form of irritation that gives a response through deep sensibility, for his consciousness of a stimulus, fails to recognize the sensory loss until it is shown him by adequate methods of examination.

In any case, when sensibility is to be tested in musculospiral injury, the skin should be shaved, as was done in all the cases reported, and tactile sensation tested with cotton, wool or a camel's hair brush with the lightest possible touch. Pin prick is usually lost as pain in a similar or somewhat smaller area, but is ordinarily recognized as pressure since, as already stated, deep sensation is not commonly lost in musculospiral injury. Sensation to cold was usually impaired or lost in an area corresponding more or less to the loss for touch and prick and sometimes in a larger area. In some cases the failure to show in the diagram any loss of sense of cold is so out of keeping with the other phenomena that it can be explained only by lack of care on the part of the examiner or by lack of intelligence and attention on the part of the patient. In only one instance was there any excessive response to cold.

Sense of the position of the joints was not lost or notably impaired in any of my cases, but vibration sense was often impaired though not lost, especially on the thumb. Discrimination of two points as two was much diminished in most of the cases, but appreciation of weight was not greatly reduced even in cases with well marked loss of superficial sensibility.

One of the problems which I encountered was to determine the sensory loss in the forearm which can fairly be ascribed to the musculospiral nerve. In several of my cases in which there was no evi-

dence that the musculocutaneous nerve had been severed there was a marked loss of sensation over the area usually ascribed to that nerve. In two instances nearly all of the dorsum of the forearm was numb, but the volar surface was not involved. In such a case it is as reasonable to assume that because the volar surface was little or not at all involved the sensory loss could not be ascribed to any injury to the musculocutaneous or the internal cutaneous nerve as to assume that because the dorsal portion of the area usually ascribed to these nerves was involved, the nerves themselves must be assumed to be involved in the injury. In such cases as Cases 7, 8 and especially 12, it may fairly be doubted that the area of sensibility loss represents only musculospiral involvement; however, it must be understood that no case was included in this series in which the motor phenomena did not clearly demonstrate musculospiral lesion or in which the motor phenomena showed any additional material lesions. A full description of these motor phenomena could only have lengthened this paper.

In those cases in which some degree of numbness was found in the forearm, there was often an area over the dorsum of the wrist in which sensation was more nearly normal or wholly normal. In ten instances in the series reported there was a strip passing from the web between the thumb and first finger in a proximal direction, of normal reaction or of very slight sensory loss separating the more deeply affected area over the dorsum of the first and second metacarpals, respectively.

As a rule, loss of sensation, if present in the thumb, extended down to the nail, but in a few instances there was a narrow band of retained sensibility proximal to the nail, and it was a common occurrence to find the dorsum of the first metacarpal less involved than the phalanges of the thumb. On the first phalanx of the thumb loss of sensibility was ordinarily bounded by the hair line on the median side, but on the external side it extended over somewhat on the palmar surface. This was much more marked over the thenar eminence where the loss of sensibility extended in a curved line on to the palmar surface. Where no evidence of this loss over the thenar eminence appears in a number of the diagrams the explanation probably lies in a lack of care on the writer's part in recording the result of the examination rather than in its actual failure to appear. Where loss of sensibility existed in the area between the thumb and first finger and between the first and second fingers the line of demarcation extended well beyond the hair line and almost to the edge of the web.

I wish to acknowledge my indebtedness to Miss Ethel Cornell, Ph.D., and to Miss Margaret Stubbs for their very valuable assistance in the examinations which form the basis of this paper.