ABSENCE OF LOBUS OLFACTORIUS AND SCLEROSIS OF CORNU AMMONIS *

FUMIO TANAKA, M.D.

Professor of Otology, Okayama Medical School

OKAYAMA, JAPAN

Cases showing absence of the olfactory lobe are rare. In 1914 Weidenreich¹ reported one case and quoted nine others from the literature on this subject. Kundrat,² in his monograph, "Arhinen-cephalie," describes brains which have no olfactory lobe, but are combined with other brain anomalies. Besides these two reports I was able to find one case reported by Valenti.³ But according to Weidenreich all reports which he quoted lack detailed information, especially in that they do not discuss the area of the brain supposedly connected with the olfactory nerve and therefore called "rhinencephalon." Valenti's report contains no statement about this area.

In Weidenreich's case there was entire absence of the bulbus and tractus olfactorius in both hemispheres. We assume that the rhinencephalon governs the sense of smell as found by Broca and Zuckerkandl in their studies in comparative anatomy, and by Retzius embryologically. If this is true also in man, we should be able to see some change in the parts regarded as the olfactory center in brains that have no olfactory lobe. Weidenreich investigated carefully his case that dealt with the bilateral absence of the bulbus and tractus olfactorius, absence (right) and reduction (left) of the tuber olfactorius. Except for these defects, no parts of the brain regarded as the rhinencephalon showed any marked change which denotes atrophy; but he was not able to study his case histologically because of the unsuitable previous treatment of the material for this purpose. He concludes the question of the olfactory center with the sentence: "Auch die Lösung diesser Frage bleibt also aufgespart, bis der Zufall eine neue Beobachtung des Defectes bringt und zugleich die Möglichkeit einer sofortigen Untersuchung am frischen und entsprechenden fixirten Material."

---

* From the Pathological Laboratory of the Massachusetts Commission on Mental Diseases, 74 Fenwood Road, Boston, Mass., U. S. A.

1. Weidenreich: Ueber partiellen Riechlappendefekt und Eunuchoidismus beim Menschen, *Ztschr. f. Morphol. u. Anthropol.* **18**:157, 1914.

2. Kundrat; Arhinen-cephalie, 1882, abstr. in Schwalbe; *Die Missbildungen des Kopfes*, 1913.

3. Valenti: Un cas d'absence unilatérale de l'appareil olfactif, *Arch. ital. de biol.* **64**:457, 1911.

It is many years since Broca and Zuckerkandl announced that the gyrus fornicatus must be regarded as the cortical olfactory area (it must be understood that under the name gyrus fornicatus or rhinencephalon are included gyrus cinguli and gyrus hippocampi with cornu ammonis). Although I am inclined to believe that certain portions of the rhinencephalon govern the smell sense, the part that subserves this function is still unknown, especially in man.

The authors quoted are arranged according to the year of publication.

Examination of animals has convinced Ferrier (cited in Bechterew's paper) that the olfactory center is located at the lower extremity of the temporal lobe. Munk (also cited by Bechterew) seeks it in the neighborhood of the cornu ammonis.

Onodi⁴ concludes, after investigation of the hitherto published clinicopathologic reports concerning the sense of smell, that the olfactory center might be traced into the gyrus hippocampi and the uncus.

Ramón y Cajal,⁵ from a histological study of the brains of several animals as well as that of man, has reached the conclusion that the focus speno-occipitalis, the subiculum, the focus pra-esubicularis and the cornu ammonis seem to have no direct connection with the olfactory nerve fibers, and that they probably represent the tertiary olfactory center.

Campbell,⁶ chiefly from histologic study of the human brain, recognizes the cross relation between the lobus pyriformis (frontal pear-shaped part of the gyrus hippocampi) and the cornu ammonis, but he doubts whether Ramón y Cajal is right in regarding the latter as a tertiary olfactory center. One of his conclusions is "Histology supports comparative anatomy in suggesting that in the human brain the lobus pyriformis must be regarded as the principal cortical center, although-not the sole one governing the olfactory sense."

Broadmann⁷ distinguishes the gyrus hippocampi from the surrounding areas on the ground of the histologic construction and names it "area ento-rhinalis."

Bechterew⁸ concludes, from the results of experimental investigation on sixteen dogs, that an olfactory center is located apparently in

4. Onodi: Die Rindencentren des Geruches und der Stimmbildung, Arch. f. Laryngol. u. Rhinol. 14:74 1903.

5. Ramón y Cajal: Studien über die Hirnrinde des Menschen, No. 4, 1903.

6. Campbell: Histological Studies on the Localization of Cerebral Function, New York, G. P. Putnam's Sons, 1905.

7. Brodman: Vergleichende Lokalisationslehre der Grosshirnrinde in ihren Prinzipien dargestellt auf Grund des zellenbaues, Leipzig, 1909. Brodman: Feinere Anatomie des Grosshirns in Lawandowsky: Handbuch der Neurologie, Vol 1.

8. Bechterew: Die Funktionen der Nervencentra, 1911.

the lobus pyriformis, and that this must be regarded as the perception center for the sense of smell, while perhaps the neighboring region (subiculum cornu ammonis, Bechterew) may have the function of its conception.

Edinger⁹ seems to approve the cornu ammonis for this center chiefly from the comparative anatomic standpoint.

As I have mentioned above, our knowledge of the olfactory center is limited. Campbell declared after covering the histologic evidence:

Above all things we seem to lack the knowledge which in the case of some other cortical regions has proved so valuable. I refer to that derived from studies in pathological history; for instance, we have much to learn of the cortical changes attending uncomplicated atrophy or lesion of the olfactory bulb and peduncle, and have virtually no conception as to what subdivisions such changes would be distributed over.

Fortunately, through the generosity of Dr. Southard, I was permitted to investigate a brain that had no olfactory tracts. I will give the macroscopic and microscopic findings of this rare and interesting case, describing particularly those parts of the brain which have been suggested as having intimate connection with the olfactory sense. I shall endeavor to throw some light on the question of the olfactory center which seems to be so obscure.

REPORT OF A CASE

History.—F. S. H. 228, S. B. I. 1916. 87 (from the record of the Foxborough [Mass.] State Hospital), R. R., a man, aged 45, had epileptic dementia, the probable cause of which was heredity. The psychosis had been of gradual onset and of over fourteen years' duration, and had been characterized by periodic attacks of an epileptiform nature during which the patient lost consciousness, had generalized clonic convulsive movements, bit his lips and tongue and had loss of sphincter control. Following these attacks the patient was weak, befogged and depressed. He also had periods of depression and at times of excitement, auditory and visual hallucinations, with delusions of persecution. His judgment and emotional fields were markedly impaired, and there was general mental deterioration with no insight. He was admitted to the hospital April 17, 1907, and died July 13, 1916. Necropsy was performed by Dr. M. M. Canavan, eleven hours after death.

Anatomic Diagnosis.—This included brown atrophy of the heart muscle; ascites; calcified lymph node in mesentery; bladder distended; chronic perisplenitis; sclerosis of the mammary arteries; hydrothorax; hydropericardium; sclerosis coronaris; tricuspid and mitral stenosis; chronic fibrous endocarditis, mitral; chronic fibrous endocarditis, aortic; acute vegetative endocarditis, aortic; possible bronchopneumonia; pulmonary infarcts; accessory spleen; uric acid deposits, kidney (?); pigmentation, fibrotic spots; focal opacities, pia; absence of lobus olfactorius; soft cerebellum.

9. Edigen: Vorlesungen über den Bau der nervösen Zentralorgane, 1911.

Fig. 1.—The base of the brain entirely bereft of bulbus and tractus olfactorius; rudimentary development of trigonum olfactorium and sulcus olfactorius are noticeable.

MACROSCOPIC STUDY OF THE BRAIN

In general form and size the brain was about normal. The pia mater showed focal opacities in the frontal and motor regions, and was nonadherent. The gyri in general were of normal richness and appearance and almost symmetrical. On the base of the brain the absence of the bulbus and tractus olfactorius first attracted our attention (Fig. 7). There was slight sclerosis in the basal arteries. On careful stripping of the pia mater the findings in the rhinencephalon were: (The nomenclature in the following description has been adopted chiefly from Retzius.¹⁰)

Fig. 2.—Mesial view of right hemisphere of the brain (see the description of macroscopic findings).

Right Hemisphere.—The weight of the right hemisphere after hardening in formaldehyd was 604 gm. The bulbus and tractus olfactorius were absent. Corresponding to the area of the trigonum olfactorium a flat and small pyramidal pointed process could be seen; both its sides and tip were free but the brain substance beneath adhered to its base. The tip of this process rested in a rudimentary sulcus olfactorius; no filia was to be seen. A rudimentary sulcus olfactorius ran forward from the top to the process described above

10. Retzius: *Das Menschenhirn*, 1896.

for about 1.7 cm., and divided into two short transverse branches. As far as this sulcus extended a gyrus rectus could be distinguished but at the end of the sulcus the gyri were bounded by a short transverse gyrus. Gyrus tuber olfactorius, gyrus olfactorius medialis and lateralis were not definite. Along the posterior ends of the gyrus rectus and gyri frontales a narrow convolution, a continuance of the area parolfactorici on the mesial surface, ran transversely into the limen insulae and separated the surface connection between the frontal lobe and the anterior perforated substance. The space of the latter was narrow. No stria of olfactoria was to be seen. Sulcus parolfactorius anterior connected upward with the sulcus cinguli, and by the sulcus

Fig. 3.—Mesial view of left hemisphere of the brain.

parolfactorius we could distinguish area parolfactoria and gyrus subcallosus, both pointing upward to gyrus cinguli (Fig. 2). The gyrus hippocampi was bounded anteriorly by incisura temporalis, above by the fissura rhinica in its anterior portion, and by sulcus collateralis on the posterior border; its length from the frontal tip to gyrus rhinencephalolingualis was about 4 cm. and seemed a little atrophied. From the inner sides of the hemisphere in which the crus cerebri was cut off could be seen clearly the whole length of the fascia dentate and the fimbria, and a part of the fornix through the dorsal surface of the gyrus hippocampi, with no pressing up of the thalamus. The gyrus semilunaris, sulcus semiannularis, gyrus ambiens and sulcus rhinica

inferior, which Retzius noted especially in the gyrus hippocampi, were easily identified. The uncus was well developed and bent sharply against the gyrus hippocampi as usual. The gyrus fusiformis was separated from the former gyrus by the fissura rhinica in its anterior portion and posteriorly by sulcus collateralis, with the exception of a narrow communication, gyrus rhinencephalofusiformis, located between the former fissura and the latter sulcus.

Figure 4

Figure 5

Fig. 4.—Section of the cornu ammonis (right), which is atrophied and the layer of gray matter remarkably narrowed. The deep branch of the alveus (*d*) which spreads into the terminal plate (into fascia dentate) is shortened and stains poorly (see also Figure 15). Compare this with Figure 5.

Fig. 5.—Section of about the same region of a normal cornu ammonis. Kulschitzky-Wolter's modification of Weigert's myelin sheath stain; thickness, 15 microns.

At its posterior extremity the gyrus hippocampi joined with the gyrus cinguli by gyrus rhinencephalolingualis, with gyrus cinguli and also with the precuneus by the isthmus cinguli as well as by precuneus anterior. The sulcus cinguli as the continuation of the sulcus parolfactorius anterior at its frontal end ran upward then backward around the genu corporis callosi and joined with the sulcus paracentralis, which separated lobulus paracentralis and precuneus. The gyrus cinguli was well developed and separated from the neighboring gyri by sulcus cinguli and subparietalis, from the corpus callosum by sulcus

Figure 6

Figure 7

Fig. 6.—An area in the dorsal plate of a section of the cornu ammonis (left); cell layer is remarkably narrowed, pyramidal cells are reduced in number and atrophied; thickened walls of blood vessels are also noticeable. Compare this with Fig. 7.

Fig. 7.—The same area as in Figure 6 of about the same region of a normal cornu ammonis. (Cresylviolet stain in paraffin section after formaldehyd and alcohol fixation; thickness 6 microns.)

corporis callosi. Its frontal end communicated with area parolfactoria and gyrus subcallosus; its posterior part was joined to the gyrus hippocampi. In the commissura anterior there was no notable change.

Left Hemisphere.—The weight of the left hemisphere after hardening in formaldehyd was 607 gm. The bulbus and tractus olfactorius were absent. In the place of the trigonum olfactorium there was a process of pyramidal form about the size of a kernel of corn, but it was not so flat as that of the right hemisphere; its tip was slightly free but the other parts adhered to the frontal lobe beneath. Gyrus tuberculi olfactorii, gyrus olfactorius medialis and lateralis could not be differentiated; neither could the filia nor the stria olfactoria. The gyrus rectus could not be differentiated from the gyri frontales, and a sulcus olfactorius did not exist. The space of substantia perforata anterior was narrowed and separated from the orbital surface by a sulcus which ran from the mesial surface as the continuation of sulcus parolfactorius posterior. The gyrus hippocampi was 5 cm. long from its frontal tip to the gyrus rhinencephalolingualis and was bounded from the gyrus fusiformis by the sulcus collateralis. The fissura rhinica, gyrus semilunaris, sulcus semiannularis, gyrus ambiens and sulcus rhinica inferior could all be recognized. Generally the gyrus appeared to be a little atrophied, especially on the dorsal surface near the fissura hippocampi, where the convolution flattened and clear strands of the fascia dentata, fimbria and a part of fornix were to be seen from the inner sides of the hemisphere (Fig. 3). The sulcus cinguli, beginning as the continuation of sulcus parolfactorius anterior at its frontal end ran as usual almost parallel to the corpus callosum and combined posteriorly with the sulcus paracentralis. Gyrus cinguli, bounded superiorly by the sulcus cinguli and sulcus subparietalis, and below by the sulcus corporis callosi, was well developed and its posterior end reached to the isthmus rhinencephalolingualis as usual. The anterior end of the gyrus cinguli joined below to the area parolfactoria and gyrus subcallosus, which were as well developed as those of the right hemisphere.

HISTOLOGIC STUDY

Although there is no definite proof of the location of the olfactory center in the human brain, it is not difficult to assume that either the gyrus hippocampi or the cornu ammonis may govern the function. Therefore I made it my chief object in the microscopic study to examine these two regions. I took out the whole gyrus hippocampi with the cornu ammonis and the fascia dentata from both hemispheres. I investigated the fine construction and the lamination of nerve cells by the cresyl violet stain in paraffin sections, the arrangement of nerve fibers by Kulschitzky-Wolter's modification of Weigert's myelin-sheath stain in celloidin sections.

To obtain exact results I prepared as many serial sections as possible, and in order to avoid any misjudgment by comparison I investigated by the same methods the same regions of four other specimens which I had obtained from four normal looking hemispheres. I also examined other important areas of the brain.

Gyrus Hippocampi.—Cajal and Campbell have studied the normal histology of the gyrus hippocampi in detail. I state the findings in my case chiefly according to the topographical subdivisions by Ramón y Cajal, at the same time taking into consideration Campbell's description.

(a) The subiculum plexiform layer was about of normal thickness; deep in this layer islets composed of numerous minute triangular cells were seen as usual. Four or five of them were found in a section of the frontal area but much fewer in the posterior area. The pyramid and polymorphous cell layers were narrowed and many nerve cells, especially the pyramidal cells, were moderately shrunken with more or less affected nuclei. There was no change in the lamination and the cells were arranged in regular parallel rows as usual. Glia cells seemed to be increased.

(b) Praesubiculum: The tangential fibers in the plexiform layer ran irregularly; some were swollen and pear-like in appearance. The thickness of the gray matter beneath was reduced and many nerve cells, especially the pyramid cells, were moderately shrunken, but we were able to recognize the small pyramid and fusiform cell layer, the deep plexiform layer, the medium sized and large pyramid cell layer and the fusiform and triangular cell layer, as described by Cajal.

(c) The Hippocampal Gyrus Proper: The tangential fibers in the plexiform layer ran irregularly and were much fewer than normal. In the layer containing a cluster of polymorphous cells, the clusters of large polymorphous cells which characterize this area of the gyrus were to be found at the lobulus pyriformis, less toward the posterior end, but their protoplasm as well as their nuclei were markedly shrunken. The deep plexiform layer or medium sized pyramid cell layer, the large pyramid cell layer and the fusiform and triangular cell layer could all be differentiated, but many nerve cells, especially the pyramid cells, were considerably shrunken. As a whole, the cortex was narrowed. In Weigert's slides, there was no notable change in the arrangement of the radial fibers in the gray matter, but the interradian fibers among the former were few in comparison with those in brains having olfactory tract intact.

Cornu Ammonis and Fascia Dentata.—The histology of these organs in the human brain has been studied carefully by Kölliker,¹¹

11. Kölliker: Handbuch der Gewebslehre, 6 te Aufl. 1896.

Fig. 8.—A gross gap of cell layer in the cornu ammonis (left). The pyramid cells have entirely disappeared in this lesion and it is filled up by neuroglia. Atrophy of pyramid cells is also noticeable. Formalin and alcohol fixation; cresyl violet stain; thickness 6 microns.

and Doinikow,¹² in his study of comparative anatomy, agrees with Kölliker.

(a) The Cornu Ammonis: The small size and the marked narrowing of the gray matter were noticeable at the first glance (Fig. 4).

Fig. 9.—Alteration of nerve cells in the terminal plate of the cornu ammonis (left); they are reduced in number and markedly atrophied. Fine construction of these nerve cells can be seen in Figure 10. Compare with Figure 11.

The microscopic investigation of the sections disclosed: the stratum zonale was about normal in thickness; the superficial tangential, middle

12. Doinikow: Beitrag zur vergleichenden Histologie des Ammonshorns, Jour. f. Psychiat. u. Neurol. **13**:166, 1908.

longitudinal and deep tangential fibers could be differentiated. The glia cells were increased and the moderate thickening of the wall of the blood vessels was visible. The cell layer, stratum cellularum pyramidarum and stratum oriens were markedly narrowed as we recognized with the naked eye, and it was a striking fact that the pyramid cells which characterized this layer by their large size as well as by their number were not only much reduced and atrophied (Fig. 6), but in some positions they had entirely disappeared and these gaps of the cell layer were filled with neuroglia (Fig. 8). Many of the pyramid cells in this layer were also markedly shrunk; their Nissl's bodies could not be differentiated and their nuclei had disappeared. Such changes of the nerve cells could be seen through the whole cornu ammonis; generally they were more noticeable in its dorsal plate than in its ventral and they were most striking in the terminal plate, as seen in Figures 9 and 10. In the area near the gaps described the change of the pyramid cells was also remarkable and in these situations the several stages of the cell alterations could be seen; some of them maintained their cell bodies only as a thread-like shell, while others kept their pyramid form with more or less affected protoplasm and nuclei. The favorite locations of the gap seemed to be the transitional regions of each plate into the other, and as far as I examined I found two spaces in the frontal part of the left cornu ammonis and individual spaces in the frontal and in the middle part of the other.

In Weigert's stain not only could the narrowing of the layer of gray matter be seen clearly, but places in which the gray matter had almost disappeared and nerve fibers were pressed together as if they connected the stratum zonale with the alveus crosswise could be seen (Figs. 4 and 13).

The Alveus: This was about normal in thickness or thicker than normal along the dorsal plate; it divided into two branches as usual—the superficial branch went into the fimbria without notable alteration, but the nerve fibers of the deep branch which spread among the nerve cells in the terminal plate of the cornu ammonis were few and fine in comparison with those of the normal sections (Figs. 15 and 16).

(*b*). **The Fascia Dentata and the Fimbria:** In the stratum zonale nothing important was found.

Stratum Granulosum and Stratum Polymorphus: The form and fine construction of each cell, as well as the number in a row, were about normal except in one area in which the ventral portions approached the surface; in this lesion many of the cells were moderately shrunk with alteration in their nuclei.

The Fimbria: There was no noticeable change in the fimbria, except that there were many blood vessels with thickened walls among the bundles of nerve fibers.

Other areas of the brain investigated were: The gyrus cinguli, gyrus temporalis superior, gyrus frontalis superior, gyrus centralis anterior, gyrus centralis posterior and gyrus occipitalis superior.

Speaking generally, the changes which I recognized in these areas, so far as I examined them, were the moderate thickening of the plexiform layer, more or less shrinkage of nerve cells in the gray matter and light thickening of the wall of the blood vessels.

SUMMARY

There were bilateral absence of the bulbus and tractus olfactorius, rudimentary development of the trigonum olfactorium in both hemi-

Fig. 10.—Alteration of nerve cells in the terminal plate of the cornu ammonis (left); they are reduced in number and markedly atrophied. Fine construction of these nerve cells can be seen in Figure 10. Compare with Figure 12.

spheres, absence (left) and partial development (right) of the sulcus olfactorius, nondevelopment of the gyrus olfactorius medialis and lateralis as well as of the gyrus tuberi olfactori in both sides, absence of the stria olfactoria and some atrophy of the gyrus hippocampi in both hemispheres. The other areas, which are regarded as the rhinencephalon, showed no notable alteration.

Comparing this case with that of Weidenreich's, I find that there is no great difference between the two cases except in the incomplete development of the sulcus olfactorius and the atrophy of the gyrus hippocampi in my case.

As already mentioned, Weidenreich is of the opinion that, although the question of the olfactory center has not been solved by his case

report, because no macroscopic change was to be seen in the rhinencephalon, perhaps the olfactory nerves connected the center in some irregular way, as the patient seemed to have had some sense of smell during life.

In my case, the absence of the gyrus olfactorius medialis and lateralis as well as of the gyrus tuberis olfactorii and the stria olfac-

Fig. 11.—Nerve cells in the terminal plate of a normal cornu ammonis. Formaldehyd alcohol fixation; cresyl violet stain; thickness 6 microns.

toria—which were also lacking in Weidenreich's case—would not be of importance in regard to the question of the olfactory center, because the development of these gyri and stria depends directly on that of the bulbus and tractus olfactorius.

The flattened gyrus hippocampi, which I have described, attracted my attention in this case, and it would be interesting to know whether this atrophy in the rhinencephalon is to be regarded as the result of the absence of the lobus olfactorius. It is to be regretted that I could not get any positive information about the sense of smell of the patient during life. Such an important question could not be solved by the macroscopic investigation alone, and therefore I discuss it later in reference to the microscopic findings.

Concerning the origin of the defect of the olfactory lobe, according to the description by Ernst,¹³ arachinencephaly, a brain without an olfactory lobe, may develop when some compression acts on the head

Fig. 12.—Finer details of these nerve cells shown in Figure 11. Formal-dehyd alcohol fixation; cresyl violet stain; thickness 6 microns.

fold of the embryo. Schwalbe¹⁴ states that the theory of compression as the cause of arhinencephaly and cyclopy has been shaken and the opinion that it is due to some embryonal defect is gaining more general acceptance.

Weidenreich is of the opinion that in his case the olfactory nerve could not combine with the olfactory lobe as usual, and consequently the latter was deserted to atrophy. The full development of the sulcus olfactorius in his case shows that the olfactory lobe might have

13. Ernst: *Missbildungen des Nervensystem*. Schwalbe: *Morphologie der Missbildungen des Menschen und der Tiere*, Teil III, Lief. 2, 1910.

14. Schwalbe und Josephy: *Die Missbildungen des Kopfes*. Schwalbe: *Morphologie der Missbildungen und der Tiere*. Tiel III, Lief. 11, 1913.

been developing for some time. He seeks the origin of the anomaly in some alteration of the embryonal connective tissue.

In the present case the sulcus olfactorius did not develop so fully but, in accord with Weidenreich's, the main anomaly has been limited to the lobus olfactorius with no remarkable change in the neighboring

Figure 13

Figure 14

Fig. 13.—An area in the dorsal plate of the cornu ammonis (right); in this lesion the gray matter has almost disappeared and nerve fibers in the narrowed layer of gray matter are pressed together between stratum zonale (Z) and alveus (A). Compare this with Figure 14

Fig. 14.—The same area of about the same region of a normal cornu ammonis as in Figure 13 (Z — stratum zonale). Kulschitzky-Wolter's modification of Weigert's myelin sheath stain; thickness 15 microns.

areas of the forebrain. This is not to be explained by the compression theory, for if so, it must have been circumscribed as well as effective, and at the same time it must have acted on both hemispheres; but such a compression is not to be considered. I would not hesitate to believe that the defect of the lobus olfactorius might have had its origin in some defect in the embryonal tissue.

In the microscopic investigation the changes were found in the cornu ammonis. There were marked reduction in the gray matter, remarkable atrophy of pyramid cells and disappearance of the cell

Figure 15

Figure 16

Fig. 15.—Fine construction of nerve fibers in the deep branch of the alveus (see also Fig. 4). The area presents itself just before the deep branch spreads among nerve cells in the terminal plate of the cornu ammonis (right). The nerve fibers are reduced in number and atrophied. Compare this with Figure 16.

Fig. 16.—Nerve fibers in the same area of about the same region of a normal cornu ammonis as in Figure 15. Kulschitzky-Wolter's modification of Weigert's myelin sheath stain; thickness 15 microns.

layer in some portions, fewer nerve fibers in the deep branch of the alveus, more or less increase of glia cells and thickenings in the walls of the blood vessels.

I believe that the change of the nerve cell denoted atrophy and not aplasia because the alteration was not the same through the whole cornu ammonis and, moreover, several stages of the change in the cells were seen.

Next to the cornu ammonis, the change in the gyrus hippocampi was most noticeable. The changes in this region were: less tangential fibers in the plexiform layer, marked atrophy of the superficial large polymorphic cells, moderate shrinkage of the pyramid cells and loss of the interradian nerve fibers.

It is interesting that to some extent these findings in the gyrus hippocampi accord with the result of the macroscopic investigation, which showed some atrophy of this gyrus. Moreover, it is extremely interesting to have found the most marked change in the cornu ammonis in my case, while the location of the olfactory center is still a question, and some authors, notably Ramon y Cajal, are suggesting the cornu ammonis as the terminal station of this sense.

According to His, quoted in Weidenreich's paper, the olfactory nerve develops separately from its central part and the connection with the brain is accomplished in a comparatively later stage. Therefore, it would not be unreasonable to expect some secondary change in the olfactory center, although it might have developed as usual. In this case in which there were no bulbus and tractus olfactorius, the cornu ammonis showed the most marked alteration in the so-called rhinencephalon.

Besides the anomaly of the olfactory lobe are there any other possible causes for this change to be considered?

Arteriosclerosis may be taken into consideration, as the thickening of the walls of the blood vessels was noted in the cornu ammonis; but the patient did not have marked arteriosclerosis and the basal arteries showed only a slight degree of the hardening. Hence an intense alteration in nervous tissue could not be explained by arteriosclerosis; moreover, the change in the nervous tissue was almost limited to the cornu ammonis.

Certain intoxications may be considered, but in the clinical history there is no evidence of any kind of intoxication as a probable cause of such a change in the nervous tissue. General intoxication would not cause the most marked change in the cornu ammonis; hence this cause may also be excluded. May I then conclude that I have eliminated doubt regarding the olfactory center, and that the cornu ammonis must be regarded as the terminal olfactory center?

Unfortunately, the patient suffered from epilepsy of long duration, as briefly described in the history. Since the sclerosis of the cornu

ammonis in epilepsy was called to attention by Meynert, Sommer,¹⁵ Bratz,¹⁶ Arzheimer,¹⁷ Worcester¹⁸ and Southard¹⁹ by histologic studies on this subject, I shall not give the results of the investigations of each one of these authors. It seems to be true that in several cases of epilepsy discussed by these physicians, some alterations — especially atrophy of pyramid cells, increase of glia cells and sclerosis of blood vessels were found in the cornu ammonis. This complicates the value of similar findings in the present case, as the defect of the lobus olfactorius cannot be excluded as a possible cause of these changes in the cornu ammonis. A discussion as to which of the two possible causes is more reasonable or was more effective in this case should be avoided, as any conclusion in this discussion appears to be only an assumption.

Although it is regrettable that this case could not bring out the soundest evidence necessary to solve the question of the olfactory center, I believe such a case and such an alteration in the cornu ammonis is a matter of enough interest to be reported and may be a contribution to further investigation on this question.

In closing this report I wish to acknowledge my great indebtedness to Dr. E. E. Southard and Dr. M. M. Canavan, whose kind help enabled me to investigate the brain discussed. I also wish to thank Mr. Herbert W. Taylor, a photographer, who prepared the photographs.

---

15. Sommer: Erkrankungen des Ammonshorn als aetiolog. Moment der Epilepsie, *Arch. f. Psychiat. u. Neurol.* **10**: 1880.

16. Bratz: Ammonshornbefunde bei Epileptikern, *Arch. f. Psychiat. u. Neurol.* **31**:1899.

17. Arzheimer: Anatomische Untersuchungen am drei und sechsig Epileptikern. Jahresversammlung des deutschen Vereins für Psychiatrie, 1907; abstr. in *Neurolog. Zentralblatt.*, 1907.

18. Worcester, W. L.: Sclerosis of the Cornu Ammonis in Epilepsy, *Tr. Nerv. & Ment. Dis.* **56**:229, 1897.

19. Southard: On the Mechanism of Gliosis in Acquired Epilepsy, *Am. J. Insanity* **64**:607, 1907-1908.