TUBEROUS SCLEROSIS *

WALTER FREEMAN, M.D.

PHILADELPHIA

Tuberous sclerosis is a relatively rare developmental anomaly of the brain and other organs, characterized clinically by idiocy and epilepsy, and pathologically by multiple sclerotic nodes over the surface of the brain, multiple subependymal tumors, and multiple tumors of the heart, kidneys, skin and other organs.

Though the disease was described by several authors before 1900, von Recklinghausen reporting a case as far back as 1862, the minute histologic alterations and their significance were appreciated first in 1912 when Bundschuh¹ and Bielschowsky² independently gave us very complete and exact studies founded on the embryological concepts laid down by Ranke.³ But little has appeared in English or American literature on the subject, although Campbell⁴ described it in his work on cerebral sclerosis, and Sailer⁵ collected twenty-eight cases in 1898. One American writer recently described a case as juvenile multiple sclerosis. It is important that these cases be recognized, for in addition to the epileptic seizures, these patients often present focal symptoms that may call for operation. It is the consensus of opinion, however, that operation has no favorable influence on their subsequent course.⁶

Because of the association of tumors of other organs with the focal sclerosis of the cortex, cases of tuberous sclerosis are recorded under many titles, and a comprehensive review of the literature is therefore difficult. Titular reference is in some instances to the cardiac tumors, as in von Recklinghausen's case; in others to the renal tumors; while

* Presented before the Philadelphia Neurological Society, March 24, 1922.

* From the Laboratory of Neuropathology, Philadelphia General Hospital, and the Graduate School of Medicine, University of Pennsylvania.

1. Bundschuh: Ein weiterer Fall von tuberöser Sklerose, Ziegler's Beitr. f. path. Anat. u. allgem. Path. **54**:278, 1912.

2. Bielschowsky and Gallus: Klinische und anatomische Studien über tuberösen Hirnsklerose, J. f. Psychol. u. Neurol. **20**: Suppl. Ergänzungsheft 1, 1913.

3. Ranke: Beitrag zur Kenntnis der normalen und pathologischen Hirnrindenbildung, Ziegler's Beitr. f. path. Anat. u. allgem. Path. **47**:51-126, 1909-1910.

4. Campbell, A. W.: Cerebral Sclerosis, Brain **28**:367-437, 1905.

5. Sailer, J.: Hypertrophic Nodular Gliosis, J. Nerv. & Ment. Dis. **25**: 402, 1898.

6. Volland: Untersuchungsergebnisse von 50 Schädeltrepanationen bei Epilepsie, Ztschr. f. d. ges. Neurol. u. Psychiat. **74**:505, 1922.

still others, in which the cutaneous tumors were prominent, are found under the name of adenoma sebaceum. On an estimate, about 100 cases have been reported.

ETIOLOGY

Degeneracy in the parents is an etiologic factor. The disease is congenital, but neither familial nor hereditary. Apparently it begins between the fourth and seventh fetal month. Ranke and Bundschuh have pointed out that the primary and secondary fissures which are completed about the fourth month are not distorted by the sclerotic patches, while the tertiary convolutions which develop later are involved. Bundschuh further called attention to the persistence of an external granule layer over the sclerotic areas which disappears in normal development about the seventh fetal month, and which is never present at birth.

CLINICAL COURSE

The disease is characterized clinically by idiocy and epilepsy. Convulsions usually are the first manifest symptom. In Hartdegen's case the attacks began a few hours after birth and continued until the end, two days later. Brückner's patient, on the other hand, was 9 years old when the first convulsion occurred. When once initiated the convulsions usually continue, although they may decrease in frequency or even cease. The first attacks are usually mild, without loss of consciousness, and they may affect only isolated groups of muscles. During a period of years, however, they become gradually more severe and may even be the direct cause of death. The attacks are quite evenly spaced, are not so frequent as genuine epileptic attacks, nor so paroxysmal as those of the atrophic sclerosis. At times equivalents of attacks are noted, fainting spells, maniacal attacks and so-called psychic epilepsy.

In cases in which the chief clinical expression is idiocy, development may be comparatively normal for several years. Usually the children are slow to walk or talk, but sometimes they are active and intelligent and may even attend school for a number of years. More characteristic than the cessation of development, however, is the retrograde process which follows it. The child ceases to take interest in playthings, prefers to sit still, loses the power of attention, becomes indifferent to everything which had previously attracted it, becomes untidy and often masturbates excessively.

Focal signs, such as localized pareses and contractures, muscular spasms and speech defects, are frequently present. Bielschowsky has reported a case in which movements on one side of the body were much retarded, with rigidity and tremor limited to that side. He records an area of sclerosis in the opposite basal ganglion.

Although status epilepticus is sometimes the direct cause of death, the patients usually die of intercurrent infection, especially of tuberculosis, or of gastro-intestinal disturbances. Occasionally the renal tumors, becoming excessively large, cause death. Patients with cardiac tumors usually die young.

DIAGNOSIS

Often the diagnosis cannot be made during life. When epilepsy and idiocy appear in a child, especially when they are progressive and when localizing signs appear, tuberous sclerosis should be considered. These signs are uncertain, however, and only when tumors of the skin are present is the diagnosis justifiable. In the Pringle⁷ type these tumors grow on the face, are small, firm papillary growths, pale to dark red, vary in size up to 1 cm., and have a "butterfly" distribution over the nose and cheeks. The Barlow type is characterized by larger nodes, sometimes 2 cm. in diameter, usually found on the trunk. They are adenomas of the sebaceous glands. Sometimes there are abnormalities in the growth of the hair. The cutaneous lesions often appear first at adolescence, giving no assistance in diagnosis during the early years of life.

Rarer malformations occasionally found are tumors of the duodenum, spleen and liver; imperfections of the heart, such as patent ductus arteriosus, cardia trilocularis, origin of the aorta from both ventricles; ectopia testis, etc.

REPORT OF A CASE

History.—M. D., Philadelphia General Hospital, service of Dr. Weisenburg, aged 6 years. The father was 29 years of age, the mother 24 at the time of conception. Both were mentally healthy and of good inheritance, but the father was tuberculous and died two months before the child was born. Birth was natural and at full term; the patient was a well developed female child. She was breast fed until 9 months of age. At seven or eight weeks of age she began to have tonic convulsions lasting about two minutes followed by stupor of several minutes' duration. The convulsions increased in frequency for six months, then decreased and ceased at the age of 5 years. During the eleven months she was in the hospital no convulsions were recorded.

Although well formed, the child was backward, could not sit up until she was 2 years old, never learned to walk, talk or feed herself; she did not play with toys until she was 4. Her chief occupation was tearing up her clothing and putting the pieces in her mouth. She masturbated constantly. She could see and hear, but apparently recognized nobody, not even her mother. She would stop tearing her clothes when spoken to, but almost immediately resume it. She was large for her age. The head was slightly enlarged; the skin was without blemish; muscular power and coordination were good. Patellar reflexes were exaggerated. Urine, blood count and spinal fluid were normal. The Wassermann test on the blood and spinal fluid was negative.

7. Pringle: A Case of Congenital Adenoma Sebaceum, *Brit. J. Dermat.* 1:64, 1891.

Before being brought to the hospital, and during her stay there she had occasional attacks of vomiting immediately after taking food, although her appetite was excellent. She became emaciated, and finally died of inanition Jan 21, 1921. She was then 6 years old.

Clinical Diagnosis: Imbecility, hydrocephalus, enteritis.

Necropsy Record (Dr. Morton McCutcheon).—The body was that of a white girl 6 years of age, weighing 40 pounds (18.14 kg.). Bony development was normal; there was marked emaciation; rigor and livor were present. The skull was slightly enlarged and normal in shape; the pupils were equal and regular, the sclerae clear. The thorax was symmetrical, the abdomen retracted. Extremities, external genitalia and hair distribution were normal.

The thoracic organs were normal. The spleen weighed 40 gm., and was normal, as were the stomach and intestines. The liver weighed 540 gm. and was of normal size and consistence. The lower border was rounded. The cut surface showed a number of yellowish white lobules, the other lobules being brown. The discolored areas did not bulge. The pancreas was normal.

The left kidney weighed 60 gm., and was of normal size and consistence. The capsule stripped readily leaving a smooth pinkish gray surface. Striations were normal. Several pale yellow nodules were present in the cortical substance, the largest being 4 mm. in diameter. Some of them bulged distinctly on section although they could not be lifted out, and one bulged through the capsule. The consistence of these structures was that of normal renal substance. The right kidney weighed 50 gm. and resembled its fellow. The suprarenals, aorta, ureters, bladder and internal genitalia were normal.

The brain was distinctly enlarged; it weighed 1,120 gm. The calvarium and dura were normal. The pia was slightly opaque in places but not markedly thickened; it was nowhere adherent to the cortex. The frontal poles of the brain were smaller than usual and unnaturally white. They felt as hard as the normal brain after fixation in liquor formaldehydi. In the temporal, parietal and occipital lobes there were similar hard areas in which the convolutions were larger than normal, projected above the general contour, were flattened or even umbilicate on the surface and unnaturally pale. These areas were fairly sharply circumscribed, irregular in outline and bounded almost everywhere by tertiary fissures. The overlying pia was lightly attached. Except for a small area in the right paracentral lobule, the central convolutions had escaped. Hippocampus, cerebellum, brain stem and cervical cord were normal.

After fixation in 10 per cent. liquor formaldehydi, the usual section was made through the basal ganglions (Fig. 1). The hypertrophic convolutions were broader at the surface than at the base and compressed the normal gyri which lay between them (Fig. 2). The cortex of the sclerotic area varied considerably in thickness; it was distinguished from the underlying white matter, not so much by color as by difference in texture, the cortex being densely hard and the white matter porous. In the fissures the tissue was softer than on the convexity. In the white matter beneath some of the sclerotic gyri were minute areas resembling the overlying cortex. Microscopic examination proved these to be heterotopias of gray matter.

In the right frontal pole, where the area of sclerosis was most extensive, there was a multilocular cyst measuring about 1 cm. in diameter, without softening, pigmentation or obvious degeneration in the neighborhood.

The posterior horn of the left lateral ventricle measured 32 mm. across. Its external surface underlay a large sclerotic area in the temporal lobe. The cerebral tissue was here reduced to 11 mm. At the point chosen for study the

Fig. 1.—Horizontal section of brain: *a*, sclerotic areas of pale color and dense consistency with depression of underlying white matter; *b*, multilocular cyst in right frontal pole formed by dilatation of perivascular lymphatic spaces; *c*, tumor at head of caudate nucleus pressing into foramen of Munro; *d*, dilated posterior horn of lateral ventricle; note the area of microgyria in left occipital lobe.

cortex was narrow, white and hard, and the white matter was soft and porous except where the fibers of the optic radiation ran, skirting the horn of the ventricle. This tract was narrow, but apparently in good condition.

In the right parietal lobe over an area 3 by 4 cm. there were numerous abnormally small and branched convolutions, in which scarcely any white matter could be seen underlying the narrow cortex. The area was not depressed below the general level as is usual in microgyria. While these convolutions were of normal consistence and relatively normal architecture, there was a sharply defined area of sclerosis immediately adjacent. A similar area appeared in the left occipital pole.

Fig. 2.—Hypertrophic convolution, broadened and umbilicate at surface, due to neuroglial overgrowth. The gliosis diminishes at the bottom of the fissure. Phosphotungstic acid-hematin stain; *a*, patch of beginning sclerosis.

At the head of the caudate nucleus on the right there was a tumor measuring 18 by 10 mm. pressing down into the foramen of Monro (Fig. 1). The tumor sprang superficially from the caudate nucleus, was of soft consistence, rather friable and darker than the surrounding tissue. It was circumscribed though not encapsulated. Its surface in the ventricle was fungoid and irregular, but at the sides it was smoother and apparently covered by ependyma. It did not invade the nervous tissue. At other points on the surface of the caudate nucleus on both sides there were smaller tumors varying from 2 to 6 mm. in diameter. In the third ventricle there were tumors up to 5 mm. in diameter, especially along the striae terminales. These smaller tumors occurred singly and in groups, and between the larger ones the ependyma was raised into cords.

Where a single tumor rose the ependyma was arranged in radial cords, resembling buttresses. The smaller tumors were much harder than the surrounding tissue and grated on the knife on sectioning. Between the tumor and the normal basal ganglion there was a layer of pale, firm tissue encapsulating the growth. In the vicinity of the tumors granular ependymitis was visible (Fig. 3). The fourth ventricle contained no tumors.

Gross Findings.—The gross findings corresponded closely to the descriptions given by Pelizzi, Vogt and Bielschowsky. They describe two varieties of tubera: the hypertrophic convolution and the sharply circumscribed node which has no definite cortical characteristics. These exist in varying proportions in different cases, but usually both are present. I could find none of the circumscribed nodes. Broadening of the convexity of the convolution and dimpling in the center were described by older authors, but Pelizzi was the first to show that the sclerotic process was confined to the external surface of the convolution and seldom or never reached the bottom of the fissure, even where two densely sclerotic gyri lay adjacent.

Fig. 3.—Granular ependymitis in vicinity of subependymal tumors.

The cyst of the frontal pole was unusual. Bundschuh found symmetrical cysts in the occipital poles in his case, and others have described them. Localized microgyria was described by Bielschowsky. Ventricular tumors were encountered as frequently as the cortical sclerosis and did not vary much from type. They were usually hard, nearly spherical, lay immediately beneath the ependyma and varied in size up to 8 mm. Their usual locations were the caudate nucleus, striae terminales and thalamus, occasionally the fourth ventricle. On account of the persistent and uncontrollable vomiting without evident local cause, the fourth ventricle was examined with particular care but no abnormalities were found. The tumor at the head of the caudate nucleus was exceptional in size and general characteristics. Whether the hydrocephalus was the result of obstruction of the foramen of Munro by this tumor is not certain. Hydrocephalus is present in a minority of cases, and without known cause. The third and fourth ventricles were not dilated.

MICROSCOPIC EXAMINATION

Ranke³ as a working hypothesis to explain the pathology of tuberous sclerosis suggests that there are two stages of differentiation of cells

of the central nervous system from germinal cells to fully developed ganglion cells and neuroglia cells. First, the cells develop to the point of differentiation when they become either neuroblasts or spongioblasts, and then they ripen into nerve cells and glia cells. This he acknowledges is purely theoretical, since none of the present histologic methods differentiate future neuroblasts from future spongioblasts. If the cells are disturbed at the time of differentiation, a great variety of forms may be produced. "Three extremes are thinkable: first, that the disturbance would result in the production from indifferent elements of nerve cells alone; second, of glia cells alone; and third, that no differentiation would take place. The first two possibilities are extreme one-sided differentiation, and the last, extreme lack of differentiation." Ranke elaborates the theory further by suggesting that harmful influences acting on the neuroblasts and spongioblasts at various periods of their development might produce further anomalies, and also that various combinations might occur.

The so-called "large cells" so characteristic of tuberous sclerosis are believed to be incompletely differentiated neuroblasts. They show the following anomalies (Fig. 4):

(a) Position: Some lie next to the surface; many are distributed through the cortex; some lie in the white matter, either singly or in heterotopic groups. They are seldom arranged in laminae.

(b) Size: The cells equal or exceed large pyramidal cells in size.

(c) Form: Occasionally a cell is seen that would be termed a ganglion cell were it not for its size or location. This is rare. More often the cell is round or oval, occasionally spindle-shaped or snakelike.

(d) Processes: There may be no processes or many processes, and they vary markedly in size. The large round cells have no processes as a rule, but some of them have scores resembling a Medusa's head. The spindle cells have two, and sometimes they dwarf the cell by comparison, extending full across the high power field, dividing, and wandering through the layers of the cortex (Compare Figs. 4, 5, 6).

(e) Nucleus: Many of the round cells have no nuclei. When present the nucleus is eccentric. Often it is lobed and sometimes two, three or more nuclei may be seen in a single cell. Some nuclei are small, but usually they are large; often grotesquely distorted; always pale and vesicular. Nucleoli are usually absent, and the chromatin network is very loose.

(f) Internal Structure: The round cells have neither extranuclear granules nor neurofibrillae, but the snakelike cells have fibrillae in their processes although seldom in the cell body. Sometimes suggestions of tigroid bodies are found in these more perfectly formed elements.

Investigators do not agree that these "large cells" are of neuroblastic origin. Several authors, especially the earlier ones, class them as

Fig. 4.—Various types of faultily differentiated cells of neuroblastic origin: *a*, simple undifferentiated “large cell”; *b*, Medusa cell; *c*, large grotesquely differentiated ganglion cell; *d*, normal ganglion cell of large pyramidal type; *e*, snakelike cell; *f*, atypical ganglion cell; *g*, twin cell; *h*, “large cell” with bilobate nucleus from heterotopia; *i*, large cell with polar nucleus. Silver diffusion stain; $\times 1,000$.

neuroglia cells. Bielschowsky indicates their neuroblastic origin by pointing out gradations from normal nerve cells to round undifferentiated ones on the one hand and snakelike ones on the other. Their staining reactions and the presence of neurofibrillae as demonstrated by Alzheimer also indicate this origin.

Fig. 5.—Low-power view of sclerotic cortex showing neuroglia margin, external granule layer, and atypical "large cells." Silver diffusion stain.

While these cells occurred usually in sclerotic areas, they were found also in the presumably normal cortex nearby; and they were more numerous in the deeper cortical strata and in the white matter than on the surface. The heterotopias consisted of groups of "large cells" interspersed with a few glia cells and fibers, and more or less perfectly formed nerve cells (Fig. 6).

NEUROGLIA

The density of the sclerotic cortex was due to innumerable glia fibers that intertwined in every direction. At the surface the fibers were sometimes collected into sheaves and bundles and stood on end above the normal margin (Fig. 5) but did not invade the pia. They were everywhere abundant, forming

a band many times the thickness of the normal "Randglia" (Fig. 5). At the surface of otherwise normal cortex, the first sign of sclerosis was thickening of the external glia margin and increase of glia nuclei, with sometimes a few large undifferentiated cells (Fig. 7).

Of more interest from the point of view of development was the external granule layer present in the most densely sclerotic areas. Beneath the broad surface layer of neuroglia lay a stratum of round cells, evidently glial in nature, divided from the deeper lying strata of nerve cells by another band of neuroglia fibers (Fig. 5).

Throughout the sclerotic areas there was a dense feltwork of glia fibers in the meshes of which lay numerous small nuclei. Most of these were in the

Fig. 6.—Higher power view of one of the heterotopias. Numerous atypical small nerve cells, some large grotesquely differentiated nerve cells and moderate gliosis.

resting stage but many were fiber formers. Where the feltwork was dense, it was impossible to trace the individual processes; but in the white matter where the tissue became spongy, the glia feltwork was much looser, and here the individual cells stood out with greater prominence (Fig. 8). They were larger than normal and possessed fairly definite cell bodies. The fibers were deeply stained, seemed to spring from the periphery of the cell, were heavy, and at a short distance from the cell divided dichotomously into larger and smaller branches which then curved away in easy paths and broke up into the usual glia fibers.

In the hypertrophic convolutions typical nerve cells were few, small and not arranged in strata; their apical dendrites pointed in various directions, and

Fig. 7.—Earliest sign of beginning sclerosis. Increase of marginal glia associated with a few large undifferentiated cells. Neuroglia stain.

Fig. 8.—Neuroglia cells from loose-meshed white matter; very many processes extending from all points of cells; giant neuroglia cells from junction of white and gray matter; $\times 1,800$.

they contained only remnants of internal structure. Myelin sheaths were almost entirely lacking. By silver impregnation, however, a surprisingly large number of nerve fibers was demonstrated even in the most densely sclerotic areas and the underlying loose-meshed white matter. Moreover, there seemed to be little interference with the projection tracts, for the spinal cord showed no degenerative changes, and such tracts as the optic radiation were well preserved.

Scarcity of myelin sheaths was quite as characteristic as overgrowth of neuroglia in the hypertrophic convolutions, yet in one area studied, nerve fibers appeared to be surrounded by multiple concentric rings, sometimes irregular or oval, depending on the angle of the section. These stain poorly for myelin, but clearly with phosphotungstic acid. They appear to be hyperplastic but poorly differentiated myelin sheaths.

Fig. 9.—Concretions in old compact tumor. They appear to originate from blood vessels.

SUBEPENDYMAL TUMORS

All the tumors in this case except the large soft one were covered by ependyma. They arose from the subependymal glia layer and projected into the ventricle. The process of tumor formation here could be seen by examining the wall of the ventricle. The subependymal neuroglia layer became thicker, the cells developed processes, and a few large undifferentiated cells appeared, thus paralleling the picture seen in the cortical sclerosis. At first the tumor was covered with ependymal cells, but as growth proceeded this layer became much flattened and rupture might have occurred so that the ependymal covering was lost. The tumors which had broken through the ependyma were more vascular and apparently of more rapid growth.

The early tumor is merely a collection of large cells with neuroglia fibers running between cells and surrounding it. In the larger tumors the cells are

more numerous and the glia fibers are proportionately diminished. They separate it into cell groups by trabeculae, and wall it off from the underlying ganglion. Many vessels in these moderate sized tumors are infiltrated with calcium salts (Fig. 9). There is granular ependymitis in the vicinity of the tumors (Fig. 3). Whether the cells of these subependymal tumors are of nervous or supporting origin is contested, some authors considering them as large cell gliomas, others as ganglioneuromas, and still others as neuroblastomas. The majority ally them with the gliomas.

The typical cells of these tumors are fusiform or retort-shaped, vary in size up to that of the largest of the Betz cells, have no tigroid bodies or neurofibrillae, and send processes away from one or both ends. Some of the

Fig. 10.—Low-power view of multilocular cyst.

cells are almost round and have no processes. The nuclei are usually large, with definite nuclear membrane, delicate chromatin network, and only occasional nucleoli. Sometimes the nuclei are grotesque in size or shape, or two or more nuclei are found in the same cell. Mitotic figures are absent. Many of the larger cells have vacuoles but otherwise do not seem to be degenerated.

The fibers sent off by the large cells run in bundles and separate the cells from one another into groups. Among these fibers lie cells of a different character, usually much elongated, with large distorted nuclei and processes following the cell outline. These take the glia stain typically whereas the large cells stain like ganglion cells.

The large tumor in this case varied in microscopic as well as gross characteristics from the others. The cells were not divided into groups, and glia

cells and fibers were almost absent. The cells were polygonal or spherical and had no processes. They stained more heavily than those of the smaller tumors, were smaller in size, and twin cells and multinucleated cells were more frequent. The extreme edge of the tumor was serrated with cell columns projecting into the ventricular lumen. There was a rich supply of vessels of large caliber and thin walls. In many respects the tumor resembled a malignant growth.

BLOOD VESSELS AND CYST

Many of the pial vessels were thickened by overgrowth of the middle coat, and the elastic lamina was split into two or more layers. There was no perivascular round cell infiltration or other sign of inflammation.

In the cortex of the sclerotic areas the blood vessels were less numerous than usual and many were abnormally thick-walled. The perivascular spaces were increased. In the white matter this increase became striking and was of two forms. In the first the adventitia was increased, filling up the space and giving the vessel the appearance of an enormously thickened wall, although the tissue was loose-meshed and nuclei were infrequent. In the other form there was further dilatation of the space to enormous proportions, without corresponding increase of adventitia, thus forming cavities (Fig. 10). This was probably the origin of the multilocular cyst of the frontal pole. Beneath a densely sclerotic cortex appeared large spaces, vacant except for an extremely delicate meshwork of fibrous tissue surrounding a small blood vessel that ran through the cavity. Between the various cavities ran trabeculae of nerve and glia fibers, with many glia nuclei in the resting stage. These septums varied in thickness down to some so fine that they were formed by only a dozen glia fibers. Moreover, in the walls of the larger spaces there were projections of glia fibers, remnants of septums which had ruptured and retracted. These cysts had no lining, and there were no degenerative changes in the adjacent tissues. They were apparently enormous dilations of perivascular lymph spaces brought about by disappearance of nervous tissue.

CONCRETIONS

Concretions were numerous in the subependymal tumors. In the sections some of these elements were circular, intensely black spots; others had central areas of lighter color; others had lumina containing red blood cells, and still others were obviously calcified capillaries and small blood vessels. This finding has been recorded by many investigators who agree that some vessels show calcareous degeneration, but the great number and uniformity of these masses in some places has led to the suggestion that some of them are calcified nervous elements. Calcium infiltration was observed in certain large cells in the sclerotic areas. These granules were most numerous in old compact tumors with dense glia capsules and few functioning vessels. On the other hand, there was no generalized degeneration of cells, and some apparently perfect tumor cells were seen lying next the calcareous bodies. Ranke ascribes both a vascular and cellular origin to these bodies.

RENAL TUMORS

These varied in character, some consisting almost entirely of fat, others of spindle cells, others of thick-walled blood vessels. Occasionally all these

elements were found in a single tumor. They were inlaid in the renal parenchyma, compressed it but little, were sharply circumscribed but not encapsulated, and gave no hint of vegetative activity.

There were no sebaceous or cardiac tumors.

SUMMARY

The cerebral changes in tuberous sclerosis are:

1. Abnormal differentiation of germinal cells during the middle fetal months with the production of neuroglial cerebral sclerosis and subependymal tumors.
2. Incomplete differentiation of ganglion cells with the production of bizarre types.
3. Persistence of the external granule layer of the cerebral cortex.
4. Agenesis of myelin sheaths in the sclerotic areas and underlying white matter.
5. Calcareous degeneration of the walls of vessels in the subependymal tumors.
6. Localized microgyria.

In addition, in this particular case there was dilatation of the perivascular lymph spaces with cavity formation. There was no involvement of the projection tracts.