American Journal of Diseases of Children

VOL. 16

AUGUST, 1918

No. 2

NEUROBLASTOMA SYMPATHICUM

REPORT OF A CASE, WITH NOTES ON TWO OTHERS, AND A BRIEF REVIEW
OF THE LITERATURE *

S. BURT WOLBACH, M.D.

Associate Professor of Pathology and Bacteriology, Harvard Medical School

AND

JOHN LOVETT MORSE, A.M., M.D.

Professor of Pediatrics, Harvard Medical School

BOSTON

CASE 1.—Charles C., 4 years and 3 months old, was admitted to the Children's Hospital, Dec. 31, 1917. His family history was good. He was born at full term, and had always been well, except for rather frequent "colds" during his infancy. His mother thought that his abdomen had always been large, but she had not noticed it especially until two weeks before, since when it had increased in size very rapidly. He began to fail rapidly about the same time and the last week had been confined to bed. Since being in bed he had been somewhat short of breath and had had an unproductive cough. His appetite was fair, he did not vomit and his bowels were regular. He had no symptoms pointing to trouble in the urinary tract.

Physical Examination.—He was pale and poorly developed, and showed some evidences of old rickets. His tongue was slightly coated. The throat, heart and lungs were normal. The abdomen was markedly distended and the superficial veins slightly enlarged. The lower border of the liver was palpable, hard and sharp, 6 cm. below the costal border in the right nipple line, 8 cm. below the ensiform and running under the left costal border in the anterior axillary line. The surface was smooth, hard and slightly tender. The spleen was not made out. The kidneys were not palpable and there was no unusual resistance in the flanks. There were no evidences of free fluid in the abdomen. The genitals were normal and the rectal examination was negative. There was slight edema of the legs. There were no evidences of any involvement of the nervous system.

The urine was acid and cloudy from urates, but contained no albumin or sugar. The sediment was negative.

The blood showed: hemoglobin, 60 per cent. (Sahli); red corpuscles, 2,000,000; white corpuscles, 5,500; mononuclears, 40 per cent.; transitionals, 4 per cent., and polynuclear neutrophils, 56 per cent.

The red corpuscles showed slight achromia, but no variation in size or shape.

The tuberculin test was negative to both the human and bovine types.

The Wassermann test was negative.

The liver increased very rapidly in size, and edema of the scrotum, back and legs became marked. He lost very rapidly in weight and strength and

* Submitted for publication June 3, 1918.

* From the Children's Hospital.

* Read at the Annual Meeting of the American Pediatric Society, Lenox, Mass., May, 1918.

died Jan. 6, 1918, one week after his admission and three weeks after the symptoms were first noted.

The diagnosis during life was sarcoma of the liver, which, in spite of the rarity of the condition, was thought to be primary rather than secondary, because no primary focus of disease could be found elsewhere.

Necropsy.—A restricted necropsy was performed by Dr. S. B. Wolbach, fifteen and a half hours after death. The head was not opened. The throat, heart, lungs, gastro-intestinal tract, pancreas and pelvic organs showed nothing abnormal. The peritoneal surfaces were smooth and glistening. The mesenteric lymph nodes were not enlarged.

Liver: The lower border of the liver extended down to the crest of the right ilium and in the median line was 14 cm. from the ensiform. The enlargement of the liver was almost symmetrical. The suspensory ligament was pushed over to the left owing to the great enlargement of the right half of

Fig. 1.—Photograph of a section of a Kaiserling specimen of the liver, Case 1. The scale is in inches.

the liver and the anterior border of the right lobe was deeply notched a few centimeters to the right of the median line of the body. The liver weighed 2,050 gm. The capsule was smooth. The color was pale red and reddish yellow, mottled with fatty, opaque areas which were not elevated above the surface. These areas varied in size from 1 mm. to 1 cm. in diameter. The liver was extremely friable. Section showed surfaces bearing very little resemblance to normal liver tissue (Fig. 1). As a whole, the cut surfaces had a translucent appearance. There were distinct areas of translucent white tissue, circular in shape, all produced by the confluence of circular areas, thickly distributed throughout the whole of the liver. There were other large areas 4 to 15 mm. in diameter which were soft, deep red and bulging slightly from the cut surfaces. Here and there were narrow bands and small areas of brownish liver tissue, but everywhere paler than normal. It was quite evident that the whole of the liver was infiltrated with a soft new growth.

The gallbladder and bile passages were patent and normal. Accompanying the cystic duct was a lymph node, 8 by 4 mm., which was soft, translucent white.

Spleen: The spleen was enlarged and measured 10 by 6 cm. The capsule was smooth; the color, salmon red. On section the cut surfaces were everywhere smooth and normal in appearance, except for a slightly paler color than is normal. The splenic corpuscles were visible and appeared normal.

Kidneys and Adrenal Glands: The kidneys were normal in size, color, markings and consistency. Partly overlying the left kidney and extending to the median line was a tumor mass (Fig. 2). The kidney itself was not connected with this tumor growth. Section of the tumor mass revealed its intimate relation with the adrenal gland. The tumor mass, as a whole, including the adrenal gland, was 8 cm. in length and 4 to 5 cm. in width. The superior end of the adrenal gland was recognizable. There was a triangular piece of adrenal present which measured 2.5 by 2.5 cm. Extending downward behind the tumor mass was a band of adrenal cortex, 4 cm. in length and from 6 to 8 mm. in width. This arm of the adrenal was not adherent to the tumor

Fig. 2.—Photograph of the primary tumor of the adrenal and the kidney, Case 1.

mass as a whole. In the center of the triangular portion of the adrenal gland first mentioned, was an elevated, white tumor nodule 8 mm. in diameter. Beneath this there bulged from the substance of the adrenal a tumor mass, 5 cm. in length, composed of three nodules, the superior one, 1 cm. in diameter, the middle one 3 cm., the inferior 2 cm. A section through the adrenal showed that the first mentioned tumor nodule extended into the substance of the adrenal and a cut section showed irregular borders. The color was grayish white. Beneath that there was a nodule, 1 cm. in diameter, soft and dark red. Section of the largest nodule, which was 3 cm. in length, presented in part firm white friable tissue, in part dark red necrotic material with many small gritty areas. The inferior nodule composing the main tumor mass was dark red, soft, and apparently composed of necrotic tumor tissue. There were numerous large lymph nodes lying between this tumor mass and the median line and grouped around the celiac axis. These lymph nodes varied in size

from 3 to 8 mm. On section they were soft, translucent and white. The right adrenal gland was normal.

Frozen Section of Tumor: Frozen sections of the tumor, after hardening in formaldehyd solution, showed in the liver almost complete replacement of the liver parenchyma by small round cells with deeply stained nuclei. In some places the destruction of the liver tissue was complete, in others the liver columns could be recognized much flattened and distorted.

Sections of the tumor of the adrenal showed a small round cell tumor with a delicate reticular framework. Portions of this tumor were necrotic and portions were fibrous, but other portions showed a rosette grouping of the cells surrounding a central granular or fibrillar mass, characteristic of ganglioma embryonale sympathicum or neuroblastoma sympathicum.

Fig. 3.—Low power photomicrograph of a section of the liver, Case 1; about 125 diameters.

Microscopic Examination.—The tissues were fixed in Zeuber's fluid and stained with eosin-methylene blue.

Liver: There is very little liver tissue remaining. The organ was largely replaced by a new growth consisting of exceedingly minute cells, smaller than lymphoid cells, most of them with densely staining homogeneous nuclei, having the appearance of pyknotic or postmortem nuclei. This appearance of the nuclei is probably due to richness in chromatin, as the preservation of other cells, that is, liver cells and bile ducts, is perfect (Fig. 3). The liver tissue is present in the form of anastomosing strands, in which the liver cells are usually much distorted, but occasionally they are almost normal in appear-

ance. The intervening tissue filled with the tumor is traversed with delicate epithelial tubules, which probably represent regenerated bile ducts. In other places, where the tumor is actively invading, there are all degrees of atrophy and degeneration of the liver cells, and it is possible to see that the tumor cells are growing between the liver columns and endothelium of the sinusoids, although in most places such details of structure are entirely obliterated. The tumor cells are small, the cytoplasm is extremely scanty in amount and difficult to demonstrate. There are occasional clusters of slightly larger cells with nuclei less densely stained and containing scattered particles of chromatin. The cells, however, are embedded in a finely granular material, which in places can be seen to have a linear striated effect suggestive of protoplasmic fibrils. In some portions of the liver the replacement by tumor cells is so complete as to obliterate all trace of liver. There are many mitotic figures among the tumor cells.

4. Photomicrograph of the primary tumor of the adrenal gland, Case 1; about 400 diameters.

Adrenal Gland: Sections through the adrenal gland containing the primary tumor showed the bulk of the tissue to be composed of minute cells similar to those in the liver metastasis. There are places, however, where there is a very perfect "rosette" formation, consisting of larger cylindrical and pyriform cells arranged about a central mass of granular material (Fig. 4). This central granular material can be resolved in places into very delicate fibrils and the latter in some instances can be traced into the pyriform tumor cells. The stroma consists of a delicate vascular fibrous tissue. The tumor is a rapidly growing neuroblastoma, of sympathetic ganglion cell origin.

Lymph Node: Sections of a lymph node adjacent to the tumor mass showed complete replacement by the exceedingly minute type of tumor cells.

Celiac Ganglia: These are almost completely replaced by the tumor cells, which are always of the very small type.

Spleen: The malpighian bodies are of normal size and shape. A few contain hyaline masses and reticular hyaline material. The pulp contains many eosinophils of the myeloid type. The venules contain groups of cells which are possibly tumor cells, since many of them were smaller than lymphocytes and have deeply stained nuclei.

Kidneys and Right Adrenal Gland: These were normal.

DISCUSSION

From the liver alone the diagnosis of neuroblastoma would have been impossible to prove, although warrantable, as numerous instances are recalled in which the cells in the metastases from neuroblastomata primary in the adrenals have had the appearances characteristic of this case. The diagnosis in this case is established by the presence of the tumor in the adrenal showing a greater degree of differentiation of the cells evidenced by the formation of protoplasmic fibrils and the arrangement into ball masses which in section produce the rosette appearance.

The essential element in the tumor for diagnosis is the fibrils. The fibrils are not to be compared with the coarser, more or less specifically staining fibrils associated with connective tissue and neuroglia cells. They are in reality delicate protoplasmic processes and are representative of axis cylinder processes. There is no special stain for their demonstration.

The earliest diagnosis based on the presence of these fibrils in malignant tumors of the adrenal gland was that of glioma. Virchow¹ and Marchand², in 1891, recognized the nervous character of a tumor of the adrenal and compared the cells with the neuroblasts of the fetal sympathetic ganglia.

J. H. Wright³, in 1910, focused attention to this group of tumors and collected twelve cases, including four of his own. He briefly and succinctly showed that the tumor cells associated with the fibrils have the "same morphology as the cells from which the sympathetic nervous system and the medulla of the adrenal develop and which are regarded by embryologists as arising from migrated primitive nerve cells." Wright offered the names neurocystoma and neuroblastoma as equally applicable.

Pick and Bielschowsky⁴, in 1911, in a classification of tumors of nerve tissue origin, gave the name ganglioma embryonale sympathicum to the tumors described by Wright. Pick⁵, in 1912, collected a total

1. Virchow: *Die Krankhaften Geschwulste*, Berlin, 1863-1865.

2. Marchand: *Festschr. f. R. Virchow; Internat. Beitr. z. wissenschaft. Med.*, Berlin, 1891, 578.

3. Wright, J. H.: *Jour. Exper. Med.*, **12**, 556.

4. Pick and Bielschowsky: *Ztschr. f. d. ges. Neurol. u. Psychiat.*, **6**, 391.

5. Pick: *Berl. klin. Wchnschr.*, **49**, 16.

of eighteen cases, including one of his own, and discussed fully and adequately the origin and the relationship of the tumors arising from the primitive nerve cells. These tumors may represent any of the potentialities of the primitive cells, which are nerve cells, glia cells and chromaffin cells. The tumors representing nerve cells with which we are now dealing may be either rapidly growing and undifferentiated or more slowly growing and more highly differentiated into ganglion cells. He completely substantiated Wright's conclusions.

Landau,⁶ independently of Pick, in the same year, published a similar classification and discussion of these tumors. He added three cases of his own and uses the name malignant sympathetic neuroblastoma. From this time on the origin and classification of these tumors has been accepted and numerous new cases have been added.

The paper of Herxheimer,⁷ in 1913, covers substantially the same ground as those of Pick and Landau. He chooses Landau's name, neuroblastoma sympathicum, in preference to Pick's ganglioma embryonale sympathicum, a choice which has proved to be in accordance with the preference of subsequent writers.

Two American papers, each introducing a new case, by Wahl⁸ and Lehman,⁹ are comprehensive and adequate to orient any one desiring more information on the subject of neuroblastoma. Lehman has collected a total of twenty-five cases of sympathetic neuroblastomas and several doubtful cases.

Hertz and Sacher's¹⁰ case is the most recent one recorded and in the rapid growth of the liver metastasis bears some resemblance to our case.

Our case is noteworthy for the rapidity of the growth of the liver metastases, which overshadowed all other signs, and the absence of apparent metastases elsewhere. As a complete necropsy was not permitted, early metastases to the bones can not be excluded. It is our opinion that these tumors are not as rare as the small number of published cases would indicate and for this reason we append the reports of two other cases from the surgical service of the Boston Children's Hospital.

CASE 2.—W. D. MacN., aged 7 months, was admitted June 12, 1917. His past history and family history are irrelevant. May 4 a physician noticed a mass in the right side of the abdomen which was not tender and which did not noticeably increase in size. The physical examination showed a well nourished infant. The skin, eyes, ears, nose, mouth and throat were all

6. Landau: *Frankfurter Ztschr. f. Path.*, **11**, 26.

7. Herxheimer: *Beitr. z. path. Anat. u. Path.*, **57**, 112.

8. Wahl: *Jour. Med. Research*, **30**, 205.

9. Lehman: *Jour. Med. Research*, **36**, 309.

10. Hertz and Secher: *Hospitalstidende*, **60**, No. 45; *abst. Jour. Am. Med. Assn.*, 1918, **70**, 278.

negative. The heart and lungs were normal. The reflexes were normal. The abdomen showed a bulging in the right upper quadrant. Palpation showed a hard irregularly shaped mass occupying the whole of the upper and a third of the lower right quadrant. It was not tender. It descended with respiration, was not attached to the abdominal wall and by rectal examination the lower end was found to be freely movable.

An exploratory operation was done on June 14. The tumor was found to be inoperable and a small piece was removed for microscopic examination. The abdomen contained a slight excess of fluid in the flanks. The tumor was hard, nodular and not cystic and extended from the right costovertebral angle to below the brim of the pelvis and inward beyond the median line. It overlay the aorta and vena cava. The ascending colon was displaced forward and inward and crossed vertically the median aspect of the tumor.

Death occurred seven hours after the operation. Permission for a necropsy was refused.

The material was fixed in Zenker's fluid and stained with eosin-methylene blue, Mallory's phosphotungstic acid hematoxylin, Van Gieson's stain and hematoxylin and eosin. The photomicrographs were made from sections stained with hematoxylin and eosin.

Microscopic Examination.—The tumor is composed mainly of small round cells irregularly arranged in masses which are separated by a delicate fibrous tissue stroma giving an alveolar appearance. The tumor cells vary in size and the nuclei contain a variable but large amount of chromatin. The cytoplasm is small in amount and usually difficult to demonstrate. There are groups of larger elongated cells in compact masses but there is no rosette grouping in the sections. Delicate fibrils are very numerous throughout the tumor between individual cells and in large masses of parallelly arranged wavy fibrils lying between the groups of tumor cells and stroma or extending between adjacent clumps of tumor cells. There are many mitotic figures and a rare type of larger cell with abundant cytoplasm suggesting a ganglion cell. The tumor is a typical malignant neuroblastoma sympathicum identical with many of the reported cases (Figs. 5 and 6).

CASE 3.—Myrtle P., aged 2 years and 2 months, was admitted to the outpatient department, March 2, 1918, for swelling and tenderness at the left knee, attributed to a fall one week previously. The knee was swollen, not red, slightly tender. There was limitation of motion to 20 degrees extension and 60 degrees flexion. A roentgenogram made March 5 shows widening of the lower end of the diaphysis of the femur with areas of rarefaction just above the epiphyseal line. A mass was noticed in the abdomen and the child was admitted to the hospital March 9.

Her past history and family history are irrelevant. The child was well nourished and well developed. The cranium, eyes, ears, nose, teeth, mouth and throat were negative. The heart and lungs were normal. The reflexes and skin were normal. The extremities with the exception of the left knee were normal. The abdomen was rounded and tympanitic, with the exception of the left upper and middle quadrants. In the left upper quadrant there was felt a hard rounded mass which did not move when the position of the child was shifted. It was smooth, not tender.

An operation, March 11, showed a large tumor of the left kidney. This was enucleated and found to be adherent posteriorly. It extended upward to the diaphragm, where there was a cyst. The cyst ruptured and a considerable quantity of brown semisolid material escaped. After removal of the tumor there was but little bleeding from the cavity, but inspection showed that there were nodules of the growth extending downward along the vertebral column and out into the broad ligament on the left side.

Death occurred March 28, presumably hastened by a low grade infection of the peritoneum.

Figure 5

Figure 6

Figs. 5 and 6.—Photomicrographs from Case 2, showing masses of fibrils extending between clumps of tumor cells; about 400 diameters.

Pathologic Report.—The specimen consisted of a kidney partly replaced by tumor. It was moderately firm and measured 13 by 7 by 5 cm. The lower pole was evenly rounded. At the upper pole there was a loss of substance involving the pelvic border of the kidney, about 3.5 cm. in diameter. There was in this location an irregularly circular yellow and red mottled crater, the base of which was formed of granular friable material. At the periphery of this crater there was attached a grayish membrane continuous with the capsule of the kidney and which probably was the remains of the capsule overlying the necrotic mass. A section through this pole of the kidney showed an irregular zone of yellowish red mottled tissue, 1 to 2 cm. deep, and inferior to this the kidney substance appeared to be normal.

Microscopic Examination.—The tissue was fixed in Zenker's fluid and 10 per cent. formaldehyd. The description is based on sections stained with eosin-

Fig. 7.—Photomicrograph, Case 3, showing bundles of fibrils between masses of the tumor cells and between the tumor and the stroma; about 400 diameters.

methylene blue, Van Gieson's stain and Mallory's phosphotungstic acid hematoxylin stain. The photomicrographs were made from sections stained with hematoxylin and eosin.

Sections through the upper pole of the kidney show a new growth consisting of large anastomosing columns of tumor cells with many mitoses and supported by a delicate connective tissue stroma. Most of the tumor cells are small and round. In many places there are elongated oval and pyriform-shaped cells. The latter have a finely granular cytoplasm which can often be traced out into long delicate processes. The nuclei are relatively large, vesicular in type and contain a large amount of chromatin in the form of a coarse network (Fig. 8). The amount of cytoplasm is very scanty and as a

rule is eccentrically situated in relation to the nucleus. The size of the cells is slightly larger than the size of lymphocytes found in the germinal centers of lymphoid follicles. Here and there throughout the tumor there are very delicate fibrils occurring in small bundles between the tumor cells and extending between adjacent clumps of tumor cells (Fig. 7). These fibrils are also found in bundles of considerable size lying between the connective tissue stroma and the tumor cells. The fibrils are exceedingly delicate and have a granular structure and bear no resemblance to the deeply staining intercellular fibrils of connective tissue and neuroglia (Fig. 8). There is no rosette formation but the diagnosis of neuroblastoma sympatheticum is established beyond question by the presence of the delicate nerve fibrils.

Fig. 8.—Higher power photomicrograph of a section from Case 3, showing the character of the nuclei and the fibrils; about 800 diameters.

No postmortem was obtained and therefore the location of the primary tumor cannot be proved. The extensive loss of substance and invasion of the kidney at the upper pole indicate, however, that the tumor took origin from the adrenal gland.

These three cases present the slight differences in the histology exhibited by this type of tumor. In the first case we have rosette formation in the primary tumor and in the metastases extreme rapidity of growth without rosette formation or fibril formation. The density of the nuclei is indicative of a large chromatin content and rapidity of growth, as has been noted by other authors. In the second and third cases no rosette formation was found. Fibril formation was abundant

in the second case, less abundant in the third, which represents a type easily overlooked. These three cases bring the number of undoubted cases of neuroblastoma sympathicum to twenty-nine, of which twenty were primary in the adrenal gland. The others have taken origin in sympathetic ganglia (three), retroperitoneal tissues (three), coccygeal gland, nose and uterus. It is highly probable that many other cases of primary tumors of the adrenal gland reported as sarcomas were neuroblastomas, as, for instance, the fourteen cases of adrenal tumors with cranial metastases collected by Tileston and Wolbach.¹¹ Of all the cases reported, but one patient survived, Lehman's case,⁹ the only successful operative removal of a tumor of this type in an infant.

Since a variety of terms have been used to designate the malignant tumors representing a primitive type of sympathetic ganglion cells, it seems advisable to indicate a choice. Wright's terms neurocytoma and neuroblastoma are too inclusive. Ganglioma embryonale sympathicum, introduced by Pick and Bielschowsky, has been regarded by several authors as unsatisfactory because of the employment of ganglioma by others to designate a more highly differentiated type of nerve cell tumor. Our own preference is for the term neuroblastoma sympathicum or sympathetic neuroblastoma.

REFERENCES

For the complete bibliography of this and related types of tumors the articles of Pick and Bielschowsky, Pick, Herxheimer, Wahl and Lehman should be consulted. The presentation of cases in the literature in tabular form by Wahl and by Lehman makes their papers particularly useful.

-
11. Tileston and Wolbach: *Am. Jour. Med. Sc.*, June, 1908.