PROGRESS IN PEDIATRICS

A REVIEW OF THE LITERATURE OF SYPHILIS IN INFANCY AND CHILDHOOD*

PARK J. WHITE, M.D. .
ST. LOUIS

Jeans¹ has reviewed the literature on this subject up to April, 1920. It is the object of the present review to bring the literature to date (December, 1921). A brief survey suffices to show the extent to which contributions from the laboratory predominate over those from clinic and practice. The reason for this is not far to seek. Syphilis, hereditary or acquired, is a disease in whose investigation and partial conquest the laboratory has played an important, an essential part. Excepting statistical reports, there are relatively few clinical observations or contributions to symptomatology. Clearly, serology, chemistry and pathology offer the widest fields for original endeavor in this disease.

INCIDENCE

Jeans and Cooke² made a study of the placentas, and of the Wassermann reactions of the cord blood of a series of 2,030 unselected infants in St. Louis. By examining the blood of 389 of these infants after 2 months of age, it was determined that the proportion of cases of hereditary syphilis that could be diagnosed with certainty by placental examination alone was 27 per cent. From the Wassermann reaction on the cord blood, 63.6 per cent. could be recognized. By applying these two methods to the series, the number of cases of hereditary syphilis in the entire group was determined to be 15 per cent. in the colored race, 1.8 per cent. in the poor of the white race, and less than 1 per cent. in the well-to-do classes. The incidence of hereditary syphilis at birth for all classes in St. Louis is 3 per cent., of which the colored population, although only 9 per cent. of the total, contributes approximately half the cases. In a somewhat similar study, Ross and Wright³ found 3.5 per cent. of the specimens of (placental) blood to be positive.

In evaluating such statistical reports, two very important factors to bear in mind are the race and the standard of living of the people

* From the Department of Pediatrics, Washington University.

1. Jeans: *Am. J. Dis. Child.* **20**:55 and 132 (July and Aug.) 1920.

2. Jeans and Cooke: *Am. J. Dis. Child.* **22**:402 (Oct.) 1921.

3. Ross and Wright: *Lancet* **1**:321 (Feb. 12) 1921.

studied. Burhans' figure ⁴ of 1.5 per cent. (for the Babies' Dispensary and Hospital of Cleveland), is rather unusually low for clinic patients. Royster's figure ⁵ of 12.5 per cent. (for a clinic in Norfolk, Va.) is decidedly high, and may be explained by the large proportion of colored patients in the clientele. The incidence of 3 per cent. for St. Louis, and 3.5 per cent. (Ross and Wright) for an English mining and industrial town, are in fairly close agreement and are based on the entire population of each town.

RELATION OF SYPHILIS TO INFANT MORTALITY

In any consideration of deaths due to syphilis, fetal mortality assumes first place. Morgan,⁶ for example, estimates that 2,000 miscarriages and stillbirths occurred in the city of Toronto in 1920, and that 30 per cent. of these were due to syphilis—thus making 670 an index of the prenatal ravages of syphilis in that city in one year. Solomon and Solomon,⁷ in a complete study of 555 families of late syphilitics, state that 29.7 per cent. gave birth to no living children, 23 per cent. being entirely sterile, 6.7 per cent. having unsuccessful pregnancies. The 29.7 per cent. of childless syphilitics should be contrasted with 11.3 per cent. childless in a similar census group. The average birth rate per family was 2.05, and the average number of living children per family was 1.62. In the same type of population taken at random, the average birth rate per family was 3.8. How large a part syphilis plays in race suicide is thus obvious.

Williams⁸ studied 302 fetal deaths (99 white, 203 black) in 4,000 deliveries. Syphilis was responsible for 34.4 per cent. of the deaths—being the most important single cause. Allowance must, of course, be made for the fact that colored people comprised more than half the clinical material reported. The high figures for syphilis given by Williams may well be compared with those given by Meyer,⁹ who, in the course of a detailed discussion of abortion, says: "Associated constitutional or venereal diseases were recorded in only seventy-six out of 697 selected histories. In fifty-two of these, other causes for the termination of the pregnancy were also recorded." Hornung's¹⁰ figures for 221 pregnant syphilitic women in Kiel are also lower than those of Williams.

Two instructive contributions to the subject of stillbirth, giving more of its pathologic aspects, are those of F. J. Browne. In the

4. Burhans: *Ohio State M. J.* **17**:821 (Dec. 1) 1921.

5. Royster: *Am. J. Syphilis* **5**:131 (Jan.) 1921.

6. Morgan: *Canad. M. A. J.* **11**:849 (Nov.) 1921.

7. Solomon and Solomon: *Social Hygiene* **6**:469 (Oct.) 1920.

8. Williams: *Bull. Johns Hopkins Hosp.* **31**:141 (May) 1920.

9. Meyer: *Am. J. Obst. & Gynec.* **2**:138 (Aug.) 1921.

10. Hornung: *Zentralbl. f. Gynäk.* **44**:1222 (Oct. 23) 1920.

first¹¹ he reports a pathologic study of 200 consecutive cases of still-birth and neonatal death in the Edinburgh Royal Maternity Hospital. Twenty-two macerated fetuses are included in the series. In fourteen of these cases syphilis was the cause of fetal death. Besides the fourteen macerated fetuses there were twenty-one other cases of syphilis. Twenty-six of the total of thirty-five cases were premature. The pathologic findings reported in this and in his later articles will be discussed under pathology. Further statistics from necropsy material are those of Warwick,¹² who found 9.5 per cent. of deaths due to syphilis in 200 necropsies performed on babies who were stillborn or who died a week after birth (in the city of St. Paul).

As reports of fetal and neonatal mortality due to syphilis vary to some extent with the factors affecting each series, so do the reports of infant mortality from this cause. In the series of Solomon and Solomon,⁷ the infant mortality rate was 124 per thousand. This is less than the rate given in Massachusetts census for 1910 (i. e., 131 per thousand). Thus, in a series of children of late syphilitics, the infant mortality rate and the deaths of children under 18 do not vary greatly from the mortality as found in the community. The authors remark that the percentage of living nonsyphilitic children is considerably greater in this study than in the studies of many other investigators, due, no doubt, to unconscious selection in other studies. It is noteworthy, however, that only 30.3 per cent. of the entire group were free from defect as to children and syphilis in the spouse.

The difficulties encountered in obtaining accurate figures on infant mortality from syphilis are well described by Morgan.⁶ He found that in one year, in the city of Toronto, only nineteen certificates mentioned hereditary syphilis as a cause of death. He mentions the physicians' fear to record syphilis "to be hawked about among relatives for the delectation of prudes." This factor of concealment is no doubt often an important one. It must be borne in mind, however, as will be seen later, that often the pathologist himself is unable definitely to prove that in a given case hereditary syphilis was actually the cause of death. Athrepsia, bronchopneumonia, etc., are such frequent "terminating complications" that it is easy, though, of course, misleading, to record any of these without mentioning the underlying syphilis.

Morgan reports that of 725 successive admissions to the medical wards of the Toronto Hospital for Sick Children, twenty-nine were syphilitic—or 4 per cent. He considers that for the whole population the figure must be considerably higher. According to this writer, 90 per cent. of deaths from syphilis occur during the first year. Burhans⁴ found that the known mortality for 141 cases was forty-nine, or 34.7

11. Browne, F. J.: *Brit. M. J.* **2**:140 (July 30) 1921.

12. Warwick: *Am. J. Dis. Child.* **21**:488 (May) 1921.

per cent. In fifteen of these the cause of death was given as hereditary syphilis. Kaufmann-Wolf and Abrahamson¹³ report their fifth series of this kind. The total mortality for the five series was at least 70 per cent. when the mother was infected; and under all conditions, it was about 50 per cent. The morbidity of the living averaged 50 per cent., so that only 25 per cent. of children in the 134 families grew up healthy.

In his valuable report of the first year's work of a clinic for syphilitic children, Jewesbury¹⁴ presents the findings in 145 children whose mothers were known to be syphilitic. Of these, nine mothers became infected after the children were born—hence the disease in the children was acquired. Of the remaining 136 mothers, forty-seven had treatment during pregnancy or before. In about 50 per cent. of the cases, the disease was most marked in the early pregnancies, and apparently became gradually attenuated. But in just as many cases, children free from infection appeared in the midst of syphilitic children and miscarriages.

HEREDITARY TRANSMISSION

To gain a complete understanding of the manner of transmission of hereditary syphilis, one should study in order the "vivo and vitro" characteristics of the *Spirochaeta pallida*, its effects on the father, the mother and, through them, on their children. A sufficiently thorough consideration of all the recent literature on the spirochete would consume too much space. Some of the work done, however, requires mention—notably the experiments with rabbits by Brown and Pearce. Their results on the dissemination of spirochetes,¹⁵ the lesions of the skin,¹⁶ mucous membranes,¹⁷ eyes,¹⁸ etc., are of much value, though without immediate bearing on the transmission of syphilis to children. Their studies on latent syphilis throw some light on the important question of how a father suffering from this form of the disease may still be capable of transmitting it to his wife and thus indirectly to the offspring. Material from the popliteal nodes of six rabbits, taken after considerable period of supposed recovery, or latency, was reinoculated into other rabbits. In each case active syphilis resulted. The authors conclude that recovered rabbits may harbor virulent spirochetes almost indefinitely, even without further manifestations of activity. The lymphoid tissues of the body are probably the chief reservoirs of the virus during latent periods. Frühwald,¹⁹ reporting

13. Kaufmann-Wolf and Abrahamson: Ztschr. f. klin. Med. **89**:274, 1920.

14. Jewesbury: Lancet **1**:962 (May 7) 1921.

15. Brown and Pearce: Arch. Dermat & Syph. **2**:470 (Oct.) 1920.

16. Brown and Pearce: J. Exper. M. **32**:445 (Oct.) 1920.

17. Brown and Pearce: J. Exper. M. **32**:447 (Nov.) 1920.

18. Brown and Pearce: J. Exper. M. **34**:167 (Aug.) 1921.

19. Frühwald: Wien. klin. Wchnschr. **33**:999 (Nov. 11) 1920.

eighty-nine gland punctures in eighty-three (acquired) syphilitics, found that as the secondary stage progresses spirochetes in the glands become rarer, but that the glands may shelter them up to the third year. In four cases, latent after the conclusion of treatment, spirochetes were demonstrated in three of fourteen gland punctures.

In an experimental study of the latent syphilitic as a carrier, Eberson and Engman²⁰ isolated the spirochete from five cases out of seventy-five, ranging in latency from one to forty years after the disappearance of syphilitic signs and symptoms. Three times the organism was found in inguinal glands (in two women and one man) and twice in semen. The strains produced typical syphilitic lesions in rabbits' testicles. *Spirochaeta pallida* was isolated from patients who gave a history of syphilis dating back eleven and thirteen years, respectively, in two instances, and one year in three instances. An inguinal gland and the semen proved positive for spirochetes in the two cases first mentioned, and the glands and semen in the last named case. The blood and other body fluids, except semen, are not infectious in latent syphilis, or if so, but rarely. These facts are clearly important from the point of view of hereditary syphilis, the persistent positivity of the semen for spirochetes even after treatment or a long period of latency, indicating the difficulties in the way of protecting the prospective mother from infection by the father.

In explanation of some of the vagaries of the spirochetes, the interesting theory has been advanced that the organism passes through an intermediary or "resting" stage like that of the protozoal parasites. One of the strongest arguments in favor of the theory has been that spirochetes have never been recovered or seen during the incubation period within the rabbit's testis. According to Eberson,²¹ the absence of organisms, as judged solely by microscopic findings, is not convincing proof that a change of form has taken place within the animal body. That the *Spirochaeta pallida* is found unaltered both in morphology and in virulence in the blood and glands, seems to him sufficient to render the "resting stage" theory untenable.

If the physical obstacles to protection against infection by the husband are great, the social obstacles are still greater. Peter,²² for example, has followed 1,000 patients who had been under treatment for venereal disease at free dispensaries, and draws a gloomy picture of the heedlessness and lack of perseverance of the great majority. Roberts²³ has had the same experience. He found that severity of symptoms at

20. Eberson and Engman: J. A. M. A. **76**:160 (Jan. 15) 1921.

21. Eberson: Arch. Dermat. & Syph. **3**:111 (Feb.) 1921.

22. Peter: Schweiz. med. Wchnschr. **50**:623 (July 15) 1920; abstr. J. A. M. A. **75**:773 (Sept. 11) 1920.

23. Roberts: Lancet **2**:277 (Aug. 6) 1921.

onset had very little influence on attendance. (This factor is more important in hereditary cases in which symptoms are more likely to be alarming.) The most important factor in relation to attendance was the positivity or negativity of the Wassermann reaction. Patients (and this applies to parents, of course) must be instructed repeatedly that a negative Wassermann reaction does not mean a cure. Letter cards are an important though often unsuccessful means of getting patients to return. It is noteworthy that in Robert's series, 8.2 per cent. gave the wrong address when registering. Persons persisting in treatment until clinical and serologic cure has been affected are not protected against reinfection. Though such reinfection undoubtedly occurs, it is no evidence of curability; and these two conceptions, according to L. Jacobi,²⁴ should be dissociated.

Widakowich²⁵ gives an extensive bibliography, together with sixty-four illustrations of anomalous forms of spermatozoa in syphilitics. The percentage of pathologic forms was much higher in syphilitic than in normal men, and in heredosyphilitics than in patients with the acquired disease. Sidler-Huguenin²⁶ states that the male sexual organs are injured more than the female by inherited syphilis, judging from the smaller number of children in the families where the syphilitic is the father. He was unable to find symptoms of syphilis in the sixty-five children of the second generation, except in one doubtful case. He believes that persons with inherited syphilis can be assured that their children will not feel the effects of it, that is, of course, if they have any children, which they often do not.

Solomon and Solomon⁷ found, as they expected, a higher incidence of positive Wassermann reactions in mates than in offspring. Thirty per cent. of the spouses were positive. From 8.4 to 16.7 per cent. of the children (depending on the method of selection) were positive. Nearly one fifth of the mothers of syphilitic children gave a negative Wassermann reaction and were without symptomatic evidence of syphilis.

Riecke and Hoernicke²⁷ cite, among others, a family in which the grandfather had tabes. His daughter and her son were syphilitic. The usual difficulty of absolutely ruling out acquired syphilis in the mother was encountered. Bruusgaard²⁸ reports the fifth thoroughly studied case of hereditary syphilis in the second generation at the Christiania Hospital. The grandmother is being treated for a gummatous tertiary

24. Jacobi, L.: *Arch. Dermat. & Syph.* **2**:493, 1920.

25. Widakowich: *Semana méd.* **27**:633 (Nov. 11) 1920; abstr. *J. A. M. A.* **76**:414 (Feb. 5) 1921.

26. Sidler Huguenin: *Schweiz. med. Wchnschr.* **51**:49 (Jan. 20) 1921.

27. Riecke and Hoernicke: *München. med. Wchnschr.* **68**:1 (Jan. 7) 1921.

28. Bruusgaard: *Norsk Mag. f. Lægevidensk.* **82**:353 (May) 1921; abstr. *J. A. M. A.* **76**:1803 (June 18) 1921.

periostitis, the mother for periostitis of both tibias, and the boy, aged 8, has fan-shaped deep scars extending from the corners of the mouth. He has a two-plus Wassermann reaction. The mother during pregnancy had tertiary ulcerations and keratitis. Another family history of interest is that given by Bernheim-Carrer.²⁹ From his records, however, and from the literature in general, it would seem that definite proof of the hereditary transmission of syphilis from one generation through the two succeeding is still lacking.

The maternal, or, more accurately, the placental transmission of syphilis to offspring, is an accepted fact. Diagnosis is much more difficult in women than in men, owing to the frequent absence of symptoms. Swayne³⁰ believes that a very large number of women contract the disease innocently; and that they are more likely than men to act as innocent carriers. This author makes the general observation that syphilitic symptoms and early lesions are not so virulent now as they were many years ago; and credits the possibility that tabes and paresis are more frequent (an observation questioned by McKenna.)

Data obtained by Fischl and Steinert³¹ confirm the placental transmission of syphilis, and also, unexpectedly, confirm Profeta's law, to the extent that seventeen apparently healthy infants failed to show any sign of syphilis, although their mothers were in the most contagious phase of florid syphilis. Ten of the infants in this group never had a positive Wassermann reaction. Five of the mothers were also constantly negative, notwithstanding their florid papules at the anus, etc. Immunization by way of the placenta before birth and by suckling afterward will have to be accepted in these cases, according to the authors, even allowing for the defective production (common to the new-born) of substances yielding the Wassermann reaction. They report several infants with pronounced lesions swarming with spirochetes, but with a constantly or frequently negative Wassermann reaction. Treatment with arsphenamin did not display any provocative action in this respect. In thirty-four infants the manifestations of inherited syphilis developed late, and the Wassermann did not become positive until some time later, but in 11.7 per cent. the Wassermann veered to positive as the precursor of the clinical manifestations.

Experimental work has shown³² that the reaction of rabbits to genital inoculations of spirochetes at the time of conception differs markedly from that of normal animals. This difference extends through the period of pregnancy and well into the period of lactation. The defensive mechanism of pregnant animals is capable of opposing

29. Bernheim-Carrer: *Monatschr. f. Kinderh.* **21**:130 (May) 1921.

30. Swayne: *Brit. M. J.* **2**:476 (Sept. 24) 1921.

31. Fischl and Steinert: *Arch. f. Kinderh.* **69**:309 (June 21) 1921; abstr. *J. A. M. A.* **77**:581 (Aug. 13) 1921.

32. Brown and Pearce: *Am. J. Syphilis* **4**:593 (Oct.) 1920.

a resistance to inoculations at the time of conception, such that little or no clinical sign of infection appears.

In inherited syphilis the placenta is likely to be exceptionally large in proportion to the size of the fetus.³³ Intense local phagocytosis explains the rarity of the discovery of spirochetes in the placenta.³⁴ All of the organisms may be incorporated in phagocytes in the blood stream or in the endothelial cells of the capillaries, even when the viscera of the fetus may be swarming with spirochetes.

McWalter³⁵ reports a case to show that infection of a mother with syphilis in the eighth month of her pregnancy results in infection of the child. The syphilis in the child is of milder form, and develops later than ordinary hereditary syphilis. Gellhorn³⁶ deplores the tendency to regard maternal syphilis as unimportant, except as it affects the fetus, and contends that it is capable of producing real obstetrical complications.

THE QUESTION OF DIFFERENT STRAINS OF THE SPIROCHETE

Whether or not there are special strains of the spirochete responsible for the various clinical forms of syphilis is a question which has given rise to considerable discussion. Many of the arguments pro and con are based on hereditary transmission. Gougerot³⁷ gives an interesting review of the subject. The "pluralistes" support their contention with the fact that cases are on record of men contaminated by the same person, all having syphilis of the central nervous system. They also emphasize the occurrence of conjugal neurosyphilis; the absence of cutaneous lesions in cases of syphilis of the central nervous system; the fact that natives of the tropics seldom have syphilis of the central nervous system; the different degrees of contagiousness; "distinct evolution." The "unicistes" dispute most of these claims, and emphasize the individual differences of the various hosts.

Levaditi and Marie³⁸ give experimental evidence in favor of two types of spirochetes: the neurotropic type on the one hand, and the skin and mucous membrane type on the other. They point to the fact that rabbits inoculated with one type are not protected against the other, and that the course of the infection differs notably according to the type inoculated.

In their article on conjugal neurosyphilis, Moore and Keidel³⁹ note that the clinical evidence favoring the existence of a neurotropic strain lies partly in the comparative frequency of neurosyphilis in both

33. Lévy-Solal: *Gynéc. et Obst.* **4**:94 (Aug.) 1921.

34. Manouélian: *Gynéc. et Obst.* **3**:1 (Jan.) 1921.

35. McWalter: *Brit. M. J.* **1**:827 (June 19) 1920.

36. Gellhorn: *Surg. Gynec. & Obst.* **32**:535 (June) 1921.

37. Gougerot: *La Médecine* **2**:85 (Nov.) 1920.

38. Levaditi and Marie: *Presse méd.* **28**:646 (Sept. 15) 1920.

39. Moore and Keidel: *J. A. M. A.* **77**:1 (July 2) 1921.

partners to a marriage and in their children. They present the results of examination of forty-two wives, two mistresses, and eight husbands, being the marital partners of fifty known neurosyphilitic patients. Forty of the partners were syphilitic, and 33 per cent. of these had neurosyphilis. Though there is a high incidence of conjugal neurosyphilis in the partners of parenchymatous neurosyphilitics, the comparatively low percentage found in the partners of cerebrospinal neurosyphilitics somewhat lessens the strength of this argument for duality of strain. Feldman⁴⁰ reports a family in which the father had taboparesis with paraplegia, the mother and the oldest boy had tabes, and a younger boy had cerebrospinal syphilis with paraplegia. The first clinical symptom in each case was nystagmus. Another boy in the same family died of cerebrosyphilis. The (blood) Wassermann reaction was positive in all. The author recalls another family in which there was no neurosyphilis, though all were syphilitic. He favors the "special strain" theory, mentioning that the hereditary transmission of a poorly resisting central nervous system is hardly logical, since inherited traits cannot be transmitted from husband to wife.

Solomon and Solomon⁷ found in their series very little difference in the various percentages obtained in families of patients with general paresis and those without central nervous involvement. The problems of syphilis are practically the same from the familial point of view, whatever course the syphilis takes in the individual patient. They add that in the past numerous articles have been written to show that the offspring of paretics were remarkably healthy and quite devoid of familial syphilis. Parallel studies indicate the contrary.

Thom⁴¹ is convinced that there is no such thing as different strains in spirochetes. The selective action of spirochetes in the body is governed entirely by the manner in which the tissues of the host react. Hutinel⁴² maintains that there is one type of hereditary syphilis in which specific dystrophic changes are produced by the spirochete or its reactions, and another in which nutritional disorders are found. The endocrine glands are the tissues principally attacked. The children of such patients may also have hereditary dystrophies. Audrain⁴³ presents the tabulated details of thirty families, through three or four generations from a progenitor with unsuspected syphilis. He finds that the law of persistent localization of the lesions is confirmed, also their periodicity and the rarity of pain.

Gordon⁴⁴ cites cases of conjugal tabes and paresis from the literature, and notes reports to the effect that either partner may develop

40. Feldman: *Med. Rec.* **100**:1107 (Dec. 24) 1921.

41. Thom: *Am. J. Syphilis* **5**:9 (Jan.) 1921.

42. Hutinel, P.: *Arch. d. med. d. enf.* **23**:5, 145, 205 (Jan.-April) 1920.

43. Audrain: *Bull. Soc. de derm. et de syph.* p. 85 (Feb. 10) 1921.

44. Gordon: *Am. J. Syphilis* **5**:248 (April) 1921.

the form of neurosyphilis from which the other has been suffering, perhaps, for years. Considerable laboratory work—particularly, of course, with the cerebrospinal fluid—is required to put familial neurosyphilis on a scientific basis. Cestan, Riser, and Stillmunkés⁴⁵ have contributed to this phase of the subject in an article in which the “colloidal benzoin” reaction of the spinal fluid is discussed.

THE WASSERMANN REACTION

Since April, 1920, the literature on this subject has been abundant—one might say, lavish. The technic, or rather the technics, of the test as employed by various observers cannot be discussed here. Studies of the test as applied to syphilitic children and their families are, of course, mostly statistical. In an analysis of 4,000 Wassermann tests on pregnant women, Williams⁴⁶ found 449 positive (11.2 per cent.). Forty-three women (1 per cent.) with negative Wassermann reactions, had syphilitic children. The author believes that the evidence shows that less than half (40 per cent.) of the positive mothers will have syphilitic children. A syphilitic placenta does not inevitably mean that the child will have the disease. It will be recalled that Jeans and Cooke² found that only 27 per cent. of cases of hereditary syphilis could be diagnosed by placental examination alone, and that 63 per cent. of the cases could be recognized by the Wassermann test on the cord blood. Williams found that a positive fetal blood Wassermann was associated with a syphilitic infant in 76 per cent. of cases. His results with the Wassermann test lead him to conclude that both paternal infection of the fetus and Colles' law have neither been proved nor disproved. Having seen families in which the father has a positive Wassermann reaction and the mother a negative one, he is not as skeptical about Colles' law as are many other observers.

In a somewhat similar Wassermann study, Nobécourt and Bonnet⁴⁷ found that in pregnant women there were more positives in the seventh than in the fourth, fifth and sixth months. There was agreement between the blood of the mother and that of the baby in 90 per cent. of the cases. The majority of discrepancies were the positive—mother-negative-child reactions—which the authors attribute to postconceptional syphilis.

Goodman⁴⁸ found that in a series of 1,320 pregnant women, 87 per cent. were Wassermann negative. Of the Wassermann negative multiparas, 37 per cent. had suffered one or more miscarriages, as

45. Cestan, Riser, and Stillmunkés: *Bull. de l'Acad. de méd., Par.* **86**:116 (July 26) 1921.

46. Williams: *Bull. Johns Hopkins Hosp.* **31**:336 (Oct.) 1920.

47. Nobécourt and Bonnet: *Presse méd.* **28**:745, 1920.

48. Goodman: *Surg. Gynec. and Obst.* **30**:368, 1920.

compared with 52 per cent. of the four plus cases. Only one woman of the entire series gave a history of having been known to be infected with syphilis.

It is becoming increasingly clear that the interpretation of the Wassermann reaction, and the recording of the degree of positivity are two factors which should not be overlooked in evaluating the results of Wassermann surveys. Yet in many such surveys they are not considered. The importance of attaching proper significance to the Wassermann report is emphasized by Heagerty.⁴⁹ Quoting the All-American Conference on Venereal Diseases (December 1920): "A frank, reliable, positive Wassermann reaction should be regarded as a sign of syphilis. In the absence of all other evidences of syphilis, a diagnosis based on a positive Wassermann alone should be made with great caution, and the Wassermann report confirmed by repetition and examination in another laboratory. A negative Wassermann cannot be regarded as evidence of the absence of syphilis." Browne⁵⁰ draws practically the same conclusions.

Thaysen⁵¹ discusses the cases in which a positive Wassermann reaction does not indicate active syphilis, and analyzes active syphilis from both the clinical and biologic points of view. He declares that notwithstanding the positive Wassermann reaction in the late latent stage, the syphilis may be regarded as inactive in a large proportion of cases. His article serves to show that the terms active, inactive, and latent, are capable of various interpretations, and that such interpretations are of considerable importance to the patient from the therapeutic point of view.

The necessity for understanding the limitations of the Wassermann test is emphasized by Rohdenburg.⁵² Three important influencing factors are named: (1) The patient. Fever and other conditions, such as recent ingestion of alcohol may give a falsely positive result. Treatment should be discontinued for at least eight weeks before a negative report is accepted. (2) The collection of the specimen. (3) The technic of the test. In 148 cases with Wassermann reaction negative for three years following treatment, the reaction again became positive in five patients. (Spinal puncture was not reported in these cases.) The late quiet cases of hereditary syphilis (six in number) have never had even a temporarily negative reaction in spite of enormous dosage and long treatment. (They apparently had no examination of the spinal fluid.)

49. Heagerty: *Canad. M. A. J.* **11**:548 (Aug.) 1921.

50. Browne, F. J.: *Edinburgh M. J.* **27**:199 (Oct.) 1921.

51. Thaysen: *Ugesk. f. Laeger* **83**:759 (June 9) 1921; abstr. *J. A. M. A.* **77**:658 (Aug. 20) 1921.

52. Rohdenburg, Garbat, Spiegel and Manheims: *J. A. M. A.* **76**:14 (Jan. 1) 1921.

Discussing the various factors responsible for the positivity of the Wassermann test, Wassermann himself says⁵³ that not only protein substances, but also lipoids may produce genuine antibodies (an observation appearing more and more frequently in the literature on anaphylaxis). The syphilitic suffers from an inversion of lipid metabolism, which explains why the Wassermann reaction is positive not only with the extracts from organs of all stillborn syphilitic children, but also with all extracts containing lipid-like substances. Epstein and Paul⁵⁴ emphasize that colloidal, electric, and surface tension reactions are involved, not changes in the chemical structure of the reacting substances.

With regard to the limitations of the Wassermann reaction in hereditary syphilis, Rolleston⁵⁵ states that a positive reaction at birth is not certain evidence of syphilis, for (McIntosh and Fildes) the reacting substance may pass from mother to child in utero. The Wassermann reaction may be negative for weeks after birth, then positive. It tends to become negative about the time of puberty; and the disappearance of a positive Wassermann does not mean the absence of spirochetes. Various organs, especially the glands of internal secretion, may become involved, and the syphilis then become inactive, the symptoms being only those of involvement of such organs. If such cases are known to be syphilitic, negativity of the Wassermann does not mean that treatment is unnecessary. Quoting the work of Jeans,⁵⁶ the author adds his own experience to show the importance of testing the spinal fluid in cases of hereditary syphilis.

Cooke⁵⁷ reports 1,855 complement fixation tests for tuberculosis on 1,556 children. He makes the important observation that nonspecific fixation reactions were found in two conditions, hereditary syphilis and diphtheria. Of seventy-five hereditary syphilitics, with strongly positive Wassermann tests, twenty-one, or nearly a third, gave positive fixation tests for tuberculosis. None of the twenty-one had any demonstrable evidence of tuberculosis, though some of the older children had the usual positive Pirquet skin test. The author considers it likely that in children with a high concentration of syphilitic antibodies in the blood, the tubercle antigen has an action similar to the other non-specific lipid antigens used in the Wassermann test. The antibodies giving the Wassermann reaction are usually present in considerably higher concentration in children with hereditary syphilis than in adults with the acquired form of the disease. None of the children under one

53. Wassermann: reported in Berlin Letter, J. A. M. A. **76**:463 (Feb. 12) 1921.

54. Epstein and Paul: Med. Klin. **17**:913 (July 24) 1921; abstr. J. A. M. A. **77**:1216 (Oct. 8) 1921.

55. Rolleston, H.: Lancet **1**:471 (March 5) 1921.

56. Jeans: Am. J. Dis. Child. **18**:173, 1919.

57. Cooke: Am. J. Dis. Child. **21**:78 (Jan.) 1921.

year reacted positively to the fixation test for tuberculosis. Similarly, infants with hereditary syphilis may not develop a positive Wassermann reaction for several months. The inability to form fixing antibodies may be associated with a lack of ability to form other protective antibodies.

Another condition requiring special interpretation of the Wassermann reaction, according to Henes,⁵⁸ is hypercholesterinemia. Pregnancy, jaundice and typhoid are three of the commonest causes of increased cholesterol content of the blood. In these conditions a strongly positive reaction should be somewhat discounted, a weakly positive one given but little weight. Conversely, in chronic nephritis and in the majority of pyrexias, the cholesterol content of the blood is diminished and corresponding allowance must be made for this in reading the Wassermann test.

To investigate the not uncommon phenomenon of so-called "Wassermann-fastness," Strickler⁵⁹ endeavored to determine whether or not some of the persistently positive reactions were due to arsphenamin treatment alone. Thirty patients with various skin diseases, whose histories and physical examinations were entirely negative for syphilis, whose first Wassermann tests were negative, were given weekly injections of arsphenamin. Only eight of the completely followed group of twenty-four remained consistently negative. Nine gave a 2 plus or greater degree of positivity, five were 1 plus, two were doubtful. As none of the patients was syphilitic, these researches could not be called experiments on the provocative Wassermann reaction. In explanation of these arsenic induced reactions, the authors suggest that the arsenic in some way acts on the liver alone or on the spleen and bone marrow also; that one or all of these organs may, as a result of the arsphenamin, elaborate a lipid substance which may yield a positive Wassermann. Commenting on these observations, Kolmer⁶⁰ says that it is commonly believed that the serum of a syphilitic patient may give a negative Wassermann reaction if tested during or immediately after intravenous treatment. According to Strickler and his associates, just the reverse is to be expected. The observations of Strickler are not to be lightly dismissed, and should be held sub judice. At present Kolmer is unconvinced that arsphenamin produces such changes in nonsyphilitic individuals as would result in their serums yielding genuine positive Wassermann reactions.

A further study of Wassermann fastness is that of Stokes and Busman,⁶¹ who found that in 458 treated patients, 22 per cent. of

58. Henes: *Am. J. Syphilis* **4**:685 (Oct.) 1920.

59. Strickler, Munson and Sidlick: *J. A. M. A.* **75**:1488 (Nov. 27) 1920.

60. Kolmer: Letter to *J. A. M. A.* **75**:1796 (Dec. 25) 1920.

61. Stokes and Busman: *Am. J. M. Sc.* **110**:658 (Nov.) 1920.

latent, late and hereditary cases remained persistently Wassermann positive. They found no evidence that Wassermann fastness is the result of any special strain. Persistent positivity seems to accompany grave, rather than mild syphilis.

There is a considerable amount of recent literature on the subject of diseases other than syphilis which give a positive Wassermann reaction. Touraine⁶² names a number of tropical diseases in this class, mentioning leprosy as giving positive tests in 47 per cent. of cases—the tuberculous form even in 80 per cent.; pulmonary tuberculosis in 23.2 per cent.; lupus in 50 per cent. Prolonged ether or chloroform narcosis will give a positive Wassermann in 25 per cent. of cases. A number of blood diseases such as hemoglobulinuria, leukemia, etc., occasionally also contribute to the sources of possible error.

Recent evidence is against malaria as a cause of Wassermann positivity. Johnson⁶³ asserts that if a case is not clinically syphilitic, a positive Wassermann in any condition requires confirmation by repetition. In the various forms of malaria, the blood does not give a positive Wassermann reaction. Other causes must be operative. Iyengar⁶⁴ tested the serums of eighty-four cases of undoubted malaria in adults, and of fourteen cases in children. Seven gave a positive Wassermann. The author believes that malaria is not one of the causes of a positive reaction, and that quinin does not affect the reaction.

Studying fifty serums from patients suffering from typhus fever, Bauer⁶⁵ found that forty-six (92 per cent.) gave a positive Wassermann reaction, though none of them had any evidence of syphilis. The serums were taken during the fever period. He concludes that the Wassermann reaction is almost always positive in typhus fever when applied to inactivated serum, if the sample is taken before the crisis. He considers this of diagnostic value, since in many control cases of typhoid and paratyphoid solely negative results were secured.

Mason⁶⁶ reports two cases of diabetes mellitus, each with a strongly positive Wassermann reaction, neither having any history or symptom or sign of syphilis. Both patients were treated for their supposed syphilitic infection over a short period of time, with marked and rapid decline of their carbohydrate tolerance. The author thinks that probably the arsenic acted as a toxic agent on the weakened pancreatic cells, altering the quality or nature of their internal secretion.

It is noteworthy that Graves⁶⁷ found, in ninety controlled cases, flat discrepancies between antemortem and postmortem Wassermann reac-

62. Touraine: *Rev. de méd.* **37**:103, 1920.

63. Johnson: *J. Path. and Bacteriol.* **24**:145 (April) 1921.

64. Iyengar: *Indian J. M. Res.* **8**:136 (July) 1920.

65. Bauer: *München. med. Wchnschr.* **68**:1251 (Sept. 30) 1921.

66. Mason: *Am. J. M. Sc.* **162**:828 (Dec.) 1921.

67. Graves: *J. A. M. A.* **75**:592 (Aug. 28) 1920.

tions in only two. The postmortem Wassermann is practically as reliable as the antemortem if the serum is taken properly and if nothing unusual appears in the serum control tube.

There is persistent difference of opinion on the "provocative Wassermann reaction." Indications are that the test is losing favor. O'Leary⁶⁸ describes the method of performing the test which is used at the Mayo Clinic. Solomon and Klauder⁶⁹ found that after intravenous (or intraspinal) treatment the cerebrospinal fluid, negative before treatment, may become positive in all routine tests, or a weakly reacting fluid may become much stronger. This is compared with the Herxheimer reaction, or the production of neurorecurrences. Clinical improvement may result, and continued treatment may again produce a negative fluid.

Rusca's experience⁷⁰ with the Wassermann reaction in human milk shows that specific antibodies are present in breast milk, even in advanced lactation. The reaction is not as sensitive as that in the blood, but a positive reaction in breast milk is strong presumptive evidence in favor of grave and still active syphilis. Under specific treatment the reaction becomes negative in the milk before it disappears from the blood. Attempts at performing the Wassermann reaction on the urine have resulted unsatisfactorily on account of the frequency of anticomplementary reactions.⁷¹

FLOCCULATION REACTIONS IN SYPHILIS

In a review of this subject, Levinson⁷² gives brief descriptions of six methods of performing flocculation tests on serum. He states that none of these has been found to be specific. The Bruck silver nitrate test, already reviewed by Jeans,¹ has also been found by Levinson to be unreliable. He quotes Vernes'⁷³ opinion that in order to produce flocculation with syphilitic serums and not with normal serums, an organic colloidal preparation, rather than a mineral suspension, is necessary. He discusses the Vernes phenomenon in some detail.

The Meinicke reaction, also discussed by Levinson, is based on the hypothesis that the reaction between serum and extract takes place when extract colloids disturb the isotonicity of salt solution permitting the union of seroglobulins and lipid extract.

The Sachs-Georgi reaction⁷⁴ is a physicochemical reaction between seroglobulins and lipid extract. The globulins in syphilitic serums

68. O'Leary: *Arch. Dermat. & Syph.* **2**:348 (Sept.) 1920.

69. Solomon and Klauder: *Arch. Dermat. & Syph.* **2**:679 (Dec.) 1920.

70. Rusca: *Pediatrics* **29**:121 (Feb. 1) 1921; abstr. *J. A. M. A.* **76**:968 (April 2) 1921.

71. Klauder and Kolmer: *J. A. M. A.* **76**:102 (Jan. 8) 1921.

72. Levinson: *Am. J. Syphilis* **5**:414 (July) 1921.

73. Vernes: *Compt. rend. Acad. d. sc.* **165**: 1917; 166, 167, 1918.

74. Sachs and Georgi: Reference given by Levinson.⁷¹

are increased in amount, and the flocculate which occurs in this reaction is a lipoglobulin aggregate. It is the simplest of all flocculation reactions. Levinson compared the Wassermann, Meinicke and Sachs-Georgi reactions, using animal serums known to give positive Wassermann reactions normally. His study gives the Sachs-Georgi reaction an advantage over the Meinicke reaction. None of the flocculation reactions can at present supplant the Wassermann test. The Sachs-Georgi reaction becomes positive earlier and remains positive longer than does the Wassermann reaction.

Epstein and Paul⁷⁵ performed the "third modification" of the Meinicke test on 1,100 serums. It proved as reliable as the Wassermann test, and, the authors believe, has a number of advantages over the latter. Gloor⁷⁶ found that the Sachs-Georgi reaction agreed with the Wassermann in a higher percentage of cases than did the Meinicke reaction.

A very large number of articles have appeared giving parallel statistics of the Sachs-Georgi and Wassermann tests.⁷⁷ Parker and Haigh⁷⁸ present many such statistics of other observers, together with data of their own. They conclude that the great simplicity of the Sachs-Georgi test, and its encouraging agreement with the Wassermann, make it worthy of further study. The consensus of opinion is well summarized by Logan,⁷⁹ when he says that unless the factors which may cause the false reactions are definitely recognized and can be controlled, the test should not be used to supplant the Wassermann reaction. It may be used profitably as a control for the Wassermann.

With regard to the Sachs-Georgi reaction in the spinal fluids, Harryman⁸⁰ finds that the results closely parallel those of the Wassermann. He believes that the test is a substitute, or may be used as a valuable addition to the spinal fluid Wassermann. It furnishes a means for an earlier diagnosis of syphilis of the central nervous system than does the Wassermann. Mandelbaum⁸¹ believes that the substance responsible for the positivity of the Sachs-Georgi test is not albumin, nor globulin, but rather a member or members of the lipid group.

75. Epstein and Paul: *Med. Klin.* **17**:1118 (Sept. 11) 1921; abstr. *J. A. M. A.* **77**:1776 (Nov. 26) 1921.

76. Gloor: *Schweiz. med. Wchnschr.* **24**:466, 1920.

77. Taniguchi: *Brit. J. Exper. Path.* **2**:41 (Feb.) 1921. Taniguchi and Yoshinare: *Brit. M. J.* **2**:239 (Aug. 13) 1921. Kurer: *Wien. klin. Wchnschr.* **33**:562, 1920. Wolfstein: *Berl. klin. Wchnschr.* **56**:1110 (Nov. 24) 1919. Hull and Faught: *J. Immunol.* **5**:521 (Nov.) 1920. Messerschmidt: *Deutsch. med. Wchnschr.* **46**:150 (Feb. 5) 1920. Kilduffe: *Arch. Dermat. & Syph.* **3**:415 (April) 1921.

78. Parker and Haigh: *Arch. Dermat. & Syph.* **4**:67 (July) 1921.

79. Logan: *Lancet* **1**:14 (Jan. 1) 1921.

80. Harryman: *Arch. Dermat. & Syph.* **4**:299 (Sept.) 1921.

81. Mandelbaum: *München. med. Wchnschr.* **67**:962, 1920.

Some of the most important work on the phenomenon of serum-flocculation in the presence of colloidal solutions has been done by Vernes.⁸² He outlines the method of the "Vernes" test, in which a colloidal solution, guinea-pig serum and red cells are used, with the formation of a colorimetric scale. Marie,⁸³ reviewing the subject, says that syphilitic serum induces flocculation a little faster than normal serum—that it is merely a question of degree in a general physico-chemic phenomenon; and that Vernes' latest reaction is based on the superflaking properties of syphilitic serums as estimated with a standard opalescent scale. In Marie's opinion, it is scarcely probable that syphilis is the only disease that induces physicochemic changes in the serum; and evidence is accumulating that serum in tuberculosis and cancer may have a special action on suspensions of colloids. Cornwall and Aronson⁸⁴ believe that if confirmed the researches of Vernes will revolutionize the laboratory diagnosis of syphilis.

MISCELLANEOUS TESTS FOR SYPHILIS

The liquor formaldehydi test has not been found to be accurate. The test consists of adding a drop of ordinary liquor formaldehydi to serum without further dilution, and allowing the whole to stand for twenty-four hours at room temperature. Coagulation of the serum is a positive result. Suffern⁸⁵ believes that the test is not without value. Ecker⁸⁶ performed both the Wassermann and the formol reaction of Gates and Papacostas, in a series of 500 serums. He concluded that the formol reaction, as it stands, is of no diagnostic value because of its failure to react in clinically and serologically clear cut cases of syphilis, and because of the occurrence of positive reactions in the absence of the disease.

An interesting new serum reaction is the "sigma" reaction of Dreyer and Ward,⁸⁷ described by them as a "simple quantitative serum reaction." The authors maintain that it is more dependable than the Wassermann. The readings are expressed in terms of standard flocculation units.

Working on the speed of sedimentation of the red cells in infancy, György⁸⁸ found that in the citrated blood of infants of about 1 month, the red cells show a normal increase in the rate of sedimentation. In infants less than 1 month old, the average rate is very much slower.

82. Vernes: *Rev. neurol.* **27**:156 (Feb.) 1920.

83. Marie, P.: *Presse méd.* **28**:856 (Nov. 20) 1920; abstr. *J. A. M. A.* **76**:144 (Jan. 8) 1921.

84. Cornwall and Aronson: *J. A. M. A.* **75**:1697 (Dec. 18) 1920.

85. Suffern: *Lancet* **2**:1107 (Nov. 26) 1921.

86. Ecker: *J. Infect. Dis.* **29**:359 (Oct.) 1921.

87. Dreyer and Ward: *Lancet* **1**:956 (May 7) 1921.

88. György: *München. med. Wchnschr.* **68**:808 (June 30) 1921.

In acute infections, tuberculosis, and especially in hereditary syphilis, a further marked rise in the rate of sedimentation is found. When other causes of increased rate have been ruled out, the test can be made of real service in the diagnosis of hereditary syphilis. It parallels the Wassermann and Sachs-Georgi reactions.

Starting with the known fact that inactivation of serum increases its viscosity, Holke⁸⁹ found that prolonged standing at 56 C. increases the viscosity of serums of syphilitics, treated or untreated, negative or positive, more readily than that of normal persons with negative Wassermann reactions.

With regard to the luetin test, Ward⁹⁰ found 87 per cent. of positive luetin reactions in syphilitics. In a second series of 200 cases there were 75 per cent. corroborative returns. He says that luetin as a measure of the allergic reaction in syphilis has a higher reaction both negatively and positively than has been emphasized before. Luetin itself is in need of standardization and could easily be improved.

COLLOIDAL AND OTHER REACTIONS IN THE CEREBROSPINAL FLUID

Unquestionably the colloidal gold test has successfully withstood the manifold investigations to which it has been subjected. Articles endeavoring to show that it is of value seem no longer to be necessary. Articles explaining and interpreting its action are commonly encountered. Such an article is that of Cruickshank,⁹¹ whose results confirm those of other workers. The substance in the spinal fluid of general paretics which causes the precipitation of colloidal gold is not dialyzable, and resides in the globulin fraction of the protein, and in this respect resembles the Wassermann reacting substance. It is not altered by heating to the coagulation point of protein. Syphilitic reactions are due to an amount of albumin sufficient to obscure partially the precipitating effect of the globulin. The precipitating action of paretic fluids cannot be ascribed solely to the increase in globulin, but is probably dependent on a specific alteration of the physical state of the globulin which is associated with a positive electrical charge.

Thompson⁹² presents the results of 8,400 colloidal gold tests, made on about 7,100 cerebrospinal fluids at the Massachusetts State Psychiatric Institute. Six hundred and seventy-seven gave paretic curves. The author believes that a paretic curve points toward a parenchymatous involvement of the brain, while the milder gold curves are obtained in meningitis, vascular disease, etc. The paretic curve is

89. Holke: *J. Path. & Bacteriol.* **24**:413 (Oct.) 1921.

90. Ward: *Am. J. Syphilis* **5**:482 (July) 1921.

91. Cruickshank: *Brit. J. Exper. Path.* **1**:71 (April) 1920.

92. Thompson: *Arch. Neurol. & Psychiat.* **5**:131 (Feb.) 1921.

also of considerable prognostic value. Bonsmann⁹³ found that a negative colloidal gold test could be relied on as excluding central nervous syphilis. Multiple sclerosis and brain tumors are likely to give misleading findings.

Keidel and Moore⁹⁴ state that in view of the complicated methods of preparing reagents for the colloidal gold test, the simplicity and uniformity of the colloidal mastic test (first described by Emmanuel in 1915) has much to recommend it. Both depend on colloidal phenomena. The suspension of finely divided colloidal mastic particles undergoes quantitative flocculation, as in the colloidal gold test, also culminating in complete precipitation. The authors' results on the spinal fluids of 311 patients show close parallelism between the two tests. When agreement is lacking, the mastic test seems to detect abnormalities more frequently than does the colloidal gold test. This fact, together with the simplicity of the test, lead the authors to conclude that the mastic test should be an indispensable part of spinal fluid examinations. According to Stanton,⁹⁵ the results of investigators using the mastic test have been discordant. Certain samples of gum mastic which have been found to be entirely unsuited for use in the mastic test may explain the conflicting reports.

Cestan et al.⁴⁵ note that the "colloidal benzoin" reaction often remains positive when the Wassermann has been rendered negative by treatment. This test is more likely to agree with the globulin tests than with the Wassermann and cell count.

Hektoen and Neymann⁹⁶ advance a test which they maintain is a definite index of the amount of protein in spinal fluid. Antihuman rabbit serum is added to various dilutions of the spinal fluid. In paresis they found an increase in the albumin and globulin, chiefly globulin. In epilepsy the increase is in albumin.

SYPHILIS OF THE CENTRAL NERVOUS SYSTEM

Fortunately for the proper understanding of hereditary syphilis, more work is being done on the disease as it affects the central nervous system. As in adults, almost any neurologic condition may be simulated. Tabulating the details of 114 cases of inherited neurosyphilis, Mensi⁹⁷ calls particular attention to a girl of 8 who presented the complete clinical picture of cerebrospinal sclerosis. A girl of 7 presented the spastic paralysis with a tendency to obesity, described by Marfan.

93. Bonsmann: *Deutsch. Arch. f. klin. Med.* **134**:20 (Oct. 26) 1920; abstr. *J. A. M. A.* **76**:901 (March 26) 1921.

94. Keidel and Moore: *Arch. Neurol. & Psychiat.* **6**:163 (Aug.) 1921.

95. Stanton: *Arch. Neurol. & Psychiat.* **4**:301, 1920.

96. Hektoen and Neymann: *J. A. M. A.* **75**:1332, 1920.

97. Mensi: *Riv. di clin. pediat.* **19**:1 (Jan.) 1921; abstr. *J. A. M. A.* **76**:1539 (May 28) 1921.

Psychic anomalies may be due to lesions of the nervous system, or to perverted functioning of the ductless glands.

Tilton⁹⁸ reviews the literature and reports three cases of cerebro-spinal syphilis in one family. Undoubtedly, hereditary syphilis is responsible for some cases of idiocy and imbecility. Thom⁹⁹ believes that it is a most important factor. Haines and Partlow¹⁰⁰ found that of 672 children in Alabama industrial schools, 118 were feeble-minded. They found no evidence of a high rate of syphilis among the mentally inferior. In a more intensive study of a smaller series, Raeder¹⁰¹ examined thirty children from 2 to 16 years old, in either or both of whose parents there was a positive Wassermann reaction (blood). "Most of the parents" had neurosyphilis, usually of the paretic type. Nineteen of the children had positive blood Wassermann reactions. Spinal puncture was done on twenty-two of the thirty children. Four of the fluids were positive. In three of these the colloidal gold curves were paretic. In the children born nearer the parental infection, not only were the physical findings and the Wassermann reactions on serum and spinal fluid positive, but these children also showed a greater degree of mental deficiency. Children born later showed less mental defect. It will be seen from the studies reported that selection of the series is an important factor in determining the extent to which inherited neurosyphilis is responsible for mental deficiency.

Virtually all workers are agreed that early examination of the spinal fluid is of paramount importance, from the point of view of therapeutics and of prognosis. It is, of course, the only means by which an asymptomatic neurosyphilis may be discovered.⁹⁹ Bigland¹⁰² says that without examination of the spinal fluid no case of organic central nervous disease should be considered syphilitic even when the blood Wassermann is positive. He also emphasizes that every case of organic central nervous disease requires spinal fluid examination even in the absence of suspected syphilis.

Kingery¹⁰³ studied the spinal fluid in fifty-two cases of hereditary syphilis. Fifteen, or 28.8 per cent. presented abnormal findings. There was a striking parallelism between the strongly positive Wassermann reactions and definite albumin and globulin increase; and an equally striking lack of parallelism between the degrees of pleocytosis and other strongly positive findings. There were four juvenile paretics and two mentally deficient cases in the series.

98. Tilton: *Northwest Med.* **20**:251 (Sept.) 1921.

99. Thom: *J. Ment. & Nerv. Dis.* **53**:8 (Jan.) 1921.

100. Haines and Partlow: *Ohio State M. J.* **16**:515 (July) 1920.

101. Raeder: *Am. J. Dis. Child.* **21**:240 (March) 1921.

102. Bigland: *Lancet* **2**:687 (Oct. 2) 1920.

103. Kingery: *J. A. M. A.* **76**:12 (Jan. 1) 1921.

In an interesting article on tuberculous meningitis and hereditary syphilis, Hutinel and Merklen¹⁰⁴ note that they have seen the two diseases associated in more than twenty cases in the past few years. They consider it proper to assume that as hereditary syphilis is an important cause of meningeal dysfunction, the implantation of tubercle bacilli as well as other germs is thereby favored. It was not until the authors began doing Wassermann tests on all cases of tuberculous meningitis that they began to realize the frequency of the association of the two conditions. (These observations should be considered in the light of Cooke's observations⁵⁷ on cross-fixation in cases of tuberculosis and of syphilis. In any event the degree of positivity of the Wassermann is of great importance in its interpretation in tuberculous patients.)

Mingazzini¹⁰⁵ reports two brothers and a sister, all of whom began to suffer at puberty from a progressive motor disturbance of the lower extremities, terminating in the syndrome of family spastic paralysis. One of the brothers also suffered from optic atrophy, choroiditis and double cataract. The father of these children was a tabetic. With the exception of the first-born (contrast with Raeder's observations¹⁰¹) all his children were more or less deficient mentally. All were syphilitic.

From five cases which Ciuffini reports,¹⁰⁶ he warns that cerebral arteritis from inherited syphilis may develop with an onset like that in the acquired disease, and the course may be the same. In eight cases, nocturnal frontal headache was the main symptom of neurosyphilis.

Parker¹⁰⁷ reviews the literature on juvenile tabes and gives a summary of seven cases. He directs attention especially to the insidious onset, lengthy and even latent course of the disease, frequency of optic atrophy, of incontinence of urine; and the relative rarity of such striking phenomena as ataxia, girdle sensations and lightning pains. There is frequently a paretic termination. The large immobile irregular pupil is more frequent with juvenile tabetic patients, in contrast to the myosis and Argyll-Robertson pupil of adult patients. Of six spinal fluid Wassermann tests, three were negative though the blood was positive. (Bigland¹⁰² would probably question the authenticity of these cases.) The observations of Rosenbeck¹⁰⁸ are in substantial agreement with those of Parker. He adds that the average age of onset of juvenile tabes is 15 years—a period equal to that between the primary lesion and tabetic manifestations in the adult. The prognosis in juvenile tabes is good for life, bad for vision.

104. Hutinel, V., and Merklen: *Arch. de méd. d. enf.* **24**:521 (Sept.) 1921.

105. Mingazzini: *Arch. Neurol. & Psychiat.* **5**:637 (June) 1921.

106. Ciuffini: *Riv. di patol. nerv.* **25**:226 (Jan.) 1921; abstr. *J. A. M. A.* **76**:1375 (May 14) 1921.

107. Parker: *Arch. Neurol. & Psychiat.* **5**:121 (Feb.) 1921.

108. Rosenbeck: *J. A. M. A.* **76**:572 (Feb. 26) 1921.

An article by Kerr,¹⁰⁹ in which he reports a case of juvenile tabes in a boy of 14, serves to illustrate how, in point of signs and laboratory findings necessary to a diagnosis of juvenile tabes, authorities differ in their criteria. The striking manifestations in Kerr's case were: (1) "painful hip," which the author believed should be interpreted as lightning pains; (2) primary optic atrophy; (3) absent knee jerks. In Kerr's opinion, these manifestations established the diagnosis of juvenile tabes in spite of the absence of a history pointing to hereditary syphilis, in spite of the negative Wassermann reaction in both blood and spinal fluid.

An important article on juvenile general paresis is that of Toni Schmidt Kräpelin.¹¹⁰ He reports thirty-nine cases of known juvenile paresis, including eight cases in which a wrong diagnosis was made, and seven cases of "preparalysis," in children who showed no clinical signs of metasymphilitic affection of the central nervous system, but in whom serologic findings characteristic of general paresis were more or less accidentally discovered. Over one fourth of the cases showed general arrest of development, one third were mentally deficient. Change in character was among the early signs. Acute onset with hemiplegia, epileptiform attacks and focal symptoms, appeared with relative frequency, suggesting a combination of epilepsy and brain syphilis. Optic atrophy without other tabetic symptoms was comparatively frequent. Total immobility of the pupil was more common than simple loss of the light reflex. In paralytic and preparalytic children, a peculiar hurried, toneless, and puffing manner of speech was noticed. Periods of excitement were common, but a persistent agitated condition was seen in only about 10 per cent. of the cases. The majority showed a typical Wassermann reaction in the blood serum and spinal fluid. No therapeutic results worth mentioning were obtained by any known treatment.

Dahlstrom¹¹¹ reports two cases of juvenile paresis, in one of which there was a misleading coincidence between trauma and the onset of the disease, with localizing neurologic signs which led to a considerable amount of brain surgery before the true condition was definitely established.

It is fortunately well-recognized that in both the acquired and hereditary forms of syphilis, infection of the central nervous system occurs early. The importance of early lumbar puncture has been urged for several years, notably by Klauder,¹¹² Fordyce,¹¹³ and in the

109. Kerr: *Lancet* **1**:598 (March 13) 1920.

110. Kräpelin, T. S.: abstr. by L. J. Pollock, *Am. J. Dis. Child.* **23**:310 (Sept.) 1920.

111. Dahlstrom: *Norsk Mag. f. Lægevidensk.* **82**:710 (Oct.) 1921.

112. Klauder: *Am. J. Syphilis* **3**:559 (Oct.) 1919.

113. Fordyce: *Am. J. M. Sc.* **161**:313 (March) 1921.

case of hereditary syphilis, by Jeans.¹¹⁴ Fordyce considers it probable that the *Spirochaeta pallida* invades the nervous system during the period of general dissemination or not at all. Wile and Hasley¹¹⁵ believe that when positive findings in the spinal fluid precede the changes in the blood, such a condition must be due to the greater delicacy of laboratory findings in the spinal fluid. Such results cannot be considered evidence for neurotropism. The authors' findings thus substantiate for the human host the observations of Brown and Pearce for dissemination in the rabbit.¹⁵

Both Bambarén¹¹⁶ and Savory¹¹⁷ urge the importance of investigating cases with epileptiform seizures for the presence of syphilis, hereditary or acquired. The former believes that the evidence is strongly against the idea of Strümpell that inherited syphilis has never been shown to be a factor in epilepsy. He cites Levy-Bing's thirteen cases of essential epilepsy, with a history of syphilis in all.

With regard to the accuracy of the blood Wassermann in neurosyphilis, Hanssen¹¹⁸ found that 13.5 per cent. of 105 patients with neurosyphilis had a negative Wassermann in the blood, but a positive one in the cerebrospinal fluid.

An experimental observation which may have some therapeutic significance is that of Weston.¹¹⁹ Working with seventeen cases of general paresis, he found that after the intraspinal injection of 1 c.c. of phenolsulphonaphthalein, it took the dye from twelve to sixty-eight minutes to appear in the urine—the time in normal subjects being from four to ten minutes. As the dye was not found in fluid drawn from the cisterna magna at any time up to five hours, the author concluded that absorption in these cases took place in the lumbar region. Whether these facts might account for the practical futility of intraspinal therapy in paretics, remains to be seen.

SYPHILIS OF THE EYES

Green's first article on this subject has been reviewed by Jeans.¹ In his later publication,¹²⁰ the literature is reviewed and complete statistics given for each form of ocular disturbance encountered in the 100 cases studied by him. Particular mention should be made of the

114. Jeans: Am. J. Dis. Child. **18**:173 (Sept.) 1919.

115. Wile and Hasley: J. A. M. A. **76**:8 (Jan. 1) 1921.

116. Bambarén: Siglo méd. **68**:485 (May 21) 1921; abstr. J. A. M. A. **77**:497 (Aug. 6) 1921.

117. Savory: Lancet **2**:296 (Aug. 7) 1920.

118. Hanssen: Acta med. Scand. **53**:748 (Jan. 14) 1921; abstr. J. A. M. A. **76**:903 (March 26) 1921.

119. Weston: Arch. Neurol. & Psychiat. **5**:58 (Jan.) 1921.

120. Green: Am. J. Dis. Child. **20**:29 (July) 1920.

nineteen cases of present or past interstitial keratitis (bilateral in seventeen cases) and of the fact that fifty-two of the 100 cases presented undoubted eyeground lesions. Fourteen different types of eyeground pathology are recorded, the most characteristic being peripheral and central punctate pigmentation. Papillary haze, or "veiling" of the optic nerve head was also quite common. Occasionally the intra-ocular lesion may be the only manifestation of the hereditary syphilis.

In the course of a comparative study of syphilis in whites and negroes, Zimmermann¹²¹ has the following to say about the hereditary cases. Only the late forms of hereditary syphilis are treated in the syphilis clinic of the Johns Hopkins Hospital Dispensary where 60 per cent. of the patients are colored. No noteworthy difference between the two races with respect to late hereditary syphilis was found. Interstitial keratitis occurred in 33.3 per cent. of the white males, and in 61.7 per cent. of the white females; in 35.7 per cent. of the colored males, and in 75 per cent. of the colored females. Such predominance of keratitis in the female has been noted frequently.

Baldino¹²² reports an unusual case of bilateral congenital cataract, with marked opacity of the crystalline lens in a case of hereditary syphilis with a positive Wassermann, saddle nose and Hutchinson's teeth.

SYPHILIS OF THE AUDITORY SYSTEM

In a Wassermann survey of twenty-three deaf mutes, thirteen were found to be positive by Ardenne.¹²³ The test was positive in eight of eleven congenital cases, and in five of twelve cases in which some disease was responsible for the deafness.

Ramadier¹²⁴ gives an illustrated description of a pneumatic (Hennebert's) test of the vestibule, a positive response being nystagmus or a slow movement of the eyeballs under compression or aspiration of the air in the external ear. This "sign of a fistula without a fistula" is evidence of some lesion in the internal ear alone, and this is comparatively common with inherited syphilis. The writer never found this test positive in the normal ear, nor in other forms of ear disease. No case has yet come to necropsy, but a specific osteitis of the bony capsule of the labyrinth would explain the clinical manifestations. The test is applied with a speculum and a rubber bulb. Hennebert¹²⁵ reports three cases of deafness due to hereditary syphilis. The first

121. Zimmermann: *Arch. Dermat. & Syph.* **4**:75 (July) 1921.

122. Baldino: *Riforma med.* **36**:833 (Sept. 11) 1920.

123. Ardenne: *J. de méd. de Bordeaux* **91**:379 (July 25) 1920.

124. Ramadier: *Presse méd.* **29**:624 (Aug. 6) 1921; abstr. *J. A. M. A.* **77**:975 (Sept. 17) 1921.

125. Hennebert: *Arch. Ital. di Otol. Rhinol. e Laringol.* **31**:198, 1920; abstr. *Am. J. Syphilis* **5**:159 (Jan.) 1921.

two constitute the classical picture of hereditary syphilitic labyrinthitis. The onset of the deafness in the first case dated back to early infancy, in the second to the age of 15 years. In one of the author's cases deafness did not begin till the age of 36 years, the case being a bona fide one of hereditary syphilis. As a rule, the deafness is very pronounced. Kay¹²⁶ reports nerve deafness due to hereditary syphilis in three children in one family, their ages being 7, 5 and 4 years. In each case the evidence of the disease was manifest between the first and second years. The Wassermann was positive in the oldest child and in the parents.

TEETH

Davidsohn¹²⁷ cites the findings of Hutchinson and Neumann as to the frequency of hypoplasia of the deciduous teeth in hereditary syphilis. Such hypoplasias are also to be found in other constitutional diseases. Knowledge of their occurrence in the milk teeth of healthy, rachitic and constitutionally sick children is essential to their correct evaluation in syphilitic children. Davidsohn is convinced that true Hutchinson's teeth are pathognomonic of the disease. Though less frequent, and seldom mentioned in the literature, the enamel defects of the permanent lower central incisors may be important manifestations of hereditary syphilis. Alterations in the entire dental morphology are noted. Whether narrow, slender teeth, with large defects, are to be regarded as in the same group as Hutchinson's teeth, is not yet determined, though this conception is not without foundation. Davidsohn and Davidsohn,¹²⁸ studying fifty cases of hereditary syphilis in children more than 6 years old, found Hutchinson's teeth in twenty-four, or 48 per cent.

Sabouraud¹²⁹ says that dental stigmata often reveal otherwise unrecognized hereditary syphilis. He emphasizes the fact that certain dental buds do not develop, and that the superior lateral incisors are more likely to be lacking—or even the four lateral incisors, both upper and lower. Sometimes the cutting edge is indented. This striking deformity has been noted on one jaw in all the teeth up to the premolars. "Fish-teeth," i. e., very narrow, sharp teeth, are among the many lesions described. Enamel defects of the molars are also considered frequent. Some molars may be found with five cusps instead of four—others, which have on the internal surface the "embryo of a fifth cusp"—the so-called "tubercle of Carabelli," or mammillary eminence. Sabouraud urges that in view of the permanence and

126. Kay: J. A. M. A. **74**:1162 (April 24) 1920.

127. Davidsohn, E.: Ztschr. f. Kinderh. **25**:249, 1920.

128. Davidsohn and Davidsohn: Deutsch. med. Wchnschr. **47**:1064 (Sept. 8) 1921; abstr. J. A. M. A. **77**:1775 (Nov. 26) 1921.

129. Sabouraud: La Médecine **2**:105 (Nov.) 1920.

importance of the dental signs of hereditary syphilis, all physicians should acquaint themselves thoroughly with them.

HAIR

In an article on the relation between alopecia areata and syphilis, Sabouraud¹³⁰ states that though many cases of alopecia areata occur without demonstrable syphilitic taint, the proportion of patients with inherited syphilis is too large for a casual coincidence. His numerous experiences with the cure of alopecia areata in children under treatment for syphilis corroborate the relation between them.

CIRCULATORY SYSTEM

There has been a marked increase in the contributions to this phase of hereditary syphilis during the past year and a half. It is interesting that in 183 Wassermann tests on eighty-four children of school age with heart disease, only one test was positive.¹³¹ (Allowance must, of course, be made for the social status of the children studied, the time between the Wassermann test and parental infection, etc.) Hahn¹³² claims to have demonstrated that the majority of cases of vascular neuroses in children are referable to hereditary syphilis and to syphilitic injuries to the fetus—a claim as yet unconfirmed by other writers.

With regard to aortitis, so common in adults with the acquired disease, Stolkind¹³³ was unable to find a single instance, in the literature, of definitely proven heredosyphilitic aortitis in older children or adults, though he does not doubt the existence of such cases. He cites a case of his own with Hutchinson's teeth, positive Wassermann, etc., with aortic regurgitation and considerable cardiac involvement, which necropsy showed to be a case of serofibrinous pericarditis with verrucous endocarditis of the aortic and mitral valves. The author insists on the difficulty of excluding cardiac lesions other than syphilitic.

Calvin¹³⁴ reviews thirty-one cases of thoracic aneurysm in children, and presents two cases of his own. He thinks the evidence far from conclusive that syphilis is the sole or even the common cause of aortic aneurysm in childhood. In fact, the greater number of these cases are due to other causes, such as infectious aortitis, erosion from without, and congenital malformations. In the first of a series of studies of aneurysm, Lucke and Rea¹³⁵ state that in 247 patients, the

130. Sabouraud: *Presse méd.* **29**:581 (July 23) 1921.

131. Donaldson: *New York State J. M.* **21**:176 (May) 1921.

132. Hahn: *Zentralbl. f. inn. Med.* **42**:601 (July 30) 1921; *abstr. J. A. M. A.* **77**:1295 (Oct. 15) 1921.

133. Stolkind: *Brit. J. Child. Dis.* **17**:126 (July-Sept.) 1920.

134. Calvin: *Am. J. Dis. Child.* **21**:327 (April) 1921.

135. Lucke and Rea: *J. A. M. A.* **77**:735 (Sept. 17) 1921.

youngest was 12 days old, the oldest 91 years. More detailed pediatric data are not given in this study.

Queyrat and Mouquin¹³⁶ found signs typical of aortic stenosis in a boy of 5, whose father had congenital cataract, considered by Cantonnet to be due to hereditary syphilis. The difficulty of definite proof that such lesions are really syphilitic has been shown by the preceding articles. The same statement is true of the mitral lesions.¹³⁷ With regard to the myocardium, Friedlander¹³⁸ is of the opinion that in many instances myocardial degeneration in infancy and early childhood rests on a basis of hereditary syphilis. There is no typical form of myocarditis which might be recognized as syphilitic. In cases of heredosyphilitic myocardial degeneration, rapid improvement has followed intensive antisiphilitic treatment.

KIDNEYS

Discussing the evolution of acute nephritis in hereditary syphilitics, Queslier¹³⁹ reports finding in the literature 101 cases in which the specificity of the nephritis was established by the avowal of the parents, stigmata of the inherited disease, and positive seroreaction. Seventy-three patients were less than 2 years of age, while the remainder were of all ages up to 23 years. In all cases other prominent causes of nephritis were excluded. Only six of the seventy-three infants survived the disease, while of the twenty-nine older individuals, eighteen succumbed to the nephritis. All varieties of nephritis were found. The acute parenchymatous type was common clinically, though, strange to say, not encountered at necropsy. Naturally, renal symptoms are often masked by extensive lesions in other organs. Rolleston⁵⁵ believes that the chronic interstitial type of nephritis in early life is not so frequently due to hereditary syphilis as was formerly thought.

BONES

Gaenslen and Thalhimer¹⁴⁰ contend that our knowledge of heredosyphilitic epiphysitis is incomplete, and question the commonly accepted conception that this condition occurs only in the early months of infancy. They report a case of epiphysitis in the greater trochanter of the femur of an adolescent boy, and attribute the lesion to hereditary

136. Queyrat and Mouquin: *Bull. et mém. Soc. méd. d. hôp. de Par.* **45**:1152 (July 21) 1921.

137. Babonnei and Denoyelle: *Bull. et mém. Soc. méd. d. hôp. de Par.* **45**:774 (June 2) 1921. Amblard: *Bull. et mém. Soc. méd. d. hôp. de Par.* **45**: (May) 1921. Merklen: *Bull. et mém. Soc. méd. d. hôp. de Par.* **45**:815 (June 9) 1921.

138. Friedlander: *Gaz. d. hôp.* **93**:64 (July 20) 1920.

139. Queslier: *abstr. in Arch. Pediat.* **38**:450 (July) 1921.

140. Gaenslen and Thalhimer: *J. Orthoped. Surg.* **3**:8, 1921.

syphilis because of (1) predominance of cartilage and the riotous appearance of the process of osteogenesis; (2) lack of completion of bone formation; (3) excess of calcium deposit in the cartilage which is in apposition to the areas of bone formation; (4) the obliterating endarteritis in the thickened periosteum.

LYMPH NODES

Fabris¹⁴¹ reports the results of examination of 2,605 infants for enlargement of the epitrochlear glands. In children who presented signs of inherited syphilis, epitrochlear enlargement was evident in 32 per cent. of those under 1 year, and in 33 per cent. of those up to 2 years of age. The figures were only 7 and 9 per cent. in tuberculous children, and 7 and 11 per cent. in those with rickets. Epitrochlear adenitis, therefore, may be regarded as a confirming sign in inherited syphilis, to judge from the author's 379 cases with the inherited taint.

DYSTROPHIES AND CONGENITAL MALFORMATIONS

Hutinel and Stevenin¹⁴² name three varieties of dystrophies which are of importance in connection with hereditary syphilis: (1) the partial, local, infectious—due directly to the lesions of the disease itself—as for example the bone lesions; (2) general dystrophies, attributable to impaired function of the organs affected—particularly the endocrine glands (a theory concurred in by Barthélémy). Such cases should be treated by organotherapy as well as by antisypilitic drugs; (3) Hereditary dystrophies due to transmission from parents to children of “organic debilities”—to the second and third generation, as for instance hyper- and hypo-thyroidism.

In the Naples children's clinic, de Stefano¹⁴³ investigated all cases with congenital anomalies. There was an overwhelming preponderance of syphilis in the antecedents of all except the cases of spina bifida.

Rios and Bisso¹⁴⁴ report a “monster from inherited syphilis.” O'Brien and Mustard¹⁴⁵ quote Socin's report of a case of phocomelia with death occurring seventeen days after birth, of hereditary syphilis. A heredosyphilitic gynandroid is also reported.¹⁴⁶ Queyrat's observations on malformations of the xiphoid in hereditary syphilis are well known. Malformed spermatozoa in heredosyphilitics have already been mentioned.²⁵ Clément Simon¹⁴⁷ also reports a case with this condition.

(To be continued)

141. Fabris: *Pediatrics* **29**:193 (March 1) 1921; abstr. *J. A. M. A.* **76**:1435 (May 21) 1921.

142. Hutinel and Stevenin: *Arch. de méd. d. enf.* **23**: Nos. 1, 2, 3, 4.

143. de Stefano: *Pediatrics* **29**:59 (Jan. 15) 1921.

144. Rios and Bisso: *Semana méd.* **28**:175 (Aug. 11) 1921.

145. O'Brien and Mustard: *J. A. M. A.* **77**:1964 (Dec. 17) 1921.

146. Laiguel, Lavastine, and Boutel: *Bull. Soc. méd.* **43**:754 (May 28) 1920.

147. Simon: *Bull. Soc. franç. de dermat. et syph.* No. 6, 193, 1921.