FURTHER OBSERVATIONS ON THE LYMPHATIC ORIGIN OF CHOLECYSTITIS, CHOLEDOCHITIS AND THE ASSOCIATED PANCREATITIS *

EVARTS A. GRAHAM, M.D., AND M. G. PETERMAN, M.D.
ST. LOUIS

The most common conceptions of the pathogenesis of infections of the gallbladder and bile tracts are based on four assumed possibilities: (1) descending infection from the liver by bacteria carried down in the bile, (2) ascending infections from the duodenum up the common bile duct, (3) hematogenous infections of the gallbladder and ducts, and (4) a spreading infection through the wall of the gallbladder from an inflamed contiguous organ. Of these possibilities, it is only the last two which take into serious consideration the actual infection of the deeper layers of the wall of the gallbladder, despite the fact that microscopic examinations of gallbladders removed at operation, at least those which we have studied, demonstrate that there are inflammatory changes not merely in the mucosa but also in the deeper layers and that usually these changes are more pronounced in the deeper layers. The other two possibilities are concerned only with the entrance of organisms into the lumen of the gallbladder, and they do not explain their entrance into its walls. They assume also that contact infection of the mucosa of the gallbladder occurs easily and frequently. But there is no direct evidence to support the idea that cholecystitis occurs, except perhaps occasionally, merely as a result of the entrance of bacteria into the lumen of the gallbladder; and there is much evidence, both clinical and experimental, that such infections are probably rare.

This evidence consists of a number of facts: (1) In experiments on ten dogs in which large amounts of pathogenic colon bacilli were injected into the normal gallbladder, no cholecystitis was produced sufficient to be demonstrable by gross appearance or clinical signs, except in one case. In order to produce it with any degree of regularity, it was necessary either to obstruct the outflow from the gallbladder by ligation of the cystic duct or to injure the blood supply by ligation of the cystic artery. In this connection a statement by Meyer, Neilson and Feusier¹ is of interest. These authors state:

* Part of this paper was read before the American Surgical Association, Toronto, June 15, 1921.

* From the Department of Surgery, Washington University School of Medicine.

1. Meyer, K. F.; Neilson, N. M., and Feusier, M. L.: The Mechanism of Gallbladder Infections in Laboratory Animals, *J. Infec. Dis.* **28**:456 (May-June) 1921.

It is generally known from the studies of Cushing and from our own, that typhoid bacilli introduced directly into the cystic bile of dogs disappear rapidly and that a cholecystitis can only be produced by considerable injury of the wall, or by placing a foreign body in the gallbladder (Marxer).

(2) Bacteria probably gain access to the gallbladder by being carried down in the bile from the liver in most, if not in all, cases of bacteriemia; yet cholecystitis is comparatively rare in such conditions. Rolleston² states that "in typhoid fever the bacilli are always present in the gallbladder, but cholecystitis is comparatively infrequent." Osler³ states that cholecystitis occurred in only nineteen of his series of 1,500 cases. (3) Theoretically, the mere presence of bacteria within the lumen of the gallbladder should be of no more significance in producing cholecystitis than that the presence of bacteria in the urine should imply an inflammation of the urinary bladder. It is well known that bacteria are frequently passed in the urine in large amounts without the coincidence of cystitis, especially in typhoid fever.

If the assumption is correct that the wall of the gallbladder beneath the mucosa is involved in most cases of cholecystitis, then it would seem reasonable to discard the idea of the serious importance of contact infections of the mucosa by organisms within the bile, particularly in view of the evidence outlined above. How then may an infection originate in the wall? It is obvious that in general only two routes are available for the transmission of organisms to the wall. One of these is the blood stream and the other is the lymph stream.

In 1909, Koch⁴ demonstrated clumps of bacilli in the wall of a gallbladder and concluded that typhoidal cholecystitis is due to capillary embolism. In the same year, also, Chiarolanza,⁵ working under Koch's direction, came to the same conclusion. Rosenow⁶ has particularly emphasized the idea of the hematogenous origin of cholecystitis. Meyer and his collaborators,¹ in their exhaustive article on the mechanism of gallbladder infections, report experiments on 500 rabbits in which they attempted to produce a condition analogous to the human typhoid carrier state. They consider that the findings of Koch and Chiarolanza of bacterial emboli in the wall of the gallbladder during

2. Rolleston, H. D.: *Diseases of the Liver, Gallbladder and Bile-Ducts*, New York, the Macmillan Company, 1912, p. 608.

3. Osler, William: *Practice of Medicine*, 1905, p. 83.

4. Koch, J.: Typhusbazillen und Gallenblase, *Ztschr. f. Hyg. u. Infektionskrankh.* **62**:1, 1909.

5. Chiarolanza: Experimentelle Untersuchungen ueber die Beziehungen der Typhusbazillen zu der Gallenblase und der Gallenwegen, *Ztschr. f. Hyg. u. Infektionskrankh.* **62**:12, 1909.

6. Rosenow, E. C.: The Etiology of Cholecystitis and Gallstones and Their Production by the Intravenous Injection of Bacteria, *J. Infec. Dis.* **19**:527 (Oct.) 1916.

Fig. 1.—A common type of hepatitis occurring in association with a subacute cholecystitis. The periportal sheaths are densely infiltrated with leukocytes, chiefly of the polymorphonuclear variety. From a study of small pieces of liver removed at operation in cases of cholecystitis, either with or without stones, it has been noted that some degree of hepatitis is a constant accompaniment of cholecystitis.

Fig. 2.—Higher power magnification of another part of the same liver showing accumulation of polymorphonuclear leukocytes around a small bile duct, indicated by line marked *A*.

Fig. 3.—Chronic hepatitis (biliary cirrhosis) in a woman who had a history of gallbladder trouble extending over a period of fifteen years. Jaundice had never been present; therefore, there had been no biliary obstruction. At operation, stones were found in the gallbladder. The marked thickening of the small bile ducts is strikingly evident.

Fig. 4. —Gallbladder removed at operation in case of cholecystitis without stones. Deep in the fibromuscular layers at a considerable distance from the mucosa were found collections of polymorphonuclear leukocytes, designated by lines marked *A*. These are usually much more conspicuous in the deeper layers than in the mucosa. Normally, the mucosa contains large numbers of round cells (lymphocytes).

Fig. 5.—Higher magnification of same gallbladder as shown in Figure 4. The leukocytic infiltration shown here is deep in the wall of the gallbladder at a considerable distance from the mucosa.

Fig. 6.—Another human gallbladder from case of cholecystitis without stones; wall extensively infiltrated with leukocytes; mucosa relatively intact.

Fig. 7.—High power magnification of Figure 6 showing polymorphonuclear leukocytes far out in the wall of the gallbladder.

Fig. 8.—Another human gallbladder: cholecystitis without stones; mucosa relatively intact, but in outer layers of wall there is rather extensive leukocytic infiltration.

typhoid to be "exceptionally rare" and concerned with an "exceedingly severe and unique form of typhoid cholecystitis." They arrive at this conclusion partly because in their own experiments, even after the intravenous injection of enormous doses of typhoid bacilli (8,000 to 24,000 million), only about one third of the rabbits presented histologic evidence of an infection through the terminal capillaries of the mucosa, and also because, as they state, examination of human gallbladders in cases of death from typhoid shows a capillary embolism only occasionally. They are inclined, however, to consider very seriously, especially in the rabbit, the possibility of contact infection of the mucosa of the gallbladder by bacteria carried down in the bile from the liver, and they hold the opinion that deep infections of the wall are frequently the result of a spread of the infection from the mucosa. In human typhoid carriers, they consider that "the embolic infection of the gallbladder wall plays an insignificant rôle." They state, furthermore, that "a condition analogous to that found in man apparently exists in the guinea-pig. The microscopic and cultural study of a limited number of gallbladders derived from infected guinea-pigs convinces us that the wall is rarely, and then only slightly, infected through the bile." This then seems to constitute another argument against the frequency of contact infection of the gallbladder in man. There can be little doubt apparently that cholecystitis not infrequently is hematogenous in origin; but yet we do not believe that an explanation based on this idea is sufficient for most cases of spontaneous cholecystitis, especially those which are not typhoidal in origin. The well known, striking tendency of cholecystitis to be associated with appendicitis and other inflammatory lesions of the portal system must be something more than a coincidence. Birch-Hirschfeld,⁷ in 1895, considered that organisms were carried to the liver and from there to the gallbladder by the bile. Since then, the "hematohepatogenous" idea of the origin of cholecystitis has been a popular one. But, as we have already shown, the idea of a contact infection of the mucosa of the gallbladder has very little to support it, at least so far as its being a frequent occurrence in man. Furthermore, it is difficult to explain the association of appendicitis and other portal infections with cholecystitis on the basis of a hematogenous infection of the wall of the gallbladder, because there is no direct path from the appendix to the gallbladder by way of the blood stream. The entire arterial supply of the gallbladder comes from the hepatic artery. The cystic artery supplies most of the blood; but there are in addition a few small branches of the hepatic artery which reach the gallbladder through its attachment to the liver. The veins of the gallbladder all empty into the portal vein. If organisms, therefore, are to be carried from the appendix to the gallbladder by the blood stream, they must

7. Birch-Hirschfeld: Quoted by Meyer et al, Footnote 1.

either travel through the whole systemic circulation back through the hepatic artery to the gallbladder or there must be a retrograde passage of them from the liver into the gallbladder through the tributaries of the portal vein. If the first alternative were correct, then there would be no more reason for cholecystitis to be associated with appendicitis than with infections elsewhere in the body, as the hand for example. The second alternative is based on the slender support of a comparatively rare possibility, a retrograde thrombosis, or at least a retrograde pylephlebitis.

There is, however, one route between the appendix and gallbladder which is comparatively direct and easy to understand, and that is a "hematolymphatic" route. Sudler⁸ has shown a very intimate lymphatic connection between the liver and gallbladder through the attachment of the gallbladder to the liver. This is shown in Figure 34, which is taken from his article. We have repeated some of Sudler's experiments and have confirmed his idea of the general arrangement and anastomosis of the hepatic and cholecystic lymphatics. The lymph vessels of the gallbladder may be seen after injections of such a substance as Prussian blue into the portal vein. Conversely, also, injections made into the wall of the gallbladder not only follow the lymphatics into the liver in the interlobular sheaths but also may be seen to pass along the lymphatics of the common duct and into the pancreas. An infection of the liver, therefore, would tend to spread to the gallbladder by way of the lymphatics, and conversely an infection of the gallbladder would tend to spread not only to the liver but also along the common duct to the pancreas. In this respect, infections in this region would merely follow the rule of infections elsewhere in the body. We should expect to find evidence, therefore, that in cases of cholecystitis there is also an associated inflammation in the liver.

In 1918, one of us⁹ showed in a series of cases in which small pieces of the liver were removed at operation for cholecystitis that microscopic evidence of hepatitis existed in every case. The inflammation was chiefly in the interlobular sheaths and appeared to be a pericholangitis. This location of inflammatory changes is in harmony with the idea of a lymphangitis, for it is here that the hepatic lymph vessels are chiefly situated.¹⁰ In another paper, we¹¹ showed that in the

8. Sudler: The Architecture of the Gallbladder, *Bull. Johns Hopkins Hosp.* **12**:126, 1901.

9. Graham, E. A.: Hepatitis: A Constant Accompaniment of Cholecystitis, *Surg., Gynec & Obst.* **26**:521 (May) 1918.

10. Mall, F. P.: On the Origin of the Lymphatics in the Liver, *Bull. Johns Hopkins Hosp.* **12**:146, 1901.

11. Peterman, M. G.; Priest, W. S., Jr., and Graham, E. A.: The Association of Hepatitis with Experimental Cholecystitis and Its Bearing on the Pathogenesis of Cholecystitis in the Human, *Arch. Surg.* **2**:92 (Jan.) 1921.

Fig. 9.—High power magnification of another human gallbladder showing small accumulations of polymorphonuclear leukocytes far out in the wall.

Fig. 10.—Cholecystitis with stones; whole wall, including mucosa, markedly inflamed.

Fig. 11.—Cholecystitis without stones; mucosa relatively normal, but leukocytic infiltration in outer layers of wall.

Fig. 12.—Cholecystitis without stones; normal mucosa but rather large collections of leukocytes (both polymorphonuclears and round cells) in deeper layers of wall, designated by lines marked A.

Fig. 13.—Hepatitis in association with chronic appendicitis; piece of liver removed at operation for chronic appendicitis; no evidence of cholecystitis. Note polymorphonuclears in lobule near its periphery.

Fig. 14.—Hepatitis with chronic appendicitis, showing leukocytic infiltration in interlobular tissue; piece of liver removed in another case of chronic appendicitis; no evidence of cholecystitis.

Fig. 15.—Hepatitis with chronic appendicitis; periportal leukocytic infiltration; piece of liver removed at operation for chronic appendicitis; no evidence of cholecystitis.

Fig. 16.—Experimental hepatitis by injection of portal vein with *Staphylococcus aureus* (Dog 1); marked periportal infiltration with polymorphonuclear leukocytes.

dog experimental cholecystitis is constantly accompanied by a hepatitis of the same sort as already observed in man. It is interesting that when this occurs secondarily to the production of a cholecystitis the right lobe of the liver is very much more affected than the left lobe, and the most marked changes of all are in the right lobe near the gallbladder. This all seems to indicate that the spread of the infection to the liver from an already inflamed gallbladder must occur through the lymphatics rather than through the veins, because in the latter instance, since the veins are tributaries of the portal vein, a more even distribution of the infection through the liver would be expected. The possibility of the establishment of a vicious circle between an infected gallbladder and the liver, whereby each may reinfect the other through the lymphatics, becomes readily apparent. This finding of the constancy of a hepatitis in association with cholecystitis has been confirmed recently by Judd,¹² who states, "A review of our studies inclines us to believe with Graham that cholecystitis rarely exists without hepatitis." In our original paper, to which reference has already been made, it was said that in a series of thirty consecutive cases coming to operation for gallbladder disease an enlargement of the liver was noted in 87 per cent. Subsequent observations have not borne out this high percentage, and we are forced to the conclusion that the actual percentage of noticeable enlargement of the liver is very much smaller than the figure given.

If we are to explain the frequent association of appendicitis with cholecystitis on the basis of a primary infection of the liver from the appendix, by way of the portal vein, and of a cholecystitis resulting from that secondarily, by way of a lymphatic spread from the liver to the gallbladder, then obviously it would seem necessary to find evidence of hepatitis in association with appendicitis. Recently, in a few cases which came to operation for chronic appendicitis, we removed small pieces of the liver for examination. Three of the accompanying photomicrographs (Figs. 13, 14 and 15) show areas of infiltration in the liver (both polymorphonuclears and round cells). In none of the cases represented here was there any evidence, either clinical or at operation, of cholecystitis. This fact seems important since in consequence the existing hepatitis could hardly be assumed to have been secondary to a cholecystitis. We have frequently noted slight liver tenderness in association with chronic appendicitis and have considered the sign of some importance in diagnosis. The question may arise as to why cholecystitis did not occur in the cases of hepatitis with appendicitis illustrated here. The answer is, of course, difficult, but perhaps the intensity of the hepatitis was not sufficient.

12. Judd, E. S.: Relation of the Liver and Pancreas to Infection of the Gallbladder, *J. A. M. A.* **77**:197 (July 16) 1921.

It is interesting that even those who favor the idea of the contact infection of the mucosa of the gallbladder by bacteria carried in the bile from the liver, the so-called "hematohepatogenous" theory, are forced to assume some degree of inflammation of the liver preceding the infection of the gallbladder. This question is discussed at some length by Meyer and his collaborators who state that "dosage, virulence and *lesions* [italics ours], and not the secretory, detoxifying activity of the liver, are the prerequisites for the passage of bacteria from the blood to the bile capillaries. . . . The conception of a purely mechanical passage deserves little consideration." Wyssokowitch¹³ and also Blachstein¹⁴ considered liver necroses prerequisites for the infection of the biliary passages in typhoid. If it is granted then that some degree of inflammation of the liver must be present before the bile can be infected, it seems only reasonable to assume that in many cases the infection will spread by means of the lymphatics and will therefore involve the gallbladder wall. Meyer states that "from an experimental standpoint the lymphatic route is of no importance," but yet he says that "in the course of peritonitis or other abdominal infections (appendicitis), streptococci can be transported through the lymphatics to the biliary passages," and again, "the so-called ascending route of infection is in all probability due to an invasion of bacteria through the lymphatics." He also states that in about 1 per cent. of the rabbits and guinea-pigs used there was a spontaneous cholecystitis due to *Bacillus coli*, streptococci or staphylococci. In these cases, "occasionally the bile appeared to be normal while the histologic picture exhibited a low-grade infection of the lymphatics of the mucosa and subserosa."

We believe that there is quite a little confusion about what is meant by the biliary lymphatics. Meyer and his collaborators speak of them as if they were entirely extrahepatic, and apparently they do not recognize that the extrahepatic lymphatics are continuous with those within the liver. This is the common conception of the lymphatics of the biliary tract. Sudler's important work is too little known. As a matter of fact, we do not see why nearly all the results on experimental cholecystitis cannot be just as easily interpreted on the basis of a lymphatic infection of the wall as on the idea either that there has been a contact infection of the mucosa or an infection of the wall directly through the blood stream. We admit, however, the possibility of infection through both of the latter means; but we are inclined to believe that the idea of a lymphatic origin secondary to a hepatitis explains the majority of cases of cholecystitis in man better than any other.

We have stated elsewhere that if a suitable substance, such as Prussian blue, be injected into the wall of the gallbladder, its course

13. Wyssokowitch: Quoted by Meyer et al, Footnote 1.

14. Blachstein: Quoted by Meyer et al., Footnote 1.

Fig. 17.—Experimental cholecystitis from Dog 1 by injection of portal vein; extensive inflammation of wall; mucosa relatively unaffected.

Fig. 18.—Higher power photomicrograph of specimen shown in Figure 17, showing leukocytic infiltration of wall of gallbladder.

Fig. 19.—Experimental hepatitis by injection of portal vein (Dog 3).

Fig. 20.—Another section from same liver as shown in Figure 19.

Fig. 21.—Experimental hepatitis (Dog 6).

Fig. 22.—Experimental hepatitis (Dog 7); polymorphonuclear leukocytes scattered throughout whole lobule.

Fig. 23.—Cholecystitis in same dog (Dog 7) ; polymorphonuclear leukocytes scattered throughout wall of gallbladder.

Fig. 24.—Same experiment as in Figure 23; polymorphonuclear leukocytes in deeper layers of wall of gallbladder.

may be followed through the lymphatics down along the common duct and into the head of the pancreas. For years, Deaver¹⁵ has strongly advocated the lymphatic origin of the chronic interstitial pancreatitis which is so common an accompaniment of cholecystitis. Maugeret,¹⁶ in 1908, wrote a thesis on this subject which offers much evidence in support of the possibility of the lymphatic spread to the pancreas. Archibald¹⁷ has raised questions with regard to the lymphatic route theory and has sought to explain the pancreatitis on a basis of an influx of bile into the pancreatic ducts in a manner similar to Opie's¹⁸ classical explanation of the origin of acute hemorrhagic pancreatitis. Deaver and Sweet¹⁵ have replied to these objections of Archibald's in their recent article.

In animal experiments, we have obtained results which we believe tend to support the idea of the lymphatic origin not only of cholecystitis but also of inflammation of the common duct and of the pancreas. These experiments have consisted of several types and will be reported in full by one of us (Peterman), in an article soon to appear. Two types are of particular interest here because of their special bearing on the idea of the lymphatic origin of cholecystitis and its complications. One of these groups of experiments has already been reported¹¹ and referred to above. It consisted in the production of hepatitis secondary to experimental cholecystitis, induced by direct inoculation of rather large amounts of organisms into the lumen of the gallbladder after ligation of the cystic duct and artery. The hepatitis produced conformed to that which had already been seen in cases of cholecystitis in man, and, for the reasons given above, it was considered as most probably representing a lymphatic spread to the liver. The other type of experiment consisted in the production of hepatitis, cholecystitis, choledochitis and pancreatitis by portal vein injections. It was felt that if cholecystitis and its common complications could be interpreted on the basis of a lymphatic spread from a liver in which the periportal spaces were the seats of inflammation, then it should be possible often in a single experiment to produce by injection of the portal vein a simultaneous hepatitis, cholecystitis, choledochitis and pancreatitis. Not only has it been possible to do this repeatedly; but the types of lesion found have agreed with those which occur in spontaneous cases in

15. Deaver, J. B., and Sweet, J. E.: Prepancreatic and Peripancreatic Disease: With a Consideration of the Anatomic Basis of Infection from the Gallbladder to the Pancreas, *J. A. M. A.* **77**:194 (July 16) 1921.

16. Maugeret, R.: *Cholécyste—Pancréatite*, Thesis, Paris, G. Steinheil, 1908.

17. Archibald, E.: The Experimental Production of Pancreatitis in Animals as the Result of the Resistance of the Common Duct Sphincter. *Surg., Gynec. & Obst.* **28**:529 (June) 1919.

18. Opie, E. L.: *Diseases of the Pancreas*, Philadelphia, J. B. Lippincott & Co., 1903.

man. It has been possible also after the injections to demonstrate the infecting organisms in the liver, in the wall of the gallbladder, in the wall of the common duct and in the pancreas. The same results occurred regardless of whether the injections were made into the main trunk of the portal vein or into one of its radicles, as the appendix vein. We believe, therefore, that the results obtained are comparable to the cases of cholecystitis in man which may be considered as following appendicitis or some other lesion of the portal system.

The experiments were conducted on seven normal dogs. After anesthetization and preparation of the skin, a median laparotomy incision was made. The portal vein was exposed at its origin, and from 4 to 6 c. c. (depending on the size of the animal) of *Staphylococcus aureus* (twenty-four hour broth culture) was injected into the portal vein, and the abdomen was closed. These animals were observed daily, and four of them were killed and examined at necropsy, from two to six days afterward. Three of the dogs were given an additional injection of hemolytic streptococci because apparently they had not been much affected by the first injection of staphylococci. From two to five days after the second injection, these dogs were killed and necropsies performed. In two additional experiments, injections of streptococci were made into a radicle of the portal vein corresponding to the appendix vein in man, and similar results were obtained to those following the injections into the portal vein itself.

In a second set of experiments in which six rabbits were used, sections were stained and examined for bacteria after injecting streptococci into the portal vein. The cystic duct, including the cystic artery, was ligated in order to eliminate the possibility of the passage of bacteria down from the liver in the bile and also to reduce to a minimum the possibility of the injected organisms reaching the gallbladder wall by way of the blood stream. After ligation of the cystic artery, the only way by which the organisms could reach the gallbladder by the blood would be either through the few small branches of the hepatic artery which pass through the attachment of the liver to the gallbladder or by a retrograde thrombosis of the portal vein into the cystic veins (which are its tributaries). In order to minimize the latter possibility, the animals were killed and examined at intervals of one, two, twelve, twenty-four and forty-eight hours, respectively, after being injected. Except perhaps in the last instance, it was felt that the time intervals were too short to permit a retrograde thrombosis. Streptococci were found in all of the last four rabbits, not only in the liver and the wall of the gallbladder but also in the pancreas. The common duct was not searched. Results of an identical sort were obtained in three other rabbits in which the injection was made into a vein corresponding to the appendix vein in man. The only exceptions,

Fig. 25.—Experimental cholecystitis (Dog 7). This shows apparently a lymphatic vessel in the outer layer of the gallbladder containing polymorphonuclear leukocytes and presenting other evidence of acute inflammation.

Fig. 26.—Experimental choledochitis by portal vein injection: extensive leukocytic infiltration of wall (Dog 3).

Fig. 27.—Experimental interstitial pancreatitis by portal vein injection (Dog 1); many leukocytes, chiefly polymorphonuclear, in trabeculae of pancreas.

Fig. 28.—Experimental interstitial pancreatitis (Dog 2); marked preponderance of polymorphonuclear leukocytes over other cells in trabeculae.

Fig. 29.—Experimental hepatitis in rabbit; portal vein injection with streptococci; demonstration, two hours later, of organisms in liver near attachment to gallbladder.

Fig. 30.—Experimental hepatitis in another rabbit after portal vein injection with streptococci; twelve hours later organisms found in liver near its attachment to the gallbladder.

Fig. 31.—Experimental cholecystitis in rabbit; injection of portal vein after ligation of cystic duct and cystic vessels; two hours later organisms found in wall of gallbladder.

Fig. 32.—Experimental cholecystitis in another rabbit; organisms demonstrated in wall twelve hours after portal vein injection.

therefore, were in the rabbits which were killed one and two hours, respectively, after the injections. It seems to us that these findings harmonize well with what would be expected on the assumption that the organisms reached the gallbladder wall from the liver through the lymphatics in the attachment of the gallbladder to the liver, because apparently there was no other route of access available. Even if they came through the very doubtful route of the few small branches of the hepatic artery which pass through the attachment of the gallbladder to the liver, one would have to assume that they had passed through the circuit of the systemic circulation after leaving the liver and had come back through the hepatic artery. Each rabbit was injected with 2 c.c. of a twenty-four hour broth culture of hemolytic streptococci.

PROTOCOLS

EXPERIMENT 1.—Five c.c. of a twenty-four hour broth culture of *Staphylococcus aureus* was injected into the portal vein of Dog 1. The dog was killed six days after the injection.

Necropsy Findings.—Liver: It was somewhat swollen, a grayish-blue, and the edge was rounded. Microscopic: There was extensive diffuse hepatitis, but the most marked infiltration of polymorphonuclears was in the interlobular sheaths.

Gallbladder: It was bound to the liver by adhesions. Microscopic: There was marked inflammation, with polymorphonuclears scattered throughout the wall. The mucosa was also involved, but the epithelium was practically intact.

Common Duct: Acute inflammation was marked. The entire wall was infiltrated with polymorphonuclears and marked vascular dilatation.

Pancreas: There was rather extensive infiltration of the trabeculae of connective tissue with polymorphonuclear leukocytes and a few red cells. There was no parenchymatous necrosis, but acute interstitial pancreatitis existed.

EXPERIMENT 2.—Four c.c. of *Staphylococcus aureus* and four days later 6 c.c. of *Staphylococcus aureus* was injected into Dog 2. The dog was killed three days after second injection.

Necropsy Findings.—Liver: It was dark and edematous. Microscopic: There were marked fatty changes, especially in the center of the lobules. There was moderate infiltration of the periportal tissues with leukocytes, mostly polymorphonuclears, extending up into parenchyma.

Gallbladder: It was pale and adherent to the liver. Microscopic: There was extensive acute inflammation, with marked infiltration of the wall with polymorphonuclears, most pronounced in the outer layers, and marked vascular dilatation. The epithelium was practically normal. *Staphylococcus* was demonstrated in the walls.

Pancreas: There was marked polymorphonuclear infiltration of the connective trabeculae, and acute interstitial pancreatitis.

EXPERIMENT 3.—Five c.c. of *Staphylococcus aureus* and three days later a second injection of 2 c.c. of hemolytic streptococci were injected in Dog 3, which was killed five days after the second injection.

Necropsy Findings.—Liver: There were several scattered abscesses. Microscopic: There were moderate fatty changes, and very extensive infiltration of polymorphonuclear leukocytes throughout the whole liver but evidently begin-

Fig. 33.—Experimental pancreatitis in rabbit; demonstration of organisms in pancreas twelve hours after injection of portal vein.

Fig. 34.—Anastomosis between lymphatics of gallbladder and liver (taken from Sudler). The extensive communication between the lymphatics of the liver and gallbladder is seen particularly at the sides of the gallbladder.

ning at the periphery of each lobule. In places there were marked accumulations of polymorphonuclear leukocytes and some mononuclear cells around the interlobular veins. There was very marked infiltration of polymorphonuclear cells beneath the liver capsule.

Gallbladder: It was adherent to the liver. The wall was gray. Microscopic: There was moderate infiltration of the wall with polymorphonuclear leukocytes and round cells, especially at the site of attachment between the liver and gallbladder.

Common Duct: The wall was extensively infiltrated with leukocytes. The mucosa was ulcerated.

Pancreas: There was polymorphonuclear infiltration into the trabeculae (moderate).

EXPERIMENT 4.—Five c.c. of *Staphylococcus aureus* and three days later a second injection of 2 c.c. of hemolytic streptococcus were injected into Dog 4, which was killed two days after second injection.

Necropsy Findings.—Liver: There was no gross change. Microscopic: There was diffuse hepatitis but again it was more marked at the peripheries of the lobules and in the interlobular tissues.

Gallbladder: It was adherent to the liver. The wall was grayish-pink. Microscopic: There was rather marked leukocytic infiltration of the wall, especially in the outer layers and more particularly in the aspect of the gallbladder away from the liver (that part covered by peritoneum). This is where most of the lymphatics are (Poirier and Charpy). Mucosa was not particularly affected.

Pancreas: The same type of lesion as already described under earlier experiments was found here.

Dogs 5, 6, and 7 showed lesions essentially similar to those already described. Dog 5 received 3 c.c. of hemolytic streptococcus three days after an initial injection of 5 c.c. of staphylococcus. In Dogs 6 and 7 there were fibrinous adhesions between the gallbladder and the liver.

SUMMARY

In many cases, probably in a majority, cholecystitis represents a direct extension to the wall of the gallbladder from a liver already inflamed. The hepatitis usually begins and is most marked in the interlobular, or periportal, tissues; and it is apparently due to infection brought to the liver by the portal vein, and, more rarely perhaps, by the hepatic artery. A pericholangitis then occurs, and because of the intimate anastomosis between the lymphatics of the intrahepatic and extrahepatic biliary systems a direct extension into the wall of the gallbladder takes place as well as into the common duct and the pancreas. From the antecedent hepatitis, therefore, a cholecystitis, choledochitis and pancreatitis can be understood to occur if in the consideration of inflammations in this locality we apply the well known fact concerning inflammation in general, namely, that it extends by way of the lymphatics. We believe that the ideas expressed in this article explain more readily than any others heretofore offered the frequent and well recognized association of biliary tract infections with

lesions of the portal system (appendicitis, peptic ulcer, typhoid fever, suppurating hemorrhoids, etc.). It has been shown also that hepatitis may follow the experimental production of cholecystitis by direct inoculation of organisms into the gallbladder. This possibility suggests that, frequently perhaps, a vicious circle between the gallbladder and liver is established whereby each may reinfect the other; and it forcibly emphasizes the desirability of cholecystectomy rather than cholecystostomy because by this means the vicious circle can most readily be broken. The ideas here expressed also furnish added evidence of the necessity of treating, in addition to the gallbladder, any coexisting lesions of the portal system.

While the opinion is expressed here that many, and perhaps the majority of, cases of cholecystitis represent a lymphatic spread from the liver, we recognize also that some are doubtless hematogenous in origin, some perhaps are contact infections from bacteria carried down in the bile and a few perhaps may have originated in an ascending infection of the common duct through its lymphatics or have been due to a chance contact of a gallbladder with an inflamed contiguous organ.

NOTE.—Since writing this article we have seen an article by George P. Muller on the "Pathology of Cholecystitis" (Brooklyn M. J. 19:11, 1905) in which he suggested in 1904 the lymphatic origin of cholecystitis. He states, "It is reasonable, therefore, to conclude that infection from the portal vein may reach the gallbladder through the superficial collecting lymphatics as well as from the bile." He apparently considers this rather a rare occurrence, however, as he states:

Clinically, attention has been called by Ochsner and others to the association of gallstones and appendicitis, but in the rather extensive series of gallbladder operations performed by Dr. Deaver, we have found but one case in which the bile infection might be directly traceable to a diseased appendix, while upon the other hand in several thousand operations upon the appendix the connection is quite as limited.